

TR-540-162

FINAL DRAFT FOR THE DRINKING WATER
CRITERIA DOCUMENT ON VINYL CHLORIDE

January 1985

Prepared Under

Program No. 1302

for

Contract 68-01-6750

Work Assignment 01

by

ICAIR
Life Systems, Inc.
Cleveland, OH 44122

for

Criteria and Standards Division
Office of Drinking Water
U.S. Environmental Protection Agency
Washington, DC 20460

SL 067119



TR-540-162

FINAL DRAFT FOR THE DRINKING WATER
CRITERIA DOCUMENT ON VINYL CHLORIDE

January 1985

Prepared Under

Program No. 1302

for

Contract 68-01-6750

Work Assignment 01

by

ICAIR
Life Systems, Inc.
Cleveland, OH 44122

for

Criteria and Standards Division
Office of Drinking Water
U.S. Environmental Protection Agency
Washington, DC 20460

SL 067120

DISCLAIMER

This document is a preliminary draft. It has not been released by the Office of Drinking Water, U.S. Environmental Protection Agency (EPA), and should not at this stage be construed to represent Agency policy. It is being circulated for comments on its technical merit.

SL 067121

TABLE OF CONTENTS

	<u>PAGE</u>
LIST OF FIGURES	ii
LIST OF TABLES	iii
I. SUMMARY	I-1
II. INTRODUCTION	II-1
III. PHYSICAL AND CHEMICAL PROPERTIES	III-1
IV. PHARMACOKINETICS	IV-1
A. Absorption and Distribution	IV-1
B. Metabolism	IV-5
C. Excretion	IV-8
V. HUMAN EXPOSURE	V-1
A. Exposure Estimation	V-1
1. Water	V-2
2. Diet	V-4
3. Air	V-6
B. Summary	V-8
VI. HEALTH EFFECTS IN ANIMALS	VI-1
A. Acute/Subchronic/Chronic Effects	VI-1
B. Teratogenicity	VI-3
C. Mutagenicity	VI-4
D. Carcinogenicity	VI-36
VII. HUMAN HEALTH EFFECTS	VII-1
A. Non-Carcinogenic Effects	VII-1
B. Carcinogenic Effects	VII-5
VIII. MECHANISMS OF TOXICITY	VIII-1
IX. QUANTIFICATION OF TOXICOLOGICAL EFFECTS FOR VINYL CHLORIDE	IX-1
A. Non-Carcinogenic Effects	IX-4
B. Quantification of Non-Carcinogenic Effects	IX-6
C. Carcinogenic Effects	IX-10
D. Quantification of Carcinogenic Effects	IX-13
X. REFERENCES	X-1
APPENDIX 1 Unit Risk Assessment for Vinyl Chloride	A1-1

LIST OF FIGURES

<u>FIGURE</u>		<u>PAGE</u>
II-1	Locations of Vinyl Chloride and Polyvinyl Chloride Plants in the United States	II-2
VII-1	Number of Cases of Vinyl Chloride/PVC Related Angiosarcomas Reported to NIOSH by Year of Diagnosis (Representing Only 63 of the 64 Cases Known to NIOSH Since Information on Diagnosis is Missing for One Case	VII-14

LIST OF TABLES

<u>TABLE</u>		<u>PAGE</u>
IV-1	Percentage of the Administered ^{14}C Activity per Gram of Tissue After Administration of (^{14}C) Vinyl Chloride by Gavage to Male Sprague-Dawley Rats	IV-3
IV-2	Percentage of ^{14}C Activity per Gram Tissue 72 Hours Following an Inhalation Exposure to (^{14}C) Vinyl Chloride for 6 Hours in Male Sprague-Dawley Rats	IV-4
IV-3	Percentage of Administered ^{14}C Activity Recovered Following a Single Oral Dose of Vinyl Chloride	IV-8
IV-4	^{14}C -Containing Urinary Metabolites from Male Sprague-Dawley Rats Given Vinyl Chloride by Gavage	IV-9
IV-5	Excretion of Radioactivity in Rats, Given a Single Dose of (^{14}C) Vinyl Chloride	IV-11
V-1	Total Estimated Cumulative Population (in Thousands) Exposed to Vinyl Chloride in Drinking Water Exceeding the Indicated Concentration	V-3
V-2	Estimated Drinking Water Intake of Vinyl Chloride	V-5
V-3	Estimated Respiratory Intake of Vinyl Chloride	V-7
V-4	Estimated Intake of Vinyl Chloride from the Environment by Adult Males in $\mu\text{g/kg/day}$ (% from Drinking Water)	V-9
VI-1	Oncogenic Effects of Inhaled Vinyl Chloride	VI-7
VI-2	Incidence of Tumors in Rats and Rabbits Exposed to Vinyl Chloride by Inhalation	VI-8
VI-3	Experiment BT1	VI-10
VI-4	Experiment BT2	VI-11
VI-5	Experiment BT6	VI-12
VI-6	Experiment BT9	VI-13
VI-7	Experiment BT15	VI-14
VI-8	Experiment BT3	VI-15
VI-9	Experiment BT10	VI-16
VI-10	Experiment BT5	VI-17
VI-11	Experiment BT14	VI-18
VI-12	Experiment BT7	VI-19
VI-13	Experiment BT17	VI-20
VI-14	Experiment BT4	VI-21
VI-15	Experiment BT8	VI-22
VI-16	Experiment BT11	VI-23
VI-17	Experiment BT27	VI-24
VI-18	Experiment BT12	VI-25
VI-19	Experiment BT13	VI-26
VI-20	Tumors Presently Correlated to VC Exposure (by Inhalation) on Experimental Rodents	VI-27
VI-21	Type and Incidence of Treatment-Related Histopathological Changes in the Liver of Rats Exposed Orally to VCM	VI-30

continued-

List of Tables - continued

<u>TABLE</u>		<u>PAGE</u>
VI-22	Site, Type and Incidence of Tumors in Organs Other than the Liver in Rats Exposed Orally to VCM for Over 2 to 5 Years . .	VI-32
VI-23	Type and Incidence of Histopathological Changes in the Liver	VI-37
VI-24	Type and Incidence of Tumors of the Mammary Glands and Incidences of Abdominal Mesotheliomas	VI-42
VII-1	Angiosarcoma of the Liver in Vinyl Chloride/PVC Worker . . .	VII-12
VIII-1	Results of Vinyl Chloride Mutagenicity Studies	VIII-4

I. SUMMARY

Almost seven billion pounds of vinyl chloride are produced in the United States annually. Most emissions into the environment originate from manufacturing plants which use the compound for the production of polyvinyl chloride resins. The predominant route of exposure to the public living near these plants is through inhalation, while the principal source of vinyl chloride exposure for most Americans is probably from polyvinyl chloride food containers. This source contributes approximately 1 ppb vinyl chloride to the diet. Vinyl chloride has also been found in drinking water. Three national surveys have demonstrated the presence of vinyl chloride at very low levels ($\mu\text{g/L}$ range) in a small number of drinking water supplies.

Upon ingestion, vinyl chloride is rapidly absorbed from the gastrointestinal tract and is distributed to the liver and other organs. Several pathways may be involved in vinyl chloride metabolism, which occurs primarily in the liver. The toxicity of vinyl chloride appears to be attributable to its enzymatic conversion to reactive polar metabolites such as chloroacetaldehyde or chloroethylene oxide. Several of these suspected metabolites are mutagenic, but vinyl chloride itself is not mutagenic, according to available information. At low doses (e.g., 1 mg/kg) of vinyl chloride the metabolites are excreted primarily in the urine. At high doses (e.g., 100 mg/kg), most of the solvent is expired as vinyl chloride.

Acute and chronic exposure to vinyl chloride can result in toxicity in experimental animals and humans. In animals, an inhalation exposure of

approximately 100,000 ppm results in death within several hours, with autopsies revealing congestion and edema of the lungs and hyperemia of the kidneys and liver. Test animals exposed to an air concentration below 100 ppm exhibit no pronounced adverse health effects. Vinyl chloride does not appear to be teratogenic in rats or rabbits, but insufficient data exist to evaluate the teratogenicity of vinyl chloride in humans.

Studies on humans working in vinyl chloride plants suggest that systemic toxic effects that are noncarcinogenic in nature can be demonstrated at exposure levels below 50 ppm. Some plant workers may have been exposed to concentrations exceeding 1,000 ppm and occasionally approaching 10,000 ppm before OSHA standards were instituted in 1974. At these levels, workers manifested dizziness, headaches and/or euphoria. Long-term exposure to these levels in vinyl chloride plants have resulted in numerous toxic effects (i.e., acroosteolysis, pulmonary insufficiency, cardiovascular and gastrointestinal manifestations and disturbances of the central nervous system). Unfortunately, data on dose-response relationships in humans are very scarce because there were few air measurements of vinyl chloride in the work environment before 1974.

Vinyl chloride is a proven carcinogen in mice, hamsters and rats. Animals studies have shown that vinyl chloride produces tumors of different types at different sites, and that the incidence and relative distribution of these tumors are influenced by dose, age of the animal and species and strain of animal used. Angiosarcomas of the liver were found in all animals studied.

whereas some types of tumors, such as brain tumors, hepatomas and lung tumors, were observed in only one species of animal. A dose-response relationship was observed in most experiments. Inhalation studies have shown that 50 ppm is the lowest vinyl chloride exposure which has a carcinogenic effect. A recently completed ingestion study demonstrated the occurrence of hepatic angiosarcomas and pulmonary angiosarcomas in rats at levels of 5.0 mg/kg bw/day or more, and the increased incidence of foci of cellular alteration and liver cell tumors at the lowest exposure level of 1.7 mg/kg bw/day.

Human data have been obtained primarily from workers exposed to vinyl chloride. A number of epidemiologic studies have linked vinyl chloride with angiosarcoma and other forms of neoplasm. The reported frequency of angiosarcoma of the liver is especially noteworthy because this is a very rare type of cancer (25 to 30 cases/year in the United States), and it is, therefore, reasonable to infer a causal relationship between exposure to vinyl chloride and the development of this tumor. Through 1977, a total of 64 cases of liver angiosarcoma have been identified worldwide among vinyl chloride-exposed industrial workers. Although rare, the carcinogenicity of vinyl chloride to humans is unambiguous.

There is insufficient data available for calculation of one-day health advisories but the ten-day health advisories can be used as conservative estimates of one-day health advisories. Ten-day health advisories of 9.0 mg/L for an adult and 2.6 mg/L for a child were developed using the subchronic toxicity data in the study by Feron et al. (1975). An adjusted ADI of 0.046 mg/L for chronic noncarcinogenic effects was calculated using the

liver lesion data in the lifetime carcinogenicity study in rats by Til et al. (1983).

The International Agency for Research on Cancer (IARC) analyzed the available data and concluded that exposure to vinyl chloride results in an increased carcinogenic risk to humans. The organs most likely to be affected are the liver, brain, lung and hemato- and lymphopoietic systems. The National Academy of Sciences (NAS) (1983) also examined the data and concluded that vinyl chloride is an established carcinogen in humans and animals, with older animals and females appearing to be more susceptible.

The NAS and EPA's Carcinogenic Assessment Group (CAG) have calculated projected incremental excess cancer risks associated with the consumption of a specific chemical via drinking water by mathematical extrapolation from high-dose animal studies. Using the risk estimates generated by the NAS (1977 to 1979), in which the linear non-threshold multi-stage model was utilized, the range of vinyl chloride concentrations were computed that would nominally increase the risk of one excess cancer per million (10^6), per hundred thousand (10^5) or per ten thousand (10^4) people over a 70-year lifetime assuming daily consumption at the stated exposure level. From the NAS model it is estimated, at the 95% confidence limit, that consuming two liters per day with a vinyl chloride concentration of 100 $\mu\text{g/L}$, 10 $\mu\text{g/L}$ or 1 $\mu\text{g/L}$ would increase the risk of one excess cancer per 10,000, 100,000 or 1,000,000 people exposed over a lifetime, respectively. Using the revised CAG approach and the multistage model, it was estimated, at the 95% confidence limit, that consuming two liters of water per day with a vinyl chloride concentration of 200 $\mu\text{g/L}$,

20 ug/L or 2 ug/L would increase the risk of one excess cancer per 10,000, 100,000 or 1,000,000 people exposed, over a lifetime respectively. The numerical differences between the NAS and CAG risk estimates are due to the selection of data for use in the model. The NAS based its calculations on an ingestion study by Maltoni et al. (1975) in which rats were exposed to vinyl chloride by gavage, while the CAG used the same Maltoni et al. (1975) study, but based its estimate upon the increased incidence of total tumors in rats exposed to vinyl chloride through inhalation.

The CAG has recalculated the cancer risk with the method described above and the Feron et al. (1981) data to estimate an increased risk of one excess cancer per 10,000, 100,000 or 1,000,000 people exposed over a lifetime to a vinyl chloride concentration of 1.5, 0.15 or 0.015 ug/L, respectively. The CAG calculation based on the Feron et al. (1981) data is under reassessment to take into account new data by Til et al. (1983).


II. INTRODUCTION

Vinyl chloride has been used for over 40 years in the production of polyvinyl chloride (PVC), the most widely used material in the manufacture of plastics in the world. About 25% of the estimated 18 billion pounds of vinyl chloride produced worldwide in 1972 was manufactured in the United State (Berk et al., 1976). Between 1968 and 1973, vinyl chloride production in the United States rose 14% annually, reaching a production level of nearly seven billion pounds in 1978 (U.S. International Trade Commission). This increase in vinyl chloride production was due to the growing dependence of virtually every branch of industry and commerce upon products and components fabricated from polyvinyl chloride (USEPA 1974). For the location of vinyl chloride and PVC manufacturing and processing plants in the United States in 1978, refer to Figure II-1.

Vinyl chloride is not known to occur in nature (NAS 1977). The compound is synthesized as chlorinated olefinic hydrocarbon monomer from petrochemical feedstock and chlorine. In 1975, vinyl chloride emissions in the United States were found to originate from three major sources: (1) 17 plants where vinyl chloride was commercially synthesized (about 11 percent); (2) 41 PVC plants where the vinyl chloride monomer was used in the production of PVC resins for various industrial purposes (about 85 percent); (3) about 8,000 PVC fabricating plants (USEPA 1975b).

Vinyl chloride and PVC are used as raw materials in the rubber, paper, glass and automotive industries. In addition, vinyl chloride and PVC are used

Figure II-1 Locations of Vinyl Chloride and Polyvinyl Chloride Plants in the United States




Source: SRI

SL 067132

00 jrtc

III. PHYSICAL AND CHEMICAL PROPERTIES

The structure of vinyl chloride is:



Vinyl chloride is highly flammable (limits of inflammability: 4.00% to 21.70%) and in sufficient concentrations (at least 1,200 to 2,000 ppm) has a sweet, pleasant odor. The compound has a boiling point of -13.3°C . Thus, at standard temperature and pressure, vinyl chloride is a gas. Vinyl chloride is only sparingly soluble in water (0.11 g/100 g water at 28°C), but is soluble in alcohol and very soluble in ether and carbon tetrachloride. The specific gravity of the chemical is 0.91. The vapor density of vinyl chloride is slightly more than twice that of air (CRC Handbook of Chemistry and Physics 1978 to 1979, Braker and Mossman 1971).

The above information indicates that vinyl chloride is volatile and readily passes from water into the gas phase under most laboratory and environmental conditions. This was confirmed in experiments where 16 mg/L vinyl chloride was added to distilled water in beakers and the concentration determined with time (USEPA 1974). The data indicate that if first order kinetics are assumed, the volatilization half-life in quiescent water (unstirred) is 290 minutes, and in continually stirred water it is 25.8 minutes. Dilling et al. (1975) found similar values for volatilization from stirred water. As Dilling et al., noted, predictions of vinyl chloride loss from water at relatively high concentrations (e.g., 1 mg/L) may not reflect the situation at very low concentrations.

in the manufacture of electrical wire insulation and cables, piping, industrial and household equipment, medical supplies, food packaging materials and building and construction products. PVC and vinyl chloride copolymers are distributed and processed in a variety of forms, including dry resins, plastisol (dispersions in plasticizers), organosol (dispersions in plasticizers plus volatile solvent), and latex (a colloidal dispersion in water used to coat paper, fabric or leather).

Volatilization appears to be the most significant process in the loss of vinyl chloride from the aquatic environment (Hill et al. 1976). Once in the atmosphere, vinyl chloride undergoes rapid photochemical oxidation (Gay et al. 1976), Lillian et al. 1975).

IV. PHARMACOKINETICS

A. Absorption and Distribution

An investigation by Duprat et al. (1977) indicated that inhaled vinyl chloride is rapidly absorbed by the lungs and immediately accumulates in the liver. In this study, rats were exposed in a chamber to 20,000 ppm ^{14}C vinyl chloride for five minutes, and then the distribution of radioactive vinyl chloride in the various body organs was determined. At ten minutes following treatment, radioactivity was found in the liver, bile duct, digestive lumen and kidneys. With increasing time (up to three hours), ^{14}C activity was detected in the urinary system, salivary and lacrimal glands, skin and thymus.

Using male Wistar rats, Withey (1976) determined that vinyl chloride is rapidly absorbed from the gastrointestinal tract following gastric intubation of aqueous solutions containing up to 2.0 mg/mL vinyl chloride. Vinyl chloride uptake by this route was extremely rapid; peak concentrations were found less than ten minutes after the dose was administered.

In a study by Watanabe et al. (1976a), rats were given single oral doses (gavage) of 0.05, 1 or 100 mg/kg of ^{14}C -vinyl chloride dissolved in corn oil, and the routes and rates of elimination of ^{14}C activity were followed for 72 hours. The percentage of the dose expired as vinyl chloride was 1%, 2% and 67%, respectively. The disposition of vinyl chloride to various organs and tissues was also determined. The liver was found to retain the greatest percentage of activity at all dose levels, three to five times the percentage

found in muscle, lung or fat (Table IV-1). The investigators concluded that the fate of vinyl chloride following oral administration is a dose-dependent saturable process, with the saturation of the vinyl chloride-metabolizing enzymes occurring at a concentration between 1 and 100 mg/kg.

In an inhalation study by this group (Watanabe et al. 1976b), rats were exposed to 10 or 1,000 ppm ^{14}C -vinyl chloride for six hours and the routes and rates of elimination of ^{14}C activity were followed for 72 hours after termination of exposure. The animals were sacrificed after 72 hours and samples of tissues collected for analysis of ^{14}C activity. Table IV-2 indicates that, as in the gavage study, the liver retained the greatest percentage of vinyl chloride (or metabolites) at the dose levels studied. In contrast to the gavage experiment, however, no saturation of vinyl chloride metabolism was discernable between 10 ppm and 1,000 ppm in this study.

Bolt et al. (1976) also studied the tissue disposition of ^{14}C -vinyl chloride in rats. Immediately after exposure by inhalation of 50 ppm vinyl chloride for five hours in a closed system, the percentage of incorporated ^{14}C -radioactivity per g tissue was highest for kidney (2.13) and liver (1.86). The percent of incorporated activity was 0.73 for the spleen and 0.17 for the brain. Forty-eight hours after the beginning of exposure, labeled material could still be detected in these tissues.

The percentage absorption of vinyl chloride from the human gastrointestinal tract has not been established. Because of the lack of data

Table IV-1 Percentage of the Administered ^{14}C Activity per Gram of Tissue After Administration of (^{14}C) Vinyl Chloride by Gavage to Male Sprague-Dawley Rats^a

Tissue	Dose (mg/kg) ^b		
	0.05	1.0	100
Liver	0.172±0.025 ^c	0.182±0.005	0.029±0.002
Skin	0.070±0.023	0.076±0.010	0.010±0.002
Carcass	0.027±0.007	0.046±0.002	0.007±0.001
Plasma	0.041±0.004	0.053±0.007	ND ^d
Muscle	0.028±0.003	0.031±0.003	0.006±0.001
Lung	0.050±0.003	0.061±0.003	0.011±0.001
Fat	0.030±0.004	0.045±0.008	0.006±0.001

^aRemaining in the body after 72 hr.

^bVinyl chloride dissolved in corn oil.

^cMean ± SE, five rats per dose.

^dNot detectable above background.

Adapted from Watanabe et al. 1976a.

Table IV-2 Percentage of ^{14}C Activity per Gram Tissue 72 Hours Following an Inhalation Exposure to (^{14}C) Vinyl Chloride for 6 Hours in Male Sprague-Dawley Rats

Tissue	Percentage ^{14}C activity Exposure concentration	
	10 ppm	1,000 ppm
Liver	0.139 ± 0.009^a (0.35) ^(b) 0.141 ± 0.009^c	0.145 ± 0.008^a (9.63) ^b 0.165 ± 0.009^c
Skin	0.072 ± 0.004 (0.18) 0.073 ± 0.004	0.115 ± 0.010 (7.64) 0.131 ± 0.011
Carcass	0.048 ± 0.004 (0.12) 0.049 ± 0.004	0.049 ± 0.004 (3.26) 0.056 ± 0.005
Plasma	0.051 ± 0.001 (0.13) 0.052 ± 0.001	ND ^d
Muscle	0.052 ± 0.005 (0.13) 0.053 ± 0.005	0.038 ± 0.003 (2.52) 0.043 ± 0.003
Lung	0.065 ± 0.007 (0.16) 0.066 ± 0.007	0.046 ± 0.001 (3.06) 0.052 ± 0.001
Fat	0.026 ± 0.006 (0.07) 0.026 ± 0.006	ND ^d
Kidney	0.079 ± 0.003 (0.20) 0.080 ± 0.003	0.057 ± 0.005 (3.79) 0.065 ± 0.006

^aExpressed as percentage of total ^{14}C activity per gram of tissue. Uncorrected for expired VC:

$$\frac{\text{dpm per g tissue}}{\text{Total dpm recovered}}$$

Mean $\pm$ SE from four rats

^bMicrogram equivalents vinyl chloride per gram of tissue.

^cExpressed as percentage metabolized ^{14}C activity per gram tissue. Corrected for expired VC:

$$\frac{\text{dpm per gram of tissue}}{\text{Total dpm recovered} - \text{dpm of expired VC}}$$

Mean $\pm$ SE from rats.

^dNot detectable, detection limit for plasma and fat was 3 $\mu\text{g/g}$ of tissue (3 ppm).


Adapted from Watanabe et al. 1976b.

on percent absorption from the gastrointestinal tract, the risk calculations in this document will assume a 100% absorption factor.

B. Metabolism

Metabolism of vinyl chloride occurs primarily by microsomal enzymes in the liver. There is strong evidence that the toxicity of this compound is attributable to its enzymatic oxidation to reactive polar metabolites. Several of these suspected metabolites are strongly mutagenic, but vinyl chloride itself is not (Bartsch and Montesano 1975). Exposure to vinyl chloride leads to the reduction of non-protein sulfhydryl levels in rat liver, suggesting that the metabolites of vinyl chloride conjugate with glutathione and/or cysteine (Hefner et al. 1975a). Hathway (1977) reported in vitro depurination of calf thymus DNA by chloroacetaldehyde identical to that observed in hepatocyte DNA following administration of vinyl chloride to rats in vivo. This suggests that vinyl chloride metabolites may interact with some purine and pyrimidine residues of DNA, providing a possible explanation for the oncogenic properties associated with vinyl chloride.

In a review of the literature, Bartsch and Montesano (1975) report two possible biotransformation schemes—one involving alcohol dehydrogenase (Scheme I) and the other involving the mixed function oxidase system (Scheme II). These are indicated below:



Evidence for biodegradation involving the alcohol dehydrogenase pathway includes data which demonstrates that pretreatment of rats with either ethanol or pyrazole (an inhibitor of alcohol dehydrogenase) inhibits the metabolism of vinyl chloride (Hefner et al. 1975a).

There is also ample evidence that the mixed function oxidase (MFO) system is involved in the metabolism of vinyl chloride. Pretreatment of rats with phenobarbital, which induces the MFO system, also enhanced liver toxicity of vinyl chloride (Jaeger et al. 1974). Rat liver microsomes catalyzed the covalent binding of vinyl chloride metabolites to protein and nucleic acids (Kappus et al. 1975, 1976). Chloroethylene oxide, which is thought to be formed by the MFO system, may be the primary microsomal metabolite capable of alkylating these intra-cellular macromolecules (Laib and Bolt 1977).

Several pathways may be involved in vinyl chloride metabolism, the predominant one depending on dose. Hefner et al. (1975) performed an inhalation study in which rats were exposed to vinyl chloride concentrations ranging from 50.5 to 1,167.0 ppm for 12-months time. The rate of metabolism, as determined by measuring the declining level of vinyl chloride in the chamber atmosphere, was three times greater for seven separate exposures ranging from 50 to 105 ppm than it was for five separate exposures ranging from 220 to 1,167 ppm. This indicated that the predominant pathway at the lower concentrations, probably involving alcohol dehydrogenase, is saturable between 105 and 220 ppm. This group also found evidence that oxidases in the microsomes may be involved in metabolism at high level exposures.

In another study, Bolt et al. (1977) subjected rats to an inspired concentration of ^{14}C -vinyl chloride ranging from 200 to 1,200 ppm in a closed system, and measured the rate of decrease of vinyl chloride levels in the chamber atmosphere. This group calculated that saturation of the vinyl chloride-metabolizing enzymes of the rat occurs at 250 ppm.

C. Excretion

Excretion of ^{14}C activity within 72 hours following a single oral dose of ^{14}C -labeled vinyl chloride (0.05, 1.0 or 100 mg/kg) is shown in Table IV-3 (Watanabe et al. 1976a). As the dose increased, a markedly greater proportion of vinyl chloride was expired unmetabolized, while the percentage of metabolite in the urine decreased substantially. Again, saturation kinetics are suggested. The table also indicates that metabolites of vinyl chloride were predominantly excreted via the urine. Administration of vinyl chloride by inhalation produced almost identical results (Watanabe et al. 1976b). Two major metabolites in the urine were identified as indicated in Table IV-4.

Buchter et al. (1980) examined the metabolic elimination of vinyl chloride in Rhesus monkeys. Rhesus monkeys were placed in a closed exposure system into which vinyl chloride was injected, and air samples were taken to determine the decline of vinyl chloride in the gas phase of the system. The results showed that the metabolic elimination of vinyl chloride in Rhesus monkeys is a dose-dependent, saturable process, as it is in rats. Elimination was shown to obey a first-order law of kinetics at air concentrations below

Table IV-3 Percentage of Administered ^{14}C Activity Recovered
Following a Single Oral Dose of Vinyl Chloride^a

	Dose (mg/kg)		
	0.05	1.0	100
Expired:			
As VC	1.43±0.13 ^b	2.13±0.22	66.64±0.67
As CO ₂	8.96±0.59	13.26±0.47	2.52±0.13
Urine	68.34±0.54	59.30±2.75	10.84±0.95
Feces	2.39±0.52	2.20±0.39	0.47±0.06
Carcass and tissues	10.13±1.93	11.10±0.47	1.83±0.14
Cage wash ^c	0	0.84±0.45	0
Total recovery	91.25±2.47	88.83±1.96	82.30±0.43

^aPercentage of dose excreted over 72 hours. Only the ^{14}C activity associated with the expired VC can be attributed to VC per se.

^bMean ± SE five rats per dose.

^cDistilled water wash of metabolism cage at termination of the study.

Adapted from Watanabe et al. 1976a.

Table IV-4 ¹⁴C-Containing Urinary Metabolites from Male
Sprague-Dawley Rats Given Vinyl Chloride by Gavage^a

Compound	Dose (mg/kg)		
	0.05(4) ^b	1.0(5)	100(5)
(A) N-acetyl-S-(2-hydroxyethyl- cysteine)	30.4±2.0 ^c	36.2±3.9	29.1±2.0
(B) Thiodiglycolic acid	25.6±1.9	23.7±1.1	25.4±0.9
(C) Unidentified	38.6±2.9	34.5±4.6	36.6±2.0
Total	94.6	94.6	91.1

^aMetabolites were separated and quantitated by high pressure liquid chromatography. Values are expressed as percentage of total urinary radioactivity.

^b() = Number of animals per dose.

^cMean ± SE

Adapted from Watanabe et al. 1976a.

200 to 300 ppm, and at high concentrations the maximal velocity of metabolic elimination of vinyl chloride was about half that of rats when related to kg body weight.

Green and Hathway (1975) measured the excretion of ^{14}C -vinyl chloride administered to rats by intragastric, intravenous (femoral vein), or intraperitoneal routes. Two doses were used: 0.25 mg/kg and 450 mg/kg. The results are shown in Table IV-5. During the first 24 hours after treatment, more than 90% was excreted from the animals irrespective route. Significant differences were noted, however, in the manner of excretion for the 0.25 mg/kg dose. For the intragastric route, 71.5% was excreted in the urine, whereas 99% was exhaled from the lungs when vinyl chloride was administered intravenously. For the intraperitoneal route, 43.2% was exhaled while 41.5% was excreted in the urine. At the higher dose (450 mg/kg), over 90% was exhaled as vinyl chloride in both intragastric and intraperitoneal administered rats. The intragastric values are consistent with the values reported in the oral studies performed by Watanabe et al. (1976a) (see Table IV-3).

Withey and Collins (1976) developed a statistical model for use in equating oral dose levels of vinyl chloride to inhalation exposure levels in rats, using blood level time curves. The authors concluded that "if the total daily liquid intake contained 20 ppm vinyl chloride, then the area generated under the blood level time curve, for rats, would be equivalent to an inhalation exposure of about 2 ppm for 24 hours." Thus, according to this model, inhalation exposure is ten times more efficient than oral exposure.

Table IV-5 EXCRETION OF RADIOACTIVITY IN RATS, GIVEN A SINGLE DOSE OF [¹⁴C]VINYL CHLORIDE

4 rats were each dosed i.g. with 250 μ g of [¹⁴C]vinyl chloride per kg in corn oil solution, and another 4 rats were each dosed similarly with 450 mg of [¹⁴C]vinyl chloride per kg. 4 rats were each injected in the femoral vein with 250 μ g of [¹⁴C]vinyl chloride per kg in *N*-(β -hydroxyethyl) lactamide. Four rats were each injected i.p. with 250 μ g of [¹⁴C]vinyl chloride per kg in *N*-(β -hydroxyethyl) lactamide, and another 4 animals were each injected similarly with 450 mg of [¹⁴C]vinyl chloride.

Size of dose	Time (h)	Radioactivity excreted (% of dose) ^a									
		Intragastric				Intravenous				Intraperitoneal	
		Exhaled air		Urine	Faeces	Exhaled air		Urine	Faeces	Exhaled air	
		Vinyl chloride	CO ₂			Vinyl chloride	CO ₂			Vinyl chloride	CO ₂
250 μ g/kg	0-24	3.7 $\pm$ 1.2	12.6 $\pm$ 1.1	71.5 $\pm$ 5.0	2.8 $\pm$ 2.5	99.0 $\pm$ 0.8	0.1	0.5	0.1	43.2 $\pm$ 4.6	10.3 $\pm$ 2.2
	24-48		0.9	3.1	1.6						0.7
	48-72			0.1	0.2						
	Total	3.7 $\pm$ 1.2	13.5 $\pm$ 1.3	75.1 $\pm$ 4.2	4.6 $\pm$ 3.0	99.0 $\pm$ 0.8	0.1	0.5	0.1	43.2 $\pm$ 4.6	11.0 $\pm$ 1.2
450 mg/kg	0-24	91.9 $\pm$ 2.5	0.6	4.5 $\pm$ 2.1	0.4					96.2 $\pm$ 4.1	0.7
	24-48		0.1	0.8	0.1						0.1
	48-72			0.1							
	Total	91.9 $\pm$ 2.5	0.7	5.4 $\pm$ 2.2	0.7					96.2 $\pm$ 4.1	0.7

^a Values shown are the means $\pm$ S.D. of those means.

Adapted from Green and Hathway (1975).

V. HUMAN EXPOSURE

Humans may be exposed to vinyl chloride in drinking water, food and air. Detailed information concerning the occurrence of and exposure to vinyl chloride in the environment is presented in another document entitled "Occurrence of Vinyl Chloride in Drinking Water, Food, and Air" (Letkiewicz et al. 1983). This section summarizes the pertinent information presented in that document in order to assess the relative source contribution from drinking water, food and air.

A. Exposure Estimation

This analysis is limited to drinking water, food and air, since these media are considered to be general sources common to all individuals. Some individuals may be exposed to vinyl chloride from sources other than the three considered here, notably in occupational settings and from the use of consumer products containing vinyl chloride. Even in limiting the analysis to these three sources, it must be recognized that individual exposure will vary widely based on many personal choices and several factors over which there is little control. Where one lives, works and travels, what one eats and physiologic characteristics related to age, sex and health status can all profoundly affect daily exposure and intake. Individuals living in the same neighborhood or even in the same household can experience vastly different exposure patterns.

Unfortunately, data and methods to estimate exposure of identifiable population subgroups from all sources simultaneously have not yet been

developed. To the extent possible, estimates are provided of the number of individuals exposed to each medium at various vinyl chloride concentrations. The 70-kg male is used for estimating intake.

1. Water

Cumulative estimates of the U.S. populations exposed to various vinyl chloride levels in drinking water from public drinking water systems are presented in Table V-1. The values in the table were obtained using Federal Reporting Data Systems data on populations served by primary water supply systems (FRDS 1983) and the estimated number of these water systems that contain a given level of vinyl chloride. An estimated 1,922,000 individuals (0.9% of the population of 214,419,000 using public water supplies) are exposed to levels of vinyl chloride in drinking water at or above 1.0 ug/L, while 591,000 individuals (0.3%) are exposed to levels above 5 ug/L. It is estimated that 118,000 individuals are exposed to levels greater than 60 ug/L. Of the approximately 1.3 million people exposed to levels ranging from 1.0 to 5 ug/L, 0.9 million (65%) obtain water from surface water supplies. All exposure to vinyl chloride in drinking water at levels above 5 ug/L is expected to be from groundwater sources.

No data were obtained on regional variations in the concentration of vinyl chloride in drinking water. The highest concentrations are expected to occur near sites of polyvinyl chloride production.

Table V-1 Total Estimated Cumulative Population (in Thousands) Exposed to Vinyl Chloride in Drinking Water Exceeding the Indicated Concentration

System type	Number of people served in U.S. (thousands)	Cumulative population (thousands) exposed to concentrations (ug/l) of:								
		>1.0	>5	>10	>20	>30	>40	>50	>60	>70
Groundwater	73,473	1,063	591	118	118	118	118	118	118	0
Surface water	<u>140,946</u>	<u>059</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>
Total	214,419	1,922	591	118	118	118	118	118	118	0
(% of total)	(100%)	(0.9%)	(0.3%)	(0.1%)	(0.1%)	(0.1%)	(0.1%)	(0.1%)	(0.1%)	(0.0%)

3-4

SL 067149

Daily intake levels of vinyl chloride from drinking water were estimated using various exposure levels and the assumptions presented in Table V-2. The data in the table suggest that the majority of the persons using public drinking water supplies would be exposed to intake levels below 0.028 $\mu\text{g}/\text{kg}/\text{day}$.

An indication of the overall exposure of the total population to vinyl chloride can be obtained through the calculation of population-concentration values. These values are a summation of the individual levels of vinyl chloride to which each member of the population is exposed. An explanation of the derivation of these values is presented in Appendix C. Population concentration estimates for vinyl chloride in drinking water were $1.1 \times 10^{-7} \mu\text{g}/\text{L} \times \text{persons}$ (best case), $1.5 \times 10^{-7} \mu\text{g}/\text{L} \times \text{persons}$ (mean best case), $2.3 \times 10^8 \mu\text{g}/\text{L} \times \text{persons}$ (mean worst case), and $2.3 \times 10^{-8} \mu\text{g}/\text{L} \times \text{persons}$ (worst case).

Assuming a consumption rate of 2 liters of water/day, population-exposure values of $2.2 \times 10^7 \mu\text{g}/\text{day} \times \text{persons}$ (best case), $3.0 \times 10^7 \mu\text{g}/\text{day} \times \text{persons}$ (mean best case), $4.6 \times 10^8 \mu\text{g}/\text{day} \times \text{persons}$ (mean worst case), and $4.6 \times 10^8 \mu\text{g}/\text{day} \times \text{persons}$ (worst case) were derived.

2. Diet

No data were obtained on levels of vinyl chloride found in foods in the United States. Therefore, no estimates of the daily intake of vinyl chloride from the U.S. diet could be made.

Table V-2 Estimated Drinking Water Intake of Vinyl Chloride

Exposure level ($\mu\text{g/L}$)	Persons using supplies exposed to indicated levels		Intake ($\mu\text{g/kg/day}$)
	Population	% of total population	
≥ 1.0	1,922,000	0.9%	≥ 0.028
> 5.0	591,000	0.3%	> 0.14
> 10	118,000	0.1%	> 0.29
> 50	118,000	0.1%	> 1.4
> 70	0	0.0%	> 2.0

Assumptions: 70-kg man, 2 liters of water/day.

3. Air

Exposure to vinyl chloride in the atmosphere varies from one location to another. The highest level of vinyl chloride reported in the atmosphere was $2,100,000 \text{ ng/m}^3$ ($2,100 \text{ ug/m}^3$) (Lillian et al. 1975 cited in Brodzinsky and Singh 1982). High levels, averaging greater than $15,000 \text{ ng/m}^3$ (15 ug/m^3), have been detected in other areas. Normal levels, however, are somewhat lower. Bordzinsky and Singh (1982) calculated a median air level of 0.0 ng/m^3 (0.0 ug/m^3) in each of three types of areas: rural/remote, urban/suburban and source dominated.

The monitoring data available are not sufficient to determine regional variations in exposure levels for vinyl chloride.

The daily respiratory intake of vinyl chloride from air was estimated using the assumptions presented in Table V-3 and median and maximum levels for vinyl chloride reported above. The estimates in Table V-3 indicate that the daily vinyl chloride intake from air for adults in rural/remote, urban/suburban, and source dominated areas is 0.0 ug/kg/day . In contrast, the intake calculated using the maximum vinyl chloride level reported is 690 ug/kg/day ; few, if any, persons are believed to be exposed at that level. The values presented do not account for variances in individual exposure or uncertainties in the assumptions used to estimate exposure.

Table V-3 Estimated Respiratory Intake of Vinyl Chloride

Exposure ($\mu\text{g}/\text{m}^3$)	Intake ($\mu\text{g}/\text{kg}/\text{day}$)
Rural/remote (0.0)	
Urban/suburban (0.0)	0.0
Source dominated (0.0)	
Maximum (2,100)	690

Assumptions: 70-kg man, 23 m^3 of air inhaled/day (ICRP 1975).

B. Summary

Table V-4 presents a general view of the total amount of vinyl chloride received by an adult male from air and drinking water. Two separate exposure levels in air and six exposure levels in drinking water are shown in the table. Since no data were obtained on levels of vinyl chloride in foods in the United States, the contribution of vinyl chloride in the diet to total vinyl chloride exposure could not be assessed.

The data presented have been selected from an infinite number of possible combinations of concentrations for the two sources. The actual exposures encountered would represent some finite subset of this infinite series of combinations. Whether exposure occurs at any specific combination of levels is not known; nor is it possible to determine the number of persons that would be exposed to vinyl chloride at any of the combined exposure levels. The data presented represent possible exposures based on the occurrence data and the estimated intakes.

The relative source contribution data are based on estimated intake and do not account for a possible differential absorption rate for vinyl chloride by route of exposure. The relative dose received may vary from the relative intake. In addition, the relative effects of the chemical on the body may vary by different routes of exposure.

Brodzinsky and Singh (1982) calculated a median urban/suburban air level of vinyl chloride of $0 \text{ } \mu\text{g}/\text{m}^3$ based on air monitoring data. Assuming an air

Table V-4 Estimated Intake of Vinyl Chloride from the Environment by Adult Males in ug/kg/day (% from Drinking Water)

Concentration in drinking water (ug/l)	Concentration in air	
	Rural/remote (0.0 ug/m ³)	Urban/suburban Source dominated Maximum (2,100 ug/m ³)
0	0.0 (--)	690 (0%)
1.0 ^a	0.028 (100%)	690 (<0.01%)
5.0 ^b	0.14 (100%)	690 (0.02%)
10 ^c	0.29 (100%)	690 (0.04%)
50 ^d	1.4 (100%)	690 (0.2%)
70 ^e	2.0 (100%)	690 (0.3%)

Intake from each source (see Sections 5.1-5.3):

Water: 1.0 ug/l: 0.028 ug/kg/day
 5.0 ug/l: 0.14 ug/kg/day
 10 ug/l: 0.29 ug/kg/day
 50 ug/l: 1.4 ug/kg/day
 70 ug/l: 2.0 ug/kg/day

Air: 0.0 ug/m³: 0.0 ug/kg/day
 2,100 ug/m³: 690 ug/kg/day

Food: Not included

^a1,922,000 individuals using public drinking water systems are estimated to be exposed to levels $\geq$ 1.0 ug/l (0.9% of population using public water supplies).

^b591,000 individuals using public drinking water systems are estimated to be exposed to levels $>$ 5.0 ug/l (0.3% of population using public water supplies).

^c118,000 individuals using public drinking water systems are estimated to be exposed to levels $>$ 10 ug/l (0.1% of population using public water supplies).

^d118,000 individuals using public drinking water systems are estimated to be exposed to levels $>$ 50 ug/l (0.1% of population using public water supplies).

^eNo individuals using public drinking water systems are estimated to be exposed to levels $>$ 70 ug/l.

level of 0 ug/m^3 , drinking water would be the predominant source of vinyl chloride exposure at all drinking water levels above 0 ug/L . An accurate assessment of the number of individuals for which drinking water is the predominant source of exposure cannot be determined from the data since specific locations containing high concentrations of vinyl chloride in drinking water and low concentrations of vinyl chloride in ambient air and food are unknown.

VI. HEALTH EFFECTS IN ANIMALS

A. Acute/Subchronic/Chronic Effects

Acute toxicity tests with vinyl chloride were performed by Patty et al. (1930) of the Bureau of Mines, Department of Commerce. Single exposures of guinea pigs to vinyl chloride gas, 10 percent in air (100,000 ppm), resulted in narcosis and death within 30 to 60 minutes. Inhalation of lower concentrations resulted in ataxia and narcosis. Pathological findings at necropsy were congestion and edema of the lungs and hyperemia of the kidneys and liver. A number of investigators have made similar observations when examining the acute inhalation effects of vinyl chloride in mice, rats, guinea pigs, rabbit, cats and dogs (Peoples and Leake 1933, Lester et al. 1963, Mastromatteo et al. 1960, Haley 1975, Prodan et al. 1975). In animal studies, LC₅₀'s at two hours ranged from 117,500 ppm for mice to 230,800 ppm for rabbits.

Feron et al. (1975) reported a subchronic toxicity study in which vinyl chloride monomer (VCM) dissolved in soybean oil was administered by gavage to male and female Wistar rats, initially weighing 44 g, at doses of 0 (controls), 30, 100 and 300 mg/kg once daily, 6 days per week for 13 weeks. Several hematological, biochemical and organ weight values were significantly ($P < 0.05$ or better) different in both mid- and high-dose animals compared to controls. The authors conservatively placed the no-effect level in this study at 30 mg/kg. However, they also estimated that the no-effect level may be

higher since it was their opinion that the effects observed with the higher doses were of doubtful toxicological significance.

Marsteller et al. (1975) reviewed and summarized the findings of previous studies on vinyl chloride exposure in laboratory animals. Torkelson et al. (1961) exposed test animals to concentrations ranging from 50 to 500 ppm. Rats exposed to 100 ppm (two hours/day for six months) were judged normal on the basis of appearance, mortality, growth, hematological examination and other factors. A slight increase in the liver weight was observed. Rats, guinea pigs, rabbits and dogs exposed to 50 ppm (7 hours/day, 130 times in 189 days) appeared to be normal in appearance, mortality and growth, and the increase in weight of the rat livers did not occur at this concentration. Basalaez et al. (1972) administered gaseous vinyl chloride to rats and rabbits at a concentration of 0.03 to 0.04 mg/L for four hours/day for six months. Cardiovascular disorders, changes in the bioelectric activity of the hypothalamus, hyperadrenalinemia, osteoporosis and resorption of bone tissue were observed.

Jaeger (1975) conducted experiments with rats to determine the interaction between vinylidene chloride (1,1,-DCE) and vinyl chloride. In this study, hepatotoxicity was measured by the elevation of serum alanine- α -ketoglutarate transaminase (AKT). When fasted rats were exposed to 0.02% (V/V), 1,1-DCE, serum AKT activity was elevated about 50-fold, two hours after the termination of a four-hour inhalation exposure. No elevation was observed when 0.1% vinyl chloride was administered alone. When the two

chemicals were administered simultaneously at the levels indicated, no elevation of serum AKT occurred. Thus, the vinyl chloride was protective. These two monomers are used together in the production of vinyl copolymers, and exposure to both agents in the workplace was reported by Kramer and Mutchler (1972).

B. Teratogenicity

John et al. (1977) examined the effects of vinyl chloride inhalation on the fetuses of mice, rats and rabbits. The pregnant animals were exposed seven hours daily to concentrations of 50 or 500 ppm for mice and 500 or 2,500 ppm for rats and rabbits. Mice and rats were exposed on days 6 to 15 of gestation, and rabbits on days 6 to 18. No teratogenic effects were observed at 2,500 ppm in rats and rabbits, except that a greater incidence of dilated ureters were noted in rats. Indeed, vinyl chloride exposure at this level actually decreased the incidence of certain skeletal anomalies in rats compared to controls (e.g., delayed ossification of the bones of the skull, and unfused centers of ossification of the skull and sternbrae). Mice were the most sensitive to vinyl chloride. No teratogenic effects were noted in the fetuses of mice exposed to 50 ppm, but a significantly greater incidence of unfused sternbrae and delayed ossification of sternbrae (no. 5) and bones of the skull were observed among litters of mice exposed to 500 ppm compared to unexposed controls. Embryotoxic effects were not generally noted, but some decrease in fetal body weight and crown-rump length was observed in rats and mice.

Radike (1977) did not observe gross (nonmicroscopic) abnormalities in the offspring of rats exposed for four hours daily on days 9 to 21 of gestation by inhalation of 600 or 6,000 ppm vinyl chloride. A small increase in the incidence of minor skeletal abnormalities, including wavy ribs, extra 14th ribs and delayed calcification of small bones, were observed in the offspring of the exposed animals. The investigators concluded that such a small incidence is difficult to distinguish from a sporadic occurrence, and, therefore, these abnormalities should be considered skeletal variants and not malformations.

Groups of pregnant CF-1 mice, Sprague-Dawley rats and New Zealand white rabbits were exposed to doses of vinyl chloride ranging from 50 to 2,500 ppm by inhalation. Exposure to these concentrations of vinyl chloride did not cause any significant embryonal or fetal toxicity and was not teratogenic in any of the three species tested (John et al. 1981).

C. Mutagenicity

Vinyl chloride is mutagenic in a number of biological systems. The mutagenic action of vinyl chloride appears to be dependent upon its metabolic conversion to chemically reactive metabolites (i.e., chloroethylene oxide, 2-chloroacetaldehyde). The mutagenic effects of vinyl chloride have been demonstrated in: (1) metabolically activated systems using Salmonella typhimurium (Bartsch et al. 1975, McCann et al. 1975, Elmore et al. 1976, Rannug et al. 1974, Garro et al. 1976) developed by Ames et al. (1973) in

which the genetic indicator refers to histidine prototrophy by base-pair substitutions, or by base-pair insertions or deletions; (2) Escherichia coli K12 bioauxotrophic strain with back mutation system arginine + (Greim et al. 1975); (3) several species of yeast inducing forward mutations and gene conversions at specific loci (Loprieno et al. 1976, 1977); (4) germ cells of Drosophila (Verburgt and Vogel 1977) and (5) Chinese hamster V79 cells (Huberman et al. 1975). The literature on the mutagenic effects of vinyl chloride was reviewed by Bartsch and Montesano (1975).

The mutagenic activity of inhaled vinyl chloride (3,000, 10,000 or 30,000 ppm for 6 hours a day for 5 days) was assessed in fertile male CD-1 strain mice with the dominant lethal assay (Anderson et al. 1976). At these concentrations, vinyl chloride was not mutagenic as judged by scoring of post-implantation fetal deaths, pre-implantation egg losses and reduction in fertility. Positive control tests indicated that the dominant lethal effect was expressed in the CD-1 mice used in these experiments.

Anderson and Richardson (1976) conducted a cytogenic study investigating mutagenic effects in the bone marrow cells of rats after exposure to paradichlorobenzene at various dose levels. In this study, benzene and vinyl chloride were used as positive controls. The results of the vinyl chloride control showed that vinyl chloride was effective in producing chromosome damage in rat bone marrow after the multiple exposure regime.

D. Carcinogenicity

Evidence has been accumulated in recent years implicating vinyl chloride as a human and animal carcinogen. The first four cases of human liver angiosarcoma in workers employed by a vinyl chloride plant were reported by Creech and Johnson in 1974. The first experimental data on the carcinogenic effect of vinyl chloride in rats were published by Viola et al. in 1971. Subsequently, preliminary results of an investigation concerned with the oncogenic potential of vinyl chloride in experimental animals were reported (Maltoni and Lefemine 1974). These initial reports spurred a series of retrospective epidemiologic investigations of workers in the vinyl chloride industry and supportive experimental studies in animals. Several comprehensive reviews and symposium proceedings have been published on the subject (e.g., Selikoff and Hammond 1975, Proceedings of the Royal Society of Medicine 1976, USEPA 1975c, Milby 1978).

Viola et al. (1971) conducted animals studies and reported the carcinogenic response of male rats (AR/IRE Wistar strain) exposed to vinyl chloride by inhalation (Table VI-1). Skin tumors were first noted at approximately ten months; tumors in the lungs and bone were observed at about 11 months.

Caputo et al. (1974) exposed male and female rats (A and IRE Wistar strain) by inhalation to various concentrations of vinyl chloride. Carcinomas and sarcomas were observed in all groups except those exposed to 50 ppm (Table VI-2). The data indicate that a dose response relationship exists

Table VI-1 Oncogenic Effects of Inhaled Vinyl Chloride

Conc. VC (ppm) 4 hours/day 5 days/week 12 months	Number rats	Skin epidermoid carcinomas	Lung adenocarcinomas and squamous cell carcinomas	Bones Osteo- chondromas
30,000	26	17	6	5
No treatment	25	--	--	--

Adapted from Viola et al. 1971).

Table VI-2 Incidence of Tumors in Rats and Rabbits Exposed to Vinyl Chloride by Inhalation

(ppm) 4 hours/day 5 days/week 12 months	Number of animals	Liver angiosarcomas cholangiomas	Lung adeno-alveolar carcinomas	Skin squa- mous cell carcinoma acanthoma	Other
<u>Rats</u>					
20,000	150	31	21	67	7
10,000	200	16	16	34	8
5,000	200	12	4	20	2
2,000	200	10	8	6	6
500	150	4	--	3	--
50	200	--	--	--	--
No treatment	200	--	--	--	--
<u>Rabbits</u>					
10,000	40	--	6	12	--
No treatment	20	--	--	--	--

Adapted from Caputo et al. (1974).

at exposures between 50 and 20,000 ppm. Tumors appeared between 8 and 13 months from the beginning of the inhalation treatment. These investigations also exposed rabbits by inhalation to 10,000 ppm vinyl chloride for 15 months (Table VI-2) and reported the occurrence of lung and skin carcinomas.

Recent inhalation studies with albino CD-1 mice and CD rats (Charles River Breeding Lab) confirmed the carcinogenicity of vinyl chloride at concentrations as low as 50 ppm (Lee et al. 1977, 1978). Liver angiosarcoma as well as other forms of cancers were found in both species.

An extensive examination of vinyl chloride in experimental animals was conducted by Maltoni (1981). A summary of these results are presented in Tables VI-3 to VI-19. Vinyl chloride caused tumors in all the animal species tested (i.e., mice, rats and hamsters) both through inhalation and ingestion exposure. A clear-cut dose-response relationship was demonstrated, with carcinogenic effects occurring at exposures as low as 50 ppm. Newborn animals appeared to be especially sensitive to the development of hepatocarcinomas and angiosarcomas and carcinogenic effects on the embryo via the placenta were demonstrated. Table VI-20 indicates the tumor types that were correlated to vinyl chloride exposure in experimental animals.

Maltoni (1981) concluded that vinyl chloride may produce tumors of different types at different sites and that the incidence and relative distribution are greatly influenced by dose, age of the animal, and species and strain of animal used.

Table VI-3

Experiment BT1.^a

Group and concentration	Tumors/100 animals		Animals with tumors, %										
	MT	BT	LAS	LA	ELAS	ELA	Hepa- tomas	Nephro- BL	Neuro- BL	Zymbal Gl.Ca	Skin EpT	Fore- stomach Pa&Ac	Mam- mary MT
I 10,000 ppm	81.7	23.3	11.7 (7/60)	-	5.0 (3/60)	5.0 (3/60)	1.7 (1/60)	8.3 (5/60)	11.7 (7/60)	26.7 (16/60)	6.0 (2/60)	-	6.0 (2/60)
II 6000 ppm	60.0	26.3	22.0 (13/60)	3.4 (2/59)	5.1 (3/59)	6.8 (4/59)	1.7 (1/59)	8.5 (5/59)	6.1 (3/59)	11.9 (7/59)	3.4 (2/59)	1.7 (1/59)	-
III 2500 ppm	63.3	20.0	21.7 (13/60)	-	5.0 (3/60)	1.1 (2/60)	1.1 (2/60)	10.0 (6/60)	6.7 (4/60)	3.3 (2/60)	1.7 (1/60)	-	3.3 (2/60)
IV 600 ppm	61.7	12.3	10.0 (6/60)	-	1.7 (1/60)	1.7 (1/60)	8.3 (5/60)	10.0 (6/60)	-	6.7 (4/60)	1.7 (1/60)	-	1.7 (1/60)
V 250 ppm	30.0	25.0	6.1 (3/59)	1.7 (1/59)	3.4 (2/59)	-	1.7 (1/59)	8.5 (5/59)	-	-	3.4 (2/59)	-	3.4 (2/59)
VI 60 ppm	16.0	36.7	1.7 (1/60)	-	1.7 (1/60)	3.3 (2/60)	-	1.7 (1/60)	-	-	1.7 (1/60)	1.7 (1/60)	3.3 (2/60)
VII No treatment (control)	13.3	43.3	-	-	-	3.4 (2/58)	-	-	-	-	1.7 (1/58)	-	-

^aExposure by inhalation to VC in air at 10,000, 6000, 2500, 500, 250, and 50 ppm; 4 hr/day, 5 days/week, for 52 weeks. Sprague-Dawley rats, M and F, 13 weeks old. Results after 135 weeks (end of experiment).

SL 067166

Adapted from Maltoni, 1981

Table VI-4

Experiment BT2.^a

Group and concentration	Tumors/100 animals		Animals with tumors, %										
	MT	BT	LAS	LA	ELAS	ELA	Hepa- tomas	Nephro- BL	Neuro- BL	Zymbal GLCs	Skin EpT	Fore- stomach Pa&Ac	Mam- mary MT
I 200 ppm	26.0	21.7	10.0 (12/120)	3.3 (4/120)	0.8 (1/120)	0.8 (1/120)	2.5 (3/120)	5.8 (7/120)	-	3.3 (4/120)	4.2 (5/120)	-	5.0 (6/120)
II 150 ppm	25.0	25.0	5.0 (6/119)	-	-	0.8 (1/119)	-	9.2 (11/119)	-	3.4 (4/119)	3.4 (4/119)	1.7 (2/119)	5.0 (6/119)
III 100 ppm	21.7	27.5	0.8 (1/120)	0.8 (1/120)	-	-	-	8.3 (10/120)	-	0.8 (1/120)	0.8 (1/120)	3.3 (4/120)	3.3 (4/120)
IV No treatment (control)	15.7	21.6	-	-	1.1 (2/185)	-	-	-	-	1.1 (2/185)	1.1 (2/185)	1.6 (3/185)	1.0 (2/185)

^aExposure by inhalation to VC in air at 200, 150, 100 ppm; 4 hr/day, 5 days/week, for 52 weeks. Sprague-Dawley rats, M and F, 13 weeks old. Results after 143 weeks (end of experiment).

SL 067167

Adapted from Maltoni, 1981

Table VI-5

Experiment BT6.*													
Group and concentration	Animals with tumors, %												
	Tumors/100 animals												
	MT	BT	LAS	LA	ELAS	ELA	Hepa- tomas	Nephro- BL	Neuro- BL	Zymbal GLCa	Skin EpT	Fore- stomach Pa&Ac	Mam- mary MT
I 30,000 ppm	100.0	50.0	30.0 (18/60)	1.7 (1/60)	1.7 (1/60)	5.0 (3/60)	1.7 (1/60)	-	1.7 (1/60)	58.3 (35/60)	1.7 (1/60)	18.3 (11/60)	2.3 (2/60)

*Exposure by inhalation to VC in air at 30,000 ppm; 4 hr/day, 5 days/week, for 52 weeks. Sprague-Dawley rats, M and F, 17 weeks old. Results after 68 weeks (end of experiment).

SL 067168

Adapted from Maltoni, 1981

Table VI-6

Experiment BT9.^a

Group and concentration	Animals with tumors, %												
	Tumors/100 animals												
	MT	BT	LAS	LA	ELAS	ELA	Hepa- tomas	Nephro- Bl.	Neuro- BL	Zymbal GI Ca	Skin EpT	Fore- stomach Pa&Ac	Mam- mary MT
I 50 ppm II	44.3	41.7	4.8 (14/294)	2.7 (8/294)	3.1 (9/294)	3.7 (11/294)	-	0.3 (1/294)	-	3.1 (9/294)	1.0 (3/294)	0.4 (1/294)	21.7 (62/294)
No treatment (control)	23.0	24.0	-	-	-	-	-	-	-	-	-	1.0 (1/98)	10.2 (10/98)

^aExposure by inhalation to VC in air at 50 ppm; 4 hr/day, 5 days/week, for 52 weeks. Sprague-Dawley rats, M and F, 18 weeks old. Results after 142 weeks (end of experiment).

SL 067169

Table VI-7

Experiment BT15.^a

Group and concentration	Animals with tumors, %												
	Tumors/100 animals												
	MT	BT	LAS	LA	ELAS	ELA	Hepa- tomas	Nephro- BL	Neuro- BL	Zymbal GLCa	Skin EpT	Fore- stomach Pad Ac	Mam- mary MT
I	83.3	68.3	4.2	0.8	-	2.5	-	0.8	-	3.3	-	-	16.0
25 ppm			(5/120)	(1/120)		(3/120)		(1/120)		(4/120)			(17/120)
II	81.7	63.3	0.8	-	1.7	2.5	-	-	-	1.7	-	-	17.6
10 ppm			(1/119)		(2/119)	(3/119)				(2/119)			(21/119)
III	85.8	66.0	-	-	-	-	-	-	-	0.8	0.8	-	18.5
5 ppm										(1/119)	(1/119)		(22/119)
IV	22.5	44.2	-	-	-	-	-	-	-	0.8	0.8	-	12.7
1 ppm										(1/118)	(1/118)		(15/118)
V													
No treatment (control)	23.3	37.6	-	-	-	0.8	-	-	-	1.7	-	-	5.8
						(1/120)				(2/120)			(7/120)

^aExposure by inhalation to VC in air at 25, 10, 5, 1 ppm; 4 hr/day, 5 days/week, for 52 weeks. Sprague-Dawley rats, M and F, 13 weeks old. Results after 147 weeks (end of experiment).

SL 067170

Table VI-8

Experiment BTJ.^a

Group and concentration	Tumors/100 animals		Animals with tumors, %										
	MT	BT	LAS	LA	ELAS	ELA	Hepa- toomas	Nephro- BL	Neuro- BL	Zymbal GI Ca	Skin EpT	Fore- stomach Pa&Ac	Mam- mary MT
I 10,000 ppm	45.0	20.0	-	-	-	1.7 (1/58)	1.7 (1/58)	1.7 (1/58)	15.6 (9/58)	15.5 (9/58)	8.6 (5/58)	1.7 (1/58)	1.7 (1/58)
II 8000 ppm	53.3	25.0	1.7 (1/60)	-	-	3.3 (2/60)	1.7 (1/60)	1.7 (1/60)	20.0 (12/60)	15.0 (9/60)	8.3 (5/60)	3.3 (2/60)	1.7 (1/60)
III 2500 ppm	41.7	35.0	1.7 (1/60)	-	-	-	3.3 (2/60)	3.3 (2/60)	8.3 (5/60)	11.7 (7/60)	3.3 (2/60)	-	6.7 (4/60)
IV 500 ppm	15.0	35.0	1.7 (1/60)	-	-	-	-	-	-	1.7 (1/60)	-	-	5.0 (3/60)
V 250 ppm	21.7	25.0	-	1.7 (1/59)	-	1.7 (1/59)	-	10.2 (6/59)	-	1.7 (1/59)	-	5.1 (3/59)	1.7 (1/59)
VI 50 ppm	10.3	25.0	-	1.7 (1/58)	-	1.7 (1/58)	-	5.2 (3/58)	-	-	1.7 (1/58)	-	1.7 (1/58)
VII No treatment (control)	14.7	20.0	-	-	0.5 (1/190)	-	-	-	-	1.0 (2/190)	0.5 (1/190)	-	2.6 (5/190)

^aExposure by inhalation to VC in air at 10,000, 6000, 2500, 500, 250, and 50 ppm; 4 hr/day, 5 days/week, for 17 weeks. Sprague-Dawley rats, M and F, 12 weeks old. Results after 156 weeks (end of experiment).

SL 067171

Adapted from Maltoni, 1981

Table VI-9

Experiment BT10.*													
Group and concentration	Tumors/100 animals		Animals with tumors, %										
	MT	BT	LAS	LA	ELAS	ELA	Hepa- tomas	Nephro- BL	Neuro- BL	Zymbal GLCa	Skin EpT	Fore- stomach Pa&Ac	Man- mary MT
I 10,000 ppm	33.3	41.7	0.8 (1/118)	-	-	0.8 (1/118)	0.8 (1/118)	-	-	7.6 (9/118)	-	2.5 (3/118)	11.0 (13/111)
II 6000 ppm	30.0	45.0	-	0.8 (1/120)	-	1.7 (2/120)	-	0.8 (1/120)	0.8 (1/120)	7.5 (9/120)	-	1.7 (2/120)	10.0 (13/120)
III 10,000 ppm	26.8	45.0	0.8 (1/119)	1.7 (2/119)	-	0.8 (1/119)	-	-	-	7.6 (9/119)	2.5 (3/119)	2.5 (3/119)	13.4 (16/119)
IV 6000 ppm	20.8	39.2	2.6 (3/118)	-	1.7 (2/118)	-	-	-	-	4.2 (5/118)	3.4 (4/118)	1.7 (2/118)	8.3 (11/118)
V 10,000 ppm	41.7	45.0	0.8 (1/119)	1.7 (2/119)	-	0.8 (1/119)	-	0.8 (1/119)	0.8 (1/119)	6.7 (8/119)	0.8 (1/119)	0.8 (1/119)	16.0 (20/119)
VI 6000 ppm	33.3	50.8	0.8 (1/120)	1.7 (2/120)	0.8 (1/120)	-	1.7 (2/120)	0.8 (1/120)	-	7.5 (9/120)	-	0.8 (1/120)	10.0 (12/120)
VII No treatment (control)	16.6	41.0	-	-	-	0.4 (1/227)	-	-	-	-	0.9 (2/227)	2.2 (5/227)	7.5 (17/227)

*Exposure by inhalation to VC in air at 10,000, 6000, ppm; 4 hr/day, 5 days/week, for 5 weeks (groups I and II) or 1 hr/day, 1 day/week, for 25 weeks (groups III and IV) or 4 hr/day, once weekly, for 25 weeks (groups V and VI) (100 hr). Sprague-Dawley rats M and F, 12 weeks old. Results after 154 weeks (end of experiment).

SL 067172

X. REFERENCES

American Public Health Association. 1975. Population residing near plants producing vinyl chloride.

Ames BN, Durston WF, Yamasaki E, Lee FD. 1975. Carcinogens are mutagens: a simple test system combining liver homogenates for activation and bacteria for detection. Proc. Natl. Acad. Sci. 8:2281-2285.

Anderson D, Hodge MCE, Purchase IFH. 1976. Vinyl chloride: dominant lethal studies in male CD-1 mice. Mutat. Res. 40:359-370.

Anderson D, Richardson CR. 1976. Paradichlorobenzene: Cytogenic study in the rat. ICI Report CTL/P/293. November 1976 (Unpublished).

Anderson D, Richardson CR, Purchase IFH, Evans, HJ, O'Riordan ML. 1981. Chromosomal analysis in vinyl chloride exposed workers: Comparisons of the standard technique with the sister chromatid exchange technique. Mutat. Res. 63:137-144.

Anderson D, Richardson CR, Weight TM, Purchase IFH, Adams WGF. 1980. Chromosomal analyses in vinyl chloride exposed workers. Results from analysis 18 and 42 months after an initial sampling. Mutat. Res. 79:151-162.

Anon. 1973. FDA to propose ban on use of PVC for liquor use. Food Chemical News, May 14, pp. 3-5.

Bartsch H, Malaveille C, Montesano R. 1975. Human, rat and mouse liver-mediated mutagenicity of vinyl chloride in S. typhimurium strains. Int. J. Cancer. 15:429-437.

Bartsch H, Montesano R. 1975. Mutagenic and carcinogenic effects of vinyl chloride. Mutat. Res. 32:93-114.

Basalae AV, Vazine AN, Kochetkov AG. 1972. On the pathogenesis of changes developing due to a long-term exposure to the effect of vinyl chloride. Gig. Tr. Prof. Zabol. 16(2):24-27 (in Russian).

Basler A, Rohrborn G. 1980. Vinyl chloride: An example of evaluating mutagenic effects in mammals in vivo after exposure to inhalation. Arch. Toxicol. 45:1-7.

Baxter PJ, Anthony PP, MacSween RNM, Schever PJ. 1977. Angiosarcoma of the liver in Great Britain 1963-1973. Br. Med. J. 2:919-921.

Becker WC. 1979. Letter from B.F. Goodrich, Chemical Division, to J.P. Lehman, Director, Office of Solid Waste, USEPA. Comments on EPA proposed additions to hazardous waste list. 44 FR 49402, August 22, 1979.

Bergman K. 1982. Reactions of vinyl chloride with RNA and DNA of various mouse tissues in vivo. Arch. Toxicol. 49:117-129.

Berk PD, Martin JF, Young RS, Creech J, Selikoff IG, Falk H, Watanabe P, Popper H, Thomas L. 1976. Vinyl chloride-associated liver disease. Ann. Intern. Med. 84:717-731.

Boettner EA, Ball GL, Weiss B. 1973. Combustion products from the incinerator of plastics. EPA Report 670/2-73-049. (PB 222 001).

Bolt HM, Kappus H, Buchter A, Bolt W. 1976. Disposition of (1,2-¹⁴C) vinyl chloride in the rat. Arch. Toxicol. 35:153-162.

Bolt HM, Laib RJ, Kappus H, Buchter A. 1977. Pharmacokinetics of vinyl chloride in the rat. Toxicol. 7:179-188.

Brady J, Liberatore F, Harper P, Greenwald P, Burnett W, Davies JNP, Bishop M, Polan A, Vianna N. 1977. Angiosarcoma of the liver: an epidemiologic survey. J. Natl. Cancer Inst. 59(5):1383-1385.

Braker, W. and A.L. Mossman. 1971. Matheson Gas Data Book, 5th Edition. East Rutherford, N.J., Matheson Gas Products, pp. 561-564.

Brodzinsky R, Singh HB. 1982. Volatile organic chemicals in the atmosphere: an assessment of available data. Prepared by SRI International for Office of Research and Development, USEPA, Research Triangle Park, N.C. Contract no. 68-02-3452.

Buchter A, Filser JG, Peter H, Bolt HM. 1980. Pharmacokinetics of vinyl chloride in rhesus monkeys. Toxicol. Letters. 6:33-36.

Byren D, Engholm G, Englund A, Westerholm P. 1976. Mortality and cancer in a group of Swedish VCM and PVC production workers. Environ. Health Perspec. 17:167-170.

Caputo A, Viola PL, Bigotti A. 1974. Oncogenicity of vinyl chloride at low concentrations in rats and rabbits. IRCS 2:1582.

Chiazze L, Nichols WE, Wong O. 1977. Mortality among employees of PVC fabricators. J. Occup. Med. 19(9):623-628.

Chiazze L Jr, Wong O, Nichols WE, Ference LD. 1980. Breast cancer mortality among PVC fabricators. J. Occup. Med. 22:677-679.

Creech JL, Johnson MN. 1974. Angiosarcoma of the liver in the manufacture of polyvinyl chloride. J. Occup. Med. 16:150-151.

Danzinger H. 1960. Accidental poisoning by vinyl chloride. Can. Med. Assoc. J. 82:828-830.

Delorme F, Theriault G. 1978. Ten cases of angiosarcoma of the liver in Shawinigan, Quebec. J. Occup. Med. 20:338-340.

Dilling WL, Tefertiller NB, Kallos GJ. 1975. Evaporation rates of methylene chloride, chloroform, 1,1,1-trichloroethane, trichloroethylene, tetrachloroethylene and other chlorinated compounds in dilute aqueous solution. Environ. Sci. Technol. 9(9):833-838.

Dinman BD, Cook WA, Whitehouse WM, Magnuson HJ. 1971. Occupational acroosteolysis. Arch. Environ. Health. 22:61-73.

Dressman RC, McFarren EF. 1978. Determination of vinyl chloride migration from polyvinyl chloride pipe into water. Am. Water Works Assoc. J. 70:29-30.

Ducatman A, Hirschhorn K, Selikoff IJ. 1975. Vinyl chloride exposure and human chromosome aberrations. Mutat. Res. 31:163-168.

Duprat P, Fabry JP, Gradiski D, Magadur JL. 1977. Metabolic approach to industrial poisoning: blood kinetics and distribution of ¹⁴C-vinyl chloride monomer (V.C.M.). Acta. Pharmacol. Toxicol. Suppl. (Kbh) 41(1):142-143.

Edmonds LD, Falk H, Nissim JE. 1975. Congenital malformations and vinyl chloride. Lancet 2:1098.

Elmore JD, Wong JL, Laumbach AD, Streips UN. 1976. Vinyl chloride mutagenicity via the metabolites chlorooxirane and chloroacetaldehyde monomer hydrate. Biochem. Biophys. Acta. 442:405-419.

Falk H, Herbert J, Crowley S, Ishak KG, Thomas LB, Popper H, Caldwell GC. 1981. Epidemiology of hepatic angiosarcoma in the United States: 1964-1974. Environ. Health Perspec. 41:107-113.

FDA. 40 FR 40529, September 3, 1975.

Federal Reporting Data System. 1983. Facilities and population served by primary water supply source (FRDS07), April 19, 1983. U.S. Environmental Protection Agency, Washington, D.C.

Feron VJ, Hendrikson CFM, Speek AJ, Til HP, Spit BJ. 1981. Lifespan oral toxicity study of vinyl chloride in rats. Fd. Cosmet. Toxicol. 19:317-331.

Feron VJ, Speek AJ, Willems MI, van Battum D, de Groot AP. 1975. Observations on the oral administration and toxicity of vinyl chloride in rats. Fd. Cosmet. Toxicol. 13:633-638.

Filatova VS, Gronsberg ES. 1957. Sanitary hygienic conditions and work in the production of polychlorvinyl tar and measures of improvement. Gig. I. Sanit. 22:38-42.

Fox AJ, Collier PF. 1977. Mortality experience of workers exposed to vinyl chloride monomer in the manufacture of polyvinyl chloride in Great Britain. Br. J. Ind. Med.

- Garro AJ, Guttanplan JB, Miluy P. 1976. Vinyl chloride dependent mutagenesis: effects of liver extracts and free radicals. *Mutat. Res.* 38:81-88.
- Gay BW, Hanst PL, Bufalini JJ, Noonan RC. 1976. Atmospheric oxidation of chlorinated ethylenes. *Environ. Science Technol.* 10:58-67.
- Green T, Hathway DE. 1975. The biological fate in rats of vinyl chloride in relation to its oncogenicity. *Chem. Biol. Interactions.* 11:545-562.
- Greim H, Bonse G, Radwan Z, Reichert D, Henschler D. 1975. Mutagenicity in vitro and potential carcinogenicity of chlorinated ethylenes as a function of metabolic oxirane formation. *Biochem. Pharmacol.* 24:2013-2017.
- Groth DH, Lynch DW, Moorman WJ, Stettler LE, Lewis TR, Wagner WD, Kouminen C. 1981. Pneumoconiosis in animals exposed to poly (vinyl chloride) dust. *Environ. Health Perspec.* 41:73-83.
- Guangerich FP, Mason P, Stott W, Fox T, Watanabe PG. 1981. Roles of 2-haloethylene oxides and 2-haloacetaldehydes derived from vinyl bromide and vinyl chloride in irreversible binding to protein and DNA. *Cancer Res.* 41:4391-4398.
- Haley TS. 1975. Vinyl chloride: how many unknown problems? *J. Toxicol. Environ. Health* 1:47-73.
- Hansteen I-L, Hillestad L, Thiss-Evensen E, Storevadt Haldos S. 1978. Effects of vinyl chloride in man a cytogenetic follow-up study. *Mutat. Res.* 51:271-278.
- Harris DK, Adams WG. 1967. Acro-osteolysis occurring in men engaged in the polymerization of vinyl chloride. *Br. Med. J.* 3:712-714.
- Hatch M, Kline J, Stein Z. 1981. Power considerations in studies of reproductive effects of vinyl chloride and some structural analogs. *Environ. Health Perspec.* 41:195-201.
- Hathway DE. 1977. Comparative mammalian metabolism of vinyl chloride and vinylidene chloride in relation to oncogenic potential. *Environ. Health Perspect.* 21:55-59.
- Heath CW, Jr, Dumont CR, Gamble J, Waxweiler RJ. 1977. Chromosomal damage in men occupationally exposed to vinyl chloride monomer and other chemicals. *Environ. Res.* 14:68-72.
- Heath CW, Jr, Falk H, Creech JL, Jr. 1975. Characteristics of cases of angiosarcoma of the liver among vinyl chloride workers in the United States. *Ann. N.Y. Acad. Sci.* 246:231-236.
- Hefner RE, Jr, Watanabe PG, Gehring PJ. 1975a. Preliminary studies on the fate of inhaled vinyl chloride monomer in rats. *Ann. NY. Acad. Sci.* 246:135-148.

- Hefner RE, Jr, Watanabe PG, Gehring PJ. 1975b. Percutaneous absorption of vinyl chloride. *Toxicol. Appl. Pharmacol.* 34:529-532.
- Hill J, Kollig HP, Parris DF, Wolfe NL, Zepp RG. 1976. Dynamic behavior of vinyl chloride in aquatic ecosystems. EPA 600/3-76-001. (PB-249 302). 63 pp.
- Hoffman D, Patrianakos C, Brunnemann KD, Gori GB. 1976. Chromatographic determination of vinyl chloride in tobacco smoke. *Anal. Chem.* 48:47-50.
- Holder B. 1974. Dow Chemical Company testimony presented at public hearings on proposed standard for occupational exposure to vinyl chloride. U.S. Dept. Labor, Washington, D.C., June 25.
- Huberman E, Bartsch H, Sachs L. 1975. Mutation induction in Chinese hamster V79 cells by two vinyl chloride metabolites, chloroethylene oxide and 2-chloroacetaldehyde. *Int. J. Cancer.* 16:639-644.
- International Agency for Research on Cancer. 1974. IARC monographs on the evaluation of carcinogenic risk of chemicals to man. Vol. 7, Lyon, France.
- International Agency for Research on Cancer. 1979. IARC monographs on the evaluation of carcinogenic risk of chemicals to man. Vol. 19. Lyon, France.
- International Commission on Radiological Protection. 1975. Report of the task group on reference man. ICRP publication 23. New York:Pergamon Press.
- Infante PF. 1976. Oncogenic and mutagenic risks in communities with polyvinyl chloride production facilities. *Ann. N.Y. Acad. Sci.* 271:49-57.
- Infante PF, Wagoner JK, Waxweiler RJ. 1976a. Carcinogenic, mutagenic, and teratogenic risks associated with vinyl chloride. *Mutat. Res.* 41:131-142.
- Infante PF, Wagoner JK, Waxweiler RJ, McMichael AJ, Falk H. 1976b. Genetic risks of vinyl chloride. *Lancet* 1:734-735.
- Jaeger RJ. 1975. Vinyl chloride monomer: comments on its hepatotoxicity and interaction with 1,1-dichloroethylene. *Ann. N.Y. Acad. Sci.* 246:150-151.
- Jaeger RJ, Reynolds ES, Conolly RB, Moslen MT, Szabo S, Murphy SD. 1974. Acute hepatic injury by vinyl chloride in rats pretreated with phenobarbital. *Nature* 252:724-726.
- John JA, Smith FA, Leong BKJ, Schwetz BA. 1977. The effects of maternally inhaled vinyl chloride on embryonal and fetal development in mice, rats, and rabbits. *Toxicol. Appl. Pharmacol.* 39:497-513.
- John JA, Smith FA, Schwetz BA. 1981. Vinyl chloride: inhalation teratology study in mice, rats and rabbits. *Environ. Health Perspectives* 41:171-179.
- Kappus H, Bolt HM, Buchter A, Bolt W. 1975. Rat liver microsomes catalyze covalent binding of ^{14}C -vinyl chloride to macromolecules. *Nature* 257:134-135.

Kappus H, Bolt HM, Buchter A, Bolt W. 1976. Liver microsomal uptake of (¹⁴C) vinyl chloride and transformation to protein alkylating metabolites in vitro. Toxicol. Appl. Pharmacol. 37:461-471.

Killian DJ, Picciano DJ, Jacobson CB. 1975. Industrial monitoring: a cytogenetic approach. Ann. N.Y. Acad. Sci. 269:4-11.

Kramer CG, Mutchler JE. 1972. The correlation of clinical and environmental measurements for workers exposed to vinyl chloride. Am. Ind. Hyg. Assoc. J. 33:19-30.

Kuzmack AM, McGaughy RE. 1975. Quantitative risk assessment for community exposure to vinyl chloride. Environ. Prot. Agency Rept., Dec. 5., Washington, D.C.

Laib RJ, Bolt HM. 1977. Alkylation of RNA by vinyl chloride metabolites in vitro and in vivo: formation of 1-N'-etheno-adenosine. Toxicology 8:185-195.

Lee CC, Bhandari JC, Winston JM, House WB, Dixon RL, Woods JS. 1978. Carcinogenicity of vinyl chloride and vinylidene chloride. J. Toxicol. Environ. Health 4:15-30.

Lee CC, Bhandari JC, Winston JM, House WB, Peters PJ, Dixon RL, Woods JS. 1977. Inhalation toxicity of vinyl chloride and vinylidene chloride. Environ. Health Perspect. 21:25-32.

Lester D, Greenberg LA, Adams WR. 1963. Effects of single and repeated exposure of humans and rats to vinyl chloride. Am. Ind. Hyg. Assoc. J. 24:265-275.

Letskiewicz F, Johnston P, Macaluso C, Elder R, Yu W, Bason C. 1983. Occurrence of vinyl chloride in drinking water, food and air. Prepared by JRB Associates for Office of Drinking Water, USEPA, EPA No. 68-01-6388.

Lillis R, Anderson H, Nicolson WJ, Daum S, Fischbein AS, Selikoff IJ. 1975. Prevalance of disease among vinyl chloride and polyvinyl chloride workers. Ann. N.Y. Acad. Sci. 246:22-41.

Lillian D, Singh HB, Appleby A, Lobban L, Arnts R, Bumpert R, Hague R, Toomey J, Kazazis J, Antell M, Hansen D, Scott B. 1975. Atmospheric fates of halogenated compounds. Environ. Sci. Technol. 9:1042-1048.

Loprieno N, Barale R, Baroncelli S, Bartsch H, Bronzetti G, Cammellini A, Corsi C, Frezza D, Nieri R, Leporini C, Rosellini D, Rossi AM. 1977. Induction of gene mutations and gene conversions by vinyl chloride metabolites in yeast. Cancer Res. 253-257.

Loprieno N, Barale R, Baroncelli S, Bauer C, Bronzetti G, Cammellini A, Cercingnani G, Corsi C, Gervasi G, Leporini C, Nieri R, Rossi AM, Stretti G, Turchi G. 1976. Evaluation of the genetic effects induced by vinyl chloride monomer (VCM) under mammalian metabolic activation: studies in vitro and in vivo. Mutat. Res. 40:85-96.

Lu PY, Metcalf RL, Plummer N, Mandel D. 1977. The environmental fate of three carcinogens: benzo(a)pyrene, benzidine, and vinyl chloride evaluated in laboratory model ecosystems. Arch. Environ. Contam. Toxicol. 6:129-142.

Makk L, Delmore F, Creech JL, Jr, Ogden LL, Fadell EH, Songster CL, Clanton J, Johnson MN, Christopherson WM. 1976. Clinical and morphologic features of hepatic angiosarcoma in vinyl chloride workers. Cancer 37:149-163.

Maltoni C. 1977. Vinyl chloride carcinogenicity: an experimental model for carcinogenesis studies. In: Hiatt HH, JD Watson, JA Winsten (eds.). Origins of Human Cancer, Book A. Cold Spring Harbor Laboratory, 119-146.

Maltoni C, Ciliberti, Gianni L, Chieco P. 1975. The oncogenic effects of vinyl chloride administered by oral route in the rat. Gli Ospedali della Vita 2(6):65-66.

Maltoni C, Lefemine G. 1974. Carcinogenicity bioassays of vinyl chloride. 1. Research plan and early results. Environ. Res. 7:387-405.

Maltoni C, Lefemine G, Ciliberti A, Cotti G, Carretti D. 1981. Carcinogenicity bioassays of vinyl chloride monomer: a model of risk assessment on an experimental basis. Environ. Health Perspec. 41:3-31.

Manusco TF. 1975. Comments for opening of discussion on "neoplastic effects." Ann. N.Y. Acad. Sci. 246:251-254.

Marsteller HJ, Leibach WK, Muller R, Gedigk P. 1975. Unusual splenomegalic liver disease as evidenced by peritoneoscopy and guided liver biopsy among polyvinyl chloride production workers. Ann. N.Y. Acad. Sci. 246:95-134.

Mastromatteo E, Fisher AM, Christie H, Danziger H. 1960. Acute inhalation toxicity of vinyl chloride to laboratory animals. Amer. Ind. Hyg. Assoc. J. 21:394-398.

McCann J, Simmon V, Streitwieser D, Ames BN. 1975. Mutagenicity of chloroacetaldehyde, a possible metabolic product of 1,2-dichloroethane (ethylene dichloride), chloroethanol (ethylene chlorohydrin), vinyl chloride, and cyclophosphamide. Proc. Natl. Acad. Sci. 72:3190-3193.

McConnell GD, Ferguson M, Pearson CR. 1975. Chlorinated hydrocarbons and the environment. Endeavor 34:13.

Milby TH (ed.). 1978. Vinyl chloride: an information resource. DHEW Pub. No. (NIH) 78-1599.

Miller A, Teirsten AS, Chuang M, Selikoff IJ, Warshaw R. 1975. Changes in pulmonary function in workers exposed to vinyl chloride and polyvinyl chloride. Ann. N.Y. Acad. Sci. 246:42-52.

Monson RR, Peters JM, Johnson MN. 1975. Proportional mortality among vinyl chloride workers. Environ. Health Perspec. 11:75-77.

National Academy of Sciences. 1977. Drinking Water and Health. Washington, DC. 783-787.

National Academy of Sciences. 1983. Drinking Water and Health, Vol. 5. National Academy Press, Washington, DC.

Newberne, PM. 1982. Assessment of the hepatocarcinogenic potential of chemicals: response of the liver, In: Plaa, G and W. R. Hewitt (eds.). Toxicology of the liver. Raven Press. New York, 266-267.

Nicholson WJ, Hammond EC, Seidman H, Selikoff IL. 1975. Mortality experience of a cohort of vinyl chloride-polyvinyl chloride workers. Ann. N.Y. Acad. Sci. 246:225-230.

Oster RH, Carr CJ, Krantz JC, Jr. 1947. Anesthesia XXVII. Narcosis with vinyl chloride. Anesthesiology 8:359-361.

Ott MG, Langner RR, Holder BB. 1975. Vinyl chloride exposure in controlled industrial environment: a long-term mortality experience in 594 employees. Arch. Environ. Health 30:333-339.

Patty FA, Yant WP, Waite CP. 1930. Acute response of guinea pigs to vapors of some new commercial organic compounds. V. Vinyl chloride. Pub. Health Reports 45:1963-1971.

Peoples AS, Leake CD. 1933. The anesthetic action of vinyl chloride. J. Pharmacol. Exp. Therapeutics 48:284.

Picciano DJ, Flake RE, Gay PC, Killian DJ. 1977. Vinyl chloride cytogenetics. J. Occup. Med. 19:527-530.

Popper H, Thomas LB. 1975. Alteration of liver and spleen among workers exposed to vinyl chloride. Ann. N.Y. Acad. Sci. 246:172-194.

Proceedings of the Royal Society of Medicine. 1976. 69:275-310.

Prodan L, Suciu I, Pislaru V, Ilea E, Pascu L. 1975. Experimental acute toxicity of vinyl chloride (monochloroethane). Ann. N.Y. Acad. Sci. 246:154-158.

Purchase IFH, Richardson CR, Anderson D. 1975. Chromosomal and dominant lethal effects of vinyl chloride. Lancet 2(7931):410-411.

Purchase IFH, Richardson CR, Anderson D, Paddle GM, Adams WGF. 1978. Chromosomal analyses in vinyl chloride-exposed workers. Mut. Res. 57:325-334.

Radike M. 1977. Transplacental effects of vinyl chloride in rats. Annual Report, pp. 183-185. USPHS-ES-00159. Center for Study of the Human Environment, Dept. Environ. Health, University of Cincinnati Medical Center.

Radike MJ, Stemmer KL, Brown PG, Larson E, Bingham E. 1977. Effect of ethanol and vinyl chloride on the induction of liver tumors: preliminary report. Environ. Health Perspect. 21:153-155.

Radike MJ, Stemmer KL, Bingham E. 1981. Effect of ethanol on vinyl chloride carcinogenesis. Environ. Health Perspect. 41:59-63.

Rowe VK. 1975. Experience in industrial exposure control. Ann. N.Y. Acad. Sci. 246:306-310.

Rannug U, Johansson A, Ramel C, Wachtmeister CA. 1974. The mutagenicity of vinyl chloride after metabolic activation. Ambio 3:194-197.

Selikoff IJ, Hammond EC, eds. 1975. Toxicity of vinyl chloride-polyvinyl chloride. Ann. N.Y. Acad. Sci., Vol. 246.

Spiro R, Kaminski R. 1978. Angiosarcoma of the liver in vinyl chloride/polyvinyl chloride workers. Update of the NIOSH Register. J. Occup. Med. 20:427-429.

Spiro R, McMichael AJ, Gamble J, Van Ert M. 1975. The association of vinyl chloride exposures with morbidity symptoms. Amer. Ind. Hyg. Assoc. J. 36:779-789.

Suciu I, Prodan L, Ilea E, Paduraru A, Pascu L. 1975. Clinical manifestations in vinyl chloride poisoning. Ann. N.Y. Acad. Sci. 246:53-69.

Tabershaw IR, Gaffey WR. 1974. Mortality study of workers in the manufacture of vinyl chloride and its polymers. J. Occup. Med. 116:509-518.

Til HP, Immel HR, Feron VJ. 1983. Lifespan oral carcinogenicity study of vinyl chloride in rats. Final report. Civo Institutes TNO. Report No. V 83.285/291099.

Torkelson RR, Oyen F, Rowe VK. 1961. The toxicity of vinyl chloride as determined by repeated exposure of laboratory animals. Amer. Ind. Hyg. Assoc. J. 22:354-361.

Tribukh SL, Tikhomirova NP, Levina SV, Kozlov LA. 1949. Working conditions and measures for their sanitation in the production and utilization of vinyl chloride plastics. Gig. Sanit. 10:38-44 (in Russian).

U.S. Department of Agriculture, Agricultural Research Service, Household Food Consumption Survey, 1965-1966, Food and Nutrient Intake of Individuals of the United States.

U.S. Environmental Protection Agency. 1974. Preliminary assessment of the environmental problems associated with vinyl chloride and polyvinyl chloride (appendices). Report on the Activities and Findings of the Vinyl Chloride Task Force. EPA 560/4-74-001, (PB-39 110). 67 pp.

U.S. Environmental Protection Agency. 1975a. Preliminary assessment of suspected carcinogens in drinking water. EPA 560/4-75-003. (PB-244 415). pp 33.

U.S. Environmental Protection Agency. 1975b. Standard Support and Environmental Impact Statement: Emission Standard for Vinyl Chloride. EPA 600/6-75-009. (PB-249 703). p. 536.

U.S. Environmental Protection Agency. 1975c. Scientific and technical assessment report on vinyl chloride and polyvinyl chloride. EPA 600/6-75-004. (PB-249 461). p. 116.

U.S. Environmental Protection Agency. 1975d. National Organics Reconnaissance Survey (Office of Drinking Water), Journal of the American Water Works Association, 67, 11, 634-647, November 1975 and 67, 12, 208.

U.S. Environmental Protection Agency. 1977a. The National Organic Monitoring Survey. Interim Report, Office of Drinking Water. p. 126.

U.S. Environmental Protection Agency. 1977b. Survey of Operating and Financial Characteristics of Community Water Systems (Temple, Barker and Sloane).

U.S. Environmental Protection Agency. 1978. Compilation of data from: A Preliminary Report on the Findings of the State Ground Water Monitoring Project and a Second Preliminary Report on the Findings of the State Ground Water Monitoring Project, State of New Jersey, Department of Environmental Protection.

U.S. Environmental Protection Agency. 1980a. Memo to Joseph Cotruvo from Robert McGaughey, September 17, 1980, Washington, D.C.: Carcinogen Risk for Pollutants in the Drinking Water -- Comparison of Results Obtained by the National Academy of Sciences and EPA's Water Criteria Program.

U.S. Environmental Protection Agency. 1980b. The Occurrence of Volatile Organics in Drinking Water (Office of Drinking Water).

U.S. Environmental Protection Agency. 1980c. Survey of EPA Regional Drinking Water Representatives to Determine the Ground Water Monitoring Data Developed by State Agencies.

U.S. Environmental Protection Agency. 1980d. Compilation of Incidents of Drinking Water Contamination with Volatile Organic Chemicals (Office of Drinking Water).

U.S. Environmental Protection Agency. 1980e. Ambient Water Quality Criteria for Vinyl Chloride (Office of Water Regulations and Standards). EPA 440/5-80-078.

U.S. Environmental Protection Agency. 1981a. Community Water Supply Survey (Office of Drinking Water).

U.S. Environmental Protection Agency. 1981b. National Organics Screening Program (SRI).

Van Esch GJ, Van Logten MJ. 1975. Vinyl chloride: a report of a European assessment. *Fd. Cosmet. Toxicol.* 13:121-124.

Verburgt FG, Vogel E. 1977. Vinyl chloride mutagenesis in Drosophila melanogaster. *Mutat. Res.* 48:327-333.

Viola PL, Bigotti A, Caputo A. 1971. Oncogenic response of rat skin, lungs, bones to vinyl chloride. *Cancer Res.* 31:516-522.

Ward AM, Udnoon S, Watkins J, Walker AE, Drake CS. 1976. Immunological mechanisms in the pathogenesis of vinyl chloride disease. *Br. Med. J.* 1:936-938.

Watanabe PG, McGowan GR, Gehring PJ. 1976a. Fate of (¹⁴C) vinyl chloride after single oral administration in rats. *Toxicol. Appl. Pharmacol.* 36:339-352.

Watanabe PG, McGowan GR, Madrid EO, Gehring PJ. 1976b. Fate of (¹⁴C) vinyl chloride following inhalation exposure in rats. *Toxicol. Appl. Pharmacol.* 37:49-59.

Waxweiler RJ, Stringer W, Wagoner JK, Jones J, Falk H, Carter C. 1976. Neoplastic risk among worker exposed to vinyl chloride. *Ann. N.Y. Acad. Sci.* 271:40-48.

Weisburger JH, Williams GM. 1981. Carcinogen testing: current problems and new approaches. *Sci.* 214:401-407.

Wilson RH, McCormick WE, Tatum CF, Creech JL. 1967. Occupational acroosteolysis. Report of 31 cases. *Amer. Med. Assoc. J.* 201:577-581.

Withey JR. 1976. Pharmacodynamics and uptake of vinyl chloride monomer administered by various routes to rats. *J. Toxicol. Environ. Health* 1:381-394.

Withey JR, Collins BT. 1976. A statistical assessment of the quantitative uptake of vinyl chloride monomer from aqueous solution. *J. Toxicol. Environ. Health* 2:311-321.

Table VI-10

Experiment BT5.*

Group and concentration	Tumors/100 animals		Animals with tumors, %										
	MT	BT	LAS	LA	ELAS	ELA	Hepa- tomas	Nephro- BL	Neuro- HL	Zymbal GL Co	Skin EpT	Fore- stomach Pa&Ar	Mam- mary MT
I 10,000 ppm	6.7	36.7	-	-	-	-	-	-	-	3.3 (1/30)	-	-	-
II 6000 ppm	6.7	21.3	-	-	-	-	-	-	-	-	-	-	-
III 10,000 ppm	29.6	22.2	-	-	-	-	-	5.9 (3/51)	-	9.8 (5/51)	-	2.0 (1/51)	2.0 (1/51)
IV 6000 ppm	21.9	46.9	-	-	-	3.1 (1/32)	-	-	-	9.4 (3/32)	3.1 (1/32)	3.1 (1/32)	6.2 (2/32)

*Exposure by inhalation to VC in air at 10,000, and 6000 ppm of breeders; 4 hr/day for 1 week (from 12th to 18th day of pregnancy) Sprague-Dawley rats, M and F, 19 weeks old (breeders). Breeders (groups I and II) and offsprings (groups III and IV). Results after 143 weeks (end of experiment).

SL 067184

Table VI-11

Group and concentration	Experiment BT14. ^a												
	Tumors/100 animals		Animals with tumors, %										
	MT	BT	LAS	LA	ELAS	ELA	Hepa- tomas	Nephro- BL	Neuro- BL	Zymbal GLCa	Skin EpT	Fore- stomach Pa&Ac	Mam- mary MIT
I 10,000 ppm (breeders)	16.7	66.7	-	-	-	-	-	-	-	-	-	-	-
II 6000 ppm (breeders)	-	-	-	-	-	-	-	-	-	-	-	-	-
III 10,000 ppm (newborn)	100.0	55.5	34.1 (15/44)	-	-	6.8 (3/44)	45.4 (20/44)	-	-	2.3 (1/44)	2.3 (1/44)	-	-
IV 6000 ppm (newborn)	109.3	58.1	40.5 (17/42)	2.4 (1/42)	2.4 (1/42)	2.4 (1/42)	47.6 (20/42)	-	-	4.8 (2/42)	4.8 (2/42)	-	2.4 (1/42)

^aExposure by inhalation to VC in air at 10,000 and 6000 ppm, 4 hr/day, 5 days/week, for 5 weeks (from 1 day to 5 weeks of age) Sprague-Dawley rats, M and F, 21 weeks old (breeders) (groups I and II) and newborn (groups III and IV). Results after 124 weeks (end of experiment).

SL 067185

Adapted from Maltoni, 1981

Table VI-12

Experiment BT7.*

Group and concentration	Tumors/100 animals		Animals with tumors, %									
	MT	BT	LAS	LA	ELAS	ELA	Hepa- tomas	Nephro- BL	Neuro- BL	Zymbal Gl. Co	Skin EpT	Fore- stomach Pa & Ac
I	50.0	10.0	29.6	-	-	-	-	3.7	11.1	7.4	-	-
10,000 ppm			(8/27)					(1/27)	(3/27)	(2/27)		
II	53.3	20.0	11.5	7.7	3.8	3.8	7.7	7.7	3.8	7.7	-	-
6000 ppm			(3/26)	(2/26)	(1/26)	(1/26)	(2/26)	(2/26)	(1/26)	(2/26)		
III	26.7	13.3	12.0	-	4.0	-	4.0	-	4.0	-	4.0	-
2500 ppm			(3/25)		(1/25)		(1/25)		(1/25)		(1/25)	
IV	20.0	10.0	10.7	3.6	-	-	-	7.1	-	-	-	-
500 ppm			(3/28)	(1/28)				(2/28)				
V	13.3	16.7	3.7	-	3.7	3.7	-	-	-	-	3.7	-
250 ppm			(1/27)		(1/27)	(1/27)					(1/27)	
VI	16.7	6.7	-	-	-	-	-	3.6	-	-	-	-
50 ppm								(1/28)				
VII												
No treatment (control)	15.0	15.0	-	-	2.6	-	-	-	-	-	-	-
					(1/38)							

*Exposure by inhalation to VC in air at 10,000, 6000, 2500, 500, 250, and 50 ppm, 4 hr/day, 5 days/week, for 52 weeks. Wistar rats, M, 11 weeks old. Results after 165 weeks (end of experiment)

SL 067186

Adapted from Maltoni, 1981

Table VI-13

Experiment BT17.^a

Group and concentration	Tumors/100 animals		Animals with tumors, %									
	MT	BT	LAS	LA	ELAS	ELA	Hepa- tomas	Nephro- BL	Neuro- BL	Zymbal GLCa	Skin EpT	Fore- stomach Ps&Ac
I 1 ppm II	24.2	29.2	-	1.0 (1/99)	3.0 (3/99)	5.0 (5/99)	1.0 (1/99)	-	-	2.0 (2/99)	-	-
No treatment (control)	20.0	18.5	-	-	-	-	-	-	-	3.2 (3/94)	-	1.1 (1/94)

^aExposure by inhalation to VC in air at 1 ppm; 4 hr/day, 5 days/week, for 52 weeks. Wistar rats, M, 13 weeks old. Results after 134 weeks (end of experiment).

SL 067187

Adapted from Maltoni, 1981

Table VI-14

Experiment HT4.^a

Group and concentration	Tumors/100 animals		Animals with tumors, %							Fore-stomach Pa&Ac
	MT	BT	LAS	LA	ELAS	ELA	Lung T	Mammary Ca	Skin EpT	
I	50.0	98.3	17.8	10.7	1.8	7.1	82.1	21.2	7.1	1.8
10,000 ppm			(10/56)	(6/56)	(1/56)	(4/56)	(46/56)	(13/56)	(4/56)	(1/56)
II	56.7	100.0	21.7	11.7	1.7	5.0	78.3	13.3	11.7	1.7
6000 ppm			(13/60)	(7/60)	(1/60)	(3/60)	(47/60)	(8/60)	(7/60)	(1/60)
III	48.3	90.0	27.1	8.5	1.5	1.7	67.8	13.5	6.8	1.7
2500 ppm			(16/59)	(5/59)	(8/59)	(1/59)	(40/59)	(8/59)	(4/59)	(1/59)
IV	68.3	103.3	23.3	8.3	1.7	5.0	83.3	13.3	3.3	-
500 ppm			(14/60)	(5/60)	(7/60)	(3/60)	(50/60)	(8/60)	(2/60)	-
V	63.3	98.3	30.0	18.3	6.0	5.0	68.3	20.0	1.7	1.7
250 ppm			(18/60)	(11/60)	(3/60)	(3/60)	(41/60)	(12/60)	(1/60)	(1/60)
VI	28.3	23.3	1.7	1.7	1.7	8.3	10.0	20.0	-	1.7
50 ppm			(1/60)	(1/60)	(1/60)	(5/60)	(6/60)	(12/60)	-	(1/60)
VII										
No treatment (control)	14.7	14.7	-	-	0.7	0.7	10.0	0.7	1.3	-
					(1/150)	(1/150)	(15/150)	(1/150)	(2/150)	

^aExposure by inhalation to VC in air at 10,000, 6000, 2500, 500, 250, and 50 ppm, 4 hr/day, 5 days/week, for 30 weeks. Swiss mice, M and F, 11 weeks old. Results after 81 weeks (end of experiment).

SL 067188

Table VI-15

Experiment BTR.^a

Group and concentration	Tumors/100 animals		Animals with tumors, %										
	MT	BT	LAS	LA	ELA	Hepa- tomas	Cholan- go-Ca	Cholan- giomas	Acoustic Duct EpT	Skin EpT	Mela- nomas	Fore- stomach Pad Ac	Leuka- mia ¹
I 10,000 ppm	50.0	73.3	-	3.3 (1/30)	6.7 (2/30)	-	6.7 (2/30)	13.3 (4/30)	3.3 (1/30)	23.3 (7/30)	3.3 (1/30)	23.3 (10/30)	16.7 (5/30)
II 6000 ppm	40.0	63.3	3.3 (1/30)	3.3 (1/30)	-	3.3 (1/30)	6.7 (2/30)	16.7 (5/30)	6.7 (2/30)	3.3 (1/30)	6.7 (2/30)	23.3 (10/30)	20.0 (6/30)
III 2500 ppm	43.3	103.3	-	6.7 (2/30)	-	-	-	26.7 (8/30)	3.3 (1/30)	10.0 (3/30)	3.3 (1/30)	66.7 (17/30)	20.0 (6/30)
IV 500 ppm	53.3	63.3	6.7 (2/30)	-	3.3 (1/30)	-	-	20.0 (6/30)	10.0 (3/30)	23.3 (7/30)	-	30.0 (9/30)	16.7 (5/30)
V 250 ppm	30.0	43.3	-	-	3.3 (1/30)	-	-	20.0 (6/30)	-	10.0 (3/30)	3.3 (1/30)	13.3 (4/30)	20.0 (6/30)
VI 50 ppm	50.0	40.0	-	-	-	-	-	23.3 (7/30)	-	30.0 (9/30)	3.3 (1/30)	10.0 (3/30)	20.0 (6/30)
VII No treatment (control)	20.0	46.7	-	-	-	-	-	36.7 (12/30)	-	5.0 (3/60)	-	5.0 (2/60)	13.3 (8/60)

^aExposure by inhalation to VC in air at 10,000, 6000, 2500, 500, 250, and 50 ppm; 4 hr/day, 5 days/week, for 30 weeks. Golden hamsters, M, 11 weeks old. Results after 109 weeks (end of experiment).

^bLatency time in weeks: Group I, 16.7; Group II, 27.2; Group III, 30.8; Group IV, 19.0; Group V, 22.5; Group VI, 35.3; Group VII 36.5.

SL 067189

Adapted from Maltoni, 1981

Table VI-16

Experiment BT11.*

Group and concentration	Tumors/100 animals		Animals with tumors, %										
	MT	BT	LAS	LA	ELAS	ELA	Hepa- tomas	Nephro UL	Neuro- BL	Zymbal Gl Ca	Skin EpT	Fore- stomach Pa&Ac	Mam mary MT
I 50.00 mg/kg	38.7	35.0	21.2 (17/80)	3.7 (3/80)	2.5 (2/80)	2.5 (2/80)	-	2.5 (2/80)	-	1.2 (1/80)	1.2 (1/80)	2.5 (2/80)	5.0 (4/80)
II 16.65 mg/kg	30.0	17.6	12.5 (10/80)	-	-	-	-	3.7 (3/80)	-	2.5 (2/80)	-	1.2 (1/80)	7.5 (6/80)
III 3.33 mg/kg	10.0	25.0	-	-	2.5 (2/80)	1.2 (1/80)	-	-	-	-	-	-	3.7 (3/80)
IV Olive oil (control)	13.7	22.5	-	-	-	-	-	-	-	1.2 (1/80)	1.2 (1/80)	-	5.0 (4/80)

*Exposure by ingestion (stomach tube) of VC in olive oil at 50.00, 16.65 and 3.33 mg/kg body weight, once daily, 4-5 days/week, for 52 weeks. Sprague-Dawley rats, M and F, 13 weeks old. Results after 136 weeks (end of experiment).

SL 067190

Table VI-17

Experiment BT27.*

Group and concentration	Tumors/100 animals		Animals with tumors, %										
	MT	BT	LAS	LA	ELAS	ELA	Hepa- tomas	Nephro- BL	Neuro- BL	Zymha- GI Ca	Skin EpT	Fore- stomach Pa&Ac	Mam- mary MT
I	24.7	35.3	2.0	-	0.7	-	0.7	-	-	3.3	-	2.0	8.0
1.0 mg/kg			(3/149)		(1/149)		(1/149)			(5/149)		(3/149)	(12/149)
II	13.3	28.0	0.7	0.7	-	-	0.7	-	-	-	0.7	1.3	2.7
0.3 mg/kg			(1/148)	(1/148)			(1/148)				(1/148)	(2/148)	(4/148)
III	18.0	31.3	-	-	-	-	-	-	-	-	0.7	0.7	9.3
0.03 mg/kg											(1/150)	(1/150)	(14/150)
IV													
Olive oil (control)	16.0	28.7	-	-	-	-	-	-	-	0.7	-	1.3	4.7
										(1/150)		(2/150)	(7/150)

*Exposure by ingestion (stomach tube) of VC in olive oil at 1.0, 0.3, 0.03 mg/kg body weight, once daily, 4-5 days/week, for 69 weeks. Sprague-Dawley rats, M and F, 10 weeks old. Results after 136 weeks (end of experiment).

SL 067191

Adapted from Maltoni, 1981

Table VI-18

Experiment BT12.^a

Group and dose	Tumors/100 animals		Animals with tumors, %										
	MT	BT	LAS	LA	ELAS	ELA	Hepa- tomas	Nephro- BL	Neuro- BL	Zymbal GI Ca	Skin EpT	Fore- stomach Pa&Ac	Mam- mary MT
I 4.25 mg x 4	18.8	25.0	-	-	-	-	-	-	-	-	-	1.8 (1/56)	1.8 (1/56)
II 4.25 mg x 3	16.7	28.9	-	-	1.9 (1/53)	-	-	-	-	-	-	1.9 (1/53)	1.9 (1/53)
III 4.25 mg x 2	11.7	18.2	-	-	1.8 (1/56)	-	-	-	-	-	1.8 (1/56)	-	6.3 (3/56)
IV 4.25 mg x 1	20.0	36.0	-	-	-	1.8 (1/55)	-	-	-	1.8 (1/55)	-	3.6 (2/55)	3.6 (2/55)
V Olive oil (control)	8.3	31.7	-	-	-	-	-	-	-	-	-	3.6 (2/55)	-

^aExposure by intraperitoneal injection of VC, 4.25 mg in olive oil (1 ml), 4, 3, 2 times, at two month intervals or once only. Sprague-Dawley rats, M and F, 17 weeks old. Results after 144 weeks (end of experiment).

SL 067192

Adapted from Maltoni, 1981

Table VI-19

Experiment DT13.^a

Group and dose	Tumors/100 animals		Animals with tumors, %										
	MT	DT	LAS	LA	ELAS	ELA	Hepa- tomas	Nephro- BL	Neuro- DL	Zymbal GI Co	Skin EpT	Fore- stomach Pa&Ac	Mam- mary MT
I 4.25 mg	16.0	17.3	-	-	-	-	-	1.3 (1/75)	-	-	-	-	4.0 (3/75)
II Olive oil (control)	13.3	26.7	-	-	-	1.3 (1/75)	-	-	-	1.3 (1/75)	-	-	1.3 (1/75)

^aExposure by subcutaneous injection of VC, 4.25 mg, in olive oil (1 ml), single dose. Sprague-Dawley rats, M and F, 21 weeks old.
Results after 146 weeks (end of experiment)

SL 067193

Adapted from Maltoni, 1981

Table VI-20 Tumors Presently Correlated to VC Exposure (by Inhalation) on Experimental Rodents

Species	Angio-sarcomas of liver	Tumors of brain	Tumors of lung	Lymphomas and leukemias	Hepa-tomas	Angio-sarcomas and an-giomas of other sites	Nephro-blasto-mas	Schia-crous cuta-neous carci-nomas	Other cuta-neous epith-e-lal tumors	Mam-mary car-cinomas	Foresto-mach pa-pillomas and acan-thomas	Mela-nomas
Rot	+	+			+	+	+	+	(+)	+	+	
Mouse	+		+			+			(+)	+	(+)	
Hamster	+			(+)		(+)			(+)		+	(+)

SL 067194

Adapted from Maltoni, 1981

Abbreviations used in tables^a

<u>Abbreviation</u>	
T	Tumor
Cs	Carcinoma
Ep ?	Epithelioma
Pp	Papilloma
Ac	Acanthoma
Ad	Adenoma
Ad †	Adenoma in malignant transformation
MT	Malignant tumors (total if not otherwise specified)
BT	Benign tumors (total if not otherwise specified)
LAS	Liver angiomas
LA	Liver angioma
ELAS	Extra-liver angiomas
ELA	Extra-liver angioma
Nephro-BL	Nephroblastoma
Neuro-BL	Neuroblastoma
A	Angioblastic hyperplasia in liver
A †	Angioblastic dysplasia in liver
Nod. nod.	Neoplastic nodules of liver
Nod. hyp.	Nodular hyperplasia of liver
Diff. hyp.	Diffused hyperplasia of liver
++	Marked
+++	Very marked

^aThe incidence of total malignant and benign tumors is given as the total number of tumors per 100 animals (one animal may bear more than one malignant or benign tumor) on the basis of the tumors observed among the animals alive, when the first tumor was observed in the experiment.

The incidence of specific tumor is given, as percent of the animals bearing the tumor considered, referred to the animals alive when the first tumor was observed (in parenthesis).

Adapted from Maltoni, 1981

SL 067195

A recently completed study by Feron et al. (1981) examined the oral toxicity of vinyl chloride in Wistar rats. This study was carried out over the lifespan of the rats, and consisted of incorporating PVC powder containing a high content of vinyl chloride monomer (VCM) in the diet, or using gastric intubation of a 10% VCM in soybean oil. The VCM doses (actual exposures) were 0, 1.7, 5.0 and 14.1 mg/kg bw through the diet or 300 mg/kg bw by gastric intubation. The results showed that rats exposed to VCM at levels of 5.0 mg/kg bw day or more demonstrated hepatic angiosarcomas, pulmonary angiosarcomas, and at the higher levels, a few primary extrahepatic abdominal angiosarcomas. At the lowest exposure level of 1.7 mg vinyl chloride monomer/kg bw/day, liver-cell tumors and an increased incidence of foci of cellular alteration were noted (Tables VI-21 and VI-22).

Feron et al. (1981) concluded that VCM is a carcinogen when administered by the oral route, and that the tumor response seems to shift from the exclusive development of angiosarcomas at very high levels to the exclusive induction of hepatocellular tumors at low levels of exposure. Feron et al. have also initiated a similar lifespan oral carcinogenicity study with vinyl chloride in rats, using three different dose levels (0.017, 0.17 and 1.7 mg VCM/kg bw day) and two control groups.

Til et al. (1983) examined the oral carcinogenicity of VCM in a lifespan study (149 weeks) with five groups of Wistar rats, each consisting of 100 males and 100 females, except for the top-dose group which comprised 50 males and 50 females. VCM was administered by incorporating PVC powder with a high VCM content into the diet. The diet was provided daily for a period of four

Table VI-21 Type and Incidence of Treatment-Related Histopathological Changes in the Liver of Rats Exposed Orally to VCM

		Incidence of change									
		Males					Females				
Type of change†	Treatment group (mg VCM/kg/day)...	0	1.7	5.0	14.1	300‡	0	1.7	5.0	14.1	300‡
Animals killed after 26 wk											
Clear-cell foci	No. of rats examined...	10	‡	—	10	9	10	—	—	10	10
		0	—	—	1	1	0	—	—	5**	2
Animals killed after 52 wk											
Clear-cell foci	No. of rats examined...	9	—	—	10	9	9	—	—	10	8
		1	—	—	8**	0	0	—	—	8**	0
Basophilic foci		0	—	—	0	0	0	—	—	4	1
Eosinophilic foci		0	—	—	2	0	0	—	—	5**	
Neoplastic nodule		0	—	—	1	0	0	—	—	2	0
Hepatocellular carcinoma		0	—	—	1	0	0	—	—	1	0
Cystic proliferation of bile ducts		0	—	—	0	0	0	—	—	4*	0
Animals found dead or killed in extremes or terminally											
Clear-cell foci	No. of rats examined...	55	58	56	59	55	57	58	59	57	54
		0	9**	16***	21***	9	4	24***	22***	36**	10*
Basophilic foci		8	18	21*	22**	12	0	33***	17	28**	19
Eosinophilic foci		3	23***	27***	33***	11	8	35***	20*	29***	6
Neoplastic nodule		0	1	7**	23***	3	2	26**	39***	44***	2
Hepatocellular carcinoma		0	1	2	8**	1	0	4	19**	29*	0
Angiosarcoma		0	0	6*	27***	27	0	0	2	9**	29

Table VI-21 - continued

Proliferation of atypical sinusoidal cells only	2	0	4	7	6	4	6	3	4	7
Extensive necrosis	4	4	8	23***	21	5	6	19***	27***	24
Cysts	2	3	4	16***	3	9	30***	41***	49**	3
Liver-cell polymorphism	4	16*	28***	42***	36	34	51*	38	41	41
Centrilobular degeneration	0	0	0	1	1	1	2	3	1	18
Focal haematopoiesis	0	1	0	10**	8	1	3	1	6	12

*Specific hepatocellular lesions were classified according to Squire & Levitt (1975).

†The figures of this group were not evaluated statistically because no corresponding control group was included in the study.

‡Not examined.

The initial number of animals was 60/sex/group. A number of rats could not be examined because of cannibalism advanced autolysis.

Values marked with asterisks differ significantly from those of the controls according to the chi-square test:

*p<0.05; **p<0.01; ***p<0.001.

Adapted from Feron et al. 1981.

Table VI-22 Site, Type and Incidence of Tumors in Organs Other than the Liver in Rats
Exposed Orally to VCM for Over 2 to 5 Years

Site and type of tumour	Treatment group (mg VCM/kg/day)...	Incidence of tumours									
		Males					Females				
		0	1.7	5.0	14.1	300,	0	1.7	5.0	14.1	30
	Effective no. of rats...	55	58	56	59	55	57	58	59	57	54
	No. of rats with primary tumours...	38	50	49	52	44	54	56	55	57	47
Large											
Angiosarcoma		0	0	4*	19 ^{ab}	19	0	0	1	5*	23
Adenoma		0	0	0	0	1	0	0	0	0	0
Salivary glands											
Squamous-cell carcinoma		0	0	2	0	1	0	0	0	0	1
Adenoma		0	0	0	0	0	0	0	0	0	1
Abdomen											
Mesothelioma		3	1	7	8	1	1	6*	3	3	0
Angiosarcoma		0	0	0	0	1	0	0	0	2	1
Fibrosarcoma		0	0	0	3	0	1	2	0	0	0
Osteosarcoma		0	0	0	0	1	0	0	0	1	0
Sarcoma		0	0	3	1	0	0	0	1	0	0
Reticulum-cell sarcoma		0	1	0	0	1	0	0	0	0	0
'Schwann-cell tumour'		0	0	0	0	0	0	1	0	0	0
Unclassified		0	0	0	0	0	2	0	0	0	0
Spleen											
Haemangioendotheliosarcoma		0	1	0	0	0	0	0	0	0	0
Lymphosarcoma		0	0	0	0	0	0	0	0	0	1
Nose											
Squamous-cell carcinoma		0	0	0	1	0	0	0	0	0	0
Brain											
Granular-cell myoblastoma		1	1	0	0	0	0	1	0	0	0
Oligodendroglioma		1	0	0	0	0	0	0	0	0	0
Plexus papilloma		0	0	0	0	0	0	1	0	0	0
Glial-cell tumour		0	0	2	0	1	0	0	0	0	0
Ependymoma		0	0	0	1	0	0	0	0	0	0
Mesodermal tumour		0	0	1	0	0	0	0	0	0	0
Pancreas											
Adenocarcinoma		0	0	0	0	0	1	2	1	0	0
Thorax											
Mesothelioma		1	0	0	0	0	0	0	0	0	0
Thyroid											
Parafollicular-cell adenoma		4	12*	10	3	3	7	10	3	2	0
Parafollicular-cell carcinoma		1	0	1	0	0	0	0	0	0	0
Follicular-cell adenoma		0	0	0	1	0	0	0	0	1	0

TABLE VI-22 (continued)

Site and type of tumour	Treatment group (mg VCH/kg/day)	Incidence of Tumours									
		Males					Females				
		0	1.7	5.0	14.1	300†	0	1.7	5.0	14.1	300†
Adrenals											
Cortical adenoma		18	25	17	10	9	26	30	20	17	14
Phaeochromocytoma		11	21	8	4	6	2	1	2	0	2
Pituitary											
Adenoma		12	25**	6	2**	0	14	16	10	5*	3
Carcinoma		1	0	1	0	0	3	0	2	0	0
Blood											
Leukemia		1	0	1	1	3	1	2	1	0	1
Heart											
Endocardial disease†		2	0	2	2	1	1	0	0	0	1
Haemangioendotheliosarcoma		1	0	0	0	0	0	0	0	0	0
Kidneys											
Nephroblastoma		1	0	0	1	0	0	0	0	0	0
Clear-cell tumour		0	0	0	0	0	0	0	0	1	0
Lipomatous tumour		0	0	1	0	0	0	0	0	0	0
Unclassified epithelial tumour		0	0	0	0	0	1	0	0	0	0
Thyroid											
Fibrosarcoma		0	1	0	0	0	0	0	0	0	0
Reticulum-cell sarcoma		0	0	0	0	1	0	1	1	0	0
Mesenteric lymph nodes											
Reticulum-cell sarcoma		0	0	0	0	1	0	0	0	0	0
Skin											
Squamous-cell carcinoma		2	3	3	1	0	0	0	0	1	0
Subcutis											
Fibroma		2	1	1	1	1	3	3	1	0	0
Fibrosarcoma		0	1	1	0	0	1	1	0	0	0
Mesenchymal tumour		0	0	0	0	0	1	0	0	0	0
Skeletal muscle											
Rhabdomyosarcoma		0	0	0	1	1	1	0	0	0	0

TABLE VI-22 (continued)

Site and type of tumour	Treatment group (mg VCM/kg/day)	Incidence of Tumours									
		Males					Females				
		0	1.7	5.0	14.1	300†	0	1.7	5.0	14.1	300†
Skull											
Osteoma		1	0	0	0	0	0	0	0	0	0
Mesenchymal tumour		0	0	0	0	0	0	1	0	0	0
Ear region											
Adenocarcinoma of unknown origin		0	0	1	0	0	0	0	0	0	0
Urinary bladder											
Unclassified epithelial tumour		0	0	1	0	0	0	0	0	0	0
Preputial glands											
Squamous-cell carcinoma		0	0	1	0	0	0	0	0	0	0
Mammary glands											
Adenoma		0	0	0	0	0	0	0	0	2	0
Fibroadenoma		0	0	0	0	0	21	25	12**	4**	7
Adenocarcinoma		0	1	0	2	0	3	2	4	7	7
Anaplastic carcinoma		0	0	0	0	0	0	0	1	0	0
Testes											
Interstitial-cell tumour		3	0	0	1	1					
Uterus											
Adenocarcinoma							6	3	1	1	0
Malignant fibroadenomatous tumour							0	1	0	0	0
Leiomyoma							0	0	1	0	0
Cervix											
Mesenchymal type of tumour							2	0	1	0	0
Adenocarcinoma							0	1	0	0	0
Ovaries											
Theca-cell tumour							0	0	1	0	0

†A small number of primary liver tumours unrelated to treatment were found in several groups. These tumours were one Kupffer-cell sarcoma, three reticulum-cell sarcomas, two fibrosarcomas, one haemangioendothelioma and one mesenchymal tumour.

‡The figures for this group were not evaluated statistically, because no corresponding control group was included in the study.

§In several cases the neoplastic character of the lesion was doubtful.

Values marked with asterisks differ significantly from those of the controls according to the chi-square test:

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

consecutive hours, and food was withdrawn during the other 20 hours. Oral VCM exposure levels were 0 (control), 0.014, 0.13 and 1.3 mg VCM/kg bw/day. An extra control group of 100 rats/sex was housed in a separate room.

Additional groups of 10 male and 10 female rats, each receiving the same treatment as the main groups, were used for determinations of glutathione levels in the liver after 9 and 18 months. Observations were made of general appearance, mortality, growth, food intake, thrombocyte count, prothrombin time, glutathione levels in the liver, gross pathology and microscopic pathology of the liver and of all grossly visible tumors or presumable tumors in the abdominal cavity, Zymbal gland and the mammary glands.

General health, behavior, body weight and food intake were not adversely affected by the test substance. In the second half of the experimental period, mortality in the extra control group was higher than in all other groups. This was most probably due to a high incidence of chronic respiratory disease in the extra control group. In the final stage of the study, the mortality in the top-dose group was slightly higher than in the lower dose groups and the controls. Thrombocyte count, prothrombin time and liver glutathione levels did not show treatment-related differences among the groups.

A clearly higher incidence of grossly visible, tumorous, liver nodules was found in both males and females of the top-dose group than in any of the

other groups. Moreover, in females of the top-dose group the incidence of hepatic cysts was considerably higher than in controls.

Microscopic examination of the liver revealed increased incidences of liver-cell polymorphism, hepatic cysts, foci of cellular alteration, neoplastic nodules and hepatocellular carcinomas in the top-dose group as compared to the control group. One mid-dose female had a hepatocellular carcinoma and neoplastic nodules were found in one low-dose female and in one mid-dose female. A control female was diagnosed with hepatocellular carcinoma. A hepatic angiosarcoma was found in one male and two females of the top-dose group, but no such tumors were encountered in any of the other groups. The number of animals bearing foci of cellular alteration in the liver was also statistically significantly increased in females of the mid-dose group as compared to controls. In addition, in females but not in males, the incidence of basophilic foci of cellular alteration in the liver was statistically significantly higher in both the low- and the mid-dose group than in the control group. Details of the histopathological examination of liver are in Table VI-23.

There was no evidence of VCM-feeding affecting the incidence of abdominal mesotheliomas or the type and incidence of mammary gland tumors (Table VI-24). No Zymbal gland tumor was found.

Table VI-23 Type and Incidence of Histopathological Changes in the Liver

Type of changes¹⁾

	Incidence of changes									
	Males					Females				
	mg VCH/kg b.w./day					mg VCH/kg b.w./day				
	0	0.017	0.17	1.7	0	0	0.017	0.17	1.7	0
Number of animals examined	99	99	99	49	100	98	100	96	49	99
1. Foci of cellular alteration										
a. clear cell foci										
I one or a few	12	8	8	16**	5	4	5	3	13***	1
II several	0	0	0	3*	0	0	0	0	6***	0
b. basophilic foci										
I one or a few	4	2	3	8*	0	9	20*	26***	19***	4
II several	0	0	0	0	0	0	1	0	2	0
c. mixed cell foci										
I one or a few	0	1	2	2	0	6	6	4	8*	0
d. eosinophilic foci										
I one or a few	1	1	2	1	0	0	0	1	10***	2
All type of foci:										
Total	17	12	13	30	5	19	32	34	58	7
Number of foci-bearing animals	16	12	15	23***	5**	19	27	31*	32***	7**

Table VI-23 - continued

Type of changes	Incidence of changes									
	Males					Females				
	mg VCH/kg b.w./day					mg VCH/kg b.w./day				
	0	0.017	0.17	1.7	0 -	0	0.017	0.17	1.7	0
2. Neoplastic nodules										
a. one	0	0	0	1	0	0	1	1	9***	0
b. a few	0	0	0	2	0	0	0	0	1	0
3. Hepatocellular carcinoma	0	0	0	3*	0	1	0	1	3	0
4. Angiosarcoma	0	0	0	1	0	0	0	0	2	0
5. Liver-cell polymorphism										
a. slight	27	23	26	19	12**	46	41	49	23	38
b. moderate	4	4	7	10**	1	14	13	8	15*	13
c. severe	1	1	1	3	0	2	3	4	9***	3
6. Prominent sinusoidal cells										
a. slight	1	1	1	0	1	4	9	3	1	6
b. moderate/marked	1	2	1	1	0	3	8	5	2	0
7. Cysts										
a. none	1	0	1	0	0	3	2	4	1	1
b. a few	4	4	3	4	1	11	11	12	7	2**
c. many	0	0	0	0	0	3	4	9	24***	1
8. Bile duct proliferation										
a. slight	37	26	35	14	21**	27	23	36	18	16*
b. moderate	9	9	8	3	1**	11	6	10	7	3*
c. severe	2	0	0	0	1	1	2	1	1	4

SL 067205

Table VI-23 - continued

Type of changes	Incidence of changes									
	Males					Females				
	mg VCM/kg b.w./day					mg VCM/kg b.w./day				
	0	0.017	0.17	1.7	0	0	0.017	0.17	1.7	0
9. Cholangiofibrosis										
a. slight	33	19*	20*	15	23	16	13	17	15	2***
b. moderate	2	6	8*	2	1	6	2	4	0	0*
10. Periportal fibrosis										
a. slight	1	2	2	3	5	0	0	0	0	1
b. moderate	2	0	0	0	2	0	0	0	0	0
11. Periportal/centrolobular infiltrates of mononuclear cells										
a. slight	14	10	8	3	7	3	2	1	2	4
b. moderate	1	1	1	0	0	0	0	2	0	0
c. severe	0	1	1	0	0	0	0	0	0	0
12. Foci of RES-cells occasionally accompanied by a few necrotic hepatocytes										
a. one or a few	18	15	27	5	12	39	28	42	12*	25*
b. several	5	8	3	0	1	6	3	5	0	3
13. Slight haematopoietic activity	2	0	0	0	1	1	0	3	0	0

Table VI-23 - continued

Type of changes	Incidence of changes									
	Males					Females				
	mg VCH/kg b.w./day					mg VCH/kg b.w./day				
	0	0.017	0.17	1.7	0	0	0.017	0.17	1.7	0
14. Single cell necrosis										
a. slight	2	3	1	0	4	2	3	13**	5*	9*
b. moderate/severe	2	1	0	0	0	1	3	4	2	2
15. Focal necrosis										
a. slight	11	3*	8	2	4*	4	4	6	6	3
b. moderate	1	2	1	0	0	3	1	2	0	3
c. severe	1	0	0	2	0	3	0	1	0	1
16. Centrilobular liver-cell degeneration	1	3	3	1	0	3	2	2	0	0
17. One or a few foci of degenerated hepatocytes	1	0	7*	3	0	1	2	0	0	0
18. Haemorrhagic areas	1	0	1	0	0	0	0	1	0	0
19. Peliosis-like changes	0	1	0	2	0	1	1	0	0	0
20. Small granuloma	0	0	0	0	0	2	0	0	0	0
21. Vacuolization of hepatocytes mainly										
a. periportal										
I slight	3	6	11*	3	4	24	22	29	4*	8***
II moderate	0	2	5*	0	0	3	2	6	0	1
III severe	0	1	1	0	0	0	0	0	0	0

Table VI-23 - continued

Type of changes	Incidence of changes									
	Males					Females				
	mg VCM/kg b.w./day					mg VCM/kg b.w./day				
	0	0.017	0.17	1.7	0	0	0.017	0.17	1.7	0
b. diffuse										
I slight	1	0	1	0	0	0	0	0	0	0
II moderate	1	0	2	0	0	2	0	0	0	0
III severe	0	2	0	0	0	1	0	0	0	2
c. centrilobular										
I slight	0	1	0	0	0	0	0	0	0	0

1) Specific hepatocellular lesions were classified according to Squire and Levitt (1975).

* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$, according to the Fishers' exact test (one tailed).

Adapted from Til et al. 1983.

SL 067208

Table VI-24 Type and Incidence of Tumors of the Mammary Glands and Incidences of Abdominal Mesothelioma

Site and type of tumours	Incidence of tumours									
	Males					Females				
	mg VCM/kg b.w./day					mg VCM/kg b.w./day				
	0	0.017	0.17	1.7	0	0	0.017	0.17	1.7	0
Number of animals examined	99	99	99	49	100	98	100	96	48	98
<u>Mammary glands</u>										
- No of tumour-bearing animals	5	8	3	0	3	41	21***	28*	21	20***
1. Adenoma										
a. single	0	0	0	0	0	6	4	2	5	2
b. two	0	0	0	0	0	1	1	1	0	0
2. Intraductal papilloma	0	0	0	0	0	0	0	0	1	0
3. Fibroadenoma										
a. single	0	1	0	0	0	25	10**	15	14	11**
b. two	0	0	0	0	0	2	3	2	2	4
c. multiple	0	0	0	0	0	3	1	1	1	1

SL 067209

Table VI-24 - continued

Site and type of tumours	Incidence of tumours									
	Males					Females				
	mg VCH/kg b.w./day					mg VCH/kg b.w./day				
	0	0.017	0.17	1.7	0	0	0.017	0.17	1.7	0
4. Fibroma										
a. single	4	6	2	0	3	1	4	1	2	2
b. two	0	0	1	0	0	0	0	0	0	0
c. multiple	0	0	0	0	0	1	0	0	0	0
5. Adenocarcinoma										
a. single	0	1	0	0	0	3	0	5	1	0
b. two	1	0	0	0	0	0	0	2	0	1
Abdomen.										
1. Mesothelioma	0	1	2	0	0	3	2	2	1	1

* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$, according to the Fisher's exact test (one tailed)

Adapted from Till et al. 1983.

SL 067210

VII. HUMAN HEALTH EFFECTS

A. Non-Carcinogenic Effects

Vinyl chloride can produce a number of pathological consequences in humans in addition to its carcinogenic effect. These effects can be caused by acute or chronic exposure to vinyl chloride. Unfortunately, data regarding dose-response relationships in humans are very scarce because few air measurements of vinyl chloride in the work environment of vinyl chloride manufacturing and polymerization plants were made before 1975 (Mancuso 1975). According to OSHA (39 Federal Register 12342, April 5, 1974), several facilities revealed vinyl chloride concentrations for some job classifications as high as 229 ppm. Rowe (1975) commented that before 1960, a few jobs resulted in exposures in the range of 100 to 385 ppm, but these measurements may be high because the method of quantification measured total halogens rather than vinyl chloride alone. Nicholson et al. (1975) reported that vinyl chloride in polymerization reactors may often have exceeded 1,000 ppm and occasionally may have approached 10,000 ppm before OSHA standards were instituted. At these levels, workers experienced dizziness, headaches and/or euphoria during work periods.

Several instances of acute exposure have occurred in vinyl chloride plants. Deaths of two Canadian workers following acute exposures to vinyl chloride gas were reported by Danziger in 1960. At autopsy, there was congestion of the liver, spleen and kidneys. In another study reported by Suciu et al. (1975), exposure of workers to high concentrations of vinyl

chloride produced euphoria, intoxication and narcosis. In this study, the investigators found a dose-response relationship for acute and subacute cases of "occupational disease" from air concentrations ranging from 2,298 mg/m³ (about 900 ppm) to about 100 mg/m³ (about 40 ppm).

In another investigation, Spirtas et al. (1975) conducted a survey of 200 vinyl chloride workers and 89 rubber plant workers (controls) in which information was sought on the frequency of eight symptoms, including dizziness, loss of consciousness, headaches, etc. The vinyl chloride workers were categorized into low and high exposure groups. Because the exposure limits had been markedly decreased a short time before the survey, the high exposure group consisted of workers who were exposed to vinyl chloride concentrations of over 200 ppm before the standard, and 20 to 30 ppm subsequently. The low exposure group consisted of workers who were exposed to 0 to 50 ppm before the standard and 0 to 10 ppm subsequent to it. Examination of the differences among the three groups indicated a statistically significant dose-response relationship for five of the eight symptoms (i.e., frequency of symptoms in the high exposure group > low exposure group > rubber workers), and a similar but non-significant trend in two of the remaining symptom categories. Thus, there appears to be a dose-response relationship between certain acute symptoms (predominantly neurological) and level of vinyl chloride exposure. The data also suggest that vinyl chloride levels below 50 ppm can produce health effects.

The earliest reports of hepatotoxicity in vinyl chloride workers were noted by Tribukh et al. (1949); however, the effects were attributed to

plasticizers added in the manufacturing process. The observed concentrations of vinyl chloride ranged from 1 to 470 ppm. Since that time, impaired liver function has been noted by other investigators (Marsteller et al. 1975, Lillis et al. 1975, Popper and Thomas 1975, Jaeger 1975).

Another effect from chronic vinyl chloride exposure is a condition known as acroosteolysis, which involves bone lesions in the distal phalanges of the hands and feet and scleroderma-like skin lesions. Also associated with this condition are Raynaud's syndrome, pseudoclubbing of fingers, and numerous other symptoms. Many cases of acroosteolysis have been reported and characterized and most involve autoclave workers in vinyl chloride plants (Wilson et al. 1967, Dinman et al 1971, Harris and Adams 1967, Lillis et al. 1975).

Other long-term effects induced by vinyl chloride are disturbances of the central nervous system, pulmonary insufficiency, cardiovascular manifestations and several gastrointestinal symptoms (Miller et al. 1975, Suciu et al. 1975). These and other vinyl chloride-induced health effects are reviewed in the New York Academy of Sciences report "Toxicity of Vinyl Chloride-Polyvinyl Chloride" (Selikoff and Hammond 1975).

Reproductive effects have also been noted. According to a study by Infante (Infante 1976, Infante et al. 1976a), the incidence of birth defects for three small communities in Ohio in which vinyl chloride polymerization plants are located was significantly higher ($P < 0.001$) than that in either the counties in which these communities are located or the State of Ohio. Significant excesses were observed for clubfoot and defects of the central

nervous system, upper alimentary tract and genital organs. A follow-up study by Edmonds et al. (1975) identified a moderate increase in central nervous system malformations, but no association could be found with vinyl chloride exposure. In another epidemiologic study by Infante et al. (1976b), there was a significant excess fetal loss ($P < 0.05$) in wives whose husbands were vinyl chloride polymerization workers compared to controls (wives of polyvinyl chloride fabrication and rubber workers). The Infante studies suggest an association between vinyl chloride and birth defects/fetal loss, but they are not yet supported by animal data. Hatch et al. (1981) examined the statistical analysis (chi-square) used in the Infante (1976a, 1976b) work and concluded that the difference in fetal loss reported as statistically significant by Infante (1976a, 1976b) was not statistically significant. Hatch et al. (1981) also concluded that the statistical test used was of rather low power, and that a negative finding with this analysis does not rule out a possible moderate effect.

Cytogenic studies have also been conducted. Picciano et al. (1977) reported no statistically significant differences in chromatid and chromosome aberrations or proportion of abnormal cells, in a group of 209 vinyl chloride exposed workers. These workers were exposed for periods ranging from 1 to 332 months to time-weighted average (TWA) levels of vinyl chloride ranging from 0.13 to 15.2 ppm. Killian et al. (1975) also reported a lack of evidence for excess chromosome breakage in a population of vinyl chloride exposed workers. In contrast, Ducatman et al. (1975) and Purchase et al. (1975) reported an increased incidence of chromosome breakage among vinyl chloride exposed workers.

Hansteen et al. (1978) and Anderson et al. (1981, 1980) showed that chromosome aberration frequencies in cultured lymphocytes from workers exposed to vinyl chloride returned to levels in the normal control range after a period without exposure to vinyl chloride or after a reduction in vinyl chloride exposure levels.

Heath et al. (1977) examined cytogenic effects in three groups of industrial workers: PVC polymerization workers (presumed high exposure), PVC processing workers (presumed low exposure) and rubber and tire manufacture workers (presumed negligible exposure). Actual vinyl chloride levels were not measured. Chromosome breakage in all three groups was significantly greater than in non-industrial controls, and overall breakage levels were similar in all three groups. The authors concluded that other agents in addition to vinyl chloride may cause cytogenic damage in workers employed in the rubber/plastics industry.

B. Carcinogenic Effects

The primary effect associated with vinyl chloride exposure in man is an increased risk of cancer in several organ systems including angiosarcoma of the liver. Human data have been obtained primarily from occupational exposure of workers to vinyl chloride.

Epidemiologic studies of vinyl chloride exposed workers have focused on cases of angiosarcoma of the liver, a type of cancer which occurs infrequently

in the general population, about 25-30 cases per year in the United States (Heath et al. 1975). Because of its rare occurrence, it is possible to infer a causal relationship between exposure to vinyl chloride and the development of this tumor. The epidemiologic evidence linking vinyl chloride to other types of cancers is more tenuous.

The first study associating vinyl chloride exposure in humans with cancer was conducted by Creech and Johnson, 1974. Three cases of angiosarcoma in workers at a polymerization plant in Louisville, Kentucky, were described. The remaining portion of this section describes some of the epidemiologic studies linking vinyl chloride with angiosarcoma and other types of cancer.

Tabershaw and Gaffey (1974) conducted a mortality study of vinyl chloride workers. Mortality calculations included only those workers who could be traced, i.e., 7,128 of 8,384 workers. These individuals were from 33 different facilities and all had been exposed to vinyl chloride for at least one year. The mean employment duration for the group of workers under study was 80 months. Among the workers, there were 854 with exposures of 20 years or longer and 1,640 exposed 15 or more years.

The overall mortality rate among vinyl chloride workers was found to be lower, i.e., 75 percent of the expected rate compared to the general male U.S. population. The favorable overall mortality rate is a phenomenon commonly observed in working populations. Standardized mortality ratios (the ratio of the number of observed deaths in the study population to the number of deaths

expected in a comparable population) for malignant neoplasms increased with increasing exposure level and/or longer duration. In the group identified as the high exposure group, there were increases in liver cancer (primarily angiosarcoma), respiratory system cancer and brain cancer. These differences were not statistically significant.

Dow Chemical Co. (Holder 1974) conducted a mortality study of 594 workers exposed to vinyl chloride in a single plant between 1942 and 1960. Workers were assigned to exposure groups based on the highest level of exposure for at least one month (low group - TWA less than 25 ppm vinyl chloride, intermediate - 25 to 200 ppm TWA, high - 200 to 300 ppm TWA). Also included in the high group were workers normally exposed to 25 to 200 ppm TWA who were also frequently exposed to excursions of 1,000 ppm. Total mortality was 91 percent of expected among the vinyl chloride exposed workers. No deaths due to liver cancer were reported, and only 13 cases of neoplasms were reported as opposed to 15.4 expected. Nine of these malignancies occurred in the high exposure group, as compared to 5.1 expected (the author stated that due to the small number of deaths, this difference was not tested for significance). Eight of these malignancies were in workers with 15 or more years of exposure.

Monson et al. (1975) conducted a proportional mortality study of workers from two vinyl chloride plants who died between 1947 and 1973. Death certificates were obtained for 142 of 161 workers (88%) who died within this time period. Deaths attributable to cancer were 50 percent higher than expected (a statistically significant difference). A 900 percent increase in

cancers of the liver and biliary tract was noted (five angiosarcomas). Excluding angiosarcoma, a 275 percent excess in the number of cancers was observed. Two brain tumors (320 percent excess) and 13 lung cancers (60 percent excess) were observed. In addition, the overall cancer death rate increased during the period.

Nicholson et al. (1975) studied a group of 257 workers (of whom 255 were traced) exposed to vinyl chloride for at least five years subsequent to 1946. Their mortality status was evaluated beginning ten years after the start of employment until 1974. Exposures were estimated to often exceed 10,000 ppm. Among the 24 deaths were three cases of angiosarcoma of the liver. Preliminary findings indicated a 25 percent increase in deaths over the expected number and a 131 percent increase in all cancer deaths, although neither of these increases was statistically significant.

The National Institute for Occupational Safety and Health (NIOSH) conducted a study which involved 1,294 individuals who were exposed to vinyl chloride for at least five years, and for whom at least ten years had elapsed since initial employment. A total of 136 deaths was reported versus 126.3 expected (not a significant difference). A 49 percent increase over the expected number of cancer deaths was noted, a statistically significant factor. A statistically significant excess number of deaths occurred for brain and CNS cancer, respiratory system cancer, and biliary and liver cancer (Waxweiler et al. 1976).

Ott et al. (1975) reexamined much of the mortality data reported by Tabershaw and Gaffey (1974) and included more clearly defined exposure levels

and follow-up of former company employees. The basic findings remain unchanged; no increase over expected in malignant neoplasms was found in the low exposure group (TWA from 10 to 100 ppm) and an increase in deaths due to malignant neoplasms was observed in the high exposure group (TWA of greater than 200 ppm).

Chiazze et al. (1977) reported a cross-sectional mortality study of 4,341 employees from 17 PVC plants who died between 1964 and 1973. The exposed employee population was compared with the entire U.S. population, specific for color and sex and adjusted for age. No angiosarcoma deaths were identified. Total cancer deaths increased in white employees (especially due to cancer of the digestive system). In white women employees, deaths from cancer of the breast and urinary organs were greater than expected. In a follow-up case-control analysis of breast cancer deaths among PVC fabricators, as an extension of their 1977 study, Chiazze et al. (1980) found no statistically significant relative risks for breast cancer in PVC fabricators and concluded that very large increases in risk for breast cancer did not exist among these workers.

In contrast, in a mortality study of 7,000 British workers exposed to vinyl chloride between 1940 and 1974, the investigators found no evidence of increased cancer mortality other than from liver cancer. In this study, four cases of malignant liver tumor were diagnosed, and two of these were confirmed as angiosarcoma. Both cases were in men exposed to high levels of vinyl chloride (Fox and Collier 1977).

In addition, Byren et al. (1976) traced 750 of 771 Swedish vinyl chloride plant workers. A four- to five-fold increase over expected in pancreas and

liver tumors was found, and two cases were diagnosed as angiosarcoma. The numbers of other tumors did not deviate significantly from expected.

Ten cases of hepatic angiosarcoma were found among the relatively small work force employed at a vinyl chloride polymerization plant in Quebec. This is the largest number of cases to be diagnosed in a single plant (Makk et al. 1976). As a result, Delorme and Theriault (1978) retrieved more detailed information on these employees. The authors suggest that the cases of hepatic angiosarcoma appear to be associated with high vinyl chloride exposure levels and overtime work hours. No correlation was found between occurrence of this tumor and alcohol consumption or cigarette smoking.

In workers engaged in the polymerization of vinyl chloride who were studied by Popper and Thomas (1975), the characteristic hepatic fibrosis was present in all cases of angiosarcoma. Although the relation of fibrotic lesions to the development of angiosarcoma requires further study, a transition from the fibrotic stage to angiosarcoma is suggested by the focal proliferation of the sinusoidal lining cells and of the hepatocytes that are seen in the fibrotic stage, but which becomes even more pronounced in the initial stages of angiosarcoma development. These findings suggest that the fibrotic lesions without angiosarcomas, frequently observed in workers to exposed to vinyl chloride (Lilis et al. 1975), might be the prestage of developing neoplastic lesions. The diagnosis of the fibrotic lesions in these workers may imply a longer latency period for tumor initiation based on a lower exposure level. The series of changes observed in the liver appear to

represent a multi-centric development of angiosarcoma and are similar to the changes induced by Thorotrast and inorganic arsenicals (Berk et al. 1976).

In the most recent update of the NIOSH register (Spirtas and Kaminaski 1978) a total of 64 cases of hepatic angiosarcoma have been identified worldwide among vinyl chloride-exposed industrial workers. A listing of all documented cases by country is presented in Table VII-1. The number of cases per year is depicted in Figure VII-1. Of the 64 cases, 23 have been reported in the United States. The authors reported that both the age at diagnosis and the latency period for cancer induction appear to be increasing. They suggested three explanations for these phenomena: (1) early cases may have heavier exposures, (2) the initial cases represented more biologically susceptible individuals and (3) random fluctuation. If the trend of increased age at diagnosis and the longer latent period for hepatic angiosarcoma induction are indeed related to lower levels of occupational exposure, then the latent period for cancer induction as a result of these low levels of exposure may be longer than previously anticipated, i.e., it would be many years before the ultimate outcome of these exposures will be known.

It has been hypothesized that inhalation of low levels of vinyl chloride by the general public in the vicinity of vinyl chloride/PVC manufacturing plants could be responsible for an increased risk of developing angiosarcoma of the liver. Brady et al. (1977) examined annual rates of hepatic angiosarcoma from 1970 through 1975 in residents of the State of New York (excluding New York City). Exposures to arsenic, vinyl chloride or thorium dioxide were suggested as significant factors in the etiology of these tumors.

Table VII-1 Angiosarcoma of the Liver in Vinyl Chloride/PVC Worker

Country	Case No.	Birth Date	1st VC of PVC Exposure	Diagnosis of Angiosarcoma	Age at Diagnosis	Years from 1st Exposure to Diagnosis	Total Years of Exposure	Date of Death
Belgium	01	00-00-00	00-00-00	00-00-00	00	00	00	06-29-76
Canada	01*	12-15-13	00-00-44	00-00-55	41	11	11	09-02-55
Canada	02*	03-06-14	00-00-43	00-00-57	43	14	14	12-21-55
Canada	03*	08-26-19	00-00-41	00-00-62	42	21	20	03-22-62
Canada	04*	04-05-19	00-00-45	00-00-67	48	22	22	01-21-68
Canada	05*	05-07-11	00-00-44	00-00-68	57	24	05	07-05-68
Canada	06*	12-15-19	00-00-47	00-00-71	51	24	23	04-10-71
Canada	07*	11-09-19	00-00-46	00-00-72	53	26	25	12-24-72
Canada	08	05-13-20	00-00-61	00-00-73	53	12	05	06-12-73
Canada	09	07-19-21	00-00-46	00-00-74	53	28	26	09-04-74
Canada	10	05-16-15	00-00-53	00-00-76	61	23	14	04-00-77
Czechoslovakia	01*	00-00-28	00-00-57	00-00-73	46	16	16	00-00-74
Czechoslovakia	02*	00-00-26	00-00-51	00-00-56	40	15	15	00-00-66
Fed Rep Germany	01*	06-04-30	10-01-56	09-19-68	38	12	12	01-25-69
Fed Rep Germany	02*	07-26-31	10-14-57	09-25-70	39	13	12	12-14-71
Fed Rep Germany	04	09-04-30	04-16-57	00-00-74	44	17	17	11-25-74
Fed Rep Germany	05*	01-01-32	12-16-62	00-00-75	43	13	12	01-09-75
Fed Rep Germany	07*	09-29-26	04-15-54	00-00-75	49	21	12	11-13-75
Fed Rep Germany	08*	10-19-17	04-19-54	00-00-75	58	22	21	12-25-75
Fed Rep Germany	09*	12-13-34	12-02-59	06-16-76	42	17	15	Alive
Fed Rep Germany	10*	07-25-29	10-10-55	06-28-77	47	22	22	06-28-77
Fed Rep Germany	11*	12-29-36	01-02-61	00-00-77	41	16	10	03-07-77
France	01*	04-15-24	01-00-46	02-18-67	43	21	19	02-19-67
France	02	06-03-11	07-06-59	01-08-75	63	15	12	01-24-75
France	03*	00-00-19	00-00-46	01-00-75	55	29	29	06-29-75
France	04*	01-27-27	10-19-49	01-04-76	49	26	26	01-04-76
France	05*	01-29-38	00-00-65	04-00-76	38	11	10	05-13-76
France	06*	04-14-34	00-00-58	09-00-76	42	18	17	09-12-76
France	07	00-00-27	07-01-50	07-00-76	49	26	23	07-02-76
France	08*	04-01-34	05-23-57	12-03-76	42	19	19	01-30-77
Great Britain	01*	04-20-01	00-00-44	12-00-72	71	28	22	12-00-72
Great Britain	03	06-02-37	02-00-66	12-00-74	37	09	04	12-24-74

VII-12

SL 067222

Table VI-1 - continued

Country	Case No.	Birth Date	1st VC of PVC Exposure	Diagnosis of Angiosarcoma	Age at Diagnosis	Years from 1st Exposure to Diagnosis	Total Years of Exposure	Date of Death
Italy	02*	11-13-29	00-00-57	12-13-72	43	15	06	12-00-72
Italy	03*	03-14-20	00-00-53	07-10-75	55	22	21	07-10-75
Japan	01	08-01-22	04-00-53	08-21-74	52	22	22	10-24-75
Norway	01*	12-23-15	03-00-50	12-20-71	56	22	21	01-04-72
Sweden	01*	06-23-27	08-14-51	08-00-74	43	19	18	10-20-70
Sweden	03*	06-10-10	05-00-47	03-19-76	65	29	21	03-19-76
Sweden	04*	11-16-14	00-00-46	05-12-77	62	31	31	05-12-77
U.S.A.	01*	10-17-23	12-09-48	03-03-73	49	24	21	03-03-73
U.S.A.	02*	08-19-33	11-15-55	05-00-70	37	14	13	09-28-71
U.S.A.	03*	05-25-15	11-28-45	12-19-73	58	28	28	12-19-73
U.S.A.	04*	01-15-24	07-06-52	08-19-67	43	15	15	01-07-68
U.S.A.	05*	01-25-12	06-19-44	04-09-64	52	20	20	04-09-64
U.S.A.	06*	11-23-28	01-17-62	02-00-74	46	12	12	07-24-75
U.S.A.	07*	05-03-22	08-27-44	00-00-68	45	24	17	03-23-68
U.S.A.	08*	05-06-20	10-07-46	08-00-61	41	15	15	08-29-61
U.S.A.	09*	11-08-31	05-28-45	03-01-74	43	29	24	03-00-75
U.S.A.	10*	08-16-13	06-12-51	05-00-68	55	17	17	05-10-68
U.S.A.	11*	05-27-09	10-14-46	03-00-70	61	23	23	03-16-70
U.S.A.	12*	11-17-18	09-13-49	05-02-69	50	20	19	05-02-69
U.S.A.	13*	12-01-21	12-11-42	05-00-74	52	32	26	07-04-74
U.S.A.	16*	11-04-27	05-08-50	00-00-69	41	19	04	03-27-69
U.S.A.	17*	05-06-31	06-23-55	10-11-74	43	19	19	Alive
U.S.A.	18*	04-22-28	09-15-54	00-00-75	46	21	11	11-02-75
U.S.A.	19*	00-00-15	00-00-43	06-19-75	60	32	22	04-06-76
U.S.A.	20*	08-31-17	00-00-55	01-30-76	58	21	18	01-30-77
U.S.A.	21*	09-02-09	12-00-46	00-00-77	67	30	21	01-02-77
U.S.A.	22*	10-02-23	07-11-47	01-00-76	52	29	28	12-04-76
U.S.A.	23*	00-00-23	09-00-58	04-06-73	50	15	14	04-06-73
U.S.A.	24*	05-07-17	00-00-39	05-27-77	60	38	26	05-27-77
U.S.A.	25*	08-07-10	02-00-47	03-10-77	67	30	20	03-10-77
Yugoslavia	01*	04-05-14	00-00-53	04-08-73	59	20	20	04-08-73
Yugoslavia	02*	11-15-31	00-00-50	07-12-73	42	23	18	07-12-73


Total Reported
Cases

64

SL 067223

Adapted from SORESEN and KAMUNSKI, 1976

Figure VII-1 Number of Cases of Vinyl Chloride/PVC Related Angiosarcomas Reported to NIOSH by Year of Diagnosis (Representing only 63 of the 64 Cases Known to NIOSH Since Information on Diagnosis is Missing for One Case)



Adapted from Spirtas and Kaminski 1978.

Direct exposure to these agents could not be demonstrated in 19 of the 26 study cases. Five of the 19 patients lived closer to vinyl chloride plants than did their matched controls. This may lend some support to the idea that "indirect modes of exposure, not specifically related to occupation, might be important in the etiology of this disorder" (Brady et al. 1971).

Falk et al. (1981) described a nationwide survey of hepatic angiosarcoma in the United States from 1964 to 1974 which identified 168 cases. Of these, 42 cases were associated with known etiologic factors including exposure to vinyl chloride, use of thorotrast, exposure to inorganic arsenic and treatment with androgenic-anabolic steroids. The remaining cases were of uncertain etiology. The authors concluded that hepatic angiosarcoma most often affects males, peaks in the sixth and seventh decades of life, and appears to occur more often in industrial areas of the Northeast and Midwest. These authors also noted a high relative risk for PVC polymerization workers.

IARC (1979) examined the available data on humans and concluded that exposure to vinyl chloride results in an increased carcinogenic risk to humans. The organ systems most likely to be affected were the liver, brain, lung and hemato- and lymphopoietic systems.

VIII. MECHANISMS OF TOXICITY

The mechanisms by which vinyl chloride causes non-carcinogenic injury are unknown. It is theorized that the toxicity of this compound is attributable to its enzymatic oxidation to reactive polar metabolites, possible chloroethylene oxide or chloroacetaldehyde (see Pharmacokinetics section).

Ward et al. (1976) hypothesized that an immunological mechanism is responsible for the non-carcinogenic pathological effects of vinyl chloride exposure. According to this model, a metabolite of vinyl chloride binds to plasma protein, producing an antibody response. The antigen and resulting immunoglobulin interact to produce a soluble complex which causes vascular occlusion, platelet aggregation and other adverse effects which explain the observed symptoms of the disease. An investigation of workers with "vinyl chloride disease" showed the presence of circulating immune complexes in 19 of 28 patients. Abnormalities were also detected in some workers exposed to vinyl chloride who had few or no overt clinical signs.

Over the past several decades, scientists have undertaken considerable research in an effort to establish the mechanism(s) by which chemical substances cause their carcinogenic effects. The somatic cell mutation theory of carcinogenicity suggests that for a carcinogenic response to occur, an irreversible change must occur in the cell which results in proliferation of a neoplasm. This change reflects a mutational event in the DNA of that cell, suggesting that the chemical carcinogen must interact directly with or

otherwise alter the DNA to initiate the change. In recent years, however, some substances have been shown to be carcinogenic, but by mechanisms with no apparent direct interaction with or alteration of the DNA of the cell by the substance. Presumably, these compounds are not capable of initiating the alteration of a normal cell to a neoplastic one, but can facilitate expression of neoplastic response in latent cells. Carcinogens now are often classified into two broad categories: genotoxic and epigenetic or nongenotoxic.

The mechanism by which a compound causes its carcinogenic effect can rarely be determined by the chronic testing of whole animals, as in the NTP bioassay. Thus, a large number of short-term in vitro and in vivo assay systems have been developed for the purpose of elucidating mechanisms. Since most of the in vitro testing systems measure mutational events, and many carcinogens are mutagens, it is suggested that positive results in certain of these test systems indicate genotoxicity. The decision as to whether a substance is genotoxic may be made qualitatively on the basis of several criteria: (1) a reliable, positive demonstration of genotoxicity in appropriate prokaryotic and eukaryotic systems in vitro, (2) studies on binding to DNA and (3) evidence of biochemical or biologic consequences of DNA damage (Weisburger and Williams 1981).

No single test system appears capable of identifying all carcinogens that are genotoxic. Therefore, a number of scientists have proposed testing batteries such that the results from each test within the battery, when evaluated in conjunction with the others, will enable researchers to determine

the mechanism of carcinogenicity of a particular compound. Vinyl chloride has not been systematically studied in any specific battery of tests, but has been evaluated in a number of test systems that have been proposed for inclusion in one or more batteries. Table VIII-1 summarizes some of the mutagenicity studies on vinyl chloride which have demonstrated the chemical's genotoxic potential. The studies have been divided according to the three criteria for genotoxicity (Weisburger and Williams 1981) outlined above.

Considering all of the data on vinyl chloride, it is probable that vinyl chloride exerts its carcinogenicity through genotoxic mechanisms.

Table VIII-1 Results of Vinyl Chloride Mutagenicity Studies

<u>A. Assay system</u>	<u>Results</u>	<u>References</u>
<u>In Vitro prokaryotic and eukaryotic systems</u>		
Metabolically activated <u><i>Salmonella typhimurium</i></u> system (Ames)	+	Bartsch <u>et al.</u> , 1975 McCann <u>et al.</u> , 1975 Elmore <u>et al.</u> , 1976 Ramsay <u>et al.</u> , 1974 Garro <u>et al.</u> , 1976
<u><i>Escherichia coli</i> K12</u> bioautotrophic strain	+	Grain <u>et al.</u> , 1975
Yeast		Loprieno <u>et al.</u> , 1976, 1977
Germ cells of <u><i>Drosophila</i></u>	+	Verburgt and Vogel, 1977
Chinese hamster V79 cells	+	Huberman <u>et al.</u> , 1975
<u>B. DNA Binding Studies</u>		
Mouse tissues (brain, lung, liver, kidney, spleen, pancreas and testes) <u>in</u> <u>vitro</u>	+(Irreversible binding to RNA and DNA)	Bergman, 1982
Rat liver microsomes, reconstituted cytochrome P-450 systems and isolated hepatocytes	+(Irreversible binding to protein and DNA)	Guengerich <u>et al.</u> , 1981
Rat liver microsomes with NADPH	+(Alkylation of RNA)	Laib and Holt, 1977
<u>C. Biochemical or biologic consequences of DNA damage</u>		
Bone marrow cells of rats (<u>in vivo</u>)	+(Chromosome damage)	Anderson and Richardson, 1976
Bone marrow cells of Chinese hamsters (<u>in vivo</u>)	+(Chromosome aberrations and sister-chromatid- exchanges)	Basler and Rohrborn, 1980
Cultured peripheral lympho- cytes in humans (vinyl chloride exposed workers)	+(Chromosomal abnormalities)	Purchase <u>et al.</u> , 1978 Purchase <u>et al.</u> , 1975 Ducatman <u>et al.</u> , 1975

APPENDIX 1

UNIT RISK ASSESSMENT FOR VINYL CHLORIDE

The data used to estimate a unit risk for oral exposure to vinyl chloride are based on the Feron et al. (1981) study. The statistically significant increases reported for liver and lung tumors were considered biologically significant. For the liver tumors, neoplastic nodules were considered a progression toward hepatocellular carcinomas, and these are included in the analysis in Tables 1 and 2. Extrapolations using the linearized multistage model show values of q_1^* for the individual tumors ranging from 8.8×10^{-2} to 1.3×10^{-1} for the males and from 5.8×10^{-2} to 1.3 for the females. The value of q_1^* based on males was 3.0×10^{-1} for liver tumors and 2.9×10^{-1} based on all tumors combined. For the females the value of q_1^* based on liver tumors was 1.9 and for all tumors combined was 2.3. All units of q_1^* are per mg/kg/day.

Before proceeding with the unit risk estimates an explanation of the total tumor counts in Tables 1 and 2 is necessary. For the liver all animals with hepatocellular carcinomas were assumed to also have the neoplastic nodules. Thus, only the neoplastic nodules and liver angiosarcomas were added to derive the total liver tumors. Otherwise, the totals would have exceeded the number of animals examined. Also, in adding the lung and liver tumors, the totals were not allowed to exceed one less than the number examined. The

Table 1 Type and Incidence of Statistically Significant Treatment-Related Changes in the Liver and Lung of Male Wistar Rats Exposed to VCM in the Diet. Values of q_1^* and Concentration from Multistage Extrapolation Model Included

	Treatment group (mg/kg/day)				q_1^* ^a (mg/kg/day) ⁻¹	95% lower-limit concentration associated with risk (ug/L) ^b		
	0	1.7	5.0	14.1		10 ⁻⁴	10 ⁻⁵	10 ⁻⁶
Number of rats examined ^c	55	58	56	59				
<u>Liver</u>								
Neoplastic nodules	0	1	7	23	2.1×10^{-1}	16.7	1.7	0.2
Hepatocellular carcinomas	0	1	2	8	8.8×10^{-2}	39.8	4.0	0.4
Angiosarcomas	0	0	6	27	1.3×10^{-1}	27.0	2.7	0.3
Total liver tumors ^d	0	2	13	50	3.0×10^{-1}	11.7	1.2	0.1
<u>Lung</u>								
Angiosarcomas	0	0	4	19	1.1×10^{-1}	31.8	3.2	0.3
Total animal with tumors ^e	0	2	17	58	2.9×10^{-1}	12.1	1.2	0.1

^a Human equivalent $q_1^* = q_1^* (a) (W_1/W_2)^{1/3}$ in (mg/kg/day)⁻¹.

^b Concentration in ug/L = $(-35,000/q_1^*) \ln(1-R)$.

^c Found dead or killed in extremis or terminally.

^d Sum of neoplastic nodules and liver angiosarcomas.

^e Total must be at least less than total examined.

SL 067231

Table 2 Type and Incidence of Statistically Significant Treatment-Related Changes in the Liver and Lung of Female Wistar Rats Exposed to VCM in the Diet. Values of q_1^* and Concentration from Multistage Extrapolation Model Included

	Treatment group (mg/kg/day)				q_1^* ^a (mg/kg/day) ⁻¹	95% lower-limit concentration associated with risk (ug/L) ^b		
	0	1.7	5.0	14.1		10 ⁻⁴	10 ⁻⁵	10 ⁻⁶
Number of rats examined ^c	57	58	59	57				
Liver								
Neoplastic nodules	2	26	39	44	1.3	2.7	0.3	0.03
Hepatocellular carcinomas	0	4	19	29	5.0 x 10 ⁻¹	70.0	0.7	0.07
Angiosarcomas	0	0	2	9	8.8 x 10 ⁻²	39.8	4.0	0.4
Total liver tumors ^d	2	26	41	53	1.9	1.8	0.2	0.02
Lung								
Angiosarcomas	0	0	1	5	5.8 x 10 ⁻²	60.3	6.0	0.6
Total animal with tumors ^e	2	26	42	56	2.3	1.5	0.2	0.02

^a Human equivalent $q_1^* = q_1^* (a) (W_1/W_2)^{1/3}$ in (mg/kg/day)⁻¹.

^b Concentration in ug/L = $(-35,000/q_1^*) \ln(1-R)$.

^c Found dead or killed in extremis or terminally.

^d Sum of neoplastic nodules and liver angiosarcomas.

^e Total must be at least less than total examined.

SL 067232

result of this latter restriction was to raise the value of q_1^* slightly due to increased variance. In fitting the response data in Tables 1 and 2 with the human equivalent dosages, the human equivalent dosages were derived by dividing the corresponding animal dosages by $(W_h/W_a)^{1/3}$. The human weight (W_h) was assumed to be 70 kg; the male rats were estimated to weigh 350 g and the female rats were estimated to weigh 200 g (Figure 1). Thus, the corresponding human equivalent dosages were 0, 0.29, 0.85, and 2.41 mg/kg/day based on the male rats, and 0, 0.24, 0.71 and 2 mg/kg/day based on the female rats.

When the response and human equivalent dose data were fit to the linearized multistage model, the 95% upper limit on the largest linear term (Table 2) was:


$$q_1^* = 2.3 \text{ (mg/kg/day)}^{-1}$$

To derive an estimate of the 95% lower level of concentration, d , corresponding to a 95% upper level of risk, R , the following equation is used:

$$R = 1 - e^{-q_1^* d}$$

where d is the lower limit on dose in mg/kg/day. To solve for d in ug/L, we use the transformation

$$1 \text{ mg/kg/day} \times (70 \text{ kg/2 L}) \times 1,000 \text{ ug/mg} = 35,000 \text{ ug/L}$$



^a The weight curves of the rats receiving 0, 1.7, 5.0 or 14.1 mg VCM/kg body weight/day from the 10% PVC diets fed for four hours each day all lie within the shaded area.

Adapted from Feron et al. 1981.

Figure 1 Average Body Weights of the Extra Controls Fed the 10%-PVC Diet Ad Libitum (-) and of the Rats Given 300 mg VCM/kg Body Weight in Oil by Gavage^a

If we set $R = 10^{-5}$ then

$$d = (-35,000/q_1^*) \ln (1-10^{-5}) \text{ (ug/L)}.$$

For the highest value of $q_1^* = 2.3 \text{ (mg/kg/day)}^{-1}$ (Table 2), setting $R = 10^{-5}$ yields a value of $d = 0.15 \text{ ug/L}$. Setting $R = 10^{-4}$ or 10^{-6} yields values of $d = 1.5 \text{ ug/L}$ and $d = 0.015 \text{ ug/L}$, respectively.

For comparison purposes only we compare the potency of vinyl chloride by the diet versus the inhalation routes. A previous memo we sent you estimated the 95% upper limit of potency for VCM as $q_1^* = 1.7 \times 10^{-2} \text{ (mg/kg/day)}^{-1}$ based on an inhalation study showing angiosarcomas and other tumors in rats. That potency estimate was derived for water quality criterion purposes. In that document an inhalation to ingestion by gavage relationship of 1 ppm inhaled = 2.28 mg/kg/day ingested was derived for 200 g rats based on VCM uptake study. Without that adjustment for route differences, a direct transformation based on a 70 kg human breathing $20 \text{ m}^3/\text{day}$ would have yielded a 1 ppm inhaled = 0.76 mg/kg/day relationship and a $q_1^* = 5.2 \times 10^{-2} \text{ mg/kg/day}$, still 44 times less than the estimate from the diet study.

In summary, the VCM potency estimates are reported in Table 3.

Table 3 VCM Potency Estimates

Route	Potency	95% lower limit concentration associated with risk (ug/L)		
	$q_1^*(\text{mg/kg/day})^{-1}$	10^{-4}	10^{-5}	10^{-6}
<u>Oral</u>				
Based on diet study	2.3	1.5	0.15	0.015
Based on inhalation study	1.7×10^{-2}	200	20.0	2.0
<u>Inhalation</u>				
Based on inhalation study	5.2×10^{-2}	67.3	6.7	0.7