

Fractions and Multiples

fraction	prefix	symbol
10^{-1}	deci	d
10^{-2}	centi	c
10^{-3}	milli	m
10^{-4}	micro	μ
10^{-5}	nano	n
10^{-10}	pico	p
10^{-16}	femto	f
10^{-18}	atto	a

multiple	prefix	symbol
10^1	deka	da*
10^2	hecto	h*
10^3	kilo	k
10^6	mega	M
10^9	giga	G
10^{12}	tera	T

*To be restricted as much as possible.

Compound prefixes should not be used. Thus 10^{-6} metre is represented by 1 μm , not 1 $\text{m}\mu\text{m}$. The attaching of a prefix to a unit in effect constitutes a new unit, so that

$$1 \text{ km}^3 = 1 (\text{km})^3 = 10^9 \text{ m}^3$$

$$\text{not } 1 \text{ k}(\text{m}^3) = 10^3 \text{ m}^3.$$

Where possible any numerical prefix should appear in the numerator of an expression.

Examples of Units Contrary to SI, with their Equivalents

physical quantity	unit	equivalent
length	angstrom	10^{-10} m
	inch	0.0254 m
	foot	0.3048 m
	yard	0.9144 m
	mile	1.609 34 km
	nautical mile	1.853 18 km
area	square inch	6.45×10^{-4} m ²
	square foot	0.092 903 m ²
	square yard	0.836 127 m ²
	square mile	2.599 00 km ²
volume	cubic inch	$1.638 71 \times 10^{-4}$ m ³
	cubic foot	0.028 316 8 m ³
	U.K. gallon	0.004 546 082 m ³
mass	pound	0.453 592 37 kg
	ounce	14.593 9 g
density	pound/cubic inch	$2.767\ 99 \times 10^4$ kg m ⁻³
	pound/cubic foot	16.0185 kg m ⁻³
force	dyne	10^{-5} N
	poundal	0.138 255 N
	pound-force	4.448 22 N
	kilogramme-force	9.806 65 N
pressure	atmosphere	$1.01\ 325$ N m ⁻²
	torr	133.322 N m ⁻²
	pound (lbf)/sq. in.	6894.76 N m ⁻²

Organochlorine Pesticides in Seals and Porpoises

by

A. V. HOLDEN

K. MARSDEN

Freshwater Fisheries Laboratory,
Pitlochry, Perthshire

Seals and porpoises in Scotland and Canada, far from the sites of application of pesticides, can accumulate high concentrations of residues in their blubber. These chemicals are spreading through the long food chain which ends with seals and porpoises and obviously cannot be confined to their place of discharge.

The presence of organochlorine residues in marine life in many areas of the world, including the Antarctic, has been reported recently¹⁻⁴. In British waters the accumulation of residue in the eggs of sea birds⁵ and the presence of concentration of pesticides through certain food chains in the marine environment⁶ have been studied. Analyses of various species of marine fish and mammals from Scottish waters during the past 5 years^{7,8} have shown that organochlorine residues are present in the tissues of marine mammals and fish. The highest concentrations have been found in organs rich in fat with the highest lipid content. Thus the muscle tissue of salmonidae, the livers of cod and the blubber of seals are all storage sites for these residues.

The aquatic mammals, such as whales, seals and porpoises, have a relatively high proportion of the body weight in the form of subcutaneous fat (blubber). Seals feed chiefly on gadoids and salmonids, and porpoises chiefly on small gadoids and clupeoids. The process of concentration of pesticides through their food chain might be expected to result in high levels of residues in blubber. The results of the analyses carried out in this laboratory confirm this expectation. In recent years, specimens of the grey seal (*Halichoerus grypus*), common seal (*Phoca vitulina*) and porpoise (*Phocoena phocaena*) have been sampled from the coasts of Scotland. Samples of blubber were available from all the specimens examined, and in a few cases other organs were also available. As well as recognized pesticide residues, other electron-capturing substances have been found in blubber, and it is difficult to say whether they are established whether such substances are widely distributed as the pesticides, blubber samples from grey seals and

harp seal (*Phoca groenlandica*) on the Canadian Atlantic coast have also been examined and the results of these analyses are reported here. Examination of samples of seal blubber from other areas is continuing in order to assess the variation between different regions of the marine environment.

All tissue samples from Scottish waters were frozen at -18°C until required for analysis. The Canadian samples were preserved in 10 per cent formalin, which showed no evidence of any influence on the analytical procedure, and in this and other cases involving fish it has not affected the efficiency of recovery of pesticides residues. The samples (g) were ground with a pestle and mortar, with anhydrous sodium sulfate as a drying agent (AR grade) and extracted in a Soxhlet apparatus with redistilled AR grade *n*-hexano (previously tested and found to be free of electron-capturing materials by gas chromatography). The hexano extracts were made up to 100 ml. and 25 ml. aliquots were subjected to clean-up by the hexane-dimethylformamide partition method of de Faubert Maudroid *et al.*, followed by column clean-up using alumina containing 5 per cent water. The loss of pesticide residues from fat extracts so treated is up to 10 per cent, as is customary in pesticide analysis; no correction has been made for such losses in the results presented here.

The cleaned-up hexano extracts were analysed by gas-liquid chromatography on a Varian 'Aerograph 205-2B' instrument employing two glass columns 5 ft. long $\times$ $\frac{1}{8}$ in. outer diameter. One column was packed with 'Chromosorb W A WIDMOS 80-100' mesh coated with 10 per cent

DC-200 silicone coated with fluoro-silicone. Nitrogen gas for dielidrin was no significant. While complete residues in p,p'-DDT interference with p,p'-TD analytical studies were evaporation and the residual alcohol in water-bath with mixing sulphate solution than analytical technique, DDE, and retained, and the original and p,p'-TD in the vials where present affected. been used.

The price examined in much a lower cost. Scottish prices of all the DC-20 by a non-25 per cent Canadian DDT price about 10 30 per cent interference material values in ference, t lyzed. A noticeable combining

Table 1.

Above
 type of
 F. Scroton
 series (?):
 N. and W.
 adult or
 N. and W.
 semi juv.
 Canada,
 juv. ad.
 (1967)
 Canada, C
 juv., in
 series (?)
 Scotland
 adult;
 E. Scot.
 porpo.

to 81,
valent
to m
254 m
716 m
144 m
10 34 km
15 18 km
10 mm²
200 003 m³
20 127 m³
20 90 km³
20 71 x 10⁻³ m³
20 310 8 m³
20 646 003 m³
20 592 37 kg
505 9 kg
0180 kg 10³ kg m⁻³
1 N
20 253 N
10 22 N
10 65 N
20 225 N m⁻³
20 225 N m⁻³
20 70 N m⁻³

on the sites of
concentrations of
feeding through
as and obviously

Canadian Atlantic
results of these
n of samples of
ing in order to
us of the marine

ers were frozen
The Canadian
formalin, which
is the analytical
involving fish
ery of pesticide
are ground to a
n sulphate (AR
aratus with re-
lated and found
by gas chroma-
le up to 100 ml.
o clean-up of
method of de-
column clean-up
or. The loss of
reated is up to
icide analyses,
on in the results
analysed by gas-
graph 205-28'
ft. long x 1 in.
with Chromo-
with 10 per cent

DC-200 silicone oil, and with the same support coated with 5 per cent J-300 + 7-5 per cent QP-1 fluorosilicone. The oven temperature was 200°C, the nitrogen gas flow 50 ml/min and the column resolution for dieldrin equivalent to 1,600 theoretical plates. There was no significant degradation of DDT on the columns. While complete separation of the common pesticide residues is possible on one or other of these columns, interference by polychlorinated biphenyl (PCB) residues^{1,2} with p,p'-TDE and p,p'-DDT peaks necessitated a further analytical stage. Aliquots (5 ml.) of some of the extracts were evaporated to dryness in a stream of cold filtered air, and the residue refluxed with 2 ml. of 2-6 per cent alcoholic potassium hydroxide for 15 min on a warm water-bath. After cooling, 5 ml. of n-hexane was added, with mixing, followed by 40 ml. of 2 per cent sodium sulphate solution. The supernatant hexane layer was then analysed by gas-liquid chromatography. By this technique, quantitative conversion of p,p'-DDT to p,p'-DDE, and of p,p'-TDE to p,p'-MDE (=DDMU), is obtained, and the conversion products confirm the identity of the original residues. The residual peaks in the p,p'-DDT and p,p'-TDE positions, where present, can be subtracted from the values of the pesticides originally determined in these positions. Hydrolysis also destroys α - and γ -BHC, where present, but polychlorinated biphenyls are unaffected. A cyanosilicone (XE-60) column³ has also been used to separate p,p'-TDE and p,p'-TDE from PCB interference, confirming the latter.

The principal pesticide residues found in all samples examined were those of the DDT group, with dieldrin in much smaller quantities. Canadian seals contained lower concentrations of dieldrin than Scottish seals, while Scottish porpoises contained remarkably high concentrations of all pesticide residues (see Table 1). Interference on the DC-200 silicone column in the p,p'-TDE position caused by the non-hydrolysable residue was responsible for about 25 per cent of the total estimated p,p'-TDE in the Canadian and Scottish samples examined. In the p,p'-DDT position, the non-hydrolysable residue constituted about 10 per cent in the Canadian samples and about 30 per cent in the Scottish samples examined for this interference, the total amount of non-hydrolysable material being smaller in the Canadian samples. The values in Table 1 have not been corrected for this interference, for only a proportion of the samples were hydrolysed. Results from the various species of seals were noticeably different in a given area, and have been combined in Table 1.

Table 1. CONCENTRATION RANGES OF ORGANOPHOSPHORUS RESIDUES IN FAT OF SEALS AND PORPOISES (p.p.m.) (Means in parentheses)

Area and type of sample	No. in sample	Dieldrin	p,p'-DDE	p,p'-TDE	p,p'-DDT
E. Scotland, adult seals (1965-66)	18	0.15-2.1 (0.79)	1.0-11.1 (3.5)	0.27-3.0 (1.2)	1.5-23.8 (7.0)
E. and W. Scotland, adult seals (1965-66)	4	0.06-0.16 (0.20)	1.0-4.0 (2.5)	0.12-0.59 (0.41)	1.3-7.7 (3.8)
Y. and W. Scotland, adult seals (1966)	9	0.06-0.44 (0.18)	1.0-4.0 (2.5)	0.12-0.59 (0.41)	1.3-7.7 (3.8)
Canada, Magdalen Is., adult seals (1967)	2	0.02-0.04 (0.03)	0.27-1.3 (0.78)	0.08-0.18 (0.12)	0.08-0.23 (0.09)
Canada, Cabot Straits, harp. seal, and adult seals (1967)	6	0.03-0.16 (0.07)	1.2-17.3 (5.5)	0.35-5.1 (2.1)	2.1-15.4 (6.4)
Orkney, adult porpoise (1967)	1	0.50	1-1	0.50	2.5
E. Scotland, adult porpoises (1965, 1967)	3	4.0-18.0 (9.0)	9.0-15.3 (12.0)	5.2-14.9 (9.0)	13.1-25.7 (21.1)

Contamination by pesticides is greater in the seals on the east coast of Scotland (taken in an area roughly from Aberdeen to the Tay estuary) than in specimens from the west and north coasts, including the Orkney Islands. Seal pups from the Orkney and Harris (west coast) breeding grounds contained only slightly lower residues than adults from the same areas. By comparison, the Canadian samples from the Magdalen Islands (Gulf of St. Lawrence)

contained very low levels of both dieldrin and the DDT group, while those from the Cabot Straits (Gulf of St. Lawrence) contained little dieldrin but higher DDT group residues, similar to those of Scottish seals. Of the few porpoises examined, that from Orkney was the least contaminated, but the three taken on the east coast of Scotland had markedly higher contents of all the residues measured. One specimen contained a total residue concentration (dieldrin and DDT group) of 73.3 p.p.m. Analyses of a few tissues other than the blubber have been made, but concentrations in the blubber have always been found to be the highest (Table 2). Concentrations in other tissues seem likely to be consistent with their lower lipid contents.

Table 2. TOTAL DDT CONCENTRATIONS (p.p.m.) IN VARIOUS TISSUES

Sample	Blubber	Liver	Brain	Kidney	Spleen	Muscle
Grey seal	5.8	0.77	0.26	—	—	—
Grey seal	9.2	0.36	0.16	—	—	—
Grey seal	14.8	0.97	0.24	—	0.57	—
Grey seal	8.4	0.48	0.17	—	0.12	—
Grey seal	2.4	0.11	0.07	—	0.08	—
Porpoise	2.8	0.58	0.02	0.04	0.10	0.56

The proportion of blubber extractable by hexane was determined for some samples. Values ranged from 72 to 86 per cent (mean 80 per cent) for the Canadian seals, and from 88 to 98 per cent (mean 93 per cent) for Scottish seals samples on the breeding grounds off the west coast. (Values for other seal samples were not estimated.) Concentrations of pesticides in terms of extractable fat for these samples would thus be up to 40 per cent greater than the values in Table 1, but the general comparison is not significantly affected. The lower extractable fat values for the Canadian samples may possibly be a result of being preserved in formalin.

Concentrations of pesticide residues in the extractable fat of various organs and tissues of one porpoise show an important relationship (Table 3). While the concentrations of the DDT group in the original tissues vary widely, those expressed in terms of extractable fat are in much closer agreement, with the sole exception of the brain. A similar anomaly for tissues of trout was described by Holden⁴. It seems likely that a large proportion of the lipid fraction of the brain may not contain pesticides, and the brain lipid composition is known to differ considerably from that of depot fat.

Table 3. CONCENTRATIONS OF TOTAL DDT (p.p.m.) IN TISSUES OF A PORPOISE

Tissue or organ	Blubber	Muscle	Liver	Spleen	Kidney	Brain
Concentration in tissue	3.8	0.50	0.26	0.13	0.04	0.08
Percentage extractable fat	87.0	9.1	13.2	6.1	1.4	0.8
Concentration in extractable fat	4.3	5.2	3.4	2.0	0.57	0.27

The unidentified peaks on the chromatograms seem to be similar to those produced by polychlorinated biphenyls, and to enable comparison to be made, the ratios (R_x) of the retention times of all regularly occurring peaks (relative to dieldrin-100) have been calculated for the operating column temperature of 200°C, these values being temperature-dependent. The R_x values of PCB compounds from commercial PCB formulations were also determined for both types of column (Table 4). At least seven PCB-type peaks were found, including those which interfere with p,p'-TDE and p,p'-DDT. Heptachlor epoxide and p,p'-MDE could not be confirmed on both columns and are not therefore recorded.

The most significant aspect of the results is that, although seals and porpoises inhabit an environment far removed from the site of application or discharge of pesticides, they can contain much greater concentrations of dieldrin and DDT group residues than those recorded for most other species, including man. Robinson and Hunter⁵ found a mean concentration of dieldrin of 0.22 p.p.m. (range 0.10-0.73 p.p.m.) and a mean total con-

centration of DDT of 4.0 p.p.m. (range 1.0–11.1 p.p.m.) in samples of human depot fat in England in 1964. No evidence of any significant change was found during 1965–66.

Table 4. RELATIVE RETENTION VALUES (R_z) OF RESIDUES (BUILDERS = 100)

Residue	Sample	DC-200 column	DC-200/0.1 column	Residue	Sample
PCB	68-5	68	70	PCB	68-5
P, DDE	150	100	100	Dieldrin	100
P, DDT	151	125	121	P, DDE	151
P, DDE	152	125	121	P, DDE	152
P, DDE	153	125	121	P, DDE	153
P, DDE	154	125	121	P, DDE	154
P, DDE	155	125	121	P, DDE	155
P, DDE	156	125	121	P, DDE	156
P, DDE	157	125	121	P, DDE	157
P, DDE	158	125	121	P, DDE	158
P, DDE	159	125	121	P, DDE	159
P, DDE	160	125	121	P, DDE	160
P, DDE	161	125	121	P, DDE	161
P, DDE	162	125	121	P, DDE	162
P, DDE	163	125	121	P, DDE	163
P, DDE	164	125	121	P, DDE	164
P, DDE	165	125	121	P, DDE	165
P, DDE	166	125	121	P, DDE	166
P, DDE	167	125	121	P, DDE	167
P, DDE	168	125	121	P, DDE	168
P, DDE	169	125	121	P, DDE	169
P, DDE	170	125	121	P, DDE	170
P, DDE	171	125	121	P, DDE	171
P, DDE	172	125	121	P, DDE	172
P, DDE	173	125	121	P, DDE	173
P, DDE	174	125	121	P, DDE	174
P, DDE	175	125	121	P, DDE	175
P, DDE	176	125	121	P, DDE	176
P, DDE	177	125	121	P, DDE	177
P, DDE	178	125	121	P, DDE	178
P, DDE	179	125	121	P, DDE	179
P, DDE	180	125	121	P, DDE	180
P, DDE	181	125	121	P, DDE	181
P, DDE	182	125	121	P, DDE	182
P, DDE	183	125	121	P, DDE	183
P, DDE	184	125	121	P, DDE	184
P, DDE	185	125	121	P, DDE	185
P, DDE	186	125	121	P, DDE	186
P, DDE	187	125	121	P, DDE	187
P, DDE	188	125	121	P, DDE	188
P, DDE	189	125	121	P, DDE	189
P, DDE	190	125	121	P, DDE	190
P, DDE	191	125	121	P, DDE	191
P, DDE	192	125	121	P, DDE	192
P, DDE	193	125	121	P, DDE	193
P, DDE	194	125	121	P, DDE	194
P, DDE	195	125	121	P, DDE	195
P, DDE	196	125	121	P, DDE	196
P, DDE	197	125	121	P, DDE	197
P, DDE	198	125	121	P, DDE	198
P, DDE	199	125	121	P, DDE	199
P, DDE	200	125	121	P, DDE	200

The seals and porpoises from the east coast of Scotland were usually more seriously contaminated than those from the north and west coasts. This, at least in respect of seals, may be because, as is believed, the populations in the two areas come from different breeding sites. The seals south of Aberdeen breed principally on the Farnes Islands, off the north-east coast of England, while the seals of north and west Scotland are primarily from the breeding grounds of the Orkney Islands in the north, and other islands off the west coast of Scotland. The sea off eastern Scotland is also likely to receive more contamination from estuarine discharges than that off north and west Scotland. Similarly, the two separate seal populations sampled off the east coast of Canada may inhabit areas with different levels of contamination. Pups contain pesticide residues in concentrations only slightly lower than in the parent population. Common seals in Holland have been found with concentrations of residue similar to those of seals on the Scottish east coast¹¹.

While the residue concentrations in subcutaneous fat are high, those of other tissues and organs are found to be much lower. This suggests that there is no risk of physiological effects, unless the metabolism of much of the fat in times of stress leads to a considerable increase in the concentration of the residue in circulating lipids, and thus in residue concentrations in other organs or tissues.

Gray seals on the coast of the Scottish mainland feed primarily on salmon (*Salmo salar*) and cod (*Gadus morhua*)¹². The highest concentrations of pesticide in salmonids in the marine environment have so far been found in sea trout (*Salmo trutta*) on the east coast with up to 0.2 p.p.m. (dieldrin + total DDT) in muscle tissue¹³, and salmon from the same area would probably have similar contents. Cod examined from the east coast have contained up to 0.35 p.p.m. (dieldrin + total DDT) in muscle tissue (mean 0.12 p.p.m.) and from the west coast up to 0.56 p.p.m. (mean 0.22 p.p.m.), while cod livers contained up to 4.0 p.p.m. and 12.0 p.p.m., respectively. Gray seals are estimated to eat about 15 lb. of food daily, and thus an adult seal could consume its own weight in food in 20–30 days. Assuming a high proportion of the pesticide intake to be stored in body fat rather than to be excreted or metabolized, an increase of two orders of magnitude in pesticide concentration from food to body fat within a few years is feasible. This degree of accumulation of residue in an environment not deliberately contaminated, and in species ecologically far distant from the target organisms of persistent pesticides, underlines the impossibility of confining such chemicals to the areas of application.

We thank Dr B. B. Rao for obtaining the Scottish samples, and Dr C. J. Kerwill for the Canadian samples.

Received December 18, 1967.

¹ Rep. President's Sci. Adv. Comm. (US Govt. Printing Office, Washington, 1963).

² Taiton, J. O'G., and Eusloka, J. R. A., *Nature*, **216**, 546 (1967).

³ Moore, N. W., and Taiton, J. O'G., *Nature*, **207**, 42 (1965).

⁴ Robinson, J., Richardson, A., Crabtree, A. N., Coulson, J. C., and Potts, G. R., *Nature*, **214**, 1307 (1967).

⁵ Department of Agriculture and Fisheries for Scotland, Fisheries of Scotland. Report for 1966.

⁶ Department of Agriculture and Fisheries for Scotland, Fisheries of Scotland. Report for 1965.

⁷ De Fabert, M., M., T. Egan, H., Gault, E. W., Hammond, E. W., Roberts, J., and Thomson, J., *Nature*, **207**, 186 (1965).

⁸ Jensen, S., *New Scientist*, **32**, 412 (1966).

⁹ Holmes, D. C., Simmons, J. H., and Taiton, J. O'G., *Nature*, **216**, 227 (1967).

¹⁰ Simmons, J. H., and Taiton, J. O'G., *Chemosphere*, **27**, 253 (1967).

¹¹ Holden, A. V., *Ann. Appl. Biol.*, **60**, 467 (1966).

¹² Robinson, J., and Hunter, C. U., *Arch. Environ. Health*, **13**, 556 (1966).

¹³ Rao, B. B., *Mar. Res.*, **2**, 50 (1966).

¹⁴ Holden, A. V., *J. Appl. Ecol.*, **3** (Suppl.), 45 (1966).

The North Pacific: an Example of Tectonics on a Sphere

by

D. P. MCKENZIE
R. L. PARKER

Institute of Geophysics and Planetary Physics,
University of California at San Diego

Individual seismic areas move as rigid plates on the surface of a sphere. Application of the Mercator projection to slip vectors shows that the paving stone theory of world tectonics is correct and applies to about a quarter of the Earth's surface.

The linear magnetic anomalies^{1,2} which parallel all active ridges can only be produced by reversal of the Earth's magnetic field³ if the oceanic crust is formed close to the ridge axis⁴. Molodtsov⁵ have shown that the anomalies cannot be observed in the North Atlantic unless most dyke intrusion, and hence crustal production, occurs within 5 km of the ridge axis. The spreading sea floor⁶ then carries these anomalies for great horizontal distances with little if any deformation. The epicentres of earthquakes also accurately follow the axis and are offset with it by transform faults^{7,8}. The structure of inland areas is less clear, though the narrow band of shallow earthquakes suggests that crust is consumed along a linear feature.

These observations are explained if the sea floor spreads as a rigid plate, and interacts with other plates in seismically active regions which also show recent tectonic activity. For the purposes of this article, ridges and trenches are respectively defined as lines along which crust is produced and destroyed. They need not also be topographic features. Transform faults conserve crust and are lines of pure slip. They are always parallel, therefore, to the relative velocity vector between two plates—a most useful property. We have tested this paving stone theory of world tectonics in the North Pacific, where it works well. Less detailed studies of other regions also support the theory.