

ACUTE TOXICITY TO AQUATIC INVERTEBRATES (E.G., DAPHNIA)

TEST SUBSTANCE

Identity: Perfluorodecanesulfonate, ammonium salt; may also be referred to as PFDS, FC-121-X, or 3M Sample E2112 and E2566-1. (1-Decanesulfonic acid, 1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,10-heneicosafuoro-, ammonium salt, CAS # 67906-42-7)

Remarks: Test sample is a mixture of the test substance in water and 2-propanol, 1-(1,1-dimethylethoxy). This mixture contains approximately 22-26% test substance, 33-37% propanol, 1-(1,1-dimethylethoxy), 0.5-1.5% perfluorononanesulfonate, ammonium salt and 38-42% water). All values reported relate to this mixture. No calculations were made to adjust for the actual concentration of the test substance in the test sample. Test sample was from 3M production lot 1.

METHOD

Method: EPA 660/3-75-009 and APHA (1980) Standard Methods for the Examination of Water and Wastewater

Test type: Static acute

GLP: Yes

Year Completed: 1988

Species: *Daphnia magna*

Analytical monitoring: pH, dissolved oxygen, temperature. Test substance concentration was not monitored.

Statistical methods: EC₅₀ values calculated using Binomial method

Test daphnid source: ABC Laboratories In-house cultures, Columbia, MO

Test daphnid age at study initiation: < 24-hours

Test conditions

Dilution water: Well water

Dilution water chemistry (from a representative sample):

hardness: 246 mg/L

alkalinity 178 mg/L

pH: 8.0

Stock and test solutions preparation: Not given. A cloudy appearance and surface foam present at preparation in the 560 and 1000 mg/L concentrations was not noted at the 24- and 48-hour observation periods.

Exposure vessels: 250 mL glass beakers containing 200 mL of test solution.

Number of replicates: 2
Number of daphnids per replicate: 10
Number of concentrations: five plus a negative control
Water chemistry during the study:
 Dissolved oxygen range (0-48 hours):
 8.7 – 9.2 mg/L (control exposure)
 8.6 – 9.2 mg/L (1000 mg/L exposure)
 pH range (0-48 hours):
 8.2 – 8.2 (control exposure)
 8.2 – 8.4 (1000 mg/L exposure)
 Test temperature range (0 – 48 hours)
 19 – 20 °C
 Element basis: mortality and sublethal effects

RESULTS

Nominal concentrations: Bk control, 100, 180, 320, 560, 1000 mg/L

Element value: 24-hour LC₅₀ = 200 (100-320) mg/L
48-hour LC₅₀ = 130 (100–180) mg/L
48-hour NOEC (sublethal effects) = < 100 mg/L
Sublethal effects included quiescence and/or daphnids tending to the bottom of test vessels.

Statistical Evaluation: The LC₅₀ value and 95% confidence interval was calculated using the binomial method.

All element values based on nominal concentrations.

Mortality of controls: None

Remarks: Values reported are for the test sample (mixture). No calculations were made to adjust for the concentration of the test substance in the test sample.

Cumulative percent mortality:

Nominal Test Conc., mg/L	24-hours	48-hours
Neg. Control	0	0
100	0	0
180	35	100
320	100	100
560	100	100
1000	100	100

CONCLUSIONS

The test sample 48-hour EC₅₀ for *Daphnia magna* was determined to be 130 mg/L with a 95% confidence interval of 100 – 180 mg/L.

Testing was conducted on the "as-sold" mixture of the subject material in water (~22-26% subject material) The values reported apply to that mixture and not directly to the subject material.

Submitter: 3M Company, Environmental Laboratory, P.O. Box 33331, St. Paul, Minnesota, 55133

DATA QUALITY

Reliability: Klimisch ranking 2. This study, while well conducted, lacks analytical data for: determination of the test substance concentration in the test solutions; and determination of the sample purity. Because a mixture was tested, care must be taken when interpreting the results. The solvent in the mixture could have enhanced or restricted biological uptake and/or toxicity of the fluorochemical component.

REFERENCES

This study was conducted by ABC Laboratories, Columbia, MO on behalf of the 3M Company.

OTHER

Last changed: 5/18/00



ANALYTICAL BIO-CHEMISTRY LABORATORIES, INC.
P.O. Box 1097 • Columbia, MO 65205
Shipping Address: 7200 East ABC Lane, Columbia, MO 65205
(314) 474-6512 • Telex: 921 814 • ABBOT 1244 4414 2244

SPONSOR

3M Company
Attn: Mr. Mohamed Elnabarawy
EE & PC Environmental Labs
Building 21-2W-05, 900 Bush Avenue
St. Paul, Minnesota 55106

STUDY TITLE

Acute Toxicity of E2566-1 to
Daphnia magna

DATA REQUIREMENT

OECD Guideline 202

AUTHOR

Scott Frazier
Biologist II

STUDY COMPLETED ON

January 9, 1988

PERFORMING LABORATORY


Analytical Bio-Chemistry Laboratories, Inc.
Aquatic Toxicology Division
7200 East ABC Lane
P.O. Box 1097
Columbia, Missouri 65205

ABC LABORATORY PROJECT ID

#36614

Submitted By: Analytical Bio-Chemistry Laboratories, Inc.
7200 East ABC Lane
P. O. Box 1097
Columbia, Missouri 65205
(314) 474-8579

Prepared By:



Scott Frazier
Biologist II/Study Director

1/18/88

Date


Approved By:



Alan D. Forbis
Supervisor, Invertebrate/
Bioconcentration Investigations

1/18/88


Date



Kent Ediger
Quality Assurance Officer

1/18/88

Date



Carl M. Thompson
Aquatic Toxicology Manager

1-18-88

Date

:dab28

TABLE OF CONTENTS

	<u>Page No.</u>
LIST OF TABLES.	36614-4
PROJECT PERSONNEL	36614-5
INVERTEBRATE ACUTE TOXICITY COMPENDIUM.	36614-6
SUMMARY	36614-7
INTRODUCTION.	36614-8
METHODS AND MATERIALS	36614-8
I. Test Organisms.	36614-8
II. Test Procedure.	36614-8
III. Test Compound	36614-9
IV. Statistics.	36614-9
RESULTS	36614-9
LITERATURE CITED.	36614-13
QUALITY ASSURANCE STATEMENT	36614-14
STUDY COMPLIANCE STATEMENT.	36614-15
APPENDIX - RAW DATA	36614-16

LIST OF TABLES

<u>Table No.</u>		<u>Page No.</u>
1	Chemical Characteristics of Well Water Used by ABC's Aquatic Toxicology Division.	36614-10
2	Acute Toxicity of E2566-1 to <u>Daphnia magna</u>	36614-11
3	Percent Mortality and Water Quality Measurements During the Acute Toxicity Test of E2566-1 to <u>Daphnia magna</u>	36614-12

PROJECT PERSONNEL

The Study Director of the project for Analytical Bio-Chemistry Laboratories, Inc. was Scott Frazier, Biologist II. Supervising the conduct of the study for 3M Company was Mr. Mohamed Elnabarawy. The following ABC personnel assisted with various phases of the study.

<u>Name</u>	<u>Title</u>	<u>Initials</u>
Jeffrey Latham	Biological Technician I	<u>JL</u>
Michael Ruddy	Biological Technician II	<u>MR</u>
Scott Frazier	Biologist II	<u>SF</u>

INVERTEBRATE ACUTE TOXICITY COMPENDIUM

Subject: "Acute Toxicity of E2566-1 to Daphnia magna,"
ABC Study Number 36614.

Sponsor: 3M Company

Testing Facility: Analytical Bio-Chemistry Laboratories, Inc.
Aquatic Toxicology Division
P.O. Box 1097
Columbia, Missouri 65205 (314-474-8579)

Location of Original Raw Data and Final Report:

3M Company
EE & PC Environmental Labs
Building 21-2W-05, 900 Bush Avenue
St. Paul, Minnesota

Test Material: E2566-1

Test Concentrations: Control, 100, 180, 320, 560 and 1000 mg/l.

Dilution Water: 246 mg/l (hardness), 178 mg/l (alkalinity),
8.0 (pH).

Test Dates: Initiation - January 7, 1988

Termination - January 9, 1988

Length of Study: 48 hours

Results: 24-hour LC_{50} = 200 mg/l (95% C.I. = 100-320 mg/l)

24-hour NOEL = 100 mg/l

48-hour LC_{50} = 130 mg/l (95% C.I. = 100-180 mg/l)

48-hour NOEL = <100 mg/l (estimated)

Test Species: Daphnia magna

Source of Organisms: ABC Laboratories in-house culture.

Age of Organisms at Study Initiation: First-instar (<24 hours old)

SUMMARY

The acute toxicity of E2566-1 to Daphnia magna was assessed using the methods outlined by the Committee on Methods for Toxicity Tests with Aquatic Organisms. Water quality parameters of temperature, dissolved oxygen and pH were measured at the termination of the test and were within acceptable limits.

The results of the 48-hour static Daphnia magna toxicity study are summarized below. All reported values were based upon nominal concentrations.

<u>Compound</u>	<u>24-Hour LC₅₀</u> <u>(95% C.I.)</u>	<u>48-Hour LC₅₀</u> <u>(95% C.I.)</u>
E2566-1	200 mg/l (100-320 mg/l)	130 mg/l (100-180 mg/l)

The no-effect level observed for E2566-1 at 24 hours was 100 mg/l based on the absence of all abnormal effects. The 48-hour no-effect level was estimated to be <100 mg/l since abnormal effects were observed in all treated test concentrations.

INTRODUCTION

This definitive static bioassay was performed by the Aquatic Toxicology Division of Analytical Bio-Chemistry Laboratories, Inc., Columbia, Missouri, for 3M Company, from January 7, 1988 to January 9, 1988. The purpose of this test was to determine the 24- and 48-hour LC_{50} levels for E2566-1 to Daphnia magna. The study was performed using ABC Protocol #7806 as approved by Mr. Mohamed Elnabarawy, 3M Company, on January 7, 1988.

METHODS AND MATERIALS

The procedures for static bioassay, as described in Methods of Acute Toxicity Tests with Fish, Macroinvertebrates and Amphibians (1) and Standard Methods for Examination of Water and Wastewater (2), were used in this experiment.

I. Test Organisms

The Daphnia magna used in the test were cultured at the ABC facilities in well water (Table 1). The adult daphnids were fed algae (Selenastrum capricornutum) at least every three days prior to testing and supplemented with a suspension of TetraMin®, cereal leaves, vitamin solution and yeast. The daphnids were identified to species using the taxonomic key presented by Pennak (3).

II. Test Procedure

The static Daphnia bioassay was conducted in 250-ml glass beakers containing 200 ml of daphnid culture/test water. Test vessels were covered with loose-fitting petri dish covers to minimize evaporation and protect against contamination during the study. These vessels were kept at 20 (± 2.0) °C in a temperature controlled area. The lighting was maintained at 50-70 footcandles on a 16-hour daylight photoperiod with 30-minute simulated dawn and dusk periods.

An initial range-finding experiment was conducted using ten Daphnia each in exposure concentrations of 1.0, 10, and 100 mg/l. From these results, five treatment concentrations in duplicate with ten Daphnia (first instar less than 24 hours old) per beaker were selected for the definitive bioassay. These concentrations were a logarithmic series ranging from 100 to 1000 mg/l and included a control. All concentrations were observed once every 24 hours for mortality and abnormal effects such as surfacing, clumping of the daphnids together and daphnids tending to the bottom of test chambers. Water chemistry parameters of temperature, dissolved oxygen and pH were measured in the control, low, middle and high concentrations at 0 and 48 hours.

III. Test Compound

The E2566-1 sample was received on December 29, 1987 as a light brown liquid and was stored at room temperature in darkness. Sample purity was unspecified. Test concentrations were prepared based on total product. All standard weights and dilutions can be found in the Appendix. Daphnid culture/test water was used in the preparation of all working stock solutions.

IV. Statistics

Statistical analysis of the concentration vs. effect data (generally mortality) was obtained by employing a computerized LC_{50} program developed by Stephan et al. (4). This program calculated the LC_{50} statistic and its 95 percent confidence limits using the binomial, the moving average and the probit tests. However, if no mortality occurred or if a dose response could not be demonstrated over a reasonable range (<37 to >63%) an LC_{50} and/or its 95 percent confidence limits could not be calculated. Three different methods of analyzing the data were used since no one method of analysis is appropriate for all possible sets of data that may be obtained. The method of calculation selected for presentation in this report was that which gave the narrowest confidence limits for the LC_{50} (4, 5) although all three models are valid.

RESULTS

Table 2 presents the calculated LC_{50} values and 95-percent confidence intervals for E2566-1. The 24- and 48-hour LC_{50} values were 200 and 130 mg/l, respectively. All results were based on the nominal concentrations of 100, 180, 320, 560 and 1000 mg/l. The no-effect concentration based on the absence of mortality and abnormal effects was 100 mg/l at 24 hours. At 48 hours the no-effect level was estimated to be <100 mg/l since all treated concentrations exhibited abnormal effects. The abnormal effects observed during the study were mortality, quiescence and/or daphnids tending to the bottom of test vessels. A cloudy appearance and surface foam present at preparation in the 560 and 1000 mg/l concentrations was not noted at the 24- and 48-hour observation periods.

Table 3 presents the percent mortality and water quality parameters measured during the test. The dissolved oxygen concentrations ranged between 8.6 and 9.2 mg/l. These values represented 99 and 103-percent saturation at 20 and 19 °C, respectively (corrected for local altitudinal pressure), and were considered adequate for testing (1). The pH values of the test solutions ranged from 8.2 to 8.4 and were consistent with the control.

The study was conducted following the intent of the Good Laboratory Practice Regulations (6) and the final report was reviewed by Analytical Bio-Chemistry Laboratories' Quality Assurance Unit. All original raw data were provided to 3M Company, with a copy retained at Analytical Bio-Chemistry Laboratories.

TABLE 1
Chemical Characteristics of Well Water Used by
ABC's Aquatic Toxicology Division.

Parameter	Concentration
Temperature	17 - 22°C ^b
Dissolved Oxygen ^a	7.4 - 9.1 ppm ^b
pH	7.6 - 8.4 ^b
Hardness (CaCO ₃)	206 - 275 ppm ^b
Alkalinity (CaCO ₃)	224 - 336 ppm ^b
Conductivity	500 - 650 μ mhos/cm ^b
Total Organic Carbon	1.4 ppm ^c (0.9 - 2.8 ppm) ^d
Suspended Solids	0 - 1.3 ppm ^b
Aluminum	<10 ppb
Arsenic	<1 ppb
Boron	2.25 ppm
Cadmium	<3 ppb
Chromium	<7 ppb
Cobalt	<3 ppb
Copper	<3 ppb
Iron	14 ppb
Lead	<30 ppb
Mercury	0.43 ppb
Nickel	<8 ppb
Silver	<4 ppb
Zinc	2 ppb
Measured organophosphorus pesticides	<0.14 ppb
Measured organochlorine pesticides plus PCB's	<1.0 ppb

^a After aeration

^b Represents seasonal variation, difference from one month to the next, will not exceed 10%

^c Average of all well water values from January-September, 1987

^d Total monthly range of organic carbon

Sample dates: Trace elements - June 3, 1987, organophosphate and organochlorine analyses - May 28, 1987.

TABLE 2
Acute Toxicity of E2566-1
to Daphnia magna^a

<u>Compound</u>	<u>LC₅₀ (mg/l)</u>	
	<u>24 hours</u>	<u>48 hours</u>
E2566-1	200 (1) (100-320) ^b	130 (1) (100-180) ^b

^aBioassay conducted at 20°C (±2.0).

^b95% confidence limits.

The 24-hour no-effect concentration was 100 mg/l, based on the absence of all abnormal effects. The 48-hour no-effect concentration was estimated to be <100 mg/l since abnormal effects were observed in all test concentrations.

All values were rounded to two significant figures following the technique provided in ABC S.O.P. #8.7.

LC₅₀ calculated using:

(1) Binomial Method

TABLE 3

Percent Mortality and Water Quality Measurements During the
Acute Toxicity Test of E2566-1 to Daphnia magna.

Nominal Concentration (mg/l)	Percent Mortality		Water Quality					
			0 hours			48 hours		
	24 hr	48 hr	Temp. ^a °C	D.O. ^b mg/l	pH ^c	Temp. °C	D.O. mg/l	pH
Control	0	0	19	9.2	8.2	20	8.7	8.2
100	0	0	19	9.2	8.2	20	8.7	8.2
180	35	100						
320	100	100	19	9.2	8.2	20	8.8	8.4
560	100	100						
1000	100	100	19	9.2	8.2	20	8.6	8.4

^aTemperature - mercury thermometer.

^bDissolved oxygen concentrations - YSI 54 ARC dissolved oxygen meter.

^cpH - Corning 140 pH/mV meter with Beckman 39831 electrode.

NOTE: Dissolved oxygen saturations at the test temperatures of 19 and 20°C are 8.9 and 8.7 mg/l, respectively (corrected for altitudinal pressure at ABC Labs).

LITERATURE CITED

- (1) Committee on Methods for Toxicity Tests with Aquatic Organisms. Methods of Acute Toxicity Tests with Fish, Macroinvertebrates and Amphibians. 1975. U.S. EPA, Ecol. Res. Ser. 660/3-75-009.
- (2) American Public Health Association. 1980. Standard Methods for the Examination of Water and Wastewater. 15th ed. Washington, DC. 1134 p.
- (3) Pennak, R. W. 1978. Freshwater Invertebrates of the United States. John Wiley & Sons, New York, 2nd ed. 803 p.
- (4) Stephan, C. E., K. A. Busch, R. Smith, J. Burke and R. W. Andrews. 1978. A Computer Program for Calculating an LC_{50} . U. S. Environmental Protection Agency, Duluth, Minnesota, pre-publication manuscript, August, 1978.
- (5) Stephan, C. 1977. Methods for Calculating an LC_{50} , p. 65-84. In F. L. Mayer and J. L. Hamelink (eds.). Aquatic Toxicology and Hazard Evaluation. ASTM Special Technical Publication 634. ASTM. Philadelphia.
- (6) Organization for Economic Cooperation and Development. 1981. OECD Guidelines for Testing of Chemicals, Principles of Good Laboratory Practice Annex 2, C(81) 30(Final):7-28.

QUALITY ASSURANCE STATEMENT

Quality Assurance Statement for report #36614 entitled, "Acute Toxicity of E2566-1 to Daphnia magna," for Mr. Mohamed Elnabarawy, 3M Company, St. Paul, Minnesota.

In accordance with ABC Laboratories' intent that all studies conducted at our facilities are designed and function in conformance with good laboratory practice regulations and the protocols for individual laboratory studies, an inspection of the final report for E2566-1 was conducted and found to be in an acceptable form by a member of our Quality Assurance Unit. A procedure audit was conducted on December 14, 1987. Results were reported to the study director on the same day and to management on December 30, 1987. A final inspection of all data and records on January 13, 1988, indicated that the report submitted to you is an accurate reflection of the study as it was conducted by ABC Laboratories.

Should you have any questions relating to the information provided in this statement or the function of our Quality Assurance Unit, please contact the Quality Assurance Unit at your convenience.

Kent Ediger
Kent Ediger
Quality Assurance Officer


1/15/88
Date

STUDY COMPLIANCE STATEMENT

Study Compliance Statement for ABC report #36614 entitled, "Acute Toxicity of E2566-1 to Daphnia magna," for Mr. Mohamed Elnabarawy, 3M Company, St. Paul, Minnesota.

In accordance with ABC Laboratories' intent that all aquatic toxicity tests conducted by our facility follow good laboratory practices, ABC's study director for the above test herein confirms that the study was conducted in compliance with the OECD Guidelines for Testing of Chemicals, Principles of Good Laboratory Practice Annex 2, C(81) 30(Final):7-28.

All original raw data were provided to 3M Company, and a copy retained at Analytical Bio-Chemistry Laboratories.


Scott Frazier

ABC Study Director

Date 11/18/87

ACTIVATED SLUDGE RESPIRATION INHIBITION TEST

TEST SUBSTANCE

Identity: Perfluorodecanesulfonate, ammonium salt; may also be referred to as PFDS, FC-121-X, or 3M Sample E2112 and E2566-1. (1-Decanesulfonic acid, 1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,10-heneicosafuoro-, ammonium salt, CAS # 67906-42-7)

Remarks: Test sample is a mixture of the test substance in water and 2-propanol, 1-(1,1-dimethylethoxy). This mixture contains approximately 22-26% test substance, 33-37% propanol, 1-(1,1-dimethylethoxy), 0.5-1.5% perfluorononanesulfonate, ammonium salt and 38-42% water). All values reported relate to this mixture. No calculations were made to adjust for the actual concentration of the test substance in the test sample. Test material was from 3M production lot 1.

Note: Only the summary page was available for this study

METHOD

Method: OECD 209

Test type: Static acute

GLP: No

Year Completed: 1987

Analytical monitoring: Dissolved oxygen concentrations.

Exposure Period: 30-minutes and 3-hours

Measurements: Dissolved oxygen depletion rate after 30-minutes and 3-hours of exposure.

RESULTS

Nominal concentrations: Blank control, 100, 180, 320, 560, 1000 mg/L, in duplicate.

Element value: 30-min $EC_{50} > 1000$ mg/L

3-hour $EC_{50} = 780$ (630-1050) mg/L

All element values based on nominal concentrations.

Statistical method: Probit analysis

Control response: The largest percent difference between control uptake rates was 7.5%; thus the two controls were within 15% of each other.

Reference substance results: Not available

Remarks: Testing was conducted on the "as-sold" mixture of the subject material (~22-26% subject substance in water). The value reported applies to that mixture and not the test material. No attempt was made to determine the impact of the presence of the organic solvents or what portion of the toxicity can be contributed to the perfluorodecanesulfonate.

CONCLUSIONS

The 3-hour EC₅₀ for FC-121 was determined to be 780 (630 – 1050) mg/L

Submitter: 3M Company, Environmental Laboratory, P.O. Box 33331, St. Paul, Minnesota, 55133

DATA QUALITY

Reliability: Klimisch ranking 4. Most of the supporting material for this study is missing. Because a mixture was tested, care must be taken when interpreting the results. The solvent in the mixture could have enhanced or restricted biological uptake and/or toxicity of the fluorochemical component. The study lacked analytical confirmation of concentrations and characterization of the sample purity.

REFERENCES

This study was conducted by the 3M Company, Environmental Laboratory, St Paul, MN, Lab Request number E2112, 10/30/87

OTHER

Last changed: 5/18/00

ENVIRONMENTAL LABORATORY
ACTIVATED SLUDGE RESPIRATION INHIBITION TEST #209
CALCULATION OF RESULTS

LR E2112
Date 10-30-87
Analyst SAB

Test Substance FC-121-X (41-2601-0429-8 Lot 1) FC Surfactant

pg 3 of 3

30 - MINUTE EXPOSURE

Test Substance Concentration (mg/L)	Oxygen Uptake Rate R	R Corrected ⁽¹⁾ to 20°C	% Inhibition in Respiration Rate ⁽²⁾
CONTROL I			
100	0.4776	0.4776	—
180	0.4248	0.4248	13
320	0.4297	0.4297	12
560	0.4273	0.4273	13
1000	0.3721	0.3721	24
	0.3642	0.3642	25
CONTROL II	0.5000	0.5000	—

Percent Difference in Controls⁽³⁾ 4.6%

EC₅₀ (95% Confidence Limits) mg/L >1000 mg/L

3 - HOUR EXPOSURE

Test Substance Concentration (mg/L)	Oxygen Uptake Rate R	R Corrected ⁽¹⁾ to 20°C	% Inhibition in Respiration Rate ⁽²⁾
CONTROL I			
100	0.5067	0.5067	—
180	0.5127	0.4897	7.0
320	0.4218	0.4218	20
560	0.4127	0.4127	22
1000	0.3285	0.3285	38
	0.2091	0.2091	60
CONTROL II	0.5461	0.5461	—

Percent Difference in Controls⁽³⁾ 7.5%

EC₅₀ (95% Confidence Limits) mg/L 780 (630-1050) mg/L Probit Analysis

(1) Temp. Correction = $\frac{R_{T,obs}}{\theta^x}$ (mg O₂/L-min.)

(3) % Difference = $\frac{|RC_1 - RC_2|}{\left(\frac{RC_1 + RC_2}{2}\right)} \times 100$

(2) % Inhibition = $\left(\frac{\left(\frac{RC_1 + RC_2}{2}\right) - R_S}{\left(\frac{RC_1 + RC_2}{2}\right)} \right) \times 100$

The two control respiration rates should be within 15% of each other

See pg 1 for test soln prep

where:

R = Oxygen uptake rate (mg O₂/L-min.)

T_{obs} = Sample temp., °C

θ = 1.047, x = T_{obs} - 20°C

C = Control

MICROBICS MICROTOX® TOXICITY TEST

TEST SUBSTANCE

Identity: Perfluorodecanesulfonate, ammonium salt; may also be referred to as PFDS, FC-121-X, or 3M Sample E2112 and E2566-1. (1-Decanesulfonic acid, 1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,10-heneicosafuoro-, ammonium salt, CAS # 67906-42-7)

Remarks: Test sample is a mixture of the test substance in water and 2-propanol, 1-(1,1-dimethylethoxy). This mixture contains approximately 22-26% test substance, 33-37% propanol, 1-(1,1-dimethylethoxy), 0.5-1.5% perfluorononanesulfonate, ammonium salt and 38-42% water). All values reported relate to this mixture. No calculations were made to adjust for the actual concentration of the test substance in the test sample. Test material was from 3M production lot 1.

Note: Only the summary page was available for this study

METHOD

Method: Microbics Microtox® Procedure

Test type: Static acute

GLP: No

Year Completed: 1987

Species: *Photobacterium phosphoreum*

Analytical monitoring: None

Statistical methods: Microbics computer software package

Test organism source: Microbics Microtox® Reagent

Test conditions: Per Microbics method

Replicates: Two

RESULTS

Nominal concentrations: 125, 250, 500, 1000 mg/L

Test Sample Element values: 5-minute EC_{50} = 640 (605-670)) mg/L
15-minute EC_{50} = 290 (260-310) mg/L

All element values based on nominal concentrations.

CONCLUSIONS

FC-121 exhibited a 15-minute EC_{50} for *Photobacterium phosphoreum* of 290 mg/L with a 95% confidence interval of 260- - 310 mg/L.

Testing was conducted on the "as-sold" mixture of the subject material in water (~22 - 26% subject material). The values reported apply to that mixture and not directly to the subject material.

Submitter: 3M Company, Environmental Laboratory, P.O. Box 33331, St. Paul, Minnesota, 55133

DATA QUALITY

Reliability: : Klimisch ranking 4. Most of the supporting material for this study is missing. Because a mixture was tested, care must be taken when interpreting the results. The solvent in the mixture could have enhanced or restricted biological uptake and/or toxicity of the fluorochemical component.

REFERENCES

This study was conducted by the 3M Company, Environmental Laboratory, St Paul, MN, Lab Request number E2112.

OTHER

Last changed: 5/18/00

Toxicity of FC-121X (41-2601-0429-8 Lot 1)
to Microbial Systems

LRE 2112
SAB

Microtox®		
Test Conc. mg/L	Percent Light Loss	
	5-MIN Exp.	15-MIN Exp.
125	24	29
250	34	44
500	45	65
1000	59	83
EC ₂₀	96	92
EC ₅₀ (95% CI)	640 (605-670)	290 (260-310)
EC ₈₀	>1000	890

OECD Activated Sludge Respiration Inhibition Test (#209)		
Test Conc. mg/L	Percent Inhibition	
	30-MIN Exp.	3-HR Exp.
100	13	7.0
180	12	20
320	13	22
560	25	38
1000	25	60
EC ₅₀ (95% CI)	>1000	780 (630-1050)

ACUTE TOXICITY TO FISH

TEST SUBSTANCE

Identity: Perfluorodecanesulfonate, ammonium salt; may also be referred to as PFDS, FC-121-X, or 3M Sample E2112 and E2566-1. (1-Decanesulfonic acid, 1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,10-heneicosafuoro-, ammonium salt, CAS # 67906-42-7)

Remarks: Test sample is a mixture of the test substance in water and 2-propanol, 1-(1,1-dimethylethoxy). This mixture contains approximately 22-26% test substance, 33-37% propanol, 1-(1,1-dimethylethoxy), 0.5-1.5% perfluorononanesulfonate, ammonium salt and 38-42% water). All values reported relate to this mixture. No calculations were made to adjust for the actual concentration of the test substance in the test sample. Test material was from 3M production lot 1.

Note: Only the summary pages of this report were available.

METHOD

Method: OECD 203

Type: Static acute

GLP: Not known

Year completed: 1988

Species: *Pimephales promelas*

Supplier: ABC Laboratories In-house cultures, Columbia, MO

Analytical monitoring: Dissolved oxygen, temperature and pH. Test substance concentration was not measured.

Exposure period: 96-hours

Statistical methods: Binomial method.

Length and weight: Average length = 18 mm.
Average weight = 0.087 g

Test conditions:

Dilution water chemistry:

hardness:	44 mg/L
alkalinity:	30 mg/L
pH:	7.5

Stock and test solution preparation: Details not available. All test vessels containing FC-121 contained a thin layer of foam at 0-

hours. The amount of foam increased with the concentration. No foam or film was present after 24-hours.

Number of fish per concentration: ten

Number of concentrations: five plus a negative control

Water chemistry during the study: within acceptable limits throughout the study.

Temperature during the study: $22 \pm 1^{\circ}\text{C}$

RESULTS

Nominal concentrations: Bk control, 3.2, 5.6, 10, 18, and 32 mg/L

Element value: 24-hour LC_{50} = 26 (18-32) mg/L

48-hour LC_{50} = 24 (18-32) mg/L

96-hour LC_{50} = 13 (10-18) mg/L

96-hour NOEL = 3.2 mg/L

All element values based on nominal concentrations

Statistical Evaluation of Mortality:

Binomial method used to calculate LC_{50} values and corresponding confidence intervals.

CONCLUSIONS

The test sample 96-hour LC_{50} for fathead minnow was determined to be 13 mg/L with a 95% confidence interval of 10 – 18 mg/L.

Testing was conducted on the "as-sold" mixture of the subject material in water (~22-26% subject material. The values reported apply to that mixture and not directly to the subject material.

Submitter: 3M Company, Environmental Laboratory, P.O. Box 33331, St. Paul, Minnesota, 55133

DATA QUALITY


Reliability: Klimisch ranking 4. A portion of the final report is missing. Because a mixture was tested, care must be taken when interpreting the results. The solvent in the mixture could have enhanced or restricted biological uptake and/or toxicity of the fluorochemical component. Characterization of the test substance purity and analytical measurement of tested concentrations was lacking.

REFERENCES

This study was conducted by ABC Laboratories, Columbia, MO on behalf of the 3M Company.

OTHER

Last changed: 5/18/00



ANALYTICAL BIO-CHEMISTRY LABORATORIES, INC.
P.O. Box 1097 • Columbia, MO 65205
Shipping Address: 7200 East ABC Lane, Columbia, MO 65205
(314) 479-6579 • Telex: ELM 514 • Analytical Bio-Chemistry Laboratories, Inc.

SPONSOR

3M Company
Attn: Mr. Mohamed Elnabarawy
EE & PC Environmental Labs
Building 21-2W-05, 900 Bush Avenue
St. Paul, Minnesota 55106

STUDY TITLE

Acute Toxicity of E2566-1
to Fathead Minnow (Pimephales promelas)

DATA REQUIREMENT

OECD Guideline #203

AUTHOR

Jane E. Bowman
Biologist II

STUDY COMPLETED ON

January 15, 1988

PERFORMING LABORATORY

Analytical Bio-Chemistry Laboratories, Inc.
Aquatic Toxicology Division
7200 East ABC Lane
P.O. Box 1097
Columbia, Missouri 65205

ABC LABORATORY PROJECT ID

#36615

Submitted By: Analytical Bio-Chemistry Laboratories, Inc.
7200 East ABC Lane
P. O. Box 1097
Columbia, Missouri 65205
(314) 474-8579

Prepared By:

Jane H. Bowman
Jane H. Bowman
Biologist II/Study Director

1-29-88
Date

Laura Mullen
Laura Mullen
Biologist I

1-26-88
Date

Approved By:

James P. Swigert
James P. Swigert, Ph.D.
Supervisor, Fish Acute/
Chronic Investigations

1-28-88
Date

Philip M. Buckler
Philip M. Buckler
Quality Assurance Supervisor

1/29/88
Date

Carl M. Thompson
Carl M. Thompson
Aquatic Toxicology Manager

1-29-88
Date

:dab31

FISH ACUTE TOXICITY COMPENDIUM

Subject: "Acute Toxicity of E2566-1 to Fathead Minnow (Pimephales promelas)," ABC Study Number 36615.

Sponsor: 3M Company

Testing Facility: Analytical Bio-Chemistry Laboratories, Inc.
Aquatic Toxicology Division
P.O. Box 1097
Columbia, Missouri 65205 (314-474-8579)

Location of Original Raw Data and Final Report:

3M Company
St. Paul, Minnesota 55106

Test Material: E2566-1

Test Concentrations: Control, 3.2, 5.6, 10, 18 and 32 mg/l

Dilution Water: 44 mg/l (hardness), 30 mg/l (alkalinity),
7.5 (pH)

Test Dates: Initiation - January 11, 1988

Termination - January 15, 1988

Length of Study: 96-hours

Results: 96-hour LC_{50} = 13 mg/l (95% C.I. = 10-18 mg/l)

96-hour NOEL = 3.2 mg/l

96-hour Slope of the Dose-Response Line was 32

Test Species: Fathead Minnow (Pimephales promelas)

Source of Organisms: ABC Laboratories in-house culture

Mean Weight: 0.087 g (± 0.027 g)

Mean Length: 18 mm (± 1.7 mm)

SUMMARY

The acute toxicity of E2566-1 to fathead minnow (Pimephales promelas) was assessed using the methods outlined by the Committee on Methods for Toxicity Tests with Aquatic Organisms. Water quality parameters of temperature, dissolved oxygen and pH were measured throughout the test and were within acceptable limits. Culture and acclimation records indicated the fish were in good condition for testing.

The study was conducted at the following nominal concentrations of E2566-1: 3.2, 5.6, 10, 18 and 32 mg/l. A thin layer of foam was present in all chambers at 0-hour. The amount of foam increased with the concentration. No foam or film was present after 24 hours. Ten fish, with a mean weight of 0.087 (± 0.027) g and a mean standard length of 18 (± 1.7) mm, were exposed to each test concentration and control.

The results of the four day static fish toxicity study using E2566-1 are summarized below. The 24-, 48- and 72-hour LC₅₀ values were also determined. The 96-hour slope of the dose response line was 32.

<u>Compound</u>	<u>96-Hour LC₅₀</u> <u>(95% C.I.)</u>
E2566-1	13 mg/l (10-18 mg/l)

Also, the results indicated a 96-hour, no-observed effect concentration could be estimated at 3.2 mg/l, which was based on the lack of mortality and abnormal effects. Abnormal effects of mortality, loss of equilibrium, light discoloration and/or fish on the bottom of test chamber were observed during the 96-hour exposure period in the 5.6, 18 and 32 mg/l test concentrations.

TABLE 2

Acute Toxicity of E2566-1 to Fathead Minnow
(Pimephales promelas)

Compound	LC ₅₀ in milligrams/liter (ppm)		
	24 hours	48 hours	96 hours
E2566-1 ^a	26 ⁽¹⁾ (18-32) ^b	24 ⁽¹⁾ (18-32) ^b	13 ⁽¹⁾ (10-18) ^b

N = 10 fish per concentration.

^aBioassay as conducted at 22°C (±1.0), mean fish weight and length, 0.087 (±0.027) g and 18 (±1.7) mm.

^b95% confidence limits.

The 96-hour no-observed effect concentration could be estimated at 3.2 mg/l, based on the lack of mortality and abnormal effects at the lowest concentration tested.

The slope of the 96-hour dose response curve as calculated by least squares regression analysis was 32.

All values were rounded to two significant figures following ABC S.O.P. #8.7.

LC₅₀ calculated using:

- (1) Binomial Method
- (2) Moving Average Method
- (3) Probit Method