

560PP

AR 226-0991

COVANCE
THE DEVELOPMENT SERVICES COMPANY

ORIGINAL

Sponsor:

3M
St. Paul, Minnesota

FINAL REPORT

Study Title:

13-Week Dietary Toxicity Study with N-Methyl Perfluorooctanesulfonamido
Ethanol (N-MeFOSE, T-6314) in Rats

Author:

Peter J. Thomford, PhD

Study Completion Date:

June 30, 2000

Performing Laboratory:

Covance Laboratories Inc.
3301 Kinsman Boulevard
Madison, Wisconsin 53704-2595

Laboratory Study Identification:

Covance 6329-225

Sponsor Study Identification:

Study No. 3M T-6314.1

Contain NO CBI

C00796

2001 JAN 18 AM 9:22

RECEIVED
OPTIC


ORIGINAL

Covance 6329-225
3M T-6314.1


QUALITY ASSURANCE STATEMENT

This report, with the exception of Appendix 7, has been reviewed by the Quality Assurance Unit of Covance Laboratories Inc., in accordance with the Food and Drug Administration (FDA) Good Laboratory Practice Regulations, 21 CFR 58. The data in Appendix 7 is generated by and is the responsibility of the Sponsor. The following inspections were conducted and findings reported to the study director and study director management.

Inspection Dates		Phase	Date Reported to Study Director and Study Director Management
From	To		
08/31/98	08/31/98	Protocol Review	08/31/98
08/31/98	08/31/98	Material Dispensation	08/31/98
04/12/99	04/29/99	Report Review	04/29/99
04/02/99	04/07/99	Data Review	05/04/99
06/02/99	06/02/99	Protocol Amendment Review	06/02/99
04/19/00	04/20/00	Report Review	04/20/00
04/28/00	04/28/00	Protocol Amendment Review	04/28/00
04/28/00	04/28/00	Protocol Amendment Review	04/28/00



Representative
Quality Assurance Unit



Date

000797

STUDY IDENTIFICATION

13-Week Dietary Toxicity Study with N-Methyl Perfluorooctanesulfonamido Ethanol (N-MeFOSE, T-6314) in Rats

Test Material	N-Methyl Perfluorooctanesulfonamido Ethanol (MeFOSE, T-6314)
Sponsor	3M Toxicology Services Building 220-2E-02, 3M Center St. Paul, Minnesota 55144-1000
Study Monitor	Andrew M. Seacat, PhD 3M Toxicology Services 651.575.3161
Alternate Study Monitor	Marvin T. Case, DVM, PhD 3M Toxicology Services 651.733.5180
Study Location	Covance Laboratories Inc. 3301 Kinsman Boulevard Madison, Wisconsin 53704-2595
Study Director	Peter J. Thomford, PhD Covance Laboratories Inc. P.O. Box 7545 Madison, Wisconsin 53707-7545 608.241.7207
Study Timetable	
Study Initiation Date	August 27, 1998
In-Life Start Date	September 1, 1998
In-Life End Date	December 2, 1998
Study Completion Date	June 30, 2000

000798

KEY PERSONNEL

Study Director	Peter J. Thomford, PhD
Study Toxicologist	Thomas E. Ryan, BS
Study Coordinator	Nii Van Bruce-Konuah
Supervisor, Small Animal Toxicology	Nathan E. Snortum, BA, BT, ALAT
Supervisor, Dose Formulation	Dixie Bushee, BS, LATG
Supervisor, Chemistry/Tox Support	Brian Schoenike, BS
Supervisor, Laboratory Animal Medicine	Donna J. Clemons, DVM Diplomate, ACLAM
Clinical Pathologist	Robert L. Hall, DVM, PhD Diplomate, ACVP (Clinical Pathology)
Supervisor, Clinical Pathology	Ronald Markevitch, BS, MT (ASCP)
Anatomical Pathologist	Sandra R. Eldridge, PhD Pathology Associates International
Anatomical Pathologist	Johnnie J. Eighmy, DVM, MS Diplomate, ACVP Diplomate, ABT
Supervisor, Anatomical Pathology	Deborah Pirkel, ALAT

000799

CONTENTS

	Page
ABSTRACT.....	8
PURPOSE.....	11
REGULATORY COMPLIANCE.....	11
TEST MATERIAL.....	11
Test Material	11
Reserve (Archive) Sample.....	11
Disposition	11
ANIMALS AND HUSBANDRY	12
Test Animal	12
Identification.....	12
Acclimation	12
Housing and Maintenance	12
Justification	13
PROCEDURES.....	13
Group Designations and Dose Levels.....	13
Dose Preparation	14
Retention Samples	15
Method of Administration.....	15
Dose Analyses	15
Clinical Observations	16
Body Weights.....	16
Food Consumption	16
Ophthalmology	17
Clinical Pathology.....	17
Serum Perfluorooctane Sulfonic Acid Level (PFOS) Determination	18
Necropsy	18
Organ Weights.....	19
Cell Proliferation Tissue Collection and Immunohistochemical Evaluation.....	19
Palmitoyl-CoA Oxidase Tissue Collection and Analyses.....	19
Liver PFOS Analyses	19
Tissue Preservation.....	20
Histopathology	20
Statistical Analyses	21
RECORD RETENTION	21

C00800

CONTENTS (Continued)

	Page
RESULTS	22
Dose Analyses	22
Clinical Observations and Survival	23
Ophthalmology	23
Body Weights	23
Food Consumption	23
Test Material Consumption	24
Clinical Pathology	24
Serum Perfluorooctane Sulfonic Acid Level (PFOS) Determination	25
Cell Proliferation Tissue Collection and Immunohistochemical Evaluation	25
Liver PFOS Determination	25
Anatomic Pathology	25
CONCLUSIONS	26
SIGNATURES	27
REFERENCES	28
OPHTHALMOLOGY REPORT	29
PATHOLOGY REPORT	30
COMMENTS ON THE DATA	37
CODES, ABBREVIATIONS, AND UNITS	39
General Codes and Abbreviations	40
Codes for Clinical Pathology	41
Abbreviations and Units for Clinical Hematology	44
Abbreviations and Units for Clinical Chemistry	46
Abbreviations and Units for Clinical Urinalysis	48
Codes for Anatomic Pathology	49
TABLES	
1 Results of Homogeneity Analyses (ppm)	51
2 Results of Stability Analyses (ppm)	52
3 Results of Dose Preparation Analyses (ppm)	55
4 Summary of Clinical Observations	56
5 Summary of Ophthalmic Observations	58
6 Summary of Body Weight Data (g)	59
7 Summary of Body Weight Change Data (g)	61
8 Summary of Food Consumption Data (g)	63

CONTENTS (Continued)

	Page
TABLES	
9 Summary of Test Material Consumption Data (mg/kg/day)	65
10 Summary of Clinical Hematology Data - Week 5	67
11 Summary of Clinical Hematology Data- Week 14	71
12 Summary of Clinical Chemistry Data - Week 5	75
13 Summary of Clinical Chemistry Data - Week 14	81
14 Summary of Clinical Urinalysis Data - Week 14	87
15 Summary of Organ Weight Data - Week 5 Sacrifice	89
16 Summary of Organ Weight Data - Week 14 Sacrifice	91
17 Incidence of Macroscopic Observations	109
18 Incidence of Microscopic Observations	112
19 Incidence of Severity of Selected Microscopic Observations	121
APPENDIX 1	124
Protocol Deviations	125
Protocol	127
Protocol Amendment No. 1	144
Protocol Amendment No. 2	146
Protocol Amendment No. 3	149
APPENDIX 2	151
Individual Animal Fate Data	152
Individual Clinical Observations	157
Individual Ophthalmic Observations	173
APPENDIX 3	189
Individual Body Weight Data (g)	190
APPENDIX 4	201
Individual Food Consumption Data (g)	202
Individual Test Material Consumption Data (mg/kg/day)	213
APPENDIX 5	223
Individual Clinical Hematology Data	224
Individual Clinical Chemistry Data	248
Individual Clinical Urinalysis Data	276
APPENDIX 6	288
Individual Anatomic Pathology Data	289
APPENDIX 7	546
Cell Proliferation Report	547

C90802

ABSTRACT

The purpose of this study was to assess the toxicity of the test material, N-Methyl Perfluorooctanesulfonamido Ethanol (N-MeFOSE, T-6314) when administered in the diet to rats for at least 13 weeks.

Four groups of CrI:CD[®](SD) IGS BR rats were exposed to N-MeFOSE in the diet for at least 13 weeks. Each group contained 20 males and 20 females. The dietary concentrations were 0 (control group), 3, 30, and 100 ppm.

Food was provided *ad libitum*, except when animals were fasted. Water was provided *ad libitum*. The animals were observed twice daily (a.m. and p.m.) for mortality and moribundity. At least once weekly, each animal was removed from its cage and examined for abnormalities and signs of toxicity. Body weight data were collected on the first day of treatment and weekly thereafter. Food consumption data were collected weekly. Ophthalmic examinations were done before initiation of treatment and during Week 14. Blood samples were collected for hematology and clinical chemistry tests during Weeks 5 and 14. Urine samples for urinalysis tests were collected during Week 14. During Weeks 5 and 14, blood was collected from five animals/sex/group for serum perfluorooctane sulfonic acid (PFOS) analyses. At the Week 5 sacrifice, samples of the liver were collected from five animals/sex/group for palmitoyl-CoA oxidase activity and liver PFOS analysis. Animals sacrificed at Week 5 were not necropsied. During Week 14, the remaining animals were anesthetized, weighed, exsanguinated, and necropsied. At necropsy, macroscopic observations were recorded, selected organs were weighed, and selected tissues were collected and preserved. Samples of liver were collected from all surviving animals for palmitoyl-CoA oxidase activity and liver PFOS analysis. Samples of the liver were collected from five animals/sex/group for proliferation cell nuclear antigen (PCNA) evaluation and microscopic examination. Microscopic examinations were done on tissues from each animal in the control and high-dose groups (terminal sacrifice only) and from the low-dose animal that was sacrificed at an unscheduled interval. Microscopic examinations were also done on the liver and stomach from each terminal sacrifice animal in the mid-dose group, and on the liver from each terminal sacrifice animal in the low-dose group.

000803

One female given 3 ppm N-MeFOSE was sacrificed on Day 59 due to severe laceration of the tail. There were no macroscopic or microscopic changes to indicate that its moribund condition was test material-related. All other animals survived to the scheduled sacrifice. No clinical observations were noted that were considered to represent primary effects of the test material. Statistically significantly lower mean body weights were noted beginning at Week 4 and continuing through the end of study for animals given 100 ppm. Statistically significantly lower mean body weights were noted beginning at Week 7 and continuing through the end of study for males given 30 ppm. Body weight changes were statistically significantly lower in a sporadic manner for animals given 30 or 100 ppm. Overall body weight changes (Weeks 1-14) were statistically significantly lower for all groups given N-MeFOSE. Although statistically significant only at Week 13, food consumption was in general lower for males given 100 ppm. For females given 100 ppm, statistically lower food consumption was apparent at Weeks 3, 4, 6, 7, 11, and 12.

Dietary administration of N-MeFOSE was associated with lower hematocrit for males and females fed 100 ppm; higher urea nitrogen for males and females fed 100 ppm; higher albumin for males fed 100 ppm; lower globulin for males fed 30 or 100 ppm; lower cholesterol and triglycerides for males fed 30 or 100 ppm and females fed 100 ppm; higher alanine aminotransferase for males fed 100 ppm; and higher hepatic palmitoyl CoA oxidase for males and females fed 100 ppm. The effects were relatively mild with the exception of those on cholesterol and triglycerides for the high-dose males; those effects were considered to be of moderate severity. All of the test material-related clinical pathology findings were more prominent at Week 14 than at Week 5.

Statistically significant and test material-related changes in terminal body weights and liver weights were seen in males and/or females in the two highest dose groups. Statistically significant decreases were seen in the terminal body weights of males fed 30 ppm and in both males and females fed 100 ppm of N-MeFOSE. Statistically significant increases were seen in the mean liver-to-body weight percentage and mean liver-to-brain weight ratio in males fed 30 ppm, in mean absolute liver weight and mean liver-to-body weight percentage in females fed 30 ppm, and in all liver weight parameters in males and females fed 100 ppm.

000004

Test material-related macroscopic findings were noted at a low incidence in the liver and glandular stomach of males and females in the 100 ppm dose group only. Liver findings included organs that were large and diffusely dark. In the glandular stomach, dark foci or areas were sometimes seen in the gastric mucosa.

Test material-related microscopic findings were seen in the liver of animals fed 30 and 100 ppm of N-MeFOSE and in the glandular stomach of males and females in the 100 ppm group only. No test material-related changes were seen in animals fed 3 ppm of N-MeFOSE. In the liver, one or more test material-related findings affected all animals fed 100 ppm and included slight-to-moderate centrilobular or diffuse hepatocellular hypertrophy, minimal-to-slight coagulative necrosis, minimal hepatocellular pigment, and minimal-to-slight centrilobular to midzonal or diffuse hepatocellular vacuolation. In animals fed 30 ppm, centrilobular hepatocellular hypertrophy was also seen, although the incidence and severity was decreased at this dose. In addition, a low incidence of minimal hepatocellular vacuolation occurred in 30 ppm dose group males, and the liver of a single female fed 30 ppm had minimal coagulative necrosis. A possible test material-related change in the glandular stomach of animals fed 100 ppm was a relatively low incidence of mucosal erosions. The erosions were of minimal to slight severity and involved a very small portion of the mucosal surface in all cases.

Based on anatomic pathology, clinical pathology, and body weight effects at 30 and 100 ppm the no-observable-adverse-effect level was determined to be less than 30 ppm when N-MeFOSE is administered at concentrations of 0, 3, 30 and 100 ppm to CrI:CD[®](SD) IGS BR rats for at least 13 weeks through dietary administration. Based on the body weight effects at 3, 30, and 100 ppm, the no-observable-effect level was determined to be less than 3 ppm when N-MeFOSE is administered at concentrations of 0, 3, 30 and 100 ppm to CrI:CD[®](SD) IGS BR rats for at least 13 weeks through dietary administration. In the absence of correlative changes in anatomic pathology and clinical pathology the body weights effects at 3 ppm were not considered to be biologically adverse.

000805

PURPOSE

The purpose of this study was to assess the toxicity of the test material, N-Methyl Perfluorooctanesulfonamido Ethanol (N-MeFOSE, T-6314) when administered in the diet to rats for at least 13 weeks.

REGULATORY COMPLIANCE

All aspects of this study were in accordance with the Food and Drug Administration Good Laboratory Practice Regulations as set forth in Title 21 of the US Code of Federal Regulations, Part 58, issued December 22, 1978 (effective June 20, 1979), and with any applicable amendments.

TEST MATERIAL

Test Material

The test material, N-Methyl Perfluorooctanesulfonamido Ethanol (MeFOSE, T-6314), Lot No. F-11364, is an off white powder. It was received at Covance on August 3, 1995.

The test material was stored at room temperature.

Information on synthesis methods, stability, purity, composition, or other characteristics that define the test material is on file with the Sponsor.

Reserve (Archive) Sample

A reserve sample (approximately 5 g) of the test material was taken and stored at room temperature. This sample will be transferred to the Sponsor after authorization from the Sponsor.

Disposition

Remaining test material will be returned after authorization from the Sponsor.

C00806

ANIMALS AND HUSBANDRY

Test Animal

Male and female Crl:CD[®](SD) IGS BR rats were obtained from the Portage, Michigan, facility of Charles River Laboratories, Inc., on August 17, 1998. The animals were 39 to 45 days old at initiation of treatment. The males weighed from 184 to 236 g, and the females weighed from 140 to 179 g at initiation of treatment.

Identification

Each animal was assigned a temporary number upon arrival. Before initiation of treatment, a microchip identification device was implanted into each animal. After randomization for placement on test, each animal was assigned a permanent number, and the microchip was coded with that number. All data for an animal are recorded under these numbers.

Acclimation

Ninety-two males and 92 females were received on August 17, 1998, and acclimated in Animal Room 348 for 15 days before initiation of treatment. In general, animals in this shipment appeared healthy. During acclimation, the animals were examined for abnormalities indicative of health problems, an ophthalmic examination was done, and body weights were recorded for all animals at randomization.

Housing and Maintenance

Animal Room 348 was used for this study. Environmental controls for the animal room were set to maintain 18 to 26°C, a relative humidity of 30 to 70%, and a 12-hour light/12-hour dark cycle.

The animals were housed individually (except for the first 7 days of acclimation when animals were group-housed) in stainless steel, screen-bottom cages.

000807

Certified rodent diet #5002 meal, (PMI Nutrition International) was provided (see Protocol Deviations for exceptions) *ad libitum*, except when animals were fasted. The diet is routinely analyzed by the manufacturer for nutritional components and environmental contaminants. The results are on file with Covance-Madison.

Water was provided *ad libitum*. Samples of the water are routinely analyzed for specified microorganisms and environmental contaminants. The results are on file with Covance-Madison.

There were no known contaminants in the diet or water at levels that would have interfered with this study.

Animals not selected for the study (12 males and 12 females) were removed from the study room and used for training procedures.

Justification

Historically rats have been used in safety evaluation studies and are recommended by appropriate regulatory agencies.

PROCEDURES

This study was conducted in accordance with the Protocol dated August 27, 1998, and Protocol Amendment Nos. 1, 2, and 3. The protocol, protocol amendments, and protocol deviations are in Appendix 1.

Group Designations and Dose Levels

The animals were examined by a laboratory animal veterinarian on August 31, 1998, and found to be suitable for study consideration. Selection of animals for the study was based on clinical observations, body weights, ophthalmic examinations, and other data as appropriate. Animals were assigned to treatment groups using a computerized blocking procedure designed to achieve body weight balance with respect to treatment group. At the time of randomization, the weight variation of the animals for each sex used did not

C00308

exceed ± 2 standard deviations of the mean weight. Group mean body weights were analyzed using Levene's test for homogeneity of variance at the 5.0% probability level and found to be homogeneous. Animals were assigned to the study according to the following design:

Group	Number of Animals		Dietary Levels (ppm MeFOSE) ^a
	Male	Female	
1 (Control) ^b	20	20	0
2 (Low)	20	20	3
3 (Mid)	20	20	30
4 (High)	20	20	100

a Dose levels were expressed as ppm of MeFOSE.

b The control animals received the basal diet only.

Dose Preparation

Dietary concentrations were based on the test material as supplied. Before initiation of treatment, diets of 1, 3, 30, 100, and 500 ppm concentrations were mixed. Diets were prepared for stability analyses at the 1-ppm and 500-ppm concentrations and for homogeneity analyses at the 3, 30, and 100 ppm concentrations. Diets were prepared once every 4 weeks for use in the in-life phase.

Preparation of Diet for Group 1. The appropriate amount of diet was weighed into a labeled container and acetone was added at the same concentration as the high-dose concentration. The combination of the diet and acetone was then mixed for 15 minutes.

Preparation of Diets for Groups 2 through 3. Each dose level was prepared independently. A specified amount of diet was weighed into a labeled container and a specified amount of diet for Group 4 was weighed for each diet. A trench was formed in the feed in the mixing bowl and the amount of diet for Group 4 was transferred to the trench. The contents of the mixing bowl were thoroughly mixed for 15 minutes.

Preparation of Diet for Group 4. A specified amount of diet was weighed into a labeled container. The required amount of test material was weighed and transferred into a labeled container. Approximately 5 mL of acetone was added to the container and mixed manually. Increments of approximately 5 mL acetone were added as necessary until the test material had dissolved. To prepare a premix, the diet was transferred into a labeled

000309

Hobart® mixing bowl. The test material and acetone were added to the mixing bowl, overlaid with a portion of diet from the mixing bowl, and thoroughly mixed. A portion of diet from the mixing bowl was transferred to a second mixing bowl, mixed manually to recover residual test material, and returned to the first mixing bowl. The contents of the mixing bowl were thoroughly mixed.

Samples for dose analyses were taken directly from the mixing bowl.

The prepared test diets were stored at room temperature in covered containers until dispensed into feeding jars.

Retention Samples

During the in-life phase, samples (approximately 100 g) were taken from each dose preparation for dose analyses and stored at room temperature. These samples were discarded on May 11, 1999 (see Protocol Deviations for exceptions).

Method of Administration

Dietary admixture was used because the potential route of exposure in humans is oral.

The dose preparations were administered *ad libitum* for at least 13 weeks, unless otherwise specified.

Dose Analyses

Analyses for the concentration of test material in the dose preparations were done by Covance using an analytical method, MP-M324-MA, supplied by the Sponsor and validated by Covance.

Homogeneity was determined for the 3-ppm, 30-ppm, and 100-ppm dose preparations once pretest. One sample (approximately 100 g) each from the top, middle, and bottom of the dose preparations mixed for homogeneity analyses were collected, divided into three subsamples for extraction and analysis, and analyzed for test material content. All samples were stored at room temperature until analyzed within 7 days of mixing.

000010

Four samples (approximately 100 g each) were taken from the 1-ppm dose level concentration of diet preparations mixed pretest. One sample was analyzed on the day of mixing and used as the baseline value. One sample was stored at room temperature for at least 19 days, then analyzed. A third sample was stored at room temperature for at least 32 days, then analyzed. The remaining sample was stored in a freezer set to maintain -10 to -30°C for 8 weeks, then analyzed. In addition, two samples (approximately 100 g each) were taken from the 500-ppm dose preparation mixed pretest. One sample was analyzed on the day of mixing. The second sample was stored at room temperature after at least 32 days, then analyzed. Two additional samples (approximately 100 g each) were taken from the 3-ppm dose preparation mixed for Weeks 9 through 12. One sample was stored at room temperature for at least 19 days, then analyzed. The second sample was stored at room temperature for at least 32 days, then analyzed. The sample collected for dose confirmation analysis was used as the baseline sample.

During the in-life phase, samples (approximately 100 g) from all dose preparations were collected and analyzed for dose confirmation. All samples were stored at room temperature until analyzed.

Clinical Observations

The animals were observed twice daily (a.m. and p.m.) for mortality and moribundity (see Protocol Deviations for exceptions). Signs of poor health or abnormal behavior were recorded as they were observed. At least once weekly, each animal was removed from its cage and examined. Any unusual or abnormal findings were recorded.

Body Weights

Individual body weight data were recorded on the first day of treatment, and weekly thereafter.

Food Consumption

Individual food consumption data were recorded weekly during treatment. Food consumption was not measured during Week 5 for animals scheduled for sacrifice during Week 5.

C00811

Ophthalmology

Ophthalmic examinations were done before initiation of treatment and during Week 14. The pupils were dilated with 0.5% Mydriacyl®, and the eyes were examined by a veterinarian with an indirect ophthalmoscope.

Clinical Pathology

Blood samples were collected from 10 animals/sex/group during Weeks 5 and 14. Urine samples were collected from 10 animals/sex/group during Week 14. Animals were fasted overnight, (see Protocol Deviations for exception) and urine was collected chilled (approximately 16 hours before blood sampling); water was provided *ad libitum*. Blood was collected from a jugular vein. Potassium EDTA was used as the anticoagulant for hematology tests. No anticoagulant was used for the clinical chemistry samples. Animals were bled in random order. The following were evaluated:

Hematology

red blood cell (erythrocyte) count	white blood cell (leukocyte) count
hemoglobin	differential blood cell count
hematocrit	segmented neutrophil count
mean corpuscular volume	lymphocyte count
mean corpuscular hemoglobin	monocyte count
mean corpuscular hemoglobin concentration	eosinophil count
platelet count	basophil count
	blood cell morphology

Reticulocyte count smears were made and held for possible future examination.

Clinical Chemistry

glucose	aspartate aminotransferase
urea nitrogen	alanine aminotransferase
creatinine	gamma glutamyltransferase
total protein	calcium
albumin	inorganic phosphorus
globulin	sodium
total bilirubin	potassium
cholesterol	chloride
triglycerides	

000312

Urinalysis

volume
specific gravity
pH
protein
glucose
ketones

bilirubin
blood
urobilinogen
microscopic examination of sediment
appearance

Serum Perfluorooctane Sulfonic Acid Level (PFOS) Determination

During Weeks 5 and 14, five animals/sex/group were fasted overnight, (see Protocol Deviations for exception) anesthetized with carbon dioxide, and blood (approximately 4 mL) was collected from a jugular vein. All samples were collected without anticoagulant, allowed to clot at room temperature, and centrifuged. Serum was harvested and stored in a freezer set to maintain -60 to -80°C until packed on dry ice and shipped to the Sponsor for analyses. The samples were analyzed for PFOS. Results of analyses will be reported separately by the Sponsor.

Necropsy

A necropsy was done on the Group 2 female that was sacrificed at an unscheduled interval. During Week 5, five animals/sex/group were fasted overnight (see Protocol Deviations for exception), anesthetized with carbon dioxide, bled for serum PFOS samples, weighed, and exsanguinated. These animals were discarded without necropsy. The liver was removed from these animals, weighed, and collected for palmitoyl CoA oxidase and PFOS analysis. During Week 14, the remaining animals were fasted overnight, anesthetized with carbon dioxide, bled for clinical pathology tests (10 animals/sex/group with the exception of Group 2 female) and serum PFOS samples (five animals/sex/group), weighed, exsanguinated, and necropsied. Liver samples were collected for palmitoyl CoA oxidase, PCNA, and PFOS analysis. Animals were necropsied in random order.

The necropsy included a macroscopic examination of the external features of the carcass; all external body orifices; the abdominal, thoracic, and cranial cavities; organs; and tissues.

000813

Organ Weights

After 13 weeks of treatment, the following organs (when present) were weighed; paired organs were weighed together:

adrenal (2)	spleen
brain	testis (2)
kidney (2)	thymus
liver	thyroid (2) with parathyroid
ovary (2)	

Organ-to-body weight percentages and organ-to-brain weight ratios were calculated.

Cell Proliferation Tissue Collection and Immunohistochemical Evaluation

At the terminal sacrifice, representative samples of the left lateral lobe of the liver were collected from five animals/sex/group and preserved in zinc formalin. After fixation, samples were embedded in paraffin and shipped to Pathology Associates International (Frederick, Maryland) for proliferation cell nuclear antigen (PCNA) evaluation. In addition, liver sections were stained with hematoxylin and eosin and examined microscopically. Results of the evaluation are in Appendix 7.

Palmitoyl-CoA Oxidase Tissue Collection and Analyses

At the scheduled sacrifices, the right lateral lobe of the liver was collected from each animal, flash-frozen in liquid nitrogen, and stored in a freezer set to maintain -60 to -80°C until analyzed by Covance for palmitoyl CoA oxidase activity.

Liver PFOS Analyses

At the scheduled sacrifices, a portion of the liver from each animal was flash-frozen in liquid nitrogen and stored in a freezer set to maintain -60 to -80°C until packed on dry ice and shipped to the Sponsor for PFOS and metabolites analyses. Results of these analyses will be reported separately by the Sponsor.

000814

Tissue Preservation

The following tissues (when present) from the animal that was sacrificed at an unscheduled interval, or from each animal sacrificed at the terminal sacrifice were collected and preserved in 10% neutral-buffered formalin, unless otherwise specified:

adrenal (2)	ovary (2)
aorta	pancreas
brain	pituitary
cecum	prostate
cervix	rectum
colon	salivary gland [mandibular (2)]
duodenum	sciatic nerve
epididymis (2)	seminal vesicle (2)
esophagus	skeletal muscle (thigh)
eye (2)	skin
femur with bone marrow (articular surface of the distal end)	spinal cord (cervical, mid-thoracic, and lumbar)
Harderian gland	spleen
heart	sternum with bone marrow
ileum with Peyer's patch (lymphoid aggregate)	stomach
jejunum	testis (2)
kidney (2)	thymus
lesions	thyroid (2) with parathyroid
liver	trachea
lung with mainstem bronchi	urinary bladder
lymph nodes (mesenteric and mandibular)	uterus
mammary glands (females only)	vagina
	Zymbal's gland

Histopathology

Tissues (as appropriate) were embedded in paraffin, sectioned, stained with hematoxylin and eosin, and examined microscopically from each animal in the control and high-dose groups and from the low-dose animal that was sacrificed at an unscheduled interval. Due to microscopic findings noted in the high-dose group, the liver and stomach from animals in the mid-dose group, and the liver from animals in the low-dose group were also embedded in paraffin, sectioned, stained with hematoxylin and eosin, and examined microscopically.

Macroscopic lesions from animals in the low- and mid-dose groups were embedded in paraffin, sectioned, stained with hematoxylin and eosin, and examined microscopically.

Bone marrow smears from the femur of each animal were prepared at the scheduled sacrifice after 13 weeks of treatment, stained with Wright's stain, and retained for possible examination.

Statistical Analyses

Levene's test (Levene, 1960) was done to test for variance homogeneity. In the case of heterogeneity of variance at $p \leq 0.05$, transformations were used to stabilize the variance. Comparison tests took variance heterogeneity into consideration.

One-way analysis of variance [ANOVA (Winer, 1971a)] was used to analyze body weights, body weight changes, food consumption, continuous clinical pathology values; organ weights, organ-to-body weight percentages, and organ-to-brain weight ratios. If the ANOVA was significant, Dunnett's t-test (Dunnett, 1964) was used for control versus treated group comparisons.

Group comparisons (Groups 2 through 4 versus Group 1) were evaluated at the 5.0%, two-tailed probability level. Only data collected on or after the first day of treatment was analyzed statistically.

RECORD RETENTION

All raw data, documentation, records, protocol, and specimens generated as a result of this study will be archived in the storage facilities of Covance-Madison for a period of at least 1 year. At least 1 year after the submission of the final report, the Sponsor will determine the final disposition of the materials. All raw data stored on magnetic media, the protocol and protocol amendments, study correspondence, and an original copy of the final report will be retained by Covance-Madison.

PCNA evaluation data, paraffin blocks, and tissue slides will be retained by Pathology Associates International.

C00816

Liver and serum samples sent to the Sponsor and analysis data generated from these samples will be retained by the Sponsor.

RESULTS

Dose Analyses

Results of the homogeneity, stability, and dose preparation analyses are in Tables 1 through 3.

Mean values of the homogeneity analyses ranged from 74.0-78.0%, 84.0-96.7%, and 77.5-84.2% of the theoretical concentrations for the diets containing 3, 30, and 100 ppm N-MeFOSE, respectively. These results indicate that the mixing procedure produced a homogeneous distribution of the test material in the dose preparations; although variability generally appeared slightly greater at the 3 and 30-ppm dietary concentrations.

Results of stability analyses of samples stored for 19 and 32 days at room temperature and 8 weeks under frozen conditions indicated that the mean concentrations ranged from 54.4-62.5% of the initial concentrations of 1 ppm for diet mixed pretest. Samples stored for 32 days at room temperature indicated that the mean concentration was 98.4% of the initial concentrations of 500 ppm. Results of stability analyses of samples stored for 19 and 32 days at room temperature indicated that the mean concentrations ranged from 93.2-94.0% of the initial concentrations of 1 ppm for diet mixed at Week 4. Samples stored for 19 days at room temperature indicated that the mean concentration was 89.0% of the initial concentrations of 3 ppm.

The mean concentrations of the dose preparation analyses for all levels ranged from 63.7 to 110.0% of the theoretical concentrations. Inherent variability and lack of sufficient sensitivity at low levels in the analytical method for measuring N-MeFOSE resulted in homogeneity, stability and routine analysis data that were in many cases outside of the standard limits of $\pm 15\%$. These results will be reevaluated in conjunction with the analytical results from the blood and liver level determinations when they are provided by the sponsor.

C00817

Clinical Observations and Survival

Clinical observations are summarized in Table 4; individual data are in Appendix 2. Individual animal fate data are also in Appendix 2.

One female given 3 ppm N-MeFOSE was sacrificed on Day 59 due to severe laceration of the tail. There were no macroscopic or microscopic changes to indicate that its moribund condition was test material-related. All other animals survived to the scheduled sacrifice.

No clinical observations were noted that were considered effects of the test material.

Ophthalmology

Ophthalmic observations are summarized in Table 5; individual data are in Appendix 2. The Ophthalmology Report contains a discussion of the data.

Animals selected for the study had no lesions at the prestudy examination. No test material-related ophthalmic observations were noted at the Week 14 examination.

Body Weights

Body weight and body weight change data are summarized in Tables 6 and 7; individual body weight data are in Appendix 3.

Test material-related lower body weights were noted for males given 30 and 100 ppm and for males and females given 100 ppm. Statistically significantly lower mean body weights were noted beginning at Week 4 and continuing through the end of study for males and females given 100 ppm. Statistically significantly lower mean body weights were noted beginning at Week 7 and continuing through the end of study for males given 30 ppm.

Body weight changes were statistically significantly lower in a sporadic manner for males given 30 ppm and for animals of both sexes given 100 ppm. Overall body weight changes (Weeks 1-14) were statistically significantly lower for all groups given N-MeFOSE.

Food Consumption

Food consumption data are summarized in Table 8; individual data are in Appendix 4.

Although statistically significant only at Week 13 for males given 100 ppm, food consumption was lower for males given 30 and 100 ppm when compared to those of controls for most of the study. Females given 100 ppm had statistically significantly lower food consumption than those of controls at Weeks 3, 4, 6, 7, 11, and 12. The overall mean food consumption for females fed the 100-ppm concentration was significantly lower by 17.5% when compared with that of the controls.

Test Material Consumption

Test material consumption data are summarized in Table 9; individual data are in Appendix 4.

Animals were fed diets containing 3, 30, or 100 ppm N-MeFOSE, T-6314. The mean amounts of test material consumed by the animals in these groups were 0.148 to 0.304 mg/kg/day; 1.390 to 3.094 mg/kg/day; and 5.054 to 10.291 mg/kg/day for males, respectively. Values for females were 0.184 to 0.319 mg/kg/day; 2.204 to 3.395; and 6.494 to 10.378, respectively. The overall mean test material consumption for males fed diets containing 3, 30, or 100 ppm N-MeFOSE, T-6314 were 0.203, 2.041, and 6.844 mg/kg/day, respectively. The overall mean test material consumption for females fed diets containing 3, 30, or 100 ppm N-MeFOSE, T-6314 were 0.240, 2.588, and 7.702, respectively.

Clinical Pathology

Hematology, clinical chemistry, and urinalysis data are summarized in Tables 10 through 14; individual data are in Appendix 5. The Pathology Report contains a discussion of the data.

Dietary administration of N-MeFOSE was associated with lower hematocrit for males and females fed 100 ppm; higher urea nitrogen for males and females fed 100 ppm; higher albumin for males fed 100 ppm; lower globulin for males fed 30 or 100 ppm; lower cholesterol and triglycerides for males fed 30 or 100 ppm and females fed 100 ppm; higher alanine aminotransferase for males fed 100 ppm; and higher hepatic palmitoyl CoA oxidase for males and females fed 100 ppm. The effects were relatively mild with the exception of those on cholesterol and triglycerides for the high-dose males; those effects

C00819

were considered to be of moderate severity. All of the test material-related clinical pathology findings were more prominent at Week 14 than at Week 5.

Serum Perfluorooctane Sulfonic Acid Level (PFOS) Determination

Results of serum PFOS level determination will be reported separately by the Sponsor.

Cell Proliferation Tissue Collection and Immunohistochemical Evaluation

Results of cell proliferation and immunohistochemical evaluation provided by Pathology Associates International are in Appendix 7.

Liver PFOS Determination

Results of liver PFOS level determination will be reported separately by the Sponsor.

Anatomic Pathology

Terminal body weights, absolute organ weights, organ-to-body weight percentages, and organ-to-brain weight ratios are summarized in Tables 15 and 16; incidences of macroscopic and microscopic observations are summarized in Tables 17 and 18; the incidence of severity of microscopic observations is summarized in Table 19. Individual data are in Appendix 6. The Pathology Report contains a discussion of the data.

Statistically significant and test material-related changes in terminal body weights and liver weights were seen in males and/or females in the two highest dose groups. Statistically significant decreases were seen in the terminal body weights of males fed 30 ppm and in both males and females fed 100 ppm of N-MeFOSE. Statistically significant increases were seen in the mean liver-to-body weight percentage and mean liver-to-brain weight ratio in males fed 30 ppm, in mean absolute liver weight and mean liver-to-body weight percentage in females fed 30 ppm, and in all liver weight parameters in males and females fed 100 ppm.

000820

Test material-related macroscopic findings were noted at a low incidence in the liver and glandular stomach of males and females in the 100 ppm dose group only. Liver findings included organs that were large and diffusely dark. In the glandular stomach, dark foci or areas were sometimes seen in the gastric mucosa.

Test material-related microscopic findings were seen in the liver of animals fed 30 and 100 ppm of N-MeFOSE and in the glandular stomach of males and females in the 100 ppm group only. No test material-related changes were seen in animals fed 3 ppm of N-MeFOSE. In the liver, one or more test material-related findings affected all animals fed 100 ppm and included slight-to-moderate centrilobular or diffuse hepatocellular hypertrophy, minimal-to-slight coagulative necrosis, minimal hepatocellular pigment, and minimal-to-slight centrilobular to midzonal or diffuse hepatocellular vacuolation. In animals fed 30 ppm, centrilobular hepatocellular hypertrophy was also seen, although the incidence and severity was decreased at this dose. In addition, a low incidence of minimal hepatocellular vacuolation occurred in 30 ppm dose group males, and the liver of a single female fed 30 ppm had minimal coagulative necrosis. A possible test material-related change in the glandular stomach of animals fed 100 ppm was a relatively low incidence of mucosal erosions. The erosions were of minimal to slight severity and involved a very small portion of the mucosal surface in all cases.

CONCLUSIONS

Based on anatomic pathology, clinical pathology, and body weight effects at 30 and 100 ppm the no-observable-adverse-effect level was determined to be less than 30 ppm when N-MeFOSE is administered at concentrations of 0, 3, 30 and 100 ppm to CrI:CD[®](SD) IGS BR rats for at least 13 weeks through dietary administration. Based on the body weight effects at 3, 30, and 100 ppm, the no-observable-effect level was determined to be less than 3 ppm when N-MeFOSE is administered at concentrations of 0, 3, 30 and 100 ppm to CrI:CD[®](SD) IGS BR rats for at least 13 weeks through dietary administration. In the absence of correlative changes in anatomic pathology and clinical pathology the body weights effects at 3 ppm were not considered to be biologically adverse.


C00821

SIGNATURES



Nii van Bruce-Konuah
Study Coordinator
Covance Laboratories Inc.

30 June 2000
Date



Peter J. Thomford, PhD
Study Director
Covance Laboratories Inc.

30 June 2000
Date

C00822

REFERENCES

Dunnett, C. W., "New Tables for Multiple Comparisons with a Control," Biometrics, 20:482-491 (1964).

Levene, H., "Robust Tests for Equality of Variances," Contributions to Probability and Statistics, (eds.) I. Olkin et al., Ch. 25, pp. 278-292, Stanford University Press: Stanford, California (1960).

Winer, B. J., "Design and Analysis of Single-Factor Experiments," Statistical Principles in Experimental Design, Second Ed., Ch. 3, pp. 149-260, McGraw-Hill: New York, New York (1971a).

000323


OPHTHALMOLOGY REPORT

Ophthalmic examinations were done before initiation of treatment and during Week 14.

Animals selected for the study had no lesions at the prestudy examination. No test material-related ophthalmic observations were noted at the Week 14 examination.



Donna J. Clemons, DVM, MS
Diplomate, ACLAM



Date

000824

PATHOLOGY REPORT

SUMMARY

The purpose of this study was to assess the toxicity of the test material, N-Methyl Perfluorooctanesulfonamido Ethanol (N-MeFOSE), when administered in the diet to rats for at least 13 weeks. The test material was administered at dose levels of 3, 30, and 100 ppm.

Dietary administration of N-MeFOSE was associated with lower hematocrit for males and females fed 100 ppm; higher urea nitrogen for males and females fed 100 ppm; higher albumin for males fed 100 ppm; lower globulin for males fed 30 or 100 ppm; lower cholesterol and triglycerides for males fed 30 or 100 ppm and females fed 100 ppm; higher alanine aminotransferase for males fed 100 ppm; and higher hepatic palmitoyl CoA oxidase for males and females fed 100 ppm. The effects were relatively mild with the exception of those on cholesterol and triglycerides for the males fed 100 ppm; those effects were considered to be of moderate severity. All of the test material-related clinical pathology findings were more prominent at Week 14 than at Week 5.

Statistically significant and test material-related changes in terminal body weights and liver weights were seen in males and/or females in the two highest dose groups. Statistically significant decreases were seen in the terminal body weights of males fed 30 ppm and in both males and females fed 100 ppm. Statistically significant increases were seen in the mean liver-to-body weight percentage and mean liver-to-brain weight ratio in males fed 30 ppm, in mean absolute liver weight and mean liver-to-body weight percentage in females fed 30 ppm, and in all liver weight parameters in males and females fed 100 ppm.

Test material-related macroscopic findings were noted at a low incidence in the liver and glandular stomach of males and females in the 100 ppm dose group only. Liver findings included organs that were large and diffusely dark. In the glandular stomach, dark foci or areas were sometimes seen in the gastric mucosa.

Test material-related microscopic findings were seen in the liver of animals fed 30 and 100 ppm of N-MeFOSE and in the glandular stomach of males and females in the 100 ppm group only. No test material-related changes were seen in animals fed 3 ppm of

C00825

N-MeFOSE. In the liver, one or more test material-related findings affected all animals fed 100 ppm, and included slight-to-moderate centrilobular or diffuse hepatocellular hypertrophy, minimal-to-slight coagulative necrosis, minimal hepatocellular pigment, and minimal-to-slight centrilobular to midzonal or diffuse hepatocellular vacuolation. In animals fed 30 ppm, centrilobular hepatocellular hypertrophy was also seen, although the incidence and severity was decreased at this dose. In addition, a low incidence of minimal hepatocellular vacuolation occurred in 30 ppm dose group males, and the liver of a single female fed 30 ppm had minimal coagulative necrosis. A possible test material-related change in the glandular stomach of animals fed 100 ppm was a relatively low incidence of mucosal erosions. The erosions were of minimal to slight severity and involved a very small portion of the mucosal surface in all cases.

METHODS

Four groups of Crl:CD®(SD) IGS BR rats (20 animals/sex/group) were fed diets containing the test material at a dose level of 0 (control group; fed basal diet only), 3, 30, or 300 ppm. During Week 5, five animals/sex/group were sacrificed to obtain liver samples for determination of hepatic palmitoyl CoA oxidase activity (by Covance) and analysis for PFOS and metabolites (by 3M). One female fed 3 ppm was sacrificed and necropsied during Week 9 because of poor health (unscheduled sacrifice). The remaining animals were sacrificed and necropsied during Week 14 (terminal sacrifice).

Blood was collected for hematology and clinical chemistry tests during Weeks 5 and 14. Urine was collected for urinalysis tests during Week 14. At necropsy, macroscopic observations were recorded, organ weights were obtained (terminal sacrifice only), and tissues were placed in fixative as specified by the protocol. In addition, samples of liver were collected at the terminal sacrifice for evaluation of proliferation cell nuclear antigen (PCNA; by Pathology Associates International; five animals/sex/group), and both scheduled sacrifices for determination of hepatic palmitoyl CoA oxidase activity (by Covance) and analysis for PFOS and metabolites (by 3M). Microscopic examinations were done on protocol-required tissues from all control and high-dose animals sacrificed during Week 14 and the 3 ppm dose group female sacrificed in a moribund condition during Week 9 (unscheduled sacrifice). In addition, livers, stomachs, and macroscopic

C00826

lesions were examined from all terminal sacrifice animals fed 30 ppm, and macroscopic lesions and livers only were examined from animals fed 3 ppm.

Statistically significant differences cited in the Results and Discussion section are based on comparisons between the control and treated groups.

RESULTS AND DISCUSSION

Mortality

One female fed 3 ppm (Animal No. C96016) was sacrificed during Week 9 (Day 59) because of poor health. All other animals survived to the respective scheduled sacrifice.

Clinical Pathology

Weeks 5 and 14. There were several statistically significant or otherwise notable differences for clinical pathology results between control and treated animals. Differences considered to be effects of the test material were lower hematocrit for males and females fed 100 ppm; higher urea nitrogen for males and females fed 100 ppm; higher albumin for males fed 100 ppm, lower globulin for males fed 30 or 100 ppm; lower cholesterol and triglycerides for males fed 30 or 100 ppm and females fed 100 ppm; higher alanine aminotransferase for males fed 100 ppm; and higher hepatic palmitoyl CoA oxidase for males and females fed 100 ppm. The effects were relatively mild with the exception of those on cholesterol and triglycerides for the males fed 100 ppm; those effects were considered to be of moderate severity. All of the test material-related clinical pathology findings were more prominent at Week 14 than at Week 5.

The mild effects on hematocrit and globulin were consistent with animals failing to gain weight appropriately. Although mildly higher urea nitrogen and albumin suggested a relative dehydration for animals given 100 ppm, there were no effects on urine volume or urine specific gravity consistent with dehydration. The mild effect on alanine aminotransferase was indicative of the hepatocellular degeneration, and mildly higher hepatic palmitoyl CoA oxidase was indicative of peroxisomal proliferation. The effects on cholesterol and triglycerides were the most prominent findings, and suggested lipid metabolism by the liver was altered.

Of uncertain relationship to administration of the test material was statistically lower mean corpuscular volume for females fed 100 ppm. The difference for mean corpuscular volume became smaller over time (i.e., from Week 5 to Week 14). Other statistically significant differences were considered incidental and unrelated to administration of the test material.

Anatomic Pathology

Unscheduled Deaths. The one female fed 3 ppm that was sacrificed in a moribund condition on Day 59 (Animal No. C96016) had no macroscopic or microscopic changes to indicate that its moribund condition was test material-related.

Terminal Sacrifice

Terminal Body Weights and Organ Weights. Test material-related changes were seen in the terminal body weights and in absolute and relative liver weights in animals fed 30 and 100 ppm N-MeFOSE.

Mean terminal body weights were statistically significantly decreased in males and females that were fed 100 ppm of N-MeFOSE and in males only fed 30 ppm. The decreased mean body weights at both dose levels were considered to be test material-related.

Statistically significant and test material-related increases were seen in the mean liver-to-body weight percentage and the mean liver-to-brain weight ratio in males fed 30 ppm, in mean absolute liver weight and mean liver-to-body weight percentage in females fed 30 ppm, and in all liver weight parameters in males and females fed 100 ppm. The liver weight increases correlated macroscopically with large livers in 3 of 15 males in the 100 ppm dose group and microscopically with the presence of hepatocellular hypertrophy in all males and females in the high-dose group.

000828

Additional statistically significant absolute or relative organ weight changes affected several other organs in both sexes and in multiple dose groups. Because there were no macroscopic or microscopic findings that correlated with the weight changes, the remaining organ weight variations were considered to be within the range of normal biologic variation or were attributed to the lower body weights in treated versus control animals.

Macroscopic Findings. Test material-related macroscopic findings were noted at a low incidence in the liver and glandular stomach of males and females fed 100 ppm N-MeFOSE. Livers that were large were seen in 3 of 15 males and in 0 females fed 100 ppm, and diffusely dark livers occurred in 2 of 15 males and 1 of 15 females fed 100 ppm. No liver changes occurred in animals fed the control material or in animals fed lower doses of N-MeFOSE. The large and diffusely dark livers correlated microscopically with hepatocellular hypertrophy in the affected animals. In the glandular stomachs, dark foci or areas occurred at an incidence of 0 of 15, 1 of 15, 0 of 15, and 2 of 15 in males, and at an incidence of 0 of 15, 0 of 14, 0 of 15, and 3 of 15 in females in the control, 3 ppm, 30 ppm, and 100 ppm groups respectively. The dark foci or areas generally correlated microscopically with small erosions in the glandular mucosa.

Diffuse reddening and mottling of the thymus and mandibular lymph nodes as well as red foci or areas in the thymus were relatively common findings and frequently correlated microscopically with hemorrhage in the affected organs. Because these findings did not occur in a dose-related manner and because rats in multiple dose groups, including control males, were sometimes affected, the findings were considered to be agonal changes that were not associated with test material administration. All remaining macroscopic findings were considered to be spontaneous and incidental and not related to test material administration.

Microscopic Findings. Test material-related microscopic findings were seen in the liver of animals fed 30 and 100 ppm of N-MeFOSE and in the glandular stomach of males and females in the 100 ppm group only. There were no test material-related findings in the livers of animals fed 3 ppm.

000829

Test material-related findings in the liver included hepatocellular hypertrophy, coagulative necrosis, hepatocellular pigment, and hepatocellular vacuolation. The incidence and severity of these findings tended to be greater in 100 ppm animals. The incidence of pertinent liver findings is presented in Text Table 1.

Text Table 1

Incidence of Selected Microscopic Findings in the Liver

Group Number/Sex	1M	2M	3M	4M	1F	2F	3F	4F
Number Examined	15	15	15	15	15	14	15	15
Hypertrophy, hepatocellular, diffuse	0	0	0	0	0	0	0	1
Hypertrophy, hepatocellular, Centrilobular	0	0	14	15	0	0	7	14
Necrosis, coagulative	0	0	0	3	0	0	1	0
Pigment, hepatocellular	0	0	0	2	0	0	0	7
Vacuolation, hepatocellular, centrilobular to midzonal	0	0	2	7	0	0	0	2
Vacuolation, hepatocellular, diffuse	0	0	2	2	0	0	0	0


As noted in the above table, hepatocellular hypertrophy generally affected centrilobular areas of the liver in animals fed 30 and 100 ppm, but the liver was more diffusely hypertrophied in one female fed 100 ppm. The hypertrophic hepatocytes were enlarged with abundant cytoplasm that often had a granular to globular eosinophilic appearance. In some animals in the high-dose group (two males and seven females), the cytoplasm of hepatocytes also contained minimal amounts of a brown granular pigment at the periphery of the cell. In addition, liver cells frequently were vacuolated (see Text Table 1), particularly in males given 100 ppm, with the vacuolation generally affecting hepatocytes in centrilobular and midzonal areas. Diffuse hepatocellular vacuolation was also seen in two males in each of the 30 and 100 ppm dose groups, but in no other animals. Lastly, minimal to slight coagulative necrosis seen in 3 of 15 males fed 100 ppm and 1 of 15 females fed 30 ppm was likely related to dietary administration of the test material since no control or 3 ppm dose group animals were similarly affected.

A possible test material-related change in the glandular stomach was the presence of mucosal erosions. The erosions were of minimal to slight severity and involved a very small portion of the mucosal surface in all cases. The erosions were seen with an

C00830


incidence of 0 of 15, 0 of 15, and 2 of 15 in males, and at an incidence of 0 of 15, 0 of 15, and 3 of 15 in females in the control, 30 ppm, and 100 ppm groups respectively. The presence of erosions microscopically generally correlated with the dark foci or areas seen in the gastric mucosa at necropsy. One male fed 3 ppm also had a slight gastric erosion that was collected as a gross lesion at necropsy. Since no erosions were seen in any of the animals' stomachs examined from animals fed 30 ppm, the lesion in the 3 ppm male was considered unlikely to be test material-related.

All remaining microscopic findings were considered to be spontaneous and incidental changes and typical of rats of this age and strain.



Robert L. Hall, DVM, PhD
Diplomate, ACVP
(Clinical Pathology)

June 19, 2000
Date



Johnnie J. Eighmy, DVM, MS
Diplomate, ACVP
Diplomate, ABT

6/19/00
Date

000831

COMMENTS ON THE DATA

Various models of calculators, computers, and computer programs were used to analyze data in this study. Because different models round off or truncate numbers differently, values in some tables (e.g., means, standard deviations, or individual values) may differ slightly from those in other tables, from individually calculated data, or from statistical analysis data. Neither the integrity nor the interpretation of the data was affected by these differences.

The number of animals listed in the heading of the summary tables for clinical observations reflects the number of animals assigned to each group at the start of the study.

The summary table for clinical observations indicate the number of animals for which a condition was observed without regard to the specific nature, severity, reversibility, number of incidences/animal, or the length of time the condition persisted.

Only observations other than normal are indicated on the summary clinical observations tables.

Each animal with observations recorded as "Normal" throughout the study has the comment "Animal has no significant findings" indicated on the individual clinical observations tables.

The specific details for comments in the individual clinical observations tables that are indicated with a "C" can be found at the end of each group for each sex.

The day of initiation of treatment is "Day 1, Week 1." Body weight data are entered at the start of a study week (e.g., a body weight recorded on Day 1 is considered a Week 1 body weight, a body weight recorded on Day 8 is considered a Week 2 body weight). Body weight change data are calculated from the first day of the study week to the first day of the following study week (e.g., Week 1 values are calculated from Day 1 through 7) and are indicated in the tables with the day being the first day of the following week (e.g., Week 1 values are indicated as "Day 8."). Weekly food consumption and test material consumption are calculated from the first day of the study week to the first day of the following study week (e.g., Week 1 values are calculated from Day 1 through 7).

The values for summary and individual test material consumption are reported with more significant figures than is appropriate for the data due to limitations of the data collection and reporting software.

000832

COMMENTS ON THE DATA (Continued)

Food consumption values are reported in whole numbers; however, PTS carries food consumption values to one place to the right of the decimal for food efficiency and test material consumption calculations. The differences in values generated do not influence the interpretation of the calculation.

The comment "SPILLED" on individual food consumption data tables indicates that food consumption was not recorded due to spillage during the interval.

The comment "NOT TAKEN" on individual food consumption data tables indicates the animal died before the end of the food consumption interval.

Differences in the population size (N) on the summary tables for clinical and anatomic pathology are explained on the individual data tables or the codes sheets.

The calculation for individual test material consumption is:

Test material consumption =

$$\frac{\text{Diet Concentration} \times \text{Food Consumption}}{\text{Food Consumption Interval} \times [\text{First Body Weight of Interval} + (\text{Daily Body Weight Gain} \times \text{Day Factor})]}$$

where: Daily Body Weight Gain =

$$\frac{\text{Last Body Weight of Interval} - \text{First Body Weight of Interval}}{\text{Day of Last Body Weight of Interval} - \text{Day of First Body Weight of Interval}}$$

and: Day Factor =

$$(\text{Food Consumption Start Day} - \text{Day of First Body Weight of Interval}) + \frac{\text{Food Consumption Interval}}{2}$$

and: Food Consumption Interval =

$$\text{Food Consumption End Day} - \text{Food Consumption Start Day}$$

000833

CODES, ABBREVIATIONS, AND UNITS

General Codes and Abbreviations

Codes for Clinical Pathology

Abbreviations and Units for Clinical Hematology

Abbreviations and Units for Clinical Chemistry

Abbreviations and Units for Clinical Urinalysis

Codes for Anatomic Pathology

Note: The following lists of codes, abbreviations, and units are used by Covance. Some, but not necessarily all, of this information may be needed for this report.

000834

General Codes and Abbreviations

WK	Week.
N	Number of measurements in a group.
Mean; MEAN	Arithmetic mean.
SD; S.D.; STAND DEV;	Standard deviation.
STANDARD DEV; sd	
*	Group mean is significantly different from the mean of the control group (Group 1) at $p \leq 0.05$.
-, NA	No value; not applicable; not present.
P	Present.
C	Comment found at the end of each group for each sex.
UNSCHED	Unscheduled.
DISPATCH	Observations transferred from the in-life module of the data collection system to the necropsy module for reference during necropsy. Observations are duplicates of the last in-life observations.
TBW	Terminal body weight.
#	Number.
CO	Clinical observation.
Animal Death Codes:	
1	Interim sacrifice 1.
T	Terminal sacrifice.
M	Sacrificed at an unscheduled interval.

000835

Codes for Clinical Pathology

GENERAL CODES

NS	No sample
QS/QNS	Quantity not sufficient
NR	No repeat (sample volume not sufficient for repeat analysis)
FS	Fibrin strands
SC	Sample clotted
SH	Slightly hemolyzed
H	Hemolyzed
SL	Slightly lipemic
L	Lipemic
SI	Slightly icteric
I	Icteric
NF	Animal not fasted
U	Unscheduled/moribund bleed
DT/DOT	Animal died on test
DB	Died during bleeding
TJ	Technician judgment to repeat test
TE	Technical error (instrument or technician error that results in unacceptable data, e.g., unacceptable instrument output, sample spilled, entry of invalid data)
RE	Recording error (recorded incorrect data, e.g., wrong number, spelling error, incorrect date)
EE	Entry error (incorrect keyboard entry)
SE	Sampling error
PC	Platelets clumped
PD	Platelets decreased
PI	Platelets increased
PL	Platelets large
PA	Platelets appear adequate
CO	Color interferes with test
HB	Heinz bodies observed
PLASMO	Plasmodium
NO AGG	No aggregation
FR	Fractionous
UTD	Unable to determine
NO COAG	No coagulation

000836

Codes for Clinical Pathology (Continued)

RESULTS NOT INCLUDED IN STATISTICAL ANALYSES

Hemolyzed clinical chemistry or coagulation samples
 Samples from animals at unscheduled intervals
 Prothrombin times (PT) greater than 50 seconds
 Activated partial thromboplastin times (PTT) greater than 110 seconds
 Bleed times (BLETIME) greater than 30 minutes

CODES FOR BLOOD CELL MORPHOLOGY

The following scale was used to measure the degree of anisocytosis (ANISO), poikilocytosis (POIK), polychromasia (POLY), hypochromasia (HYPO), or basophilic stippling (BASTIP) or the presence of Howell-Jolly bodies (HJBODY), toxic neutrophils (TOXNEUT), or atypical lymphocytes (ATYPLYM):

Scale	Degree	Presence
-	Normal for the species	Not present
1	Slight	Rare
2	Moderate	Few
3	Marked	Moderate
4	Not applicable	Many

URINE APPEARANCE

Color			Clarity	Miscellaneous
A Pale	E Amber	I Black	J Clear	M Debris
B Straw	F Brown	P Blue/green	K Hazy	O Feces
C Yellow	G Red	Q Blue	L Cloudy	
D Dark yellow	H Green	R Orange		

000337

Codes for Clinical Pathology (Continued)

URINE CHEMISTRY MULTISTIX® STRIP

Urine Glucose		Urine Ketone		Urine Blood	
-	Negative	-	Negative	-	Negative
+	100 mg/dL	+	5 mg/dL	+	Small
++	250 mg/dL	++	15 mg/dL	++	Moderate
+++	500 mg/dL	+++	40 mg/dL	+++	Large
++++	1,000 mg/dL	++++	80 mg/dL		
+++++	≥2,000 mg/dL	+++++	160 mg/dL		

Urine Urobilinogen		Urine Bilirubin	
-	0.2 mg/dL	-	Negative
+	1 mg/dL	+	Small
++	2 mg/dL	++	Moderate
+++	4 mg/dL	+++	Large
++++	8 mg/dL		

(1 mg = approximately 1 Ehrlich unit)

URINE SEDIMENT

Cells, Crystals, Casts, and Comments		Bacteria
A Amorphous urates	Q Sperm	0 Not present
B Amorphous phosphates	R Fecal contamination	1 Few
C Uric acid	S Pinworm ova found	2 Moderate
D Triple phosphates	T Pinworm larvae found	3 Many
E Calcium oxalate	U Parasite ova found	
F Calcium carbonate		
G Granular casts		
H Hyaline casts	0 Not present	
I Cellular casts	1 1-5 per field	
J Waxy casts	2 6-10 per field	
K Unknown crystal	3 11-20 per field	
P Mucous threads	4 >20 per field	

C00838

Abbreviations and Units for Clinical Hematology

Test	Abbreviation (Units)
Red blood cell count	RBC (E6/UL or $\times 10^6/\mu\text{L}$)
Hemoglobin	HGB (G/DL)
Hematocrit	HCT (%)
Mean corpuscular volume	MCV (FL)
Mean corpuscular hemoglobin	MCH (PG)
Mean corpuscular hemoglobin concentration	MCHC (%)
Platelet count	PLT (E3/UL or $\times 10^3/\mu\text{L}$)
Mean platelet volume	MPV (FL)
Reticulocyte count	RETIC (%)
Absolute reticulocyte count	RETIC (E3/UL or $\times 10^3/\mu\text{L}$)
Heinz body count	HEINZ (%)
Erythrocyte sedimentation rate	ESR (MM/HR)
Prothrombin time	PT (SEC)
Activated partial thromboplastin time	PTT (SEC)
Thrombin time	TT (SEC)
Activated coagulation time	ACT (SEC)
Fibrinogen	FBR (MG/DL)
Fibrin/fibrinogen degradation products	FDP (UG/ML)
Platelet aggregation	
Collagen	PAGG/COL (%)
Adenosine diphosphate	PAGG/ADP (%)
Alpha 2-antiplasmin	ANTIPLAS (%)
Bleeding time	BLE TIME (SEC)
Methemoglobin	METHGB (%)
Plasma hemoglobin	PLA HGB (MG/DL)
Myeloid/erythroid ratio	M/E RATIO
Estimated myeloid/erythroid ratio	EST M/E RATIO
White blood cell count	WBC (E3/UL or $\times 10^3/\mu\text{L}$)
Differential blood cell count	
Nucleated red blood cell count	NRBC (/100 WBC)
Corrected white blood cell count	COR WBC (E3/UL or $\times 10^3/\mu\text{L}$)
Segmented neutrophil count	N-SEG (E3/UL or $\times 10^3/\mu\text{L}$) and %
Band neutrophil count	N-BAND (E3/UL or $\times 10^3/\mu\text{L}$) and %
Lymphocyte count	LYMPH (E3/UL or $\times 10^3/\mu\text{L}$) and %
Monocyte count	MONO (E3/UL or $\times 10^3/\mu\text{L}$) and %
Eosinophil count	EOSIN (E3/UL or $\times 10^3/\mu\text{L}$) and %
Basophil count	BASO (E3/UL or $\times 10^3/\mu\text{L}$) and %
Anisocytosis	ANISO (-,1,2,3)
Polychromasia	POLY (-,1,2,3)

000339

Abbreviations and Units for Clinical Hematology (Continued)

Test	Abbreviation (Units)
Poikilocytosis	POIK (-,1,2,3)
Hypochromasia	HYPO (-,1,2,3)
Howell-Jolly bodies	HJBODY (-,1,2,3,4)
Basophilic stippling	BASTIP (-,1,2,3)
Toxic neutrophils	TOXNEUT (-,1,2,3,4)
Atypical lymphocytes	ATYPLYM (-,1,2,3,4)
Aqueous white blood cell count (right eye)	R EYE (WBC/UL)
Aqueous white blood cell count (left eye)	L EYE (WBC/UL)

C00820

Abbreviations and Units for Clinical Chemistry

Test	Abbreviation (Units)
Glucose	GLU (MG/DL)
Urea nitrogen	UN (MG/DL)
Urea	UREA (MG/DL)
Creatinine	CREAT (MG/DL)
Total protein	T PRO (G/DL)
Albumin	ALB (G/DL)
Globulin	GLOB (G/DL)
Albumin/globulin ratio	A/G RATIO
Total bilirubin	T BILI (MG/DL)
Direct bilirubin	D BILI (MG/DL)
Indirect bilirubin	I BILI (MG/DL)
Cholesterol	CHOL (MG/DL)
Triglyceride	TRIG (MG/DL)
Urea nitrogen/creatinine ratio	UN/CREAT (RATIO)
Total lipids	T LIPIDS (MG/DL)
Phospholipids	P LIPIDS (MG/DL)
High-density lipoprotein cholesterol	HDL (MG/DL)
Low-density lipoprotein cholesterol	LDL (MG/DL)
Uric acid	UA (MG/DL)
Aspartate aminotransferase	AST/SGOT (IU/L)
Alanine aminotransferase	ALT/SGPT (IU/L)
Alkaline phosphatase	ALK PHOS (IU/L)
Gamma glutamyl transferase	GGT (IU/L)
Sorbitol dehydrogenase	SDH (IU/L)
Lactate dehydrogenase	LDH (IU/L)
Creatine kinase	CK (IU/L)
Amylase	AMYLASE (IU/L)
Lipase	LIPASE (IU/L)
Palmitoyl CoA oxidase	PCOAO (IU/G)
Calcium	CA (MG/DL)
Ionized calcium	ION CA (MG/DL)
Inorganic phosphorus	I PHOS (MG/DL)
Sodium	NA (MMOL/L)
Potassium	K (MMOL/L)
Chloride	CL (MMOL/L)
Magnesium	MG (MEQ/L or MG/DL)
Zinc	ZN (MG/L or PPM)
Strontium	SR (MG/L or PPM)
Iron	FE (UG/DL)

000341

Abbreviations and Units for Clinical Chemistry (Continued)

Test	Abbreviation (Units)
Excess iron	EX FE (UG/DL)
Total iron binding capacity	TIBC (UG/DL)
Unbound iron binding capacity	UIBC (UG/DL)
Percent iron saturation	FE %SAT (%)
Plasma cholinesterase	CHEP (MU/ML)
Red blood cell cholinesterase	CHER (MU/ML)
Brain cholinesterase	CHEB (MU/ML)
Caudate putamen	CAUD PUT (UMOL/G)
Hippocampus	HIPPOCAM (UMOL/G)
Frontal cortex	F CORTEX (UMOL/G)
Cerebellum	CEREBELL (UMOL/G)
Bicarbonate	BICARB (MMOL/L)
Serum hemoglobin	SER HGB (MG/DL)
Serum bile acids	SBA (UMOL/L or MG/DL)
Fecal bile acids	FBA (UG/ML)
Average fecal weight	FCC WGT (G)
Fecal bile acids (calculation)	FBA (MG/Day)
Osmolality	OSMO (MOSM/KG)
Electrophoresis	
Albumin	E ALB (G/DL)
Alpha-1-globulin	E A-1 (G/DL)
Alpha-2-globulin	E A-2 (G/DL)
Beta globulin	E BETA (G/DL)
Gamma globulin	E GAMMA (G/DL)
High-density lipoprotein	E-HDL (%)
Low-density lipoprotein	E-LDL (%)
Very-low-density lipoprotein	E-VLDL (%)
Insulin	INSULIN (UU/ML)
Adrenocorticotrophic hormone	ACTH (PG/ML)
Cortisol	CORTISOL (UG/ML)
Glucagon	GLUCAGON (PG/ML)
Triiodothyronine	T3 (NG/DL)
Thyroxine	T4 (UG/DL)
Creatine kinase isoenzymes	
BB	CK-BB (U/L)
MB	CK-MB (U/L)
MM	CK-MM (U/L)

000342

Abbreviations and Units for Clinical Urinalysis

Test	Abbreviation (Units)
Urine volume	U VOL (ML)
8 hour urine volume	8 HR VOL (ML)
Specific gravity	SP GR
Urine osmolality	U OSMO (MOSM/KG)
Quantitative urinary/cerebrospinal fluid protein	QUAN PRO (MG/DL)
Urine protein excretion	PRO EXC (MG)
Urine chemistry Multistix® strip	
Urine pH	U PH
Urine protein	U PRO (MG/DL)
Urine glucose	U GLU
Urine ketones	U KET
Urine bilirubin	U BILI
Urine blood	U BLOOD
Urine urobilinogen	UROBILI
Urine reducing substances	U RE SUB
Microscopic examination of urine sediment	
Red blood cells per high-power field	RBC (PER HPF)
White blood cells per high-power field	WBC (PER HPF)
Epithelial cells per high-power field	EPITH (PER HPF)
Bacteria per high-power field	BACT (PER HPF)
Casts per low-power field	CASTS (PER LPF)
Crystals per low-power field	CRYSTALS (PER LPF1 or PER LPF2)
Urine appearance	URINE APP1 or URINE APP2
Comments	COMMENTS

Miscellaneous Codes and Abbreviations for Clinical Pathology

Fecal occult blood	Not applicable
Fecal parasite detection	Not applicable
Hemolytic potential	Not applicable
Osmolality	OSMO (MOSM/KG)

Codes for Anatomic Pathology

Code	Definition
ANIMAL DEATH CODES	
I	Interim sacrifices 1
T	Terminal sacrifice
M	Sacrificed at an unscheduled interval

MACROSCOPIC CODES

EX	Indicates that organ weight is excluded from calculations
NOT TAKEN	Organ weight not taken; explanation given in necropsy notes
MISSING	Organ missing or lost
UNSUITABLE	Organ technically unsuitable for weighing
AUTOLYTIC	Organ autolyzed and could not be weighed
EXCLUDE	Weight was taken, but was excluded from all calculations

MICROSCOPIC CODES

Codes Prefacing Neoplastic Findings

B-	Primary, benign neoplasm
M-	Primary, malignant neoplasm
N-	Metastatic neoplasm
I-	Locally invasive neoplasm
X-	Other neoplasm

Code	Definition
------	------------

Distribution of Findings

Focal
Diffuse
Multifocal

000314

Codes for Anatomic Pathology (Continued)

Grades for Severity or Amount

1	Minimal - the least amount of change that can be observed with the light microscope
2	Slight - less than average amount of change, but readily discernible as abnormal
3	Moderate - the average amount of change that is expected for a lesion
4	Moderately severe (marked) - a marked amount of change with possible loss of function of the affected cells or organs
5	Severe - a great amount of change with probable loss of function of the affected cell or organs and frequently involves large areas of the organ

Other Microscopic Codes

TL	Total
P	Finding present
-	Finding not present
MN	Mean

TISSUE ABBREVIATIONS

Abbreviation	Definition
LN	Lymph node
GL	Gland
STOMACH, GL	Glandular stomach
STOMACH, NONGL	Nonglandular
SALIV GL, MANDIB	Mandibular salivary gland
LN, ANT MES/PANC	Anterior mesenteric/pancreatic lymph node
AUDITORY SEB GL	Auditory sebaceous gland
LACRIMAL GLAND, EX	Exorbital lacrimal gland
HEMATO NEOPLASIA	Hematopoietic neoplasia
LACRIMAL GL, INT	Internal lacrimal gland
CAVITY, ABDOM	Abdominal cavity
SALIV GL, PAROTID	Parotid salivary gland
LN, TRACHEOBRON	Tracheobronchial lymph node

000845

Table 1

Results of Homogeneity Analyses (ppm)
Mixed 8/20/98

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MeFOSE, T6314) IN RATS

Samples Location	Replicate	T6314 (ppm)		
		3	30	100
Top	1	2.19	33.7	79.4
	2	2.43	27.8	76.5
	3	2.09	25.6	76.7
	Mean	2.24 (74.7) ^a	29.0 (96.7)	77.5 (77.5)
Middle	1	2.24	25.2	75.0
	2	2.27	25.3	79.7
	3	2.14	25.1	89.5
	Mean	2.22 (74.0)	25.2 (84.0)	81.4 (81.4)
Bottom	1	2.03	23.7	83.2
	2	3.04	27.6	81.6
	3	1.94	29.4	87.8
	Mean	2.34 (78.0)	26.9 (89.7)	84.2 (84.2)

^a Each value in parenthesis is the percent of theoretical.

C00346

Table 2

Results of Stability Analyses (ppm)
Mixed 8/20/98

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MeFOSE, T6314) IN RATS

Storage Conditions	T6314 (ppm)		
	Replicate	1	500
Initial	1	1.88	415
	2	0.954	468
	Mean	1.42 (142)	442 (88.4) ^a
19 day, room temperature	1	0.733	-
	2	0.811	-
	Mean	0.772 (77.2)	-
32 day, room temperature	1	-	444
	2	-	425
	Mean	-	435 (87.0)
32 day, room temperature ^b	1	0.709	-
	2	0.850	-
	Mean	0.780 (78.0)	-
8 week, frozen	1	0.779	-
	2	0.995	-
	Mean	0.887 (88.7)	-

a Each value in the parenthesis is the percent of theoretical.

b Reinjecting

C00847

Table 2 (Continued)

Results of Stability Analyses (ppm)
Mixed 9/24/98

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MeFOSE, T6314) IN RATS

Storage Conditions	Replicate	T6314 (ppm)	
		500	1
Initial	1	-	0.788
	2	-	0.656
	Mean	-	0.722 (72.2) ^a
19 day, room temperature	1	-	0.707
	2	-	0.651
	Mean	-	0.679 (67.9)
32 days, room temperature	1	-	0.688
	2	-	0.657
	Mean	-	0.673 (67.3)

^a Each value in the parenthesis is the percent of theoretical.

000348

Table 2 (Continued)

Results of Stability Analyses (ppm)
Mixed 10/22/98

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MeFOSE, T6314) IN RATS

Storage Conditions	Replicate	T6314 (ppm)
		3
Initial	1	2.24
	2	2.49
	Mean	2.37 (79.0) ^a
19 day, room temperature	1	1.89
	2	2.33
	Mean	2.11 (70.3)

^a Each value in the parenthesis is the percent of theoretical.

000849

Table 3

Results of Dose Preparation Analyses (ppm)

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

Week	Replicate	T-6314 (ppm)			
		0	3	30	100
1-4	1	a	2.16	29.7	87.3
	2	a	2.12	27.0	85.2
	Mean	-	2.14 (71.3) ^b	28.4 (94.7)	86.3 (86.3)
1-4 ^c	1	-	1.98	-	-
	2	-	1.84	-	-
	Mean	-	1.91 (63.7)	-	-
1-4 ^d	1	-	2.48	-	-
	2	-	2.28	-	-
	Mean	-	2.38 (79.3)	-	-
5-8	1	a	3.55	27.9	89.3
	2	a	3.05	27.0	82.3
	Mean	a	3.30 (110)	27.5 (91.7)	85.8 (85.8)
9/12	1	a	2.24	27.6	89.7
	2	a	2.49	26.8	87.2
	Mean	a	2.37 (79.0)	27.2 (90.7)	88.5 (88.5)
13/14	1	a	2.51	26.2	98.1
	2	a	2.42	26.4	98.7
	Mean	a	2.47 (82.3)	26.3 (87.7)	98.4 (98.4)

a Below the limit of quantitation (<0.5 ppm).

b Each value in parenthesis is the percent of theoretical.

c Retest.

d Retention.

000850

Table 4

Summary of Clinical Observations

PAGE: 1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEPOSE, T-6314) IN RATS

DAYS 1-93	NUMBER OF ANIMALS AFFECTED											
	SEX:						FEMALE					
CATEGORY	1	2	3	4	1	2	3	3	3	3	4	100
KEYWORD	0	3	30	100	0	0	0	0	0	0	0	100
QUALIFIER	20	20	20	20	20	20	20	20	20	20	20	20
*** TOP OF LIST ***												
APPEARANCE												
LIMITED USE	0	0	0	0	0	1	0	0	0	0	0	0
LIMB-HIND-LEFT	3	6	3	4	1	1	3	3	1			
MALOCCLUSION	0	0	0	0	0	0	0	0	1			
MISSING	0	0	0	0	1	0	0	0	0			
TAIL-DISTAL	0	0	0	0	0	0	0	0	0			
TEETH	0	0	0	0	0	1	0	0	0			
SWOLLEN	0	1	0	0	0	0	0	0	0			
LIMB-HIND-LEFT												
THIN												
DISCHARGE												
GENITAL	1	0	0	0	0	0	0	0	0			
RED IN COLOR												
NASAL	0	0	0	1	0	0	0	0	0			
RED IN COLOR												
EYES												
RED DISCHARGE	0	0	0	1	0	0	0	0	0			
EYES												
RESPIRATION												
AUDIBLE	0	0	0	1	0	0	0	0	0			

000351

Table 4

Summary of Clinical Observations

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

DAYS 1-93	NUMBER OF ANIMALS AFFECTED											
	SEX:			MALE			FEMALE					
CATEGORY	GROUP:	1	2	3	4	1	2	3	4			
KEYWORD	DOSE:	0	3	30	100	0	3	30	100			
QUALIFIER	NUMBER:	20	20	20	20	20	20	20	20	20	20	20
SKIN & PELAGE												
BLACK SKIN		1	0	0	0	0	0	0	0	0	0	0
TAIL-DISTAL												
SORE/SCAB		0	0	0	0	0	0	0	1	0	0	0
MOUTH		0	0	0	0	0	0	1	0	0	0	0
TAIL		0	0	0	0	0	0	1	0	0	0	0
YELLOW HAIR COAT												
PENIS		0	0	1	0	0	0	0	0	0	0	0

*** END OF LIST ***

000352

Table 5

Summary of Ophthalmic Observations
Week 14

PAGE: 1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

		NUMBER OF ANIMALS AFFECTED											
WEEK 14		SEX:						MALE					
		GROUP:						DOSE:					
		1						2					
		3						4					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					
		15						15					

Table 6

Summary of Body Weight Data (g)

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 1

WEEK	SEX:		MALE				FEMALE			
	GROUP:		1		2		1		2	
	DOSE:	UNITS:	0	3	3	4	0	3	3	4
			PPM	PPM	PPM	PPM	PPM	PPM	PPM	PPM
1	N	MEAN	20	20	20	20	20	20	20	20
		S.D.	210	211	211	208	159	159	160	158
			11.2	10.2	11.7	11.9	10.0	11.6	9.2	10.3
2	N	MEAN	20	20	20	20	20	20	20	20
		S.D.	263	263	263	258	185	180	183	178
			16.0	13.4	13.8	13.1	11.9	15.5	11.8	12.6
3	N	MEAN	20	20	20	20	20	20	20	20
		S.D.	313	314	313	303	203	200	203	194
			21.5	16.6	17.1	15.9	16.0	25.0	15.2	17.5
4	N	MEAN	20	20	20	20	20	20	20	20
		S.D.	355	355	351	335*	224	222	223	207*
			25.1	19.9	20.5	18.5	17.4	23.3	14.1	16.6
5	N	MEAN	20	20	20	20	20	20	20	20
		S.D.	385	382	378	359*	236	230	233	215*
			28.6	20.5	21.6	21.4	21.7	23.1	14.6	16.2
6	N	MEAN	15	15	15	15	15	15	15	15
		S.D.	410	401	393	378*	247	237	235	223*
			24.4	23.9	21.1	19.2	22.9	27.3	14.7	21.0
7	N	MEAN	15	15	15	15	15	15	15	15
		S.D.	435	424	414*	403*	255	244	245	229*
			24.5	26.8	23.2	21.8	25.5	25.3	14.5	21.9

000354

Table 6

Summary of Body Weight Data (g)

PAGE: 2

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

WEEK	SEX:	MALE				FEMALE			
		GROUP:		DOSE:		GROUP:		DOSE:	
		1	2	3	4	1	2	3	4
UNITS:		PPM	PPM	PPM	PPM	PPM	PPM	PPM	PPM
8	N	15	15	15	15	15	15	15	15
	MEAN	452	439	428*	416*	259	245	248	231*
	S.D.	26.9	27.0	28.6	22.9	26.8	26.5	14.9	22.5
9	N	15	15	15	15	15	15	15	15
	MEAN	475	461	448*	430*	267	256	257	238*
	S.D.	28.1	25.8	28.4	24.6	26.3	26.9	17.6	22.3
10	N	15	15	15	15	15	14	15	15
	MEAN	490	475	460*	442*	275	259	263	241*
	S.D.	29.7	28.4	30.2	23.7	29.2	23.6	16.8	22.6
11	N	15	15	15	15	15	14	15	15
	MEAN	503	485	471*	453*	281	264	266	245*
	S.D.	33.0	31.9	31.9	24.9	31.3	26.9	16.1	23.7
12	N	15	15	15	15	15	14	15	15
	MEAN	516	496	480*	462*	287	266	270	246*
	S.D.	34.1	30.6	33.8	24.8	33.2	25.7	18.0	24.3
13	N	15	15	15	15	15	14	15	15
	MEAN	532	510	490*	471*	293	273	274	250*
	S.D.	36.1	30.0	30.1	25.1	36.2	29.3	19.6	24.5
14	N	15	15	15	15	15	14	15	15
	MEAN	547	521	491*	481*	300	280	279	254*
	S.D.	38.6	27.6	37.1	25.3	35.9	26.7	19.3	24.5

000855

Table 7

Summary of Body Weight Change Data (g)

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 1

WEEK	GROUP:	SEX:				MALE				FEMALE			
		DOSE:		PPM		DOSE:		PPM		DOSE:		PPM	
		UNITS:	PPM	UNITS:	PPM	UNITS:	PPM	UNITS:	PPM	UNITS:	PPM		
1-2	N	20	20	20	20	20	20	20	20	20	20	20	20
	MEAN	53	52	54	53	50	50	50	50	50	50	50	50
	S.D.	6.5	5.4	5.4	4.6	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8
2-3	N	20	20	20	20	20	20	20	20	20	20	20	20
	MEAN	49	51	51	49	45*	45*	45*	45*	45*	45*	45*	45*
	S.D.	9.3	5.6	5.6	5.0	4.6	4.6	4.6	4.6	4.6	4.6	4.6	4.6
3-4	N	20	20	20	20	20	20	20	20	20	20	20	20
	MEAN	42	41	41	39	32*	32*	32*	32*	32*	32*	32*	32*
	S.D.	10.4	5.5	5.5	5.2	4.1	4.1	4.1	4.1	4.1	4.1	4.1	4.1
4-5	N	20	20	20	20	20	20	20	20	20	20	20	20
	MEAN	31	27	27	26	24*	24*	24*	24*	24*	24*	24*	24*
	S.D.	7.6	3.6	3.6	4.9	6.1	6.1	6.1	6.1	6.1	6.1	6.1	6.1
5-6	N	15	15	15	15	15	15	15	15	15	15	15	15
	MEAN	19	17	17	17	14	14	14	14	14	14	14	14
	S.D.	7.8	3.9	3.9	4.7	4.9	4.9	4.9	4.9	4.9	4.9	4.9	4.9
6-7	N	15	15	15	15	15	15	15	15	15	15	15	15
	MEAN	26	23	23	21*	24	24	24	24	24	24	24	24
	S.D.	3.3	4.4	4.4	3.6	6.2	6.2	6.2	6.2	6.2	6.2	6.2	6.2
7-8	N	15	15	15	15	15	15	15	15	15	15	15	15
	MEAN	16	15	15	14	13	13	13	13	13	13	13	13
	S.D.	4.3	4.9	4.9	8.1	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8

Table 7

Summary of Body Weight Change Data (g)

PAGE: 2

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

WEEK	SEX:	MALE				FEMALE			
		1	2	3	4	1	2	3	4
GROUP:		0	3	30	100	0	3	30	100
DOSE:		PPM	PPM	PPM	PPM	PPM	PPM	PPM	PPM
UNITS:									
MEAN									
S.D.									
8-9	N	15	15	15	15	15	15	15	15
	MEAN	23	22	20	14*	8	11	9	7
	S.D.	3.9	4.3	3.7	10.3	7.1	3.5	4.9	3.1
9-10	N	15	15	15	15	15	14	15	15
	MEAN	16	15	12	12	7	5	6	3
	S.D.	5.4	4.9	5.7	7.5	6.3	3.8	4.4	2.5
10-11	N	15	15	15	15	15	14	15	15
	MEAN	12	10	11	11	6	5	3	3
	S.D.	4.7	19.0	4.1	3.1	4.7	5.5	5.2	2.8
11-12	N	15	15	15	15	15	14	15	15
	MEAN	13	10	9*	9*	6	2	4	2
	S.D.	5.1	11.5	4.3	2.5	5.9	4.3	4.5	4.8
12-13	N	15	15	15	15	15	14	15	15
	MEAN	16	15	10*	9*	6	8	4	4
	S.D.	4.2	6.3	10.6	4.0	7.2	5.6	5.7	2.8
13-14	N	15	15	15	15	15	14	15	15
	MEAN	15	10	1	10	6	6	5	3
	S.D.	5.0	5.0	29.5	3.0	7.7	7.7	5.4	4.1
1-14	N	15	15	15	15	15	14	15	15
	MEAN	38	307*	282*	270*	141	121*	120*	96*
	S.D.	35.8	27.0	35.5	21.5	30.7	17.8	14.9	17.0

000357

Table 8

Summary of Food Consumption Data (g)

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 1

WEEK	UNITS	SEX: MALE				FEMALE			
		GROUP:		DOSE:		GROUP:		DOSE:	
		1	2	3	4	1	2	3	4
		PPM	PPM	PPM	PPM	PPM	PPM	PPM	PPM
1	N	20	20	20	20	19	16	17	18
	MEAN	169	168	171	168	126	127	138	122
	S.D.	12.3	9.6	13.8	8.6	13.5	16.5	24.9	19.5
2	N	19	19	19	20	19	16	13	16
	MEAN	176	177	178	176	130	124	137	116
	S.D.	12.9	11.6	11.3	9.1	15.6	24.3	17.1	14.3
3	N	20	19	20	19	19	16	16	19
	MEAN	181	183	186	173	138	135	144	115*
	S.D.	16.2	16.5	14.6	8.0	16.7	16.8	18.7	13.3
4	N	18	19	20	20	17	16	13	20
	MEAN	186	186	191	176	135	134	138	121*
	S.D.	14.3	13.6	15.7	10.8	17.7	14.6	10.7	17.2
5	N	15	14	14	15	14	12	8	15
	MEAN	177	176	172	166	129	122	122	115
	S.D.	20.5	14.7	13.4	9.4	16.1	17.1	7.4	15.6
6	N	15	15	15	15	12	13	12	15
	MEAN	191	190	190	182	138	131	150	121*
	S.D.	14.7	17.5	22.8	9.4	18.8	13.7	20.5	14.1
7	N	14	15	15	15	12	12	6	14
	MEAN	191	192	184	182	144	138	136	117*
	S.D.	17.7	15.4	23.2	8.5	15.1	21.1	6.9	13.8

C000858

Table 8

Summary of Food Consumption Data (g)

PAGE: 2

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

WEEK	SEX:	MALE				FEMALE			
		1	2	3	4	1	2	3	4
		GROUP: DOSE: PPM	GROUP: DOSE: PPM	GROUP: DOSE: PPM	GROUP: DOSE: PPM	GROUP: DOSE: PPM	GROUP: DOSE: PPM	GROUP: DOSE: PPM	GROUP: DOSE: PPM
UNITS:		PPM	PPM	PPM	PPM	PPM	PPM	PPM	PPM
8	N	13	12	14	15	12	12	11	14
	MEAN	185	188	179	176	134	136	138	120
	S.D.	10.9	12.6	15.2	10.0	19.8	12.0	11.0	15.6
9	N	15	15	15	15	11	12	9	14
	MEAN	190	189	180	175	130	131	146	118
	S.D.	20.8	16.6	19.6	9.2	17.1	13.4	32.1	15.7
10	N	14	15	15	15	13	13	12	14
	MEAN	187	180	176	174	139	128	143	115
	S.D.	24.3	27.5	18.8	10.2	36.5	19.2	29.6	13.9
11	N	14	15	14	14	10	12	11	13
	MEAN	191	183	176	173	136	124	145	112*
	S.D.	29.7	17.5	22.1	9.7	22.8	11.9	32.6	12.5
12	N	13	15	15	15	13	13	15	14
	MEAN	185	181	175	173	137	129	146	114*
	S.D.	14.6	19.4	19.4	11.3	29.3	15.2	38.0	13.3
13	N	13	15	15	15	11	11	13	14
	MEAN	187	178	160	168*	133	120	143	119
	S.D.	20.7	12.3	46.1	8.9	26.4	17.2	34.8	26.4
1-13	N	9	10	13	13	8	4	4	10
	MEAN	2397	2329	2271	2283	1769	1667	1683	1505*
	S.D.	102.5	115.5	145.8	77.0	182.0	184.7	24.1	173.4

000859

Table 9

Summary of Test Material Consumption Data (mg/kg/day)

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 1

WEEK	SEX: MALE				FEMALE			
	GROUP:		DOSE:		GROUP:		DOSE:	
	UNITS:	PPM	PPM	PPM	UNITS:	PPM	PPM	PPM
1	N	20	20	20	N	16	17	18
	MEAN	0.304	3.094	10.291		0.319	3.395	10.378
	S.D.	0.017	0.242	0.471		0.040	0.601	1.400
2	N	19	19	20	N	16	13	16
	MEAN	0.263	2.649	8.941		0.273	3.028	8.867
	S.D.	0.012	0.125	0.331		0.042	0.512	1.027
3	N	19	20	19	N	16	16	19
	MEAN	0.235	2.394	7.794		0.269	2.861	8.142
	S.D.	0.015	0.126	0.276		0.027	0.393	0.679
4	N	19	20	20	N	16	13	20
	MEAN	0.216	2.243	7.238		0.252	2.570	8.219
	S.D.	0.010	0.137	0.335		0.015	0.170	0.997
5	N	14	14	15	N	12	8	15
	MEAN	0.192	1.920	6.399		0.226	2.262	7.464
	S.D.	0.014	0.132	0.350		0.012	0.188	0.685
6	N	15	15	15	N	13	12	15
	MEAN	0.198	2.017	6.660		0.236	2.662	7.675
	S.D.	0.017	0.176	0.231		0.030	0.337	0.703
7	N	15	15	15	N	12	6	14
	MEAN	0.191	1.873	6.343		0.247	2.348	7.325
	S.D.	0.015	0.152	0.186		0.051	0.140	0.627

C00860

Table 9
Summary of Test Material Consumption Data (mg/kg/day)

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

WEEK	N	MEAN	S.D.	MALE			FEMALE		
				2	3	4	2	3	4
				PPM	PPM	PPM	PPM	PPM	PPM
8	12	0.178	0.010	14	15	15	12	11	14
				1.764	5.946	5.946	0.232	2.335	7.243
				0.097	0.210	0.210	0.023	0.212	0.492
9	15	0.173	0.014	15	15	15	12	9	14
				1.694	5.749	5.749	0.218	2.394	6.995
				0.116	0.216	0.216	0.016	0.686	0.545
10	15	0.160	0.023	15	15	15	13	12	14
				1.618	5.554	5.554	0.209	2.310	6.721
				0.112	0.211	0.211	0.021	0.545	0.481
11	15	0.160	0.014	14	14	14	12	11	13
				1.582	5.369	5.369	0.197	2.274	6.529
				0.138	0.281	0.281	0.016	0.516	0.456
12	15	0.155	0.015	15	15	15	13	15	14
				1.544	5.301	5.301	0.206	2.319	6.494
				0.122	0.248	0.248	0.022	0.680	0.484
13	15	0.148	0.012	15	15	15	11	13	14
				1.390	5.054	5.054	0.184	2.204	6.700
				0.387	0.196	0.196	0.020	0.601	1.173
1-13				2.041	6.844	6.844	0.240	2.588	7.702

000861

Table 10
Summary of Clinical Hematology Data
Males Week 5
13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

DOSE ppm MeFOS	RBC X10 ⁶ /μL	HGB G/DL	HCT %	MCV FL	MCH PG	MCHC %	PLT X10 ³ /μL
0	MEAN 8.18 S.D. .424 N 10	16.2 .66 10	47.0 2.04 10	57.4 1.97 10	19.8 .42 10	34.5 .61 10	1173 61.5 10
3	MEAN 8.14 S.D. .427 N 10	16.4 .47 10	47.5 1.70 10	58.4 1.77 10	20.1 .73 10	34.4 .51 10	1151 136.0 10
30	MEAN 8.24 S.D. .477 N 10	16.3 .38 10	47.1 1.64 10	57.2 1.93 10	19.8 .81 10	34.7 .52 10	1038 * 96.9 10
100	MEAN 7.99 S.D. .303 N 10	15.9 .71 10	46.0 1.71 10	57.6 1.86 10	19.9 .76 10	34.5 .45 10	1081 112.7 10

000862

Table 10
Summary of Clinical Hematology Data
Males Week 5
13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDE
ETHANOL (N-MEFOSE, T-6314) IN RATS

DOSE ppm MeFOS	WBC $\times 10^3/\mu\text{L}$	N-SEG $\times 10^3/\mu\text{L}$	LYMPH $\times 10^3/\mu\text{L}$	MONO $\times 10^3/\mu\text{L}$	EOSIN $\times 10^3/\mu\text{L}$	BAO $\times 10^3/\mu\text{L}$	N-SEG%	LYMPH%	MONO%	EOSIN%	BAO%
0	MEAN 9.1 S.D. 1.96 N 10	.6 .14 10	7.9 1.86 10	.4 .15 10	.1 .03 10	.0 .00 10	.0 2.1 10	87 2.2 10	5 1.3 10	1 1.3 10	0 .3 10
3	MEAN 9.1 S.D. 1.74 N 10	.6 .10 10	8.0 1.64 10	.4 .20 10	.1 .05 10	.0 .00 10	.0 1.6 10	88 3.5 10	5 1.9 10	1 .7 10	0 .0 10
30	MEAN 10.0 S.D. 2.48 N 10	.5 .15 10	9.0 2.27 10	.4 .13 10	.1 .06 10	.0 .04 10	.0 1.4 10	90 1.8 10	4 1.1 10	1 .3 10	0 .4 10
100	MEAN 10.7 S.D. 3.51 N 10	.9 .40 10	9.0 3.17 10	.6 .20 10	.1 .03 10	.0 .05 10	.0 3.6 10	84 4.2 10	6 1.3 10	1 .3 10	0 .5 10

000863

Table 10
Summary of Clinical Hematology Data
Females Week 5
13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

DOSE ppm MeFOS	RBC x10 ⁶ /μL	HGB G/DL	HCT %	MCV FL	MCH PG	MCHC %	PLT x10 ³ /μL	
0	MEAN S.D. N	7.70 .214 10	15.7 .51 10	45.4 1.31 10	59.1 1.45 10	20.4 .68 10	34.5 .43 10	1114 104.4 10
3	MEAN S.D. N	8.03 .488 10	16.0 .61 10	46.1 2.08 10	57.6 1.95 10	20.0 .76 10	34.7 .64 10	1178 82.3 10
30	MEAN S.D. N	7.98 .357 10	16.0 .55 10	45.8 1.91 10	57.4 1.48 10	20.0 .56 10	35.0 .37 10	1123 121.8 10
100	MEAN S.D. N	7.89 .326 10	15.5 .54 10	44.7 1.62 10	56.6 * 1.28 10	19.6 .34 10	34.7 .71 10	1058 217.2 10

000864

Table 10

Summary of Clinical Hematology Data

Females Week 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

DOSE ppm MeFOS	WBC X10 ³ /μL	N-SEG X10 ³ /μL	LYMPH X10 ³ /μL	MONO X10 ³ /μL	EOSIN X10 ³ /μL	BASO X10 ³ /μL	N-SEG%	LYMPH%	MONO%	EOSIN%	BASO%	
0	MEAN S.D. N	6.3 1.59 10	.4 .29 10	5.4 1.32 10	.3 .15 10	.0 .05 10	.0 .00 10	7 3.1 10	87 4.5 10	5 1.5 10	1 .3 10	0 .3 10
3	MEAN S.D. N	7.4 2.21 10	.5 .20 10	6.5 2.20 10	.2 .12 10	.1 .07 10	.0 .00 10	7 3.2 10	88 4.0 10	4 1.6 10	1 .7 10	0 .0 10
30	MEAN S.D. N	7.3 2.42 10	.5 .19 10	6.4 2.24 10	.3 .19 10	.1 .05 10	.0 .00 10	8 4.4 10	87 5.4 10	4 1.8 10	1 .3 10	0 .0 10
100	MEAN S.D. N	7.3 2.09 10	.5 .18 10	6.4 1.96 10	.4 .27 10	.1 .04 10	.0 .03 10	7 3.7 10	87 4.5 10	5 2.3 10	1 .3 10	0 .3 10

000865

Table 11
Summary of Clinical Hematology Data
Males Week 14
13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

DOSE ppm MeFOS	RBC x10 ⁶ /μL	HGB G/DL	HCT %	MCV FL	MCH PG	MCHC %	PLT x10 ³ /μL
0	MEAN S.D. N	15.3 .71 10	47.0 2.00 10	54.9 1.43 10	17.8 .40 10	32.5 .40 10	1122 119.5 10
3	MEAN S.D. N	15.3 .67 10	46.7 2.02 10	55.6 2.31 10	18.2 .77 10	32.7 .32 10	1060 73.0 10
30	MEAN S.D. N	15.2 .56 10	46.8 1.99 10	54.0 1.79 10	17.6 .82 10	32.7 .71 10	1000 140.3 10
100	MEAN S.D. N	14.6 .71 10	44.3 * 1.92 10	53.7 1.39 10	17.7 .56 10	33.0 .30 10	1011 138.4 10

000366

Table 11

Summary of Clinical Hematology Data

Males Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

DOSE	WBC	N-SEG	LYMPH	MONO	EOSIN	BAO	N-SEG%	LYMPH%	MONO%	EOSIN%	BAO%
ppm MeFOS	X10 ³ /μL	X10 ³ /μL	X10 ³ /μL	X10 ³ /μL	X10 ³ /μL	X10 ³ /μL					
0	6.9 .88 10	.9 .28 10	5.4 .85 10	.5 .14 10	.1 .03 10	.0 .00 10	13 4.0 10	78 4.3 10	7 1.9 10	2 .5 10	0 .4 10
3	6.7 1.48 10	1.0 .23 10	5.1 1.35 10	.4 .18 10	.1 .05 10	.0 .00 10	15 3.4 10	76 5.6 10	7 2.8 10	2 .6 10	0 .3 10
30	7.7 2.68 10	.9 .24 10	6.2 2.57 10	.4 .14 10	.1 .05 10	.0 .00 10	13 4.7 10	80 5.6 10	5 2.1 10	1 .7 10	0 .0 10
100	7.2 2.63 10	1.1 .35 10	5.6 2.13 10	.4 .23 10	.1 .05 10	.0 .03 10	16 2.8 10	77 2.9 10	5 1.7 10	1 .4 10	0 .3 10

000867

Table 11

Summary of Clinical Hematology Data

Females Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

DOSE ppm MeFOS	RBC X10 ⁶ /μL	HGB G/DL	HCT %	MCV FL	MCH PG	MCHC %	PLT X10 ³ /μL
0	MEAN S.D. N	14.8 .45 10	45.1 .95 10	58.1 1.13 10	19.0 .54 10	32.7 .40 10	1016 103.3 10
3	MEAN S.D. N	15.1 .75 9	46.1 1.97 9	57.1 1.70 9	18.7 .63 9	32.7 .49 9	1073 108.7 9
30	MEAN S.D. N	15.1 .56 10	46.2 1.77 10	57.2 1.57 10	18.7 .52 10	32.7 .22 10	980 97.8 10
100	MEAN S.D. N	14.2 .67 10	43.1 * 1.91 10	56.1 * 1.26 10	18.5 .49 10	33.0 .37 10	1056 110.7 10

000868

Table 11

Summary of Clinical Hematology Data

Females Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDE
ETHANOL (N-MEFOSE, T-6314) IN RATS

DOSE ppm MeFOS	WBC x10 ³ /μL	N-SEG x10 ³ /μL	LYMPH x10 ³ /μL	MONO x10 ³ /μL	EOSIN x10 ³ /μL	BASO x10 ³ /μL	N-SEG%	LYMPH%	MONO%	EOSIN%	BASO%	
0	MEAN S.D. N	4.0 .95 10	.8 .47 10	3.0 .69 10	.2 .12 10	.1 .03 10	.0 .00 10	19 8.5 10	74 8.8 10	5 2.8 10	2 .6 10	0 .4 10
3	MEAN S.D. N	4.7 1.27 9	.7 .18 9	3.7 1.27 9	.3 .05 9	.1 .00 9	.0 .00 9	15 5.3 9	77 6.4 9	6 1.3 9	2 .8 9	0 .0 9
30	MEAN S.D. N	5.7 1.08 10	1.0 .49 10	4.4 1.10 10	.3 .10 10	.1 .00 10	.0 .03 10	17 7.6 10	76 8.0 10	5 1.7 10	1 .5 10	0 .3 10
100	MEAN S.D. N	4.7 2.03 10	.6 .28 10	3.8 1.89 10	.2 .09 10	.1 .04 10	.0 .00 10	14 7.0 10	80 8.4 10	4 1.8 10	2 .7 10	0 .3 10

000869

Table 12

Summary of Clinical Chemistry Data

Males Week 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-NEFOSE, T-6314) IN RATS

DOSE ppm MeFOS	GLU MG/DL	UN MG/DL	CREAT MG/DL	T PRO G/DL	ALB G/DL	GLOB G/DL	T BILI MG/DL	CHOL MG/DL	TRIG MG/DL	
0	MEAN S.D. N	103 5.7 10	11 1.9 10	.6 .07 10	6.9 .39 10	4.7 .23 10	2.2 .26 10	.2 .00 10	59 15.4 10	45 18.3 10
3	MEAN S.D. N	114 14.1 10	11 1.4 10	.6 .05 10	6.9 .23 10	4.7 .19 10	2.1 .14 10	.2 .00 10	58 12.1 10	55 18.8 10
30	MEAN S.D. N	106 15.6 10	12 1.7 10	.6 .03 10	6.6 .22 10	4.7 .14 10	2.0 * .19 10	.2 .03 10	54 10.6 10	40 10.4 10
100	MEAN S.D. N	106 17.5 10	13 * 1.6 10	.6 .07 10	6.7 .29 10	4.7 .21 10	2.0 * .18 10	.2 .00 10	44 14.1 10	25 * 9.9 10

000870

Table 12
Summary of Clinical Chemistry Data
Males Week 5
13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

DOSE ppm MeFOS	AST/SGOT IU/L	ALT/SGPT IU/L	GGT IU/L	CA MG/DL	I PHOS MG/DL	NA MMOL/L	K MMOL/L	CL MMOL/L
0	MEAN 103	32	0	10.9	9.5	147	6.2	105
	S.D. 18.9	3.6	.5	.47	.61	1.3	.39	2.3
	N 10	10	10	10	10	10	10	10
3	MEAN 96	33	0	10.8	9.8	147	6.2	105
	S.D. 16.1	4.6	.5	.19	.71	1.6	.42	1.6
	N 10	10	10	10	10	10	10	10
30	MEAN 101	32	0	10.5	9.0	148	6.1	106
	S.D. 18.3	3.8	.7	.30	.78	1.2	.52	1.2
	N 10	10	10	10	10	10	10	10
100	MEAN 97	40 *	0	10.6	9.0	147	6.1	105
	S.D. 12.2	4.5	.3	.42	1.02	2.5	.46	2.5
	N 10	10	10	10	10	10	10	10

000371

Table 12
Summary of Clinical Chemistry Data
Females Week 5
13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

DOSE ppm MeFOS	GLU MG/DL	UN MG/DL	CREAT MG/DL	T PRO G/DL	ALB G/DL	GLOB G/DL	T BILI MG/DL	CHOL MG/DL	TRIG MG/DL	
0	MEAN S.D. N	108 9.2 10	12 1.2 10	.7 .03 10	7.4 .31 10	5.3 .21 10	2.1 .24 10	.2 .05 10	68 11.9 10	30 8.1 10
3	MEAN S.D. N	108 8.9 10	12 2.1 10	.7 .04 10	7.4 .47 10	5.3 .39 10	2.1 .24 10	.2 .00 10	69 12.0 10	28 8.9 10
30	MEAN S.D. N	108 10.0 10	12 1.5 10	.7 .03 10	7.3 .34 10	5.3 .29 10	1.9 .19 10	.2 .03 10	64 9.7 10	28 6.9 10
100	MEAN S.D. N	95 * 12.0 10	14 1.5 10	.6 .07 10	7.2 .41 10	5.3 .28 10	1.9 .23 10	.2 .00 10	53 21.7 10	26 9.3 10

000872

Table 12
Summary of Clinical Chemistry Data

Females Week 5
13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDE
ETHANOL (N-MEFOSE, T-6314) IN RATS

DOSE ppm MeFOS	AST/SGOT IU/L	ALT/SGPT IU/L	GGT IU/L	CA MG/DL	I PHOS MG/DL	NA MMOL/L	K MMOL/L	CL MMOL/L
0	MEAN	99	28	0	10.8	7.9	146	105
	S.D.	15.3	4.8	.5	.52	.65	1.3	1.2
	N	10	10	10	10	10	10	10
3	MEAN	93	27	0	10.7	8.2	147	105
	S.D.	7.5	4.9	.7	.40	.77	1.6	1.4
	N	10	10	10	10	10	10	10
30	MEAN	93	28	1	10.8	7.7	146	104
	S.D.	14.0	3.5	1.1	.38	1.07	1.9	1.9
	N	10	10	10	10	10	10	10
100	MEAN	93	30	1	10.6	8.2	147	106
	S.D.	16.0	5.8	.8	.32	1.06	2.0	2.6
	N	10	10	10	10	10	10	10

000873

Table 12
Summary of Clinical Chemistry Data
Males Week 5
13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

DOSE ppm MeFOS	PCOAO IU/G	
0	12	MEAN
	3.6	S.D.
	5	N
3	13	MEAN
	2.1	S.D.
	5	N
30	8	MEAN
	4.3	S.D.
	5	N
100	9	MEAN
	1.9	S.D.
	5	N

000874

Table 12
Summary of Clinical Chemistry Data
Females Week 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

DOSE ppm MeFOS	PCOAO IU/G
-----	-----
0	MEAN 7 S.D. 2.5 N 5
3	MEAN 5 S.D. 2.9 N 5
30	MEAN 7 S.D. 3.3 N 5
100	MEAN 11 S.D. 2.7 N 5

000375

Table 13

Summary of Clinical Chemistry Data

Males Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

DOSE ppm MeFOS	GLU MG/DL	UN MG/DL	CREAT MG/DL	T PRO G/DL	ALB G/DL	GLOB G/DL	T BILI MG/DL	CHOL MG/DL	TRIG MG/DL
0	MEAN	117	13	.8	7.2	4.7	2.5	.2	69
	S.D.	8.2	2.1	.05	.34	.24	.21	.05	15.2
	N	10	10	10	10	10	10	10	10
3	MEAN	119	12	.7	7.0	4.7	2.3	.1	57
	S.D.	7.9	2.1	.05	.34	.20	.21	.05	17.4
	N	10	10	10	10	10	10	10	10
30	MEAN	113	14	.8	6.8	4.6	2.1 *	.2	49 *
	S.D.	9.7	2.0	.07	.57	.35	.30	.05	15.1
	N	10	10	10	10	10	10	10	10
100	MEAN	128 *	15 *	.7	7.2	5.2 *	2.0 *	.2	26 *
	S.D.	11.3	1.4	.05	.39	.29	.26	.05	12.1
	N	10	10	10	10	10	10	10	10

000376

Table 13

Summary of Clinical Chemistry Data

Males Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

DOSE ppm MeFOS	AST/SGOT IU/L	ALT/SGPT IU/L	GGT IU/L	CA MG/DL	I PHOS MG/DL	NA MMOL/L	K MMOL/L	CL MMOL/L
0	MEAN	89	38	2	10.7	6.6	150	6.1
	S.D.	18.4	5.1	1.3	.49	.65	2.5	.31
	N	10	10	10	10	10	10	10
3	MEAN	86	41	2	10.6	6.5	150	6.1
	S.D.	9.7	6.4	.7	.26	.29	2.2	.28
	N	10	10	10	10	10	10	10
30	MEAN	91	38	2	10.2 *	6.5	149	6.2
	S.D.	16.7	7.0	1.3	.40	.58	2.2	.36
	N	10	10	10	10	10	10	10
100	MEAN	91	60 *	2	10.4	6.9	150	6.4
	S.D.	14.3	21.6	.7	.30	.39	2.6	.48
	N	10	10	10	10	10	10	10

000377

Table 13
Summary of Clinical Chemistry Data
Females Week 14
13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

DOSE ppm MeFOS	GLU MG/DL	UN MG/DL	CREAT MG/DL	T PRO G/DL	ALB G/DL	GLOB G/DL	T BILI MG/DL	CHOL MG/DL	TRIG MG/DL	
0	MEAN S.D. N	119 10.6 10	12 2.0 10	.8 .07 10	7.9 .50 10	5.8 .41 10	2.1 .31 10	.2 .03 10	73 12.6 10	43 16.8 10
3	MEAN S.D. N	121 8.0 9	13 2.1 9	.8 .07 9	7.7 .53 9	5.6 .42 9	2.1 .24 9	.2 * .05 9	69 9.4 9	38 21.5 9
30	MEAN S.D. N	110 7.0 10	14 1.8 10	.8 .07 10	8.0 .42 10	5.8 .24 10	2.2 .28 10	.2 .03 10	70 9.9 10	36 12.7 10
100	MEAN S.D. N	110 9.2 10	17 * 2.1 10	.8 .06 10	8.2 .50 10	6.0 .36 10	2.1 .23 10	.2 .00 10	51 * 17.7 10	25 4.2 10

000378

Table 13
Summary of Clinical Chemistry Data
Females Week 14
13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

DOSE ppm MeFOS	AST/SGOT IU/L	ALT/SGPT IU/L	GGT IU/L	CA MG/DL	I PHOS MG/DL	NA MMOL/L	K MMOL/L	CL MMOL/L	
0	MEAN S.D. N	84 10.1 10	34 5.7 10	2 1.2 10	11.0 .32 10	4.8 .54 10	150 1.9 10	5.7 .29 10	111 2.7 10
3	MEAN S.D. N	107 32.6 9	43 20.7 9	3 1.2 9	10.8 .41 9	5.1 .93 9	150 3.6 9	6.1 .70 9	111 3.3 9
30	MEAN S.D. N	128 126.7 10	64 105.5 10	3 1.7 10	11.0 .26 10	5.6 .74 10	151 2.6 10	6.2 .50 10	109 2.6 10
100	MEAN S.D. N	80 9.3 10	36 6.1 10	3 1.0 10	10.9 .27 10	5.4 .90 10	150 2.7 10	5.7 .43 10	109 3.0 10

000379

Table 13
Summary of Clinical Chemistry Data
Males Week 14
13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

DOSE ppm MeFOS	PCOAO IU/G
-----	-----
0	MEAN 10 S.D. 2.3 N 15
3	MEAN 9 S.D. 2.7 N 15
30	MEAN 9 S.D. 2.0 N 15
100	MEAN 15 * S.D. 8.1 N 15

000330

Table 13
Summary of Clinical Chemistry Data
Females Week 14
13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

DOSE ppm MeFOS	PCOAO IU/G
0	MEAN 7 S.D. 2.5 N 15
3	MEAN 7 S.D. 2.9 N 14
30	MEAN 9 S.D. 2.8 N 15
100	MEAN 20 * S.D. 6.7 N 15

C00381

Table 14
Summary of Clinical Urinalysis Data

Males Week 14
13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

DOSE ppm MeFOS	U VOL ML	SP GR	U PH
0	9.9 7.15 10 10	1.036 .0179 10	6.9 .32 10
3	10.6 5.52 10 10	1.026 .0093 10	6.9 .39 10
30	13.8 13.29 10 10	1.026 .0162 10	6.8 .42 10
100	21.0 16.52 10 10	1.024 .0148 10	6.6 .24 10

000832

Table 14
Summary of Clinical Urinalysis Data
Females Week 14
13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

DOSE ppm MeFOS	U VOL ML	SP GR	U PH
0	MEAN 17.0 S.D. 14.13 N 10	1.016 .0112 10	6.7 .26 10
3	MEAN 11.4 S.D. 10.11 N 9	1.020 .0102 9	6.8 .44 9
30	MEAN 16.7 S.D. 14.62 N 10	1.016 .0084 10	6.8 .26 10
100	MEAN 15.7 S.D. 12.05 N 10	1.019 .0113 10	6.6 .21 10

000383

TABLE 15
Summary of Organ Weight Data
Week 5 Sacrifice

PAGE: 1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

TABLE INCLUDES:
SEX=ALL; GROUP=ALL; WEEKS=ALL
DEATH=1; SUBSET=T

LIVER

SEX	DOSE GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO-BODY WT (%)	SEX	DOSE GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO-BODY WT (%)

M 1					F 1				
NUMBER IN GROUP:					NUMBER IN GROUP:				
MEAN:					MEAN:				
STANDARD DEV:					STANDARD DEV:				

M 2					F 2				
NUMBER IN GROUP:					NUMBER IN GROUP:				
MEAN:					MEAN:				
STANDARD DEV:					STANDARD DEV:				

M 3					F 3				
NUMBER IN GROUP:					NUMBER IN GROUP:				
MEAN:					MEAN:				
STANDARD DEV:					STANDARD DEV:				

000834

TABLE 15
Summary of Organ Weight Data
Week 5 Sacrifice

PAGE: 2

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

TABLE INCLUDES:
SEX=ALL; GROUP=ALL; WEEKS=ALL
DEATH=1; SUBSET=T

LIVER

SEX	DOSE GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO- BODY WT (%)	SEX	DOSE GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO- BODY WT (%)
M	4				F	4			
NUMBER IN GROUP: 5 MEAN: 311.6 STANDARD DEV: 22.0					NUMBER IN GROUP: 5 MEAN: 188.1 STANDARD DEV: 9.5				
14.7069 1.1550					8.4256 0.4870				
4.7214 0.1919					4.4836 0.2373				

000885

TABLE 16
Summary of Organ Weight Data
Week 14 Sacrifice

PAGE: 1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

TABLE INCLUDES:
SEX=ALL; GROUP=ALL; WEEKS=ALL
DEATH=T; SUBSET=ALL

ADRENAL

SEX	DOSE	GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO- BODY WT (%)	ORGAN-TO- BRAIN WT RATIO	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO- BODY WT (%)	ORGAN-TO- BRAIN WT RATIO
F 1										
NUMBER IN GROUP:			15	15	15	15	15	15	15	15
MEAN:			505.3	0.0562	0.0112	0.0258	270.5	0.0709	0.0265	0.0365
STANDARD DEV:			35.7	0.0092	0.0020	0.0044	33.3	0.0123	0.0054	0.0062
F 2										
NUMBER IN GROUP:			15	15	15	15	14	14	14	14
MEAN:			482.1	0.0575	0.0120	0.0267	251.3	0.0675	0.0268	0.0343
STANDARD DEV:			27.7	0.0083	0.0020	0.0038	24.0	0.0114	0.0035	0.0057
F 3										
NUMBER IN GROUP:			15	15	15	15	15	15	15	15
MEAN:			460.1*	0.0551	0.0121	0.0254	253.4	0.0700	0.0278	0.0348
STANDARD DEV:			31.8	0.0088	0.0022	0.0040	17.8	0.0089	0.0040	0.0050

000886

TABLE 16
Summary of Organ Weight Data
Week 14 Sacrifice

PAGE: 2

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

TABLE INCLUDES:
SEX=ALL; GROUP=ALL; WEEKS=ALL
DEATH=T; SUBSET=ALL

		ADRENAL									
SEX	DOSE GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO- BODY WT (%)	ORGAN-TO- BRAIN WT RATIO	SEX	DOSE GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO- BODY WT (%)	ORGAN-TO- BRAIN WT RATIO
M		F									
4		4									
NUMBER IN GROUP:		15	15	15	15	NUMBER IN GROUP:		15	15	15	15
MEAN:		445.8*	0.0560	0.0126	0.0259	MEAN:		229.8*	0.0672	0.0293	0.0330
STANDARD DEV:		24.4	0.0073	0.0014	0.0039	STANDARD DEV:		22.7	0.0113	0.0047	0.0052

000837

TABLE 16
Summary of Organ Weight Data
Week 14 Sacrifice

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 3

TABLE INCLUDES:
SEX=ALL; GROUP=ALL; WEEKS=ALL
DEATH=T; SUBSET=ALL

BRAIN

SEX	DOSE	GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO- BODY WT (%)	ORGAN-TO- BRAIN WT RATIO	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO- BODY WT (%)	ORGAN-TO- BRAIN WT RATIO
F 1										
NUMBER IN GROUP:			15	15	15	15	15	15	15	15
MEAN:			505.3	2.1771	0.4326	1.0000	270.5	1.9434	0.7256	1.0000
STANDARD DEV:			35.7	0.0642	0.0283	0.0000	33.3	0.1049	0.0712	0.0000
F 2										
NUMBER IN GROUP:			15	15	15	15	14	14	14	14
MEAN:			482.1	2.1519	0.4480	1.0000	251.3	1.9678	0.7891*	1.0000
STANDARD DEV:			27.7	0.0910	0.0354	0.0000	24.0	0.0694	0.0733	0.0000
F 3										
NUMBER IN GROUP:			15	15	15	15	15	15	15	15
MEAN:			460.1*	2.1680	0.4731*	1.0000	253.4	2.0183	0.7995*	1.0000
STANDARD DEV:			31.8	0.0768	0.0323	0.0000	17.8	0.0826	0.0538	0.0000

000838

TABLE 16
Summary of Organ Weight Data
Week 14 Sacrifice

PAGE: 4

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

TABLE INCLUDES:
SEX=ALL; GROUP=ALL; WEEKS=ALL
DEATH=T; SUBSET=ALL

		BRAIN									
SEX	DOSE GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO- BODY WT (%)	ORGAN-TO- BRAIN WT RATIO	SEX	DOSE GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO- BODY WT (%)	ORGAN-TO- BRAIN WT RATIO
M	4					F	4				
NUMBER IN GROUP:		15	15	15	15	NUMBER IN GROUP:		15	15	15	15
MEAN:		445.8*	2.1678	0.4878*	1.0000	MEAN:		229.8*	2.0372*	0.8910*	1.0000
STANDARD DEV:		24.4	0.0884	0.0360	0.0000	STANDARD DEV:		22.7	0.1113	0.0599	0.0000

600389

TABLE 16
Summary of Organ Weight Data
Week 14 Sacrifice13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 5

TABLE INCLUDES:
SEX=ALL; GROUP=ALL; WEEKS=ALL
DEATH=T; SUBSET=ALL

KIDNEY

SEX	DOSE GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO-BODY WT (%)	ORGAN-TO-BRAIN WT RATIO	SEX	DOSE GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO-BODY WT (%)	ORGAN-TO-BRAIN WT RATIO
F 1											
NUMBER IN GROUP:		15	15	15	15	NUMBER IN GROUP:		15	15	15	15
MEAN:		270.5	2.1241	0.7893	1.0931	MEAN:		270.5	2.1241	0.7893	1.0931
STANDARD DEV:		33.3	0.2283	0.0711	0.1021	STANDARD DEV:		33.3	0.2283	0.0711	0.1021
F 2											
NUMBER IN GROUP:		14	14	14	14	NUMBER IN GROUP:		14	14	14	14
MEAN:		251.3	2.1259	0.8469	1.0809	MEAN:		251.3	2.1259	0.8469	1.0809
STANDARD DEV:		24.0	0.2488	0.0664	0.1239	STANDARD DEV:		24.0	0.2488	0.0664	0.1239
F 3											
NUMBER IN GROUP:		15	15	15	15	NUMBER IN GROUP:		15	15	15	15
MEAN:		253.4	2.2559	0.8936*	1.1190	MEAN:		253.4	2.2559	0.8936*	1.1190
STANDARD DEV:		17.8	0.1509	0.0771	0.0819	STANDARD DEV:		17.8	0.1509	0.0771	0.0819

SEX	DOSE GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO-BODY WT (%)	ORGAN-TO-BRAIN WT RATIO	SEX	DOSE GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO-BODY WT (%)	ORGAN-TO-BRAIN WT RATIO
M 1											
NUMBER IN GROUP:		15	15	15	15	NUMBER IN GROUP:		15	15	15	15
MEAN:		505.3	3.8199	0.7549	1.7543	MEAN:		270.5	2.1241	0.7893	1.0931
STANDARD DEV:		35.7	0.4337	0.0520	0.1893	STANDARD DEV:		33.3	0.2283	0.0711	0.1021
M 2											
NUMBER IN GROUP:		15	15	15	15	NUMBER IN GROUP:		14	14	14	14
MEAN:		482.1	3.7300	0.7744	1.7358	MEAN:		251.3	2.1259	0.8469	1.0809
STANDARD DEV:		27.7	0.3858	0.0762	0.1924	STANDARD DEV:		24.0	0.2488	0.0664	0.1239
M 3											
NUMBER IN GROUP:		15	15	15	15	NUMBER IN GROUP:		15	15	15	15
MEAN:		460.1*	3.8085	0.8308*	1.7609	MEAN:		253.4	2.2559	0.8936*	1.1190
STANDARD DEV:		31.8	0.3063	0.0819	0.1761	STANDARD DEV:		17.8	0.1509	0.0771	0.0819

000390

TABLE 16
Summary of Organ Weight Data
Week 14 Sacrifice

PAGE: 6

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

TABLE INCLUDES:
SEX=ALL; GROUP=ALL; WEEKS=ALL
DEATH=T; SUBSET=ALL

KIDNEY

SEX	DOSE GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO- BODY WT (%)	ORGAN-TO- BRAIN WT RATIO	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO- BODY WT (%)	ORGAN-TO- BRAIN WT RATIO
M	4								
		15	15	15	15	15	15	15	15
		445.8*	4.1850*	0.9389*	1.9357*	229.8*	2.1983	0.9576*	1.0773
		24.4	0.4155	0.0787	0.2325	22.7	0.2606	0.0797	0.0948
		NUMBER IN GROUP:	NUMBER IN GROUP:	NUMBER IN GROUP:	NUMBER IN GROUP:	NUMBER IN GROUP:	NUMBER IN GROUP:	NUMBER IN GROUP:	NUMBER IN GROUP:
		MEAN:	MEAN:	MEAN:	MEAN:	MEAN:	MEAN:	MEAN:	MEAN:
		STANDARD DEV:	STANDARD DEV:	STANDARD DEV:	STANDARD DEV:	STANDARD DEV:	STANDARD DEV:	STANDARD DEV:	STANDARD DEV:

000391

TABLE 16
Summary of Organ Weight Data
Week 14 Sacrifice

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 7

TABLE INCLUDES: SEX=ALL; GROUP=ALL; WEEKS=ALL DEATH=T; SUBSET=ALL											
LIVER											
SEX	DOSE GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO-BODY WT (%)	ORGAN-TO-BRAIN WT RATIO	SEX	DOSE GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO-BODY WT (%)	ORGAN-TO-BRAIN WT RATIO
M 1						F 1					
NUMBER IN GROUP:		15	15	15	15	NUMBER IN GROUP:		15	15	15	15
MEAN:		505.3	14.5017	2.8663	6.6577	MEAN:		270.5	8.0512	2.9775	4.1397
STANDARD DEV:		35.7	1.6903	0.2214	0.7163	STANDARD DEV:		33.3	1.3405	0.3423	0.6146
M 2						F 2					
NUMBER IN GROUP:		15	15	15	15	NUMBER IN GROUP:		14	14	14	14
MEAN:		482.1	14.3465	2.9803	6.6774	MEAN:		251.3	7.8673	3.1298	3.9983
STANDARD DEV:		27.7	1.3494	0.2896	0.6795	STANDARD DEV:		24.0	0.9141	0.2035	0.4464
M 3						F 3					
NUMBER IN GROUP:		15	15	15	15	NUMBER IN GROUP:		15	15	15	15
MEAN:		460.1*	15.9014	3.4572*	7.3423*	MEAN:		253.4	9.2512*	3.6610*	4.5878
STANDARD DEV:		31.8	1.5170	0.2312	0.7472	STANDARD DEV:		17.8	0.7652	0.3281	0.3853

000882

TABLE 16
Summary of Organ Weight Data
Week 14 Sacrifice

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

TABLE INCLUDES:
SEX=ALL; GROUP=ALL; WEEKS=ALL
DEATH=T; SUBSET=ALL

LIVER

SEX	DOSE	GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO- BODY WT (%)	ORGAN-TO- BRAIN WT RATIO	SEX	DOSE	GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO- BODY WT (%)	ORGAN-TO- BRAIN WT RATIO
M		4					F		4				
NUMBER IN GROUP:			15	15	15	15	NUMBER IN GROUP:			15	15	15	15
MEAN:			445.8*	24.5652*	5.5083*	11.3342*	MEAN:			229.8*	12.2021*	5.3304*	5.9827*
STANDARD DEV:			24.4	2.7276	0.4975	1.1828	STANDARD DEV:			22.7	1.4877	0.6283	0.5773

000393

TABLE 16
Summary of Organ Weight Data
Week 14 Sacrifice

PAGE: 9

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

TABLE INCLUDES: SEX=ALL; GROUP=ALL; WEEKS=ALL DEATH=T; SUBSET=ALL									
OVARY									
SEX	DOSE	TERMINAL	ORGAN	ORGAN-TO-	ORGAN-TO-	TERMINAL	ORGAN	ORGAN-TO-	ORGAN-TO-
GROUP	GROUP	BODY WT	WEIGHT	BODY WT	BODY WT	BODY WT	WEIGHT	BODY WT	BODY WT
		(g)	(g)	RATIO		(g)	(g)	(%)	RATIO
F 1									
NUMBER IN GROUP:		15	15	15	15	15	15	15	15
MEAN:		505.3				270.5	0.1280	0.0480	0.0658
STANDARD DEV:		35.7				33.3	0.0205	0.0097	0.0096
F 2									
NUMBER IN GROUP:		15	14	14	14	14	14	14	14
MEAN:		482.1				251.3	0.1225	0.0489	0.0624
STANDARD DEV:		27.7				24.0	0.0227	0.0081	0.0117
F 3									
NUMBER IN GROUP:		15	15	15	15	15	15	15	15
MEAN:		460.1*				253.4	0.1221	0.0486	0.0605
STANDARD DEV:		31.8				17.8	0.0215	0.0109	0.0109

000894

TABLE 16
Summary of Organ Weight Data
Week 14 Sacrifice

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 10

TABLE INCLUDES:
SEX=ALL; GROUP=ALL; WEEKS=ALL
DEATH=T; SUBSET=ALL

OVARY

SEX	DOSE GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO- BODY WT (%)	ORGAN-TO- BRAIN WT RATIO	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO- BODY WT (%)	ORGAN-TO- BRAIN WT RATIO
M	4								
NUMBER IN GROUP:		15	15	15	15	15	15	15	15
MEAN:		445.8*				229.8*	0.1206	0.0527	0.0592
STANDARD DEV:		24.4				22.7	0.0238	0.0099	0.0107

000895

TABLE 16
Summary of Organ Weight Data
Week 14 Sacrifice

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 11

TABLE INCLUDES:
SEX=ALL; GROUP=ALL; WEEKS=ALL
DEATH=T; SUBSET=ALL

SPLEEN

SEX	DOSE GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO-BODY WT (%)	ORGAN-TO-BRAIN WT RATIO	SEX	DOSE GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO-BODY WT (%)	ORGAN-TO-BRAIN WT RATIO
F 1											
NUMBER IN GROUP:		15	15	15	15	NUMBER IN GROUP:		15	15	15	15
MEAN:		505.3	0.8041	0.1592	0.3691	MEAN:		270.5	0.5153	0.1907	0.2649
STANDARD DEV:		35.7	0.0904	0.0149	0.0371	STANDARD DEV:		33.3	0.1324	0.0408	0.0644
F 2											
NUMBER IN GROUP:		15	15	15	15	NUMBER IN GROUP:		14	14	14	14
MEAN:		482.1	0.7666	0.1597	0.3558	MEAN:		251.3	0.5235	0.2090	0.2658
STANDARD DEV:		27.7	0.1431	0.0332	0.0604	STANDARD DEV:		24.0	0.0643	0.0239	0.0290
F 3											
NUMBER IN GROUP:		15	15	15	15	NUMBER IN GROUP:		15	15	15	15
MEAN:		460.1*	0.7341	0.1602	0.3390	MEAN:		253.4	0.5166	0.2049	0.2568
STANDARD DEV:		31.8	0.0802	0.0203	0.0390	STANDARD DEV:		17.8	0.0506	0.0251	0.0306

000336

TABLE 16
Summary of Organ Weight Data
Week 14 Sacrifice

PAGE: 12

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

TABLE INCLUDES:
SEX=ALL; GROUP=ALL; WEEKS=ALL
DEATH=T; SUBSET=ALL

SPLEEN

SEX	DOSE	GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO- BODY WT (%)	ORGAN-TO- BRAIN WT RATIO	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO- BODY WT (%)	ORGAN-TO- BRAIN WT RATIO
M			15	15	15	15	15	15	15	15
NUMBER IN GROUP:			445.8*	0.6707*	0.1501	0.3101*	229.8*	0.4675	0.2037	0.2294
MEAN:			24.4	0.1266	0.0251	0.0620	22.7	0.0678	0.0232	0.0296
STANDARD DEV:										
F			4							
NUMBER IN GROUP:										
MEAN:										
STANDARD DEV:										

600897

TABLE 16
Summary of Organ Weight Data
Week 14 Sacrifice

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 13

TABLE INCLUDES:
SEX=ALL; GROUP=ALL; WEEKS=ALL
DEATH=T; SUBSET=ALL

TESTIS

SEX	DOSE GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO-BODY WT (%)	ORGAN-TO-BRAIN WT RATIO	SEX	DOSE GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO-BODY WT (%)	ORGAN-TO-BRAIN WT RATIO
M 1						F 1					
NUMBER IN GROUP:		15	15	15	15	NUMBER IN GROUP:		15			
MEAN:		505.3	3.3907	0.6733	1.5579	MEAN:		270.5			
STANDARD DEV:		35.7	0.2211	0.0559	0.0976	STANDARD DEV:		33.3			
M 2						F 2					
NUMBER IN GROUP:		15	15	15	15	NUMBER IN GROUP:		14			
MEAN:		482.1	3.3852	0.7028	1.5752	MEAN:		251.3			
STANDARD DEV:		27.7	0.3024	0.0565	0.1496	STANDARD DEV:		24.0			
M 3						F 3					
NUMBER IN GROUP:		15	15	15	15	NUMBER IN GROUP:		15			
MEAN:		460.1*	3.4481	0.7535*	1.5922	MEAN:		253.4			
STANDARD DEV:		31.8	0.3067	0.0874	0.1509	STANDARD DEV:		17.8			

000338

TABLE 16
Summary of Organ Weight Data
Week 14 Sacrifice

PAGE: 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

TABLE INCLUDES:
SEX=ALL; GROUP=ALL; WEEKS=ALL
DEATH=T; SUBSET=ALL

TESTIS

SEX	DOSE GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO- BODY WT (%)	ORGAN-TO- BRAIN WT RATIO	SEX	DOSE GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO- BODY WT (%)	ORGAN-TO- BRAIN WT RATIO
M		F									
4		4									
NUMBER IN GROUP:		15	15	15	15	NUMBER IN GROUP:		15	15	15	15
MEAN:		445.8*	3.4789	0.7824*	1.6042	MEAN:		229.8	229.8	22.7	22.7
STANDARD DEV:		24.4	0.2974	0.0756	0.1126	STANDARD DEV:		22.7	22.7		

000399

TABLE 16
Summary of Organ Weight Data
Week 14 Sacrifice

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 15

TABLE INCLUDES:
SEX=ALL; GROUP=ALL; WEEKS=ALL
DEATH=T; SUBSET=ALL

THYMUS

SEX	DOSE GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO-BODY WT (%)	ORGAN-TO-BRAIN WT RATIO	SEX	DOSE GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO-BODY WT (%)	ORGAN-TO-BRAIN WT RATIO

M 1						F 1					
NUMBER IN GROUP:						NUMBER IN GROUP:					
MEAN:						MEAN:					
STANDARD DEV:						STANDARD DEV:					
15 15 505.3 0.3454 0.0684 0.1586						15 15 270.5 0.2814 0.1053 0.1451					
35.7 0.0847 0.0160 0.0383						33.3 0.0627 0.0259 0.0330					

M 2						F 2					
NUMBER IN GROUP:						NUMBER IN GROUP:					
MEAN:						MEAN:					
STANDARD DEV:						STANDARD DEV:					
15 15 482.1 0.3117 0.0651 0.1451						14 14 251.3 0.2951 0.1176 0.1499					
27.7 0.0630 0.0152 0.0301						24.0 0.0491 0.0179 0.0236					

M 3						F 3					
NUMBER IN GROUP:						NUMBER IN GROUP:					
MEAN:						MEAN:					
STANDARD DEV:						STANDARD DEV:					
15 15 460.1* 0.2725* 0.0592 0.1260*						15 15 253.4 0.2978 0.1174 0.1476					
31.8 0.0678 0.0147 0.0324						17.8 0.0627 0.0225 0.0310					

000800

TABLE 16
Summary of Organ Weight Data
Week 14 Sacrifice

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

TABLE INCLUDES:													
SEX=ALL; GROUP=ALL; WEEKS=ALL													
DEATH=T; SUBSET=ALL													
THYMUS													
SEX	DOSE	GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO- BODY WT (%)	ORGAN-TO- BRAIN WT RATIO	SEX	DOSE	GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO- BODY WT (%)	ORGAN-TO- BRAIN WT RATIO
M							F			4			
NUMBER IN GROUP:			15	15	15	15	NUMBER IN GROUP:			15	15	15	15
MEAN:			445.8*	0.2610*	0.0586	0.1206*	MEAN:			229.8*	0.2654	0.1152	0.1300
STANDARD DEV:			24.4	0.0515	0.0115	0.0246	STANDARD DEV:			22.7	0.0525	0.0190	0.0229

C00901

TABLE 16
Summary of Organ Weight Data
Week 14 Sacrifice

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 17

TABLE INCLUDES: SEX=ALL; GROUP=ALL; WEEKS=ALL DEATH=T; SUBSET=ALL											
THYROID/PARATHYR											
SEX	DOSE GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO-BODY WT (%)	ORGAN-TO-BRAIN WT RATIO	SEX	DOSE GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO-BODY WT (%)	ORGAN-TO-BRAIN WT RATIO
F 1											
NUMBER IN GROUP:		15	15	15	15	NUMBER IN GROUP:		15	15	15	15
MEAN:		505.3	0.0255	0.0051	0.0117	MEAN:		270.5	0.0203	0.0075	0.0104
STANDARD DEV:		35.7	0.0063	0.0013	0.0029	STANDARD DEV:		33.3	0.0062	0.0024	0.0029
F 2											
NUMBER IN GROUP:		15	15	15	15	NUMBER IN GROUP:		14	14	14	14
MEAN:		482.1	0.0254	0.0053	0.0118	MEAN:		251.3	0.0180	0.0072	0.0092
STANDARD DEV:		27.7	0.0078	0.0017	0.0034	STANDARD DEV:		24.0	0.0066	0.0027	0.0034
F 3											
NUMBER IN GROUP:		15	15	15	15	NUMBER IN GROUP:		15	15	15	15
MEAN:		460.1*	0.0244	0.0053	0.0112	MEAN:		253.4	0.0190	0.0075	0.0094
STANDARD DEV:		31.8	0.0048	0.0010	0.0019	STANDARD DEV:		17.8	0.0046	0.0019	0.0024

000902

TABLE 16
Summary of Organ Weight Data
Week 14 Sacrifice

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

TABLE INCLUDES:
SEX=ALL; GROUP=ALL; WEEKS=ALL
DEATH=T; SUBSET=ALL

THYROID/PARATHYR

SEX	DOSE GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO- BODY WT (%)	ORGAN-TO- BRAIN WT RATIO	SEX	DOSE GROUP	TERMINAL BODY WT (g)	ORGAN WEIGHT (g)	ORGAN-TO- BODY WT (%)	ORGAN-TO- BRAIN WT RATIO
M		4				F		4			
NUMBER IN GROUP:		15	15	15	15	NUMBER IN GROUP:		15	15	15	15
MEAN:		445.8*	0.0252	0.0057	0.0117	MEAN:		229.8*	0.0203	0.0088	0.0100
STANDARD DEV:		24.4	0.0033	0.0009	0.0017	STANDARD DEV:		22.7	0.0052	0.0020	0.0024

000903

TABLE 17

Incidence of Macroscopic Observations

PAGE: 1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

TABLE INCLUDES:													
SEX=ALL; GROUP=ALL; WEEKS=ALL													
DEATH=ALL; SUBSET=ALL													
ORGAN AND KEYWORD(S) OR PHRASE													
SEX:													
GROUP: -1- -2- -3- -4- -1- -2- -3- -4-													
NUMBER: 20 20 20 20 20 20 20 20 20 20 20 20 20 20													
NUMBER EXAMINED: 20 20 20 20 20 20 20 20 20 20 20 20 20 20													
BONE MARROW SMEAR TAKEN													
MALOCCLUSION													
NO MACROSCOPIC LESIONS													
TAIL - BENT/MISSING													
STAINS-PERINEUM/PERIANAL													
LIVER (LI)													
NUMBER EXAMINED: 20 20 20 20 20 20 20 20 20 20 20 20 20 20													
NOT REMARKABLE: 20 20 20 20 16 20 20 20 20 20 19													
LIGHT FOCUS (I)/AREA (S)													
LARGE													
DIFFUSELY DARK													
KIDNEY (KD)													
NUMBER EXAMINED: 20 20 20 20 20 20 20 20 20 20 20 20 20 20													
NOT REMARKABLE: 19 19 20 18 20 20 20 20 20 20 20 20 20 20													
LARGE PELVIS(ES)													
GRANULAR MATERIAL													
SPLEEN (SP)													
NUMBER EXAMINED: 20 20 20 20 20 20 20 20 20 20 20 20 20 20													
NOT REMARKABLE: 20 20 20 20 19 20 20 20 20 20 20 20 20 20													
ADHESION (S)													

000904

TABLE 17
Incidence of Macroscopic Observations

PAGE: 2

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

		--- N U M B E R - O F - A N I M A L S - A F F E C T E D ---											
TABLE INCLUDES:		SEX:				MALE				FEMALE			
SEX=ALL; GROUP=ALL; WEEKS=ALL DEATH=ALL; SUBSET=ALL		GROUP: -1- -2- -3- -4-				-1- -2- -3- -4-				-1- -2- -3- -4-			
ORGAN AND KEYWORD(S) OR PHRASE		NUMBER:				20 20 20 20				20 20 20 20			
		NUMBER EXAMINED:				20 20 20 20				20 20 20 20			
		NOT REMARKABLE:				17 18 16 18				17 19 19 20			
THYMUS (TH)		2 1 3 1				0 0 0 0				0 0 0 0			
RED FOCUS(I)/AREA(S)		0 1 1 1				0 0 0 0				0 0 0 0			
DIFUSELY RED		1 0 0 0				0 0 0 0				0 0 0 0			
MOTTLED													
PITUITARY (PI)		20 20 20 20				20 20 20 20				20 20 20 20			
		NOT REMARKABLE:				20 20 20 20				20 20 20 20			
LARGE		0 0 0 0				0 0 0 0				0 0 0 0			
STOMACH, GL (ST)		20 20 20 20				20 20 20 20				20 20 20 20			
		NOT REMARKABLE:				20 18 20 18				20 20 19 17			
DARK FOCUS(I)/AREA(S)		0 1 0 2				0 0 0 0				0 0 0 0			
LIGHT FOCUS(I)/AREA(S)		0 1 0 0				0 0 0 0				0 0 0 0			
LN, MANDIBULAR (MN)		20 20 20 20				20 20 20 20				20 20 20 20			
		NOT REMARKABLE:				16 18 15 16				20 20 17 20			
DIFUSELY RED		3 2 4 3				0 0 0 0				0 0 0 0			
MOTTLED		1 0 1 1				0 0 0 0				0 0 0 0			

000905

TABLE 17
Incidence of Macroscopic Observations

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 3

TABLE INCLUDES:		--- N U M B E R - O F - A N I M A L S - A F F E C T E D ---											
SEX=ALL; GROUP=ALL; WEEKS=ALL		SEX: ---MALE---						---FEMALE---					
DEATH=ALL; SUBSET=ALL													
ORGAN AND KEYWORD(S) OR PHRASE													
UTERUS (UT)	NUMBER EXAMINED:	0	0	0	0	0	0	20	20	20	20	20	20
	NOT REMARKABLE:	0	0	0	0	0	0	19	17	19	19	19	19
LUMEN FILLED WITH FLUID													
OVARY (OV)	NUMBER EXAMINED:	0	0	0	0	0	0	1	3	1	1	1	1
	NOT REMARKABLE:	0	0	0	0	0	0	20	20	20	20	19	18
DIFFUSELY RED													
TAIL (TI)	NUMBER EXAMINED:	20	20	20	20	20	20	20	20	20	20	20	20
	NOT REMARKABLE:	20	20	20	20	20	20	19	20	19	20	20	20
LACERATION(S)													
** END OF LIST **		0	0	0	0	0	0	0	1	0	0	0	0

000906

TABLE 18

Incidence of Microscopic Observations

PAGE: 1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

--- N U M B E R - O F - A N I M A L S - A F F E C T E D ---												
TABLE INCLUDES:												
SEX=ALL; GROUP=ALL; WEEKS=ALL												
DEATH=ALL; FIND=ALL; SUBSET=ALL												
ORGAN AND FINDING DESCRIPTION	SEX:		---MALE---				---FEMALE---					
	GROUP:	-1-	-2-	-3-	-4-	-1-	-2-	-3-	-4-			
	NUMBER:	20	20	20	20	20	20	20	20	20	20	20
** TOP OF LIST **												
BONE, FEMUR (FE)	NUMBER EXAMINED:	15	0	0	15	15	1	0	15	1	0	15
	NOT REMARKABLE:	15	0	0	15	15	1	0	15	1	0	15
MARROW, FEMUR (FM)	NUMBER EXAMINED:	15	0	0	15	15	1	0	15	1	0	15
	NOT REMARKABLE:	15	0	0	15	15	1	0	15	1	0	15
MARROW, STERNUM (SE)	NUMBER EXAMINED:	15	0	0	15	15	1	0	15	1	0	15
	NOT REMARKABLE:	15	0	0	15	15	1	0	15	1	0	15
BONE, STERNUM (SB)	NUMBER EXAMINED:	15	0	0	15	15	1	0	15	1	0	15
	NOT REMARKABLE:	15	0	0	15	15	1	0	15	1	0	15
EYE (EY)	NUMBER EXAMINED:	15	0	0	15	15	1	0	15	1	0	15
	NOT REMARKABLE:	15	0	0	15	15	1	0	15	1	0	15
BRAIN (BR)	NUMBER EXAMINED:	15	0	0	15	15	1	0	15	1	0	15
	NOT REMARKABLE:	15	0	0	15	15	1	0	15	1	0	15
--INFILTRATE, LYMPHOHISTIOCYTIC												
SPINAL CORD (SC)	NUMBER EXAMINED:	15	0	0	15	15	1	0	15	1	0	15
	NOT REMARKABLE:	15	0	0	15	15	1	0	15	1	0	15

000907

TABLE 18

Incidence of Microscopic Observations

PAGE: 2

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

TABLE INCLUDES:		--- N U M B E R - O F - A N I M A L S - A F F E C T E D ---											
SEX=ALL; GROUP=ALL; WEEKS=ALL		SEX:		MALE		FEMALE							
DEATH=ALL; FIND=ALL; SUBSET=ALL		GROUP: -1- -2- -3- -4- -1- -2- -3- -4-		20 20 20 20 20 20 20 20		20 20 20 20 20 20 20 20							
ORGAN AND FINDING DESCRIPTION		NUMBER:		20 20 20 20 20 20 20 20		20 20 20 20 20 20 20 20							
LUNG (LU)		NUMBER EXAMINED:		15 0 0 0 15 15 15 15		1 0 0 0 1 0 1 0							
		NOT REMARKABLE:		6 0 0 0 7 12 1 0 12									
--HEMORRHAGE				1 0 0 0 0 0 0 0		0 0 0 0 0 0 0 0							
--INFILTRATE, MACROPHAGE, ALVEOLAR				4 0 0 0 5 1 0 0		0 0 0 0 0 0 0 0							
--INFLAMMATION, SUBACUTE				1 0 0 0 0 0 0 0		0 0 0 0 0 0 0 0							
--MINERALIZATION, VASCULAR				5 0 0 0 3 2 0 0		0 0 0 0 0 0 0 0							
TRACHEA (TR)		NUMBER EXAMINED:		15 0 0 0 15 15 15 15		1 0 1 0 1 0 1 0							
		NOT REMARKABLE:		15 0 0 0 15 15 15 15									
ESOPHAGUS (ES)		NUMBER EXAMINED:		15 0 0 0 15 15 15 15		1 0 1 0 1 0 1 0							
		NOT REMARKABLE:		15 0 0 0 15 15 15 15									
LIVER (LI)		NUMBER EXAMINED:		15 15 15 15 15 15 15 15		15 15 15 15 15 15 15 15							
		NOT REMARKABLE:		5 0 1 0 5 4 2 0									
--FIBROSIS, CAPSULAR				0 0 1 0 1 0 0 0		0 0 0 0 0 0 0 0							
--HYPERTROPHY, HEPATOCELLULAR, DIFFUSE				0 0 0 0 0 0 0 0		0 0 0 0 0 0 0 0							
--HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR				0 0 14 15 0 0 0 0		0 0 0 0 0 0 0 0							
--INFILTRATE, LYMPHOHISTIOCYTIC				10 15 13 10 10 11 12 9		0 0 0 0 0 0 0 0							
--NECROSIS, COAGULATIVE				0 0 0 0 3 0 0 0		0 0 0 0 0 0 0 0							
--NECROSIS, INDIVIDUAL HEPATOCYTE				6 9 5 6 3 6 4 5		0 0 0 0 0 0 0 0							
--PIGMENT, HEPATOCELLULAR				0 0 0 0 2 0 0 0		0 0 0 0 0 0 0 0							
--VACUOLATION, HEPATOCELLULAR, CENTRILLOBULAR TO MIDZONAL				0 0 0 0 2 0 0 0		0 0 0 0 0 0 0 0							
--VACUOLATION, HEPATOCELLULAR, DIFFUSE				0 0 0 0 2 2 0 0		0 0 0 0 0 0 0 0							

** CONTINUED ON NEXT PAGE **

000908

TABLE 18

Incidence of Microscopic Observations

PAGE: 3

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

--- N U M B E R - O F - A N I M A L S - A F F E C T E D ---									
TABLE INCLUDES:									
SEX=ALL; GROUP=ALL; WEEKS=ALL									
DEATH=ALL; FIND=ALL; SUBSET=ALL									
SEX: -----MALE-----FEMALE-----									
GROUP: -1- -2- -3- -4- -1- -2- -3- -4-									
NUMBER: 20 20 20 20 20 20 20 20 20									
ORGAN AND FINDING DESCRIPTION									
** FROM PREVIOUS PAGE **									
LIVER (LI)	NUMBER EXAMINED:	15	15	15	15	15	15	15	15
	NOT REMARKABLE:	5	0	1	0	5	4	2	0
--VACUOLATION, HEPATOCELLULAR, PERIportal									
KIDNEY (KD)	NUMBER EXAMINED:	15	1	0	15	15	1	0	15
	NOT REMARKABLE:	11	0	0	12	14	1	0	12
--BASOPHILIC TUBULES, CORTEX									
--DILATATION, PELVIC		2	0	0	0	0	0	0	0
--INFLAMMATION, CHRONIC		1	1	0	2	0	0	0	0
--MINERALIZATION, TUBULAR		0	0	0	0	0	0	0	1
--THICKENING, BASEMENT MEMBRANE		1	0	0	1	1	0	0	2
		1	0	0	0	0	0	0	0
HEART (HT)	NUMBER EXAMINED:	15	0	0	15	15	1	0	15
	NOT REMARKABLE:	12	0	0	12	15	1	0	14
--DEGENERATION/NECROSIS									
MUSCLE, SKELETAL (SM)		3	0	0	3	0	0	0	1
	NUMBER EXAMINED:	15	0	0	15	15	1	0	15
	NOT REMARKABLE:	15	0	0	15	15	1	0	15
NERVE, SCIATIC (SN)	NUMBER EXAMINED:	15	0	0	15	15	1	0	15
	NOT REMARKABLE:	15	0	0	15	15	1	0	15

TABLE 18
Incidence of Microscopic Observations

PAGE: 4

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

TABLE INCLUDES: SEX=ALL; GROUP=ALL; WEEKS=ALL DEATH=ALL; FIND=ALL; SUBSET=ALL	--- N U M B E R - O F - A N I M A L S - A F F E C T E D ---											
	SEX: ---MALE---						---FEMALE---					
ORGAN AND FINDING DESCRIPTION	GROUP: -1- -2- -3- -4- -1- -2- -3- -4-											
	NUMBER: 20 20 20 20 20 20											
SPLEEN (SP)	NUMBER EXAMINED: 15	0	0	15	15	1	0	15	1	0	15	
	NOT REMARKABLE: 15	0	0	15	14	1	0	15				
--INFLAMMATION, CHRONIC, CAPSULAR	0	0	0	0	1	0	0	0				
THYMUS (TH)	NUMBER EXAMINED: 15	2	4	15	15	2	1	15				
	NOT REMARKABLE: 9	0	0	6	11	1	0	10				
--HEMORRHAGE	6	2	4	9	4	1	1	4				
--INVOLUTION	0	0	1	1	1	1	0	2				
LN, MESENTERIC (MS)	NUMBER EXAMINED: 15	0	0	15	15	1	0	15				
	NOT REMARKABLE: 15	0	0	13	15	1	0	15				
--HEMORRHAGE	0	0	0	2	0	0	0	0				
THYROID (TY)	NUMBER EXAMINED: 15	0	0	15	15	1	0	15				
	NOT REMARKABLE: 4	0	0	8	6	1	0	9				
--CYST, ULTIMOBRANCHIAL	9	0	0	7	9	0	0	6				
--ECTOPIC THYMUS	1	0	0	0	0	0	0	0				
--INFILTRATE, LYMPHOHISTIOCYTIC	1	0	0	0	0	0	0	0				
--ONE EXAMINED	1	0	0	0	0	0	0	1				
PARATHYROID (PT)	NUMBER EXAMINED: 15	0	0	15	15	1	0	15				
	NOT REMARKABLE: 13	0	0	15	15	1	0	14				
--CYST	1	0	0	0	0	0	0	0				

** CONTINUED ON NEXT PAGE **

C00910

TABLE 18

Incidence of Microscopic Observations

PAGE: 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

--- N U M B E R - O F - A N I M A L S - A F F E C T E D ---												
SEX: ---MALE---FEMALE---												
GROUP: -1- -2- -3- -4- -1- -2- -3- -4-												
NUMBER: 20 20 20 20 20 20 20 20												
ORGAN AND FINDING DESCRIPTION												
** FROM PREVIOUS PAGE **												
PARATHYROID (PT)	15	0	0	15	15	1	0	15	1	0	15	14
	NOT EXAMINED:	13	0	0	15	15	1	0	14	0	15	14
--ONE EXAMINED												
ADRENAL, CORTEX (AC)	15	0	0	15	15	1	0	15	1	0	15	15
	NOT EXAMINED:	15	0	0	14	15	1	0	15	1	15	15
--INFILTRATE, LYMPHOHISTIOCYTIC												
ADRENAL, MEDULLA (MA)	0	0	0	1	0	0	0	0	0	0	0	0
	NUMBER EXAMINED:	15	0	0	15	15	1	0	15	1	0	15
	NOT EXAMINED:	15	0	0	15	15	1	0	15	1	0	15
AORTA (AO)												
	NUMBER EXAMINED:	15	0	0	15	15	1	0	15	1	0	15
	NOT EXAMINED:	15	0	0	15	15	1	0	15	1	0	15
PITUITARY (PI)												
	NUMBER EXAMINED:	15	0	0	15	15	1	0	15	1	0	15
	NOT EXAMINED:	14	0	0	14	14	1	0	15	1	0	15
--CYST												
--HYPERPLASIA												
	1	0	0	1	0	0	0	0	0	0	0	0
	0	0	0	0	0	1	0	0	0	0	0	0
STOMACH, GL (ST)												
	NUMBER EXAMINED:	14	2	15	15	15	1	15	15	1	15	15
	NOT EXAMINED:	14	1	15	13	15	1	15	15	1	15	12
--EROSION												
	0	1	0	2	0	0	0	0	0	0	0	3

TABLE 18

Incidence of Microscopic Observations

PAGE: 6

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

TABLE INCLUDES:		--- N U M B E R - O F - A N I M A L S - A F F E C T E D ---											
SEX=ALL; GROUP=ALL; WEEKS=ALL		SEX:		-----MALE-----				-----FEMALE-----					
DEATH=ALL; FIND=ALL; SUBSET=ALL		GROUP:		1-	2-	3-	4-	1-	2-	3-	4-		
ORGAN AND FINDING DESCRIPTION		NUMBER:		20	20	20	20	20	20	20	20		
STOMACH, NONGL (SU)		NUMBER EXAMINED:		14	0	15	15	15	1	15	15		
		NOT REMARKABLE:		14	0	15	15	15	1	15	15		
PANCREAS (PA)		NUMBER EXAMINED:		15	0	0	15	15	1	0	15		
		NOT REMARKABLE:		11	0	0	13	14	1	0	13		
--ATROPHY, ACINAR				0	0	0	0	1	0	0	1		
--INFILTRATE, LYMPHOHISTIOCYTIC				2	0	0	0	1	0	0	2		
--INFLAMMATION, CHRONIC-ACTIVE				1	0	0	0	2	0	0	0		
--PIGMENT, BROWN				2	0	0	0	2	0	0	0		
DUODENUM (DU)		NUMBER EXAMINED:		14	0	0	15	15	1	0	15		
		NOT REMARKABLE:		14	0	0	15	15	1	0	15		
JEJUNUM (JE)		NUMBER EXAMINED:		15	0	0	15	15	1	0	15		
		NOT REMARKABLE:		15	0	0	15	15	1	0	15		
ILEUM (IL)		NUMBER EXAMINED:		15	0	0	15	15	1	0	15		
		NOT REMARKABLE:		15	0	0	15	15	1	0	15		
CECUM (CE)		NUMBER EXAMINED:		15	0	0	15	15	1	0	15		
		NOT REMARKABLE:		15	0	0	15	15	1	0	15		

000912

TABLE 18

Incidence of Microscopic Observations

PAGE: 7

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

--- N U M B E R - O F - A N I M A L S - A F F E C T E D ---												
ORGAN AND FINDING DESCRIPTION	SEX:		---MALE---			---FEMALE---						
	GROUP:		-1-	-2-	-3-	-4-	-1-	-2-	-3-	-4-		
	NUMBER:		20	20	20	20	20	20	20	20		
COLON (CO)	NUMBER EXAMINED:	15	0	0	0	15	15	1	0	15		
	NOT REMARKABLE:	15	0	0	0	15	15	1	0	15		
RECTUM (RE)	NUMBER EXAMINED:	15	0	0	0	15	15	1	0	15		
	NOT REMARKABLE:	13	0	0	0	15	14	1	0	15		
--PARASITISM		2	0	0	0	0	1	0	0	0		
LN, MANDIBULAR (MN)	NUMBER EXAMINED:	15	2	5	15	15	1	1	3	15		
	NOT REMARKABLE:	11	0	0	7	12	1	0	0	14		
--HEMORRHAGE		4	2	5	8	2	0	0	3	1		
--HYPERPLASIA, PLASMACYTIC		0	0	0	0	1	0	0	0	0		
SALIV GL, MANDIB (SG)	NUMBER EXAMINED:	15	0	0	0	15	15	1	0	15		
	NOT REMARKABLE:	15	0	0	0	15	15	1	0	15		
HARDERIAN GLAND (HG)	NUMBER EXAMINED:	15	0	0	0	15	15	1	0	15		
	NOT REMARKABLE:	14	0	0	0	13	15	1	0	14		
--INFILTRATE, LYMPHOHISTIOCYTIC		1	0	0	0	2	0	0	0	1		
AUDITORY SEB GL (AS)	NUMBER EXAMINED:	15	0	0	0	15	15	1	0	15		
	NOT REMARKABLE:	15	0	0	0	12	11	1	0	13		
--ONE EXAMINED		0	0	0	0	3	4	0	0	2		

TABLE 18

Incidence of Microscopic Observations

PAGE: 8

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

TABLE INCLUDES:		--- N U M B E R - O F - A N I M A L S - A F F E C T E D ---											
SEX=ALL; GROUP=ALL; WEEKS=ALL DEATH=ALL; FIND=ALL; SUBSET=ALL		SEX:		MALE				FEMALE					
		GROUP: -1- -2- -3- -4-		-1- -2- -3- -4-		-1- -2- -3- -4-		-1- -2- -3- -4-					
ORGAN AND FINDING DESCRIPTION		NUMBER: 20 20 20 20		20 20 20 20		20 20 20 20		20 20 20 20					
SKIN (SK)		NUMBER EXAMINED: 15 0 0 15		15 15 15 15		1 0 1 0		1 0 1 0					
--EXUDATE, EPIDERMAL SURFACE		NOT REMARKABLE: 14 0 0 15		15 15 15 15		1 0 1 0		1 0 1 0					
--INFLAMMATION, SUBACUTE		1 0 0 0		0 0 0 0		0 0 0 0		0 0 0 0					
MAMMARY, FEMALE (MF)		1 0 0 0		0 0 0 0		0 0 0 0		0 0 0 0					
		NUMBER EXAMINED: 0 0 0 0		0 15 15 15		1 0 1 0		1 0 1 0					
		NOT REMARKABLE: 0 0 0 0		0 15 15 15		1 0 1 0		1 0 1 0					
URINARY BLADDER (UB)		NUMBER EXAMINED: 15 0 0 15		15 15 15 15		1 0 1 0		1 0 1 0					
		NOT REMARKABLE: 15 0 0 13		15 15 15 15		1 0 1 0		1 0 1 0					
--INFILTRATE, LYMPHOHISTIOCYTIC		0 0 0 2		0 0 0 0		0 0 0 0		0 0 0 0					
UTERUS (UT)		NUMBER EXAMINED: 0 0 0 0		0 15 4 1		15 4 1 1		15 4 1 1					
		NOT REMARKABLE: 0 0 0 0		0 14 1 1		10 10 10 10		10 10 10 10					
--DILATATION		0 0 0 0		0 1 3 0		5 5 5 5		5 5 5 5					
CERVIX (CV)		NUMBER EXAMINED: 0 0 0 0		0 15 1 0		15 1 0 15		15 1 0 15					
		NOT REMARKABLE: 0 0 0 0		0 15 1 0		15 1 0 15		15 1 0 15					
OVARY (OV)		NUMBER EXAMINED: 0 0 0 0		0 15 1 1		15 1 1 15		15 1 1 15					
		NOT REMARKABLE: 0 0 0 0		0 15 1 1		15 1 1 15		15 1 1 15					

000014

TABLE 18

Incidence of Microscopic Observations

PAGE: 9

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

--- N U M B E R - O F - A N I M A L S - A F F E C T E D ---												
TABLE INCLUDES:												
SEX=ALL; GROUP=ALL; WEEKS=ALL												
DEATH=ALL; FIND=ALL; SUBSET=ALL												
ORGAN AND FINDING DESCRIPTION												
SEX: -----MALE-----FEMALE-----												
GROUP: -1- -2- -3- -4- -1- -2- -3- -4-												
NUMBER: 20 20 20 20 20 20 20 20												
VAGINA (VA)	NUMBER EXAMINED:	0	0	0	0	15	1	0	15			
	NOT REMARKABLE:	0	0	0	0	15	1	0	15			
PROSTATE (PR)	NUMBER EXAMINED:	15	0	0	15	0	0	0	0	0	0	0
	NOT REMARKABLE:	3	0	0	9	0	0	0	0	0	0	0
--INFILTRATE, LYMPHOHISTIOCYTIC		12	0	0	6	0	0	0	0	0	0	0
--INFLAMMATION, CHRONIC-ACTIVE		0	0	0	1	0	0	0	0	0	0	0
SEMINAL VESICLES (SV)	NUMBER EXAMINED:	15	0	0	15	0	0	0	0	0	0	0
	NOT REMARKABLE:	15	0	0	15	0	0	0	0	0	0	0
TESTIS (TE)	NUMBER EXAMINED:	15	0	0	15	0	0	0	0	0	0	0
	NOT REMARKABLE:	15	0	0	15	0	0	0	0	0	0	0
EPIDIDYMIDES (EP)	NUMBER EXAMINED:	15	0	0	15	0	0	0	0	0	0	0
	NOT REMARKABLE:	15	0	0	15	0	0	0	0	0	0	0
TAIL (TI)	NUMBER EXAMINED:	0	0	0	0	0	0	1	0	0	0	0
	NOT REMARKABLE:	0	0	0	0	0	0	0	0	0	0	0
--HEMORRHAGE		0	0	0	0	0	0	1	0	0	0	0
--ULCERATION		0	0	0	0	0	0	1	0	0	0	0
** END OF LIST **												

000915

TABLE 19

Incidence of Severity of Selected Microscopic Observations

Covance 6329-225
3M T-6314.113-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 1

--- N U M B E R - O F - A N I M A L S - A F F E C T E D ---												
SEX:		MALE			FEMALE							
GROUP:	-1-	-2-	-3-	-4-	-1-	-2-	-3-	-4-				
NUMBER:	20	20	20	20	20	20	20	20				
--- ** TOP OF LIST ** ---												
LIVER (LI) NUMBER EXAMINED: 15 15 15 15 15 15 15 15												
--FIBROSIS, CAPSULAR												
-> 15 15 15 14 15 15 14 15 15 15 15 15												
1> 0 0 1 0 1 0 1 0 0 0 0												
TL> 15 15 15 15 15 15 15 15 15 15 15 15												
MN> 0.0 0.0 0.1 0.0 0.1 0.0 0.1 0.0 0.0 0.0 0.0												
--HYPERTROPHY, HEPATOCELLULAR, DIFFUSE												
-> 15 15 15 15 15 15 15 15 15 15 14												
2> 0 0 0 0 0 0 0 0 0 0 1												
TL> 15 15 15 15 15 15 15 15 15 15 15												
MN> 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.1												
--HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR												
-> 15 15 1 0 15 15 8 1												
1> 0 0 11 0 0 0 7 0												
2> 0 0 3 1 0 0 0 7												
3> 0 0 0 14 0 0 0 7												
TL> 15 15 15 15 15 15 15 15 15 15 15												
MN> 0.0 0.0 1.1 2.9 0.0 0.0 0.5 2.3												
--INFILTRATE, LYMPHOHISTIOCYTIC												
-> 5 0 2 5 5 4 3 6												
1> 10 14 13 10 10 11 12 9												
2> 0 1 0 0 0 0 0 0												
TL> 15 15 15 15 15 15 15 15 15 15 15												
MN> 0.7 1.1 0.9 0.7 0.7 0.7 0.8 0.6												

** CONTINUED ON NEXT PAGE **

000916

TABLE 19

Incidence of Severity of Selected Microscopic Observations

PAGE: 2

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

--- N U M B E R - O F - A N I M A L S - A F F E C T E D ---												
TABLE INCLUDES:												
SEX=ALL; GROUP=ALL; WEEKS=ALL												
DEATH=ALL; FIND=ALL; SUBSET=T												

ORGAN/TISSUE EXAMINED												

SEX: -----MALE-----FEMALE-----												
GROUP: -1- -2- -3- -4- -1- -2- -3- -4-												
NUMBER: 20 20 20 20 20 20 20 20												

** FROM PREVIOUS PAGE ** LIVER (LI) --NECROSIS, COAGULATIVE	NUMBER EXAMINED:	15	15	15	15	15	15	15	15	15	15	15
	->	15	15	15	12	15	15	15	14	15		
	1>	0	0	0	2	0	0	0	1	0		
	2>	0	0	0	1	0	0	0	0	0		
	TL>	15	15	15	15	15	15	15	15	15	15	
MN>	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.1	0.0	0.0		

--NECROSIS, INDIVIDUAL HEPATOCYTE	->	9	6	10	9	12	9	11	10			
	1>	6	9	5	5	3	6	4	5			
	2>	0	0	0	1	0	0	0	0			
	TL>	15	15	15	15	15	15	15	15	15		
	MN>	0.4	0.6	0.3	0.5	0.2	0.4	0.3	0.3			

--PIGMENT, HEPATOCELLULAR	->	15	15	15	13	15	15	15	8			
	1>	0	0	0	2	0	0	0	7			
	TL>	15	15	15	15	15	15	15	15	15		
	MN>	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.5			

--VACUOLATION, HEPATOCELLULAR, CENTRIOBLULAR TO MIDZONAL	->	15	15	13	8	15	15	15	13			
	1>	0	0	2	5	0	0	0	2			
	2>	0	0	0	2	0	0	0	0			
	TL>	15	15	15	15	15	15	15	15	15		
	MN>	0.0	0.0	0.1	0.6	0.0	0.0	0.0	0.0	0.1		
** CONTINUED ON NEXT PAGE **												

000917

TABLE 19

Incidence of Severity of Selected Microscopic Observations

Covance 6329-225
3M T-6314.113-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 3

--- N U M B E R - O F - A N I M A L S - A F F E C T E D ---												
SEX:		MALE				FEMALE						
GROUP:	-1-	-2-	-3-	-4-	-1-	-2-	-3-	-4-	-1-	-2-	-3-	-4-
NUMBER:	20	20	20	20	20	20	20	20	20	20	20	20

** FROM PREVIOUS PAGE **												

ORGAN/TISSUE EXAMINED												
LIVER (LI) NUMBER EXAMINED:												
--VACUOLATION, HEPATOCELLULAR, DIFFUSE												
	>	15	15	15	15	15	15	15	15	15	15	15
	1>	0	0	2	0	0	0	0	0	0	0	0
	2>	0	0	0	2	0	0	0	0	0	0	0
	TL>	15	15	15	15	15	15	15	15	15	15	15
	MN>	0.0	0.0	0.1	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0

--VACUOLATION, HEPATOCELLULAR, PERIPORTAL												
	>	15	15	14	15	13	12	15	15	15	15	15
	1>	0	0	1	0	2	3	0	0	0	0	0
	TL>	15	15	15	15	15	15	15	15	15	15	15
	MN>	0.0	0.0	0.1	0.0	0.1	0.2	0.0	0.0	0.0	0.0	0.0

STOMACH, GL (ST) NUMBER EXAMINED:												
	>	14	2	15	15	15	1	15	15	1	15	15
	1>	14	1	15	13	15	1	15	12	1	15	12
	2>	0	0	0	0	1	0	0	0	0	0	2
	TL>	14	1	0	1	0	1	0	0	1	15	15
	MN>	0.0	1.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.3

--EROSION												
	>	14	1	15	13	15	1	15	12	1	15	12
	1>	0	0	0	0	1	0	0	0	0	0	2
	2>	0	1	0	1	0	1	0	0	1	15	15
	TL>	14	2	15	15	15	1	15	1	15	15	15
	MN>	0.0	1.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.3

** END OF LIST **												

000913

APPENDIX 1

Protocol Deviations

Protocol

Protocol Amendment No. 1

Protocol Amendment No. 2

Protocol Amendment No. 3

Protocol Deviations

Protocol. Housing. Diet. "Certified Rodent Diet #5002, meal (PMI Nutrition International) *ad libitum*, unless otherwise specified."

Actual Procedure. The animals were given Teklad 8728C pelleted food from August 17, 1998 to August 23, 1998.

Protocol. Dose Preparation. Retention Samples. "During the in-life phase, samples (approximately 100 g) will be taken from each dose preparation sampled for dose analyses and stored at room temperature. Unless used for analyses, these samples will be discarded approximately 1 month after completion of the in-life phase."

Actual Procedure. Retention samples were discarded after approximately 1 month of the completion of the in-life phase.

Protocol. Observation of Animals. Clinical Observations. "Each animal will be observed twice daily (a.m. and p.m.) for mortality and moribundity, recording findings as they are observed."

Actual Procedure. On Days 16 through 30, the a.m. and p.m. mortality and moribundity checks were not documented.

Protocol. Termination. Scheduled Sacrifices. "After at least 4 weeks of treatment, five animals/sex/group will be fasted overnight, bled for serum samples, anesthetized with carbon dioxide, weighed, and exsanguinated."

Actual Procedure. Fasting of animals at the Week 5 scheduled sacrifice was not documented.

000920

Protocol Deviations (Continued)

Protocol. Experimental Design. Postmortem Procedures. "The following tissues (when present) or representative samples will be collected from each animal and preserved in 10% phosphate-buffered formalin, unless otherwise specified."

Actual Procedure. Some tissues, required by the protocol, were not available for histopathologic examination. Missing tissues are listed with appropriate comments in the pathology data sheets for individual animals. Summary tables do not include them as having been examined.

These deviations are not expected to have affected the results of the study.

000921



Sponsor:

3M
St. Paul, Minnesota

PROTOCOL

Study Title:

13-Week Dietary Toxicity Study with N-Methyl Perfluorooctanesulfonamido
Ethanol (N-MeFOSE, T-6314) in Rats

Date:

August 27, 1998

Performing Laboratory:

Covance Laboratories Inc.
3301 Kinsman Boulevard
Madison, Wisconsin 53704-2595

Laboratory Study Identification:

Proposal No. 90545D

Covance 6329-225

Sponsor Project Identification:

3M T-6314.1

000932

Study

13-Week Dietary Toxicity Study with N-Methyl Perfluorooctanesulfonamido Ethanol
(N-MeFOSE, T-6314) in Rats

Purpose

To assess the toxicity of the test material when administered in the diet to rats for at least
13 weeks

Sponsor

3M
Toxicology Services
Building 220-2E-02, 3M Center
St. Paul, Minnesota 55144-1000

Study Monitor

Andrew M. Seacat, PhD
3M Toxicology Services
Telephone No.: 651.575.3161
Facsimile No.: 651.733.1773

Alternate Study Monitor

Marvin T. Case, DVM, PhD
3M Toxicology Services
Telephone No.: 651.733.5180
Facsimile No.: 651.733.1773

Study Location

Covance Laboratories Inc.
3301 Kinsman Boulevard
Madison, Wisconsin 53704-2595

Mailing Address: PO Box 7545
Madison, Wisconsin 53707-7545

000923

Study Director

Peter J. Thomford, PhD
Covance Laboratories Inc.
Telephone No.: 608.241.7207
Facsimile No.: 608.242.2736

Toxicologist

Thomas E. Ryan, BS
Covance Laboratories Inc.

Proposed Study Timetable

In-Life Start Date: September 1, 1998
In-Life End Date: December 3, 1998
Audited Draft Report Date: May 26, 1998

Regulatory Compliance

This study will be conducted in compliance with the Food and Drug Administration Good Laboratory Practice Regulations as set forth in Title 21 of the US Code of Federal Regulations, Part 58, issued December 22, 1978 (effective June 20, 1979), and with any applicable amendments.

Animal Care and Use Statement

All procedures in this protocol are in compliance with the Animal Welfare Act Regulations, 9 CFR 1-4. In the opinion of the Sponsor and study director, the study does not unnecessarily duplicate any previous work.

Quality Assurance

The protocol, study conduct, and final report will be audited by the Covance Quality Assurance Unit (QAU). The proliferation cell nuclear antigen (PCNA) evaluation, data, and report will be audited by the QAU of Pathology Associates International. Liver and serum analyses, data, and report will be audited by the QAU of 3M Environmental Technology and Safety Services.

000924

Test Material

Identification

N-Methyl Perfluorooctanesulfonamido Ethanol (MeFOS, T-6314)

Lot Number

The lot numbers will be maintained in the raw data.

Purity

Responsibility of the Sponsor

Stability

Responsibility of the Sponsor

Storage Conditions

At room temperature

Characteristics

Information on synthesis methods, composition, or other characteristics that define the test material is on file with the Sponsor.

Reserve (Archive) Samples

A reserve sample (approximately 5 g) of each lot will be taken and stored at room temperature. These samples will be transferred to the Sponsor after completion of the in-life phase.

Disposition of Test Material

After authorization from the Sponsor, any remaining test material will be returned to:

Andrew M. Seacat, PhD
3M
Toxicology Services
Building 220-2E-02, 3M Center
St. Paul, Minnesota 55144-1000
Telephone No.: 651.575.3161
Facsimile No.: 651.733.1773

000925

Animals**Species**

Rat

Strain

Crl:CD®(SD) IGS BR

Source

Charles River Laboratories, Inc., Portage, Michigan

Age at Initiation of Treatment

Preferably less than 6 weeks of age but not more than 8 weeks of age

Weight at Initiation of Treatment

100 to 300 g

Number and Sex

80 males and 80 females

Identification

Implantable microchip identification device

Husbandry**Housing**

Individual (may be group-housed during acclimation). Animals will be housed in suspended, stainless steel cages.

Diet

Certified Rodent Diet #5002, meal (PMI Nutrition International) *ad libitum*, unless otherwise specified. The diet is routinely analyzed by the manufacturer for nutritional components and environmental contaminants. Specified nutrient and contaminant analyses are on file at Covance-Madison.

000936

Water

Ad libitum. Samples of the water are routinely analyzed for specified microorganisms and environmental contaminants. The results are on file at Covance-Madison.

Contaminants

There are no known contaminants in the diet or water at levels that might interfere with this study.

Environment

Environmental controls for the animal room will be set to maintain 18 to 26°C, a relative humidity of 30 to 70%, and a 12-hour light/12-hour dark cycle. The light/dark cycle may be interrupted to accommodate in-life procedures.

Acclimation

At least 1 week

Randomization

Selection of animals for the study will be based on clinical observations, ophthalmic examinations, and other data as appropriate. Animals will be assigned to treatment groups using a computerized blocking procedure designed to achieve body weight balance with respect to treatment groups. At the time of randomization, the weight variation of the animals of each sex used will not exceed ± 2 standard deviations of the mean weight, and the mean body weight for each group of each sex will not be statistically different at the 5.0% probability level.

Justification

Rats historically have been used in safety evaluation studies and are recommended by appropriate regulatory agencies.

000927

Group Designations and Dietary Levels

Group	Number of Animals		Dietary Levels (ppm MeFOS) ^a
	Male	Female	
1 (Control) ^b	20	20	0
2 (Low)	20	20	3
3 (Mid)	20	20	30
4 (High)	20	20	100

a Dose levels are expressed as ppm of MeFOS.

b The control animals will receive the basal diet only.

Dosing Procedures**Method of Administration**

Dietary, 7 days/week for at least 13 weeks. Treatment will continue through the day before necropsy.

Reason for Dosing Route

The potential route of exposure in humans is oral.

Dose Preparation

All dose preparations will be mixed according to the study-specific mixing procedure developed by Covance. Dose concentrations will be based on the MeFOS content as supplied. All dose preparations will be stored at room temperature.

Before initiation of treatment, dose preparations of 1 ppm, 3 ppm, 30 ppm, 100 ppm and 500 ppm will be mixed.

Dose preparations will be mixed at least once every 4 weeks during the in-life phase.

Retention Samples

During the in-life phase, samples (approximately 100 g) will be taken from each dose preparation sampled for dose analyses and stored at room temperature. Unless used for analyses, these samples will be discarded approximately 1 month after completion of the in-life phase.

000928

Dose Analyses

By Covance, using a method supplied by the Sponsor and validated by Covance

Homogeneity

Homogeneity will be determined for 3-ppm, 30-ppm, and 100-ppm dose preparations once pretest. One sample (approximately 100 g) each from the top, middle, and bottom of the dose preparations mixed for homogeneity analyses will be collected, divided into three subsamples for extraction and analysis, and analyzed for test material content. All samples will be stored at room temperature until analyzed within 7 days after mixing. Homogeneity analysis will be repeated if batch size changes by more than 30%.

Stability

Four samples (approximately 100 g each) will be taken from the 1-ppm dose level concentration of diet preparations mixed pretest. One sample will be analyzed on the day of mixing and used as the baseline value. One sample will be stored at room temperature for at least 19 days, then analyzed. A third sample will be stored at room temperature after at least 32 days, then analyzed. The remaining sample will be stored in a freezer set to maintain -10 to -30°C for 8 weeks, then analyzed.

In addition, two samples (approximately 100 g each) will be taken from the 500-ppm dose preparation mixed pretest. One sample will be analyzed on the day of mixing. The second sample will be stored at room temperature after at least 32 days, then analyzed.

Dose Confirmation

During the in-life phase, samples (approximately 100 g) from all dose preparations will be analyzed. All samples will be stored at room temperature until analyzed.

600929

Observation of Animals**Clinical Observations**

Each animal will be observed twice daily (a.m. and p.m.) for mortality and moribundity, recording findings as they are observed. At least once weekly, each animal will be observed (cage will be opened, and the animal will be removed); abnormal findings or an indication of normal will be recorded. Additional findings will be recorded as they are observed.

Body Weights

Each animal will be weighed at least once prior to treatment, on the first day of treatment, and weekly thereafter.

Food Consumption

Food consumption will be recorded weekly during treatment. Food consumption will not be measured during Week 5 for animals scheduled for sacrifice during Week 5.

Ophthalmic Examinations

Ophthalmic examinations will be done for each animal before initiation of treatment and before the Week 14 scheduled sacrifice. The eyes of each animal will be examined by a veterinarian using an indirect ophthalmoscope. A mydriatic agent will be instilled into the eyes prior to examination.

Clinical Pathology**Frequency**

Hematology and clinical chemistry during Weeks 5 and 14; urinalysis during Week 14

Number of Animals

Ten animals/sex/group (the same animals will be used at each interval, if possible)

Method of Collection

Animals will be fasted overnight; blood will be collected from a jugular vein. The anticoagulant will be potassium EDTA for hematology tests. Urine will be collected chilled overnight (approximately 16 hours).

000930

Tests

Hematology

red blood cell (erythrocyte) count	platelet count
hemoglobin	white blood cell (leukocyte) count
hematocrit	differential blood cell count
mean corpuscular volume	blood cell morphology
mean corpuscular hemoglobin	reticulocyte smear (made, but not examined)
mean corpuscular hemoglobin concentration	

Clinical Chemistry

glucose	alanine aminotransferase
urea nitrogen	gamma glutamyltransferase
creatinine	aspartate aminotransferase
total protein	calcium
albumin	inorganic phosphorus
globulin	sodium
cholesterol	potassium
triglycerides	chloride
total bilirubin	

Urinalysis

appearance	glucose
volume	ketones
specific gravity	bilirubin
pH	blood
protein	microscopic examination of sediment
urobilinogen	

Serum Perfluorooctane Sulfonic Acid (PFOS) Analyses

Frequency and Number of Animals

Five animals/sex/group after at least 4 and 13 weeks of treatment. Samples after 4 weeks of treatment will be collected from animals selected for the interim sacrifice.

000931

Method of Collection

Animals will be fasted overnight; blood (approximately 4 mL) will be collected from a jugular vein. Samples will be collected without anticoagulant.

Sample Handling

Blood samples will be allowed to clot at room temperature and centrifuged. Serum samples will be harvested and stored in a freezer set to maintain -60 to -80°C. Samples will be packed on dry ice and shipped to:

Kris J. Hansen, PhD
3M Environmental Technology and Safety Services
935 Bush Avenue
Building 2-3E-09
St. Paul, Minnesota 55133-3331
Telephone No.: 651.778.6018
Facsimile No.: 651.778.6176

Serum samples will be analyzed for PFOS and metabolites by the Sponsor. Results will be reported separately by the Sponsor.

Termination**Unscheduled Sacrifices and Deaths**

Necropsies will be done. Animals to be sacrificed will be anesthetized with carbon dioxide, weighed, and exsanguinated.

Scheduled Sacrifices

After at least 4 weeks of treatment, five animals/sex/group will be fasted overnight, bled for serum samples, anesthetized with carbon dioxide, weighed, and exsanguinated. The abdominal cavity of each animal will be opened, the liver will be removed and weighed, and liver samples will be collected for palmitoyl CoA and PFOS analysis. Animals will be discarded after liver collection.

000032

After at least 13 weeks of treatment, all surviving animals will be fasted overnight, bled for serum samples (five animals/sex/group) and clinical pathology tests (10/sex/group), anesthetized with carbon dioxide, weighed, exsanguinated, and necropsied. Liver samples will be collected for palmitoyl CoA, PCNA, and PFOS analysis.

Postmortem Procedures

Necropsy

The necropsy will include an examination of the external features of the carcass; all external body orifices; the abdominal, thoracic, and cranial cavities; organs; and tissues.

Organ Weights

At the scheduled sacrifice after 13 weeks of treatment, the following organs (when present) will be weighed; paired organs will be weighed together:

adrenal (2)	spleen
brain	testis (2)
kidney (2)	thymus
liver	thyroid (2) with parathyroid
ovary (2)	

Organ-to-body weight percentages and organ-to-brain weight ratios will be calculated.

Bone Marrow Smear

From the femur of each animal at the scheduled sacrifice after 13 weeks of treatment only; made but not examined

Cell Proliferation Tissue Collection and Immunohistochemical Evaluation

At the scheduled sacrifice after 13 weeks of treatment, representative samples of left lateral lobe of the liver from each animal will be collected and preserved in zinc formalin.

After fixation, each sample of liver will be embedded in paraffin, and the paraffin blocks will be shipped to:

000933

Sandra R. Eldridge, PhD
Pathology Associates International
15 Worman's Mill Court, Suite I
Frederick, Maryland 21701
Telephone No.: 301.663.1644, ext. 2201
Facsimile No: 301.663.8994

Proliferation cell nuclear antigen (PCNA) evaluation will be done on the samples. Results will be provided for inclusion in the final report.

Palmitoyl-CoA Oxidase Tissue Collection and Analyses

At the scheduled sacrifices, the right lateral lobe of the liver will be collected from each animal and flash-frozen in liquid nitrogen. The liver tissue will be stored in a freezer set to maintain -60 to -80°C until analyzed by Covance for palmitoyl-CoA oxidase activity.

Liver PFOS Analysis

At scheduled sacrifices, the remaining portion of the liver from each animal after other required collections will be stored in a freezer set to maintain -60 to -80°C. Samples will be packed on dry ice and shipped to Kris J. Hansen, PhD, 3M Environmental Technology and Safety Services. Liver samples will be analyzed for PFOS and metabolites by the Sponsor. Results will be reported separately by the Sponsor.

Tissue Preservation

The following tissues (when present) from each animal that is found dead, sacrificed at an unscheduled interval, or sacrificed at the scheduled sacrifice after 13 weeks of treatment will be preserved in 10% neutral-buffered formalin:

000934

adrenal (2)	ovary (2)
aorta	pancreas
brain	pituitary
cecum	prostate
cervix	rectum
colon	salivary gland [mandibular (2)]
duodenum	sciatic nerve
epididymis (2)	seminal vesicle (2)
esophagus	skeletal muscle (thigh)
eye (2)	skin
femur with bone marrow (articular surface of the of the distal end)	spinal cord (cervical, mid-thoracic, and lumbar)
Harderian gland	spleen
heart	sternum with bone marrow
ileum with Peyer's patch (lymphoid aggregate)	stomach
jejunum	testis (2)
kidney (2)	thymus
lesions	thyroid (2) with parathyroid
liver	trachea
lung with mainstem bronchi	urinary bladder
lymph nodes (mesenteric and mandibular)	uterus
mammary glands (females only)	vagina
	Zymbal's gland

Histopathology

Tissues (as appropriate) from each animal in the control and high-dose groups and each animal that dies or is sacrificed at an unscheduled interval will be embedded in paraffin, sectioned, stained with hematoxylin and eosin, and examined microscopically.

Macroscopic lesions from animals in the low- and mid-dose groups will be embedded in paraffin, sectioned, stained with hematoxylin and eosin, and examined microscopically.

Suspected target organs noted at the high dose will be examined microscopically from each animal (at the Sponsor's request and added by amendment).

Reports

One copy of the draft report will be sent to the Sponsor. The report will include the following information:

000935

Experimental Design and Methods

Results

dose analyses
mortality
clinical observations
body weights
body weight changes
food consumption
test material consumption
ophthalmic findings
clinical pathology results
palmitoyl CoA oxidase activities
macroscopic observations
microscopic observations
cell proliferation assessments (provided by the Sponsor's designee)

Statistical Evaluation

Levene's test will be done to test for variance homogeneity. In the case of heterogeneity of variance at $p \leq 0.05$, transformations will be used to stabilize the variance.

Comparison tests will take variance heterogeneity into consideration.

One-way analysis of variance (ANOVA) will be used (if applicable) to analyze body weights, body weight changes, food consumption, continuous clinical pathology values, and organ weight data. If the ANOVA is significant, Dunnett's t-test will be used for control versus treated group comparisons.

If the ANOVA shows significance for body weights at Week 1, one-way analysis of covariance (ANCOVA) will be used to analyze body weights, with initial body weights as the covariate. If the ANCOVA is significant, covariate-adjusted means will be used for control versus treated group comparisons.

Group comparisons (Groups 2 through 4 versus Group 1) will be evaluated at the 5.0%, two-tailed probability level. Only data collected on or after the first day of treatment will be analyzed statistically.

000936

At the end of 1 year after issuance of the audited draft report, if no requested revisions or instructions to finalize have been communicated by the Sponsor, then the audited draft report will be considered 'final' and issued as the final report, signed by the study director, and submitted to the Sponsor.

Any modifications or changes to the audited draft report requested 1 year after issuance will be performed at additional cost to the Sponsor.

Two copies of the signed final report (one unbound and one bound) will be sent to the client.

Record Retention

All raw data, documentation, records, protocol, specimens, and final report generated as a result of this study will be archived in the storage facilities of Covance-Madison for a period of 1 year following submission of the final report to the Sponsor. One year after submission of the final report, the aforementioned materials will be sent to the Sponsor, and a return fee will be charged; all raw data stored on magnetic media, protocol, study correspondence, and the original final report will be retained by Covance. The Sponsor may elect to have the materials retained in the Covance archives for an additional period of time, and Covance will charge a storage fee. If the Sponsor chooses to have Covance dispose of the materials, a disposal fee will be charged.

PCNA antigen evaluation data will be retained by Pathology Associates International.


Liver and serum samples sent to the Sponsor and analysis data will be retained by the Sponsor.

000337


PROTOCOL APPROVAL




Andrew M. Seacat, PhD
Study Monitor
3M



Date



Peter J. Thomford, PhD
Study Director
Covance Laboratories Inc.



Date

000938



PROTOCOL AMENDMENT NO. 1

Covance 6329-225

13-Week Dietary Toxicity Study with N-Methyl Perfluorooctanesulfonamido Ethanol (N-MeFOSE, T-6314) in Rats

Sponsor:	3M, St. Paul, Minnesota
Study Monitor:	Andrew M. Seacat, PhD
Testing Facility:	Covance Laboratories Inc., Madison, Wisconsin
Study Director:	Peter J. Thomford, PhD

This amendment modifies the following portion of the protocol:

Effective August 27, 1998

1. **Page 3, Proposed Study Timetable.** To correct the audited draft report date, delete the text in this section and replace with the following:

In-Life Start Date: September 1, 1998
In-Life End Date: December 3, 1998
Audited Draft Report Date: May 26, 1999


Effective October 1, 1998

2. **Page 12, Postmortem Procedures, Cell Proliferation Tissue Collection and Immunohistochemical Evaluation, Paragraph 1.** To reflect the decision to perform proliferation cell nuclear antigen evaluation for five animals/sex/group only, delete this paragraph and replace with the following:

At the scheduled sacrifice after 13 weeks of treatment, representative samples of the left lateral lobe of the liver from five animals/sex/group will be collected and preserved in zinc formalin.

000939


AMENDMENT APPROVAL



Andrew M. Seacat, PhD
Study Monitor
3M

11/5/98

Date



Peter J. Thomford, PhD
Study Director
Covance Laboratories Inc.

10/26/98

Date

070340

PROTOCOL AMENDMENT NO. 2

Covance 6329-225

13-Week Dietary Toxicity Study with N-Methyl Perfluorooctanesulfonamido
Ethanol (N-MeFOSE, T-6314) in Rats

Sponsor:	3M, St. Paul, Minnesota
Study Monitor:	Andrew M. Seacat, PhD
Testing Facility:	Covance Laboratories Inc., Madison, Wisconsin
Study Director:	Peter J. Thomford, PhD

This amendment modifies the following portion of the protocol:

Effective October 21, 1998

1. **Page 8, Dose Preparation, Dose Analysis, Stability.** To reflect the decision to perform additional stability analysis, add the following to this section:

Two additional samples (approximately 100 g each) will be taken from the 3-ppm dose preparation mixed for Week 9 through 12. One sample will be stored at room temperature for at least 19 days, then analyzed. The second sample will be stored at room temperature for at least 32 days, then analyzed. The sample collected for dose confirmation analysis will be used as the baseline sample.

000941

Effective December 4, 1998

2. **Page 13, Postmortem Procedures, Liver PFOS Analysis, Sentence 1.** To clarify collection of liver sections for PFOS analysis, delete this sentence and replace with the following:

At scheduled sacrifices, the remaining portion of the liver from each animal after other required collections will be flash-frozen in liquid nitrogen and stored in a freezer set to maintain -60 to -80°C.

Effective December 15, 1998

3. **Page 13, Postmortem Procedures, Cell Proliferation Tissue Collection and Immunohistochemical Evaluation, Paragraph 3.** To include examination of liver sections will be stained with hematoxylin and eosin as part of the cell proliferation evaluation, delete this paragraph and replace with the following:

Proliferation cell nuclear antigen (PCNA) evaluation will be done on the samples. In addition, liver sections will be stained with hematoxylin and eosin and examined microscopically. Results will be provided for inclusion in the final report.

Effective January 25, 1999

4. **Page 14, Postmortem Procedures, Histopathology, Paragraph 3.** Liver and stomach were identified as target organs in animals in the high-dose group and will be examined in animals from the mid-dose group. Therefore delete this paragraph and replace with the following:

Liver and stomach from each animal in the mid-dose group will be embedded in paraffin, sectioned, stained with hematoxylin and eosin, and examined microscopically.


000942

Effective February 8, 1999


5. **Page 14, Postmortem Procedures, Histopathology, Paragraph 3.** Liver and was identified as target organ in animals in the mid-dose group and will be examined in animals from the low-dose group. Therefore delete this paragraph and replace with the following:

Liver and stomach from each animal in the mid-dose group and liver from each animal in the low-dose group will be embedded in paraffin, sectioned, stained with hematoxylin and eosin, and examined microscopically.


AMENDMENT APPROVAL




Andrew M. Seacat, PhD
Study Monitor
3M



Date



Peter J. Thomford, PhD
Study Director
Covance Laboratories Inc.



Date

000913



PROTOCOL AMENDMENT NO. 3

Covance 6329-225

13-Week Dietary Toxicity Study with N-Methyl Perfluorooctanesulfonamido
Ethanol (N-MeFOSE, T-6314) in Rats

Sponsor:	3M, St. Paul, Minnesota
Study Monitor:	Andrew M. Seacat, PhD
Testing Facility:	Covance Laboratories Inc., Madison, Wisconsin
Study Director:	Peter J. Thomford, PhD

This amendment modifies the following portion of the protocol:

Effective August 28, 1998

1. **Page 4.** To include the vehicle used to dissolve the test material before mixing with the diet, add the following section.

Vehicle

Identification

Acetone

Lot Numbers

The lot numbers will be maintained in the raw data.

Purity

On file with the manufacturer

Stability

On file with the manufacturer

090944

Storage Conditions
At room temperature

Characteristics
Information on synthesis methods, composition, or other characteristics that define the vehicle is on file with the manufacturer.

AMENDMENT APPROVAL

Andrew M. Seacat
Andrew M. Seacat, PhD
Study Monitor
3M

June 10th 1999
Date

Peter J. Thomford
Peter J. Thomford, PhD
Study Director
Covance Laboratories Inc.

01 June 1999
Date

① To correct an error.

PJT 17 Feb 2000

000345

APPENDIX 2

Individual Animal Fate Data
Individual Clinical Observations
Individual Ophthalmic Observations

000946

APPENDIX 2
Individual Animal Fate Data

PAGE: 1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	DOSE GROUP	SEX	DEATH CODE	TYPE OF DEATH	DESCRIPTION OF DEATH	DATE OF DEATH	DAY OF STUDY	WEEK OF STUDY
C95901	1	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95902	1	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95903	1	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95904	1	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95905	1	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95906	1	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95907	1	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95908	1	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95909	1	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95910	1	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95911	1	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95912	1	MALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C95913	1	MALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C95914	1	MALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C95915	1	MALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C95916	1	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95917	1	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95918	1	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95919	1	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95920	1	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95921	2	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95922	2	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95923	2	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95924	2	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95925	2	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95926	2	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95927	2	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95928	2	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95929	2	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95930	2	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95931	2	MALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C95932	2	MALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C95933	2	MALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C95934	2	MALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5

0009/17

APPENDIX 2
Individual Animal Fate Data

PAGE: 2

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	DOSE GROUP	SEX	DEATH CODE	TYPE OF DEATH	DESCRIPTION OF DEATH	DATE OF DEATH	DAY OF STUDY	WEEK OF STUDY
C95935	2	MALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C95936	2	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95937	2	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95938	2	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95939	2	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95940	2	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95941	3	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95942	3	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95943	3	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95944	3	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95945	3	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95946	3	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95947	3	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95948	3	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95949	3	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95950	3	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95951	3	MALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C95952	3	MALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C95953	3	MALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C95954	3	MALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C95955	3	MALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C95956	3	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95957	3	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95958	3	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95959	3	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95960	3	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95961	4	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95962	4	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95963	4	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95964	4	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95965	4	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95966	4	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95967	4	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95968	4	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14

000948

APPENDIX 2

Individual Animal Fate Data

PAGE: 3

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	DOSE GROUP	SEX	DEATH CODE	TYPE OF DEATH	DESCRIPTION OF DEATH	DATE OF DEATH	DAY OF STUDY	WEEK OF STUDY
C95969	4	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95970	4	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95971	4	MALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C95972	4	MALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C95973	4	MALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C95974	4	MALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C95975	4	MALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C95976	4	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95977	4	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95978	4	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95979	4	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95980	4	MALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95981	1	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95982	1	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95983	1	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95984	1	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95985	1	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95986	1	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95987	1	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95988	1	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95989	1	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95990	1	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95991	1	FEMALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C95992	1	FEMALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C95993	1	FEMALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C95994	1	FEMALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C95995	1	FEMALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C95996	1	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95997	1	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95998	1	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C95999	1	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96000	1	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96001	2	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96002	2	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14

APPENDIX 2

Individual Animal Fate Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL-
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 4

ANIMAL NUMBER	DOSE GROUP	SEX	DEATH CODE	TYPE OF DEATH	DESCRIPTION OF DEATH	DATE OF DEATH	DAY OF STUDY	WEEK OF STUDY
C96003	2	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96004	2	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96005	2	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96006	2	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96007	2	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96008	2	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96009	2	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96010	2	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96011	2	FEMALE	1	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96012	2	FEMALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C96013	2	FEMALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C96014	2	FEMALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C96015	2	FEMALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C96016	2	FEMALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C96017	2	FEMALE	M	UNSCHEDULED	UNSCHEDULED DEATH	10/29/98	59	9
C96018	2	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96019	2	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96020	2	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96021	3	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96022	3	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96023	3	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96024	3	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96025	3	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96026	3	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96027	3	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96028	3	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96029	3	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96030	3	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96031	3	FEMALE	1	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96032	3	FEMALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C96033	3	FEMALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C96034	3	FEMALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C96035	3	FEMALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C96036	3	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14

APPENDIX 2

Individual Animal Fate Data

PAGE: 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	DOSE GROUP	SEX	DEATH CODE	TYPE OF DEATH	DESCRIPTION OF DEATH	DATE OF DEATH	DAY OF STUDY	WEEK OF STUDY
C96037	3	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96038	3	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96039	3	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96040	3	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96041	4	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96042	4	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96043	4	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96044	4	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96045	4	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96046	4	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96047	4	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96048	4	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96049	4	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96050	4	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96051	4	FEMALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C96052	4	FEMALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C96053	4	FEMALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C96054	4	FEMALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C96055	4	FEMALE	1	SCHEDULED	FIRST INTERIM SACRIFICE	10/01/98	31	5
C96056	4	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96057	4	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96058	4	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96059	4	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14
C96060	4	FEMALE	T	SCHEDULED	TERMINAL SACRIFICE	12/02/98	93	14

000951

Appendix 2

Individual Clinical Observations

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 1

ANIMAL DEATH NUMBER CODE	WK OF DEATH	CATEGORY KEYWORD QUALIFIER	GROUP: M1	DOSE: 0 PPM	DAYS 1-93 - OBSERVED DURING THE DAYS LISTED BELOW; 'C' - COMMENTS LISTED AT END OF OBSERVATIONS
C95901	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95902	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95903	T	14	APPEARANCE MALOCCLUSION	29, 36	
			SKIN & PELAGE BLACK SKIN TAIL-DISTAL	50, 57, 64, 71	
C95904	T	14	DISCHARGE GENITAL RED IN COLOR	92	
C95905	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95906	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95907	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95908	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95909	T	14	APPEARANCE MALOCCLUSION	71, 78, 85, 92-93	

000952

Appendix 2

Individual Clinical Observations

PAGE: 2

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL DEATH WK OF NUMBER CODE DEATH	CATEGORY KEYWORD QUALIFIER	GROUP: M1	DOSE: 0 PPM	DAYS 1-93 - OBSERVED DURING THE DAYS LISTED BELOW: 'C' - COMMENTS LISTED AT END OF OBSERVATIONS
C95910	T 14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95911	1 5	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95912	1 5	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95913	1 5	APPEARANCE MALOCCLUSION		22, 29, 31
C95914	1 5	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95915	1 5	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95916	T 14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95917	T 14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95918	T 14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95919	T 14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95920	T 14	ANIMAL HAS NO SIGNIFICANT FINDINGS		

000953

Appendix 2

Individual Clinical Observations

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 3

ANIMAL DEATH WK OF NUMBER CODE DEATH	CATEGORY KEYWORD QUALIFIER	GROUP: M2	DOSE: 3 PPM	DAYS 1-93 - OBSERVED DURING THE DAYS LISTED BELOW; 'C' - COMMENTS LISTED AT END OF OBSERVATIONS
C95921	T 14 APPEARANCE MALOCCLUSION			22, 29
C95922	T 14 ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95923	T 14 ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95924	T 14 ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95925	T 14 APPEARANCE MALOCCLUSION THIN			71, 78, 85, 92-93 71
C95926	T 14 ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95927	T 14 ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95928	T 14 ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95929	T 14 ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95930	T 14 APPEARANCE MALOCCLUSION			64, 71, 78
C95931	1 5 ANIMAL HAS NO SIGNIFICANT FINDINGS			

000954

Appendix 2

Individual Clinical Observations

PAGE: 4

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL DEATH NUMBER CODE	WK OF DEATH	CATEGORY KEYWORD QUALIFIER	GROUP: M2	DOSE: 3 PPM	DAYS 1-93 - OBSERVED DURING THE DAYS LISTED BELOW: 'C' - COMMENTS LISTED AT END OF OBSERVATIONS
C95932	1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95933	1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95934	1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95935	1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95936	T	14	APPEARANCE MALOCCLUSION		78, 85, 92-93
C95937	T	14	APPEARANCE MALOCCLUSION		78, 85, 92-93
C95938	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95939	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95940	T	14	APPEARANCE MALOCCLUSION		29, 36, 43, 50, 57, 64, 71, 78, 85, 92-93

000955

Appendix 2

Individual Clinical Observations

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 5

ANIMAL DEATH WK OF NUMBER CODE DEATH	CATEGORY KEYWORD QUALIFIER	GROUP: M3	DOSE: 30 PPM	DAYS 1-93 - OBSERVED DURING THE DAYS LISTED BELOW; 'C' - COMMENTS LISTED AT END OF OBSERVATIONS
C95941	T 14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95942	T 14	APPEARANCE MALOCCLUSION		50, 57, 64, 71, 78, 85, 92-93
C95943	T 14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95944	T 14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95945	T 14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95946	T 14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95947	T 14	APPEARANCE MALOCCLUSION		43, 50, 57, 64, 71
C95948	T 14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95949	T 14	APPEARANCE MALOCCLUSION		85, 92-93
		SKIN & PELAGE YELLOW HAIR COAT PENIS		92-93

936003

Appendix 2

Individual Clinical Observations

PAGE: 6

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL DEATH NUMBER CODE	WK OF DEATH	CATEGORY KEYWORD QUALIFIER	GROUP: M3	DOSE: 30 PPM	DAYS 1-93 - OBSERVED DURING THE DAYS LISTED BELOW; 'C' - COMMENTS LISTED AT END OF OBSERVATIONS
C95950	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95951	1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95952	1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95953	1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95954	1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95955	1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95956	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95957	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95958	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95959	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95960	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		

C00957

Appendix 2

Individual Clinical Observations

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 7

ANIMAL DEATH WK OF NUMBER CODE DEATH	CATEGORY KEYWORD QUALIFIER	GROUP: M4	DOSE: 100 PPM	DAYS 1-93 - OBSERVED DURING THE DAYS LISTED BELOW; 'C' - COMMENTS LISTED AT END OF OBSERVATIONS
C95961	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS	
C95962	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS	
C95963	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS	
C95964	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS	
C95965	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS	
C95966	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS	
C95967	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS	
C95968	T	14	APPEARANCE MALOCCLUSION MISSING TEETH	55, 57, 64, 71, 78, 85, 92-93 55
			DISCHARGE NASAL RED IN COLOR	55
			EYES RED DISCHARGE EYES	55
			RESPIRATION AUDIBLE	55

000958

Appendix 2

Individual Clinical Observations

PAGE: 8

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL DEATH NUMBER	DEATH CODE	WK OF DEATH	CATEGORY KEYWORD QUALIFIER	GROUP: M4	DOSE: 100 PPM	DAYS 1-93 - OBSERVED DURING THE DAYS LISTED BELOW; 'C' - COMMENTS LISTED AT END OF OBSERVATIONS
C95969	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95970	T	14	APPEARANCE MALOCCLUSION			64, 71, 78, 85, 92-93
C95971	1	5	APPEARANCE MALOCCLUSION			29, 31
C95972	1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95973	1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95974	1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95975	1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95976	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95977	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95978	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95979	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95980	T	14	APPEARANCE MALOCCLUSION			29, 36

000959

Appendix 2

Individual Clinical Observations

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 9

ANIMAL NUMBER	DEATH CODE	WK OF DEATH	CATEGORY KEYWORD QUALIFIER	GROUP: F1	DOSE: 0 PPM	DAYS 1-93 - OBSERVED DURING THE DAYS LISTED BELOW; 'C' - COMMENTS LISTED AT END OF OBSERVATIONS
C95981	T	14	APPEARANCE LIMITED USE LIMB-HIND-LEFT SWOLLEN LIMB-HIND-LEFT			64 64
C95982	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95983	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95984	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95985	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95986	T	14	APPEARANCE MALOCCLUSION			85, 92-93
C95987	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95988	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95989	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95990	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS			

630960

Appendix 2

Individual Clinical Observations

PAGE: 10

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL DEATH WK OF NUMBER CODE DEATH	CATEGORY KEYWORD QUALIFIER	GROUP: F1	DOSE: 0 PPM	DAYS 1-93 - OBSERVED DURING THE DAYS LISTED BELOW: 'C' - COMMENTS LISTED AT END OF OBSERVATIONS
C95991 1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95992 1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95993 1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95994 1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95995 1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95996 T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95997 T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95998 T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C95999 T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C96000 T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		

000961

Appendix 2

Individual Clinical Observations

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 11

ANIMAL DEATH NUMBER CODE	WK OF DEATH	CATEGORY KEYWORD	GROUP: F2	DOSE: 3 PPM	DAYS 1-93 - OBSERVED DURING THE DAYS LISTED BELOW; 'C' - COMMENTS LISTED AT END OF OBSERVATIONS
C96001	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C96002	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C96003	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C96004	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C96005	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C96006	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C96007	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C96008	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C96009	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C96010	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C96011	1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C96012	1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C96013	1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS		

000962

Appendix 2

Individual Clinical Observations

PAGE: 12

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL DEATH WK OF NUMBER CODE DEATH	CATEGORY KEYWORD QUALIFIER	GROUP: F2	DOSE: 3 PPM	DAYS 1-93 - OBSERVED DURING THE DAYS LISTED BELOW; 'C' - COMMENTS LISTED AT END OF OBSERVATIONS
C96014	1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS	
C96015	1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS	
C96016	M	9	APPEARANCE MALOCCLUSION	22, 29
			SKIN & PELAGE SORE/SCAB TAIL	59
C96017	T	14	APPEARANCE MALOCCLUSION	78, 85, 92-93
C96018	T	14	APPEARANCE MALOCCLUSION	15
C96019	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS	
C96020	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS	

000963

Appendix 2

Individual Clinical Observations

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 13

ANIMAL DEATH WK OF NUMBER CODE DEATH	CATEGORY KEYWORD QUALIFIER	GROUP: F3	DOSE: 30 PPM	DAYS 1-93 - OBSERVED DURING THE DAYS LISTED BELOW; 'C' - COMMENTS LISTED AT END OF OBSERVATIONS
C96021	T 14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C96022	T 14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C96023	T 14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C96024	T 14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C96025	T 14	APPEARANCE MALOCCLUSION		78, 85, 92-93
C96026	T 14	APPEARANCE MALOCCLUSION		57, 64
C96027	T 14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C96028	T 14	APPEARANCE MALOCCLUSION SKIN & PELAGE SORE/SCAB MOUTH		78, 85, 92-93 78
C96029	T 14	ANIMAL HAS NO SIGNIFICANT FINDINGS		

000964

Appendix 2
Individual Clinical Observations

PAGE: 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL DEATH NUMBER CODE	WK OF DEATH	CATEGORY KEYWORD QUALIFIER	GROUP: F3	DOSE: 30 PPM	DAYS 1-93 - OBSERVED DURING THE DAYS LISTED BELOW; 'C' - COMMENTS LISTED AT END OF OBSERVATIONS
C96030	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C96031	1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C96032	1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C96033	1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C96034	1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C96035	1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C96036	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C96037	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C96038	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C96039	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		
C96040	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS		

000965

Appendix 2

Individual Clinical Observations

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 15

ANIMAL NUMBER	DEATH CODE	WK OF DEATH	CATEGORY KEYWORD QUALIFIER	GROUP: F4	DOSE: 100 PPM	DAYS 1-93 - OBSERVED DURING THE DAYS LISTED BELOW; 'C' - COMMENTS LISTED AT END OF OBSERVATIONS
C96041	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C96042	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C96043	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C96044	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C96045	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C96046	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C96047	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C96048	T	14	APPEARANCE MALOCCLUSION		50	
C96049	T	14	APPEARANCE MISSING TAIL-DISTAL			
C96050	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS			36, 43, 50, 57, 64, 71, 78, 85, 92-93
C96051	1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS			

000066

Appendix 2

Individual Clinical Observations

PAGE: 16

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL DEATH WK OF		CATEGORY	GROUP: F4	DOSE: 100 PPM	DAYS 1-93 - OBSERVED DURING THE DAYS LISTED BELOW:	'C' - COMMENTS LISTED AT END OF OBSERVATIONS
NUMBER	CODE					
DEATH	DEATH	KEYWORD				
		QUALIFIER				
C96052	1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C96053	1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C96054	1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C96055	1	5	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C96056	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C96057	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C96058	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C96059	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C96060	T	14	ANIMAL HAS NO SIGNIFICANT FINDINGS			

000967

Appendix 2

Individual Ophthalmic Observations
Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL,
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 1

ANIMAL NUMBER	CATEGORY KEYWORD QUALIFIER	GROUP: M1	DOSE: 0 PPM	WEEK 14
C95901	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95902	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95903	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95904	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95905	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95906	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95907	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95908	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95909	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95910	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95916	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95917	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95918	ANIMAL HAS NO SIGNIFICANT FINDINGS			

000968

Appendix 2

Individual Ophthalmic Observations
Week 14

PAGE: 2

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	CATEGORY		GROUP: M1	DOSE: 0 PPM	WEEK 14
	KEYWORD	QUALIFIER			
C95919	ANIMAL HAS NO SIGNIFICANT FINDINGS				
C95920	ANIMAL HAS NO SIGNIFICANT FINDINGS				

640969

Appendix 2

Individual Ophthalmic Observations

Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 3

ANIMAL NUMBER	CATEGORY KEYWORD QUALIFIER	GROUP: M2	DOSE: 3 PPM	WEEK 14
C95921	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95922	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95923	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95924	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95925	CONJUNCTIVA CONJUNCTIVITIS EYES			14
C95926	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95927	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95928	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95929	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95930	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95936	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95937	ANIMAL HAS NO SIGNIFICANT FINDINGS			

000970

Appendix 2
Individual Ophthalmic Observations
Week 14

PAGE: 4

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

GROUP: M2 DOSE: 3 PPM WEEK 14

CATEGORY
KEYWORD
QUALIFIER

ANIMAL
NUMBER

C95938 ANIMAL HAS NO SIGNIFICANT FINDINGS
C95939 ANIMAL HAS NO SIGNIFICANT FINDINGS
C95940 ANIMAL HAS NO SIGNIFICANT FINDINGS

000971

Appendix 2

Individual Ophthalmic Observations
Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 5

ANIMAL NUMBER	CATEGORY KEYWORD QUALIFIER	GROUP: M3	DOSE: 30 PPM	WEEK 14
C95941	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95942	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95943	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95944	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95945	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95946	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95947	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95948	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95949	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95950	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95956	RETINA RETINAL DEGENERATION-FOCAL EYE-RIGHT			14
C95957	ANIMAL HAS NO SIGNIFICANT FINDINGS			

000972

Appendix 2
Individual Ophthalmic Observations
Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 6

GROUP: M3 DOSE: 30 PPM WEEK 14

CATEGORY
KEYWORD
QUALIFIER

ANIMAL
NUMBER

C95958 ANIMAL HAS NO SIGNIFICANT FINDINGS
C95959 ANIMAL HAS NO SIGNIFICANT FINDINGS
C95960 ANIMAL HAS NO SIGNIFICANT FINDINGS

000973

Appendix 2

Individual Ophthalmic Observations
Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 7

GROUP: M4 DOSE: 100 PPM
WEEK 14

CATEGORY
KEYWORD
QUALIFIER

C95961 ANIMAL HAS NO SIGNIFICANT FINDINGS
C95962 ANIMAL HAS NO SIGNIFICANT FINDINGS
C95963 ANIMAL HAS NO SIGNIFICANT FINDINGS
C95964 ANIMAL HAS NO SIGNIFICANT FINDINGS
C95965 ANIMAL HAS NO SIGNIFICANT FINDINGS
C95966 ANIMAL HAS NO SIGNIFICANT FINDINGS
C95967 ANIMAL HAS NO SIGNIFICANT FINDINGS
C95968 CONJUNCTIVA
CONJUNCTIVITIS
EYE-LEFT
14
C95969 ANIMAL HAS NO SIGNIFICANT FINDINGS
C95970 ANIMAL HAS NO SIGNIFICANT FINDINGS
C95976 ANIMAL HAS NO SIGNIFICANT FINDINGS
C95977 ANIMAL HAS NO SIGNIFICANT FINDINGS

000974

Appendix 2

Individual Ophthalmic Observations
Week 14

PAGE: 8

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	CATEGORY KEYWORD QUALIFIER	GROUP: M4	DOSE: 100 PPM	WEEK 14
------------------	----------------------------------	-----------	---------------	---------

C95978	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95979	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C95980	ANIMAL HAS NO SIGNIFICANT FINDINGS			

000975

Appendix 2

Individual Ophthalmic Observations

Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 9

GROUP: F1 DOSE: 0 PPM WEEK 14

CATEGORY
KEYWORD
QUALIFIER

ANIMAL
NUMBER

C95981	ANIMAL HAS NO SIGNIFICANT FINDINGS
C95982	ANIMAL HAS NO SIGNIFICANT FINDINGS
C95983	ANIMAL HAS NO SIGNIFICANT FINDINGS
C95984	ANIMAL HAS NO SIGNIFICANT FINDINGS
C95985	ANIMAL HAS NO SIGNIFICANT FINDINGS
C95986	ANIMAL HAS NO SIGNIFICANT FINDINGS
C95987	ANIMAL HAS NO SIGNIFICANT FINDINGS
C95988	ANIMAL HAS NO SIGNIFICANT FINDINGS
C95989	ANIMAL HAS NO SIGNIFICANT FINDINGS
C95990	ANIMAL HAS NO SIGNIFICANT FINDINGS
C95996	ANIMAL HAS NO SIGNIFICANT FINDINGS
C95997	ANIMAL HAS NO SIGNIFICANT FINDINGS
C95998	ANIMAL HAS NO SIGNIFICANT FINDINGS

000976

Appendix 2

Individual Ophthalmic Observations
Week 14

PAGE: 10

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	CATEGORY KEYWORD QUALIFIER	GROUP: F1	DOSE: 0 PPM	WEEK 14
C95999	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C96000	ANIMAL HAS NO SIGNIFICANT FINDINGS			

000977

Appendix 2

Individual Ophthalmic Observations
Week 14

PAGE: 11

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

GROUP: F2 DOSE: 3 PPM WEEK 14

CATEGORY

KEYWORD

QUALIFIER

ANIMAL
NUMBER

C96001	ANIMAL HAS NO SIGNIFICANT FINDINGS
C96002	ANIMAL HAS NO SIGNIFICANT FINDINGS
C96003	ANIMAL HAS NO SIGNIFICANT FINDINGS
C96004	ANIMAL HAS NO SIGNIFICANT FINDINGS
C96005	ANIMAL HAS NO SIGNIFICANT FINDINGS
C96006	ANIMAL HAS NO SIGNIFICANT FINDINGS
C96007	ANIMAL HAS NO SIGNIFICANT FINDINGS
C96008	ANIMAL HAS NO SIGNIFICANT FINDINGS
C96009	ANIMAL HAS NO SIGNIFICANT FINDINGS
C96010	ANIMAL HAS NO SIGNIFICANT FINDINGS
C96016	ANIMAL HAS NO SIGNIFICANT FINDINGS
C96017	ANIMAL HAS NO SIGNIFICANT FINDINGS
C96018	ANIMAL HAS NO SIGNIFICANT FINDINGS

000978

Appendix 2
Individual Ophthalmic Observations
Week 14

PAGE: 12

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	CATEGORY KEYWORD QUALIFIER	GROUP: F2	DOSE: 3 PPM	WEEK 14
C96019	ANIMAL HAS NO SIGNIFICANT FINDINGS			
C96020	ANIMAL HAS NO SIGNIFICANT FINDINGS			

000979

Appendix 2

Individual Ophthalmic Observations
Week 14

PAGE: 13

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

GROUP: F3 DOSE: 30 PPM WEEK 14

CATEGORY
KEYWORD
QUALIFIER

C96021 ANIMAL HAS NO SIGNIFICANT FINDINGS
C96022 ANIMAL HAS NO SIGNIFICANT FINDINGS
C96023 ANIMAL HAS NO SIGNIFICANT FINDINGS
C96024 ANIMAL HAS NO SIGNIFICANT FINDINGS
C96025 ANIMAL HAS NO SIGNIFICANT FINDINGS
C96026 ANIMAL HAS NO SIGNIFICANT FINDINGS
C96027 ANIMAL HAS NO SIGNIFICANT FINDINGS
C96028 ANIMAL HAS NO SIGNIFICANT FINDINGS
C96029 ANIMAL HAS NO SIGNIFICANT FINDINGS
C96030 ANIMAL HAS NO SIGNIFICANT FINDINGS
C96036 ANIMAL HAS NO SIGNIFICANT FINDINGS
C96037 ANIMAL HAS NO SIGNIFICANT FINDINGS
C96038 ANIMAL HAS NO SIGNIFICANT FINDINGS

000980

Appendix 2
Individual Ophthalmic Observations
Week 14

PAGE: 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

GROUP: F3 DOSE: 30 PPM WEEK 14

CATEGORY
KEYWORD
QUALIFIER

ANIMAL
NUMBER

ANIMAL HAS NO SIGNIFICANT FINDINGS

ANIMAL HAS NO SIGNIFICANT FINDINGS

C96039

C96040

000981

Appendix 2

Individual Ophthalmic Observations

Week 14

PAGE: 15

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

GROUP: F4 DOSE: 100 PPM WEEK 14

CATEGORY
KEYWORD
QUALIFIER

ANIMAL
NUMBER

C96041	ANIMAL HAS NO SIGNIFICANT FINDINGS
C96042	ANIMAL HAS NO SIGNIFICANT FINDINGS
C96043	ANIMAL HAS NO SIGNIFICANT FINDINGS
C96044	ANIMAL HAS NO SIGNIFICANT FINDINGS
C96045	ANIMAL HAS NO SIGNIFICANT FINDINGS
C96046	ANIMAL HAS NO SIGNIFICANT FINDINGS
C96047	ANIMAL HAS NO SIGNIFICANT FINDINGS
C96048	ANIMAL HAS NO SIGNIFICANT FINDINGS
C96049	ANIMAL HAS NO SIGNIFICANT FINDINGS
C96050	ANIMAL HAS NO SIGNIFICANT FINDINGS
C96056	ANIMAL HAS NO SIGNIFICANT FINDINGS
C96057	ANIMAL HAS NO SIGNIFICANT FINDINGS
C96058	ANIMAL HAS NO SIGNIFICANT FINDINGS

000932

Appendix 2

Individual Ophthalmic Observations
Week 14

PAGE: 16

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	CATEGORY		GROUP: F4	DOSE: 100 PPM	WEEK 14
	KEYWORD	QUALIFIER			
C96059	ANIMAL HAS NO SIGNIFICANT FINDINGS				
C96060	ANIMAL HAS NO SIGNIFICANT FINDINGS				

000383

APPENDIX 3

Individual Body Weight Data (g)

C00934

Appendix 3

Individual Body Weight Data (g)

PAGE: 1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WEEK 1	WEEK 2	WEEK 3	WEEK 4	WEEK 5	WEEK 6	WEEK 7	WEEK 8	WEEK 9	WEEK 10
GROUP: MALE 1 - 0 PPM										
C95901	208	261	314	352	380	391	420	430	455	473
C95902	216	271	321	362	392	407	433	444	473	491
C95903	204	254	308	355	374	398	424	437	459	472
C95904	206	259	315	361	395	409	436	453	472	485
C95905	204	257	303	341	376	386	411	433	453	466
C95906	197	250	304	355	391	411	445	461	485	507
C95907	208	253	298	342	366	383	406	417	438	455
C95908	202	262	316	364	399	417	442	461	480	497
C95909	203	254	326	377	415	429	457	477	505	533
C95910	215	272	337	389	411					
C95911	225	281	334	389	402					
C95912	222	281	345	353	377					
C95913	234	299	349	377	397					
C95914	202	256	302	349	377					
C95915	185	219	245	276	293	462	487	506	531	545
C95916	226	284	342	389	425	428	454	474	502	510
C95917	205	264	324	370	400	418	438	458	476	485
C95918	209	262	321	371	395	427	452	465	491	506
C95919	213	270	326	370	405	422	443	464	481	504
C95920	216	267	317	360	397					
GROUP: MALE 2 - 3 PPM										
C95921	215	267	312	350	372	389	411	422	449	466
C95922	207	256	301	345	374	389	413	421	453	466
C95923	201	253	306	346	374	390	418	437	465	479
C95924	232	281	327	360	388	406	427	445	465	479
C95925	227	282	334	360	396	418	439	461	480	494
C95926	206	250	294	324	350	366	380	401	421	433
C95927	202	251	299	336	359	373	397	417	434	454
C95928	199	257	314	359	391	404	429	447	466	477

Appendix 3

Individual Body Weight Data (g)

PAGE: 2

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WEEK 1	WEEK 2	WEEK 3	WEEK 4	WEEK 5	WEEK 6	WEEK 7	WEEK 8	WEEK 9	WEEK 10
GROUP: MALE 2 - 3 PPM										
C95929	208	259	311	351	377	393	418	436	461	475
C95930	220	275	329	374	398	413	441	449	472	477
C95931	210	258	316	367	398					
C95932	203	252	303	342	365					
C95933	206	260	308	353	381					
C95934	191	238	283	322	352					
C95935	208	261	318	361	392					
C95936	224	290	355	408	436	465	496	511	528	554
C95937	217	273	321	357	378	395	412	421	444	459
C95938	221	283	335	380	406	426	451	466	486	505
C95939	209	255	299	341	363	384	404	413	433	442
C95940	214	263	311	350	383	398	418	432	455	470
GROUP: MALE 3 - 30 PPM										
C95941	199	245	284	318	342	357	373	387	403	407
C95942	203	248	294	325	347	360	380	372	400	408
C95943	230	283	331	371	397	408	429	452	466	486
C95944	209	263	307	347	366	382	404	413	430	438
C95945	205	261	315	355	385	395	418	443	462	477
C95946	204	253	294	336	369	386	410	425	450	464
C95947	202	252	299	337	368	381	397	415	433	451
C95948	216	265	316	358	380	397	419	433	457	467
C95949	219	276	332	383	413	428	457	478	502	513
C95950	215	268	323	364	389	407	431	445	463	465
C95951	236	297	351	392	416					
C95952	205	250	294	329	347					
C95953	207	258	308	342	362					
C95954	225	279	332	374	404					
C95955	198	259	314	350	381					
C95956	209	264	318	366	396	425	448	469	489	505

C009386

Appendix 3

Individual Body Weight Data (g)

PAGE: 3

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WEEK 1	WEEK 2	WEEK 3	WEEK 4	WEEK 5	WEEK 6	WEEK 7	WEEK 8	WEEK 9	WEEK 10
GROUP: MALE 3 - 30 PPM										
C95957	226	279	326	367	389	411	427	440	460	474
C95958	205	261	308	343	369	389	406	420	438	459
C95959	190	244	290	323	353	372	392	401	421	436
C95960	214	264	317	348	383	399	420	426	446	452
GROUP: MALE 4 - 100 PPM										
C95961	227	284	334	373	402	417	442	456	476	489
C95962	205	252	292	321	345	356	381	397	412	423
C95963	217	264	310	343	369	389	412	427	452	463
C95964	222	266	317	351	382	389	418	434	447	456
C95965	195	245	283	313	333	344	371	382	393	411
C95966	197	245	284	317	349	360	386	405	426	438
C95967	204	256	301	334	363	374	399	408	423	438
C95968	206	258	310	349	377	390	422	437	447	463
C95969	219	277	317	351	377	386	417	432	451	463
C95970	232	281	329	362	385	394	427	434	453	447
C95971	203	252	298	336	340					
C95972	200	253	304	341	367					
C95973	184	234	271	295	311					
C95974	197	246	293	320	343					
C95975	214	262	306	334	357	350	361	367	380	388
C95976	200	244	283	312	332	376	400	416	432	448
C95977	209	256	299	330	358	391	410	422	439	449
C95978	212	268	313	344	367	379	398	407	419	429
C95979	204	255	300	334	359	379	397	409	429	437
C95980	219	269	313	342	369	382				

Appendix 3

Individual Body Weight Data (g)

PAGE: 4

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WEEK 1	WEEK 2	WEEK 3	WEEK 4	WEEK 5	WEEK 6	WEEK 7	WEEK 8	WEEK 9	WEEK 10
GROUP: FEMALE 1 - 0 PPM										
C95981	156	191	220	240	256	260	273	287	286	288
C95982	142	175	213	216	224	234	241	246	256	261
C95983	146	173	186	195	214	211	224	219	229	231
C95984	169	192	215	241	247	243	251	254	263	276
C95985	169	195	213	234	240	254	262	267	270	286
C95986	169	190	218	230	236	247	260	262	262	273
C95987	146	173	182	213	211	218	229	229	240	250
C95988	150	171	182	215	217	232	239	234	250	251
C95989	153	189	204	228	249	252	254	262	286	283
C95990	172	208	225	256	280	279	292	301	302	322
C95991	168	193	214	219	240					
C95992	148	167	177	203	213					
C95993	158	180	184	207	218					
C95994	172	194	217	229	245					
C95995	152	162	181	201	207	258	260	264	271	276
C95996	165	194	210	237	244	223	233	238	249	252
C95997	152	176	192	211	219	241	241	246	244	250
C95998	167	194	202	226	237	300	318	318	332	345
C95999	173	194	222	260	286	248	261	261	268	275
C96000	161	180	206	225	232					
GROUP: FEMALE 2 - 3 PPM										
C96001	143	161	187	204	201	214	224	228	243	249
C96002	142	160	176	178	200	205	209	209	222	231
C96003	153	168	190	201	196	207	217	215	219	226
C96004	176	153	124	198	218	236	239	239	251	260
C96005	165	187	205	221	224	232	236	241	252	254
C96006	150	170	187	210	218	220	232	234	244	248
C96007	164	188	202	226	234	245	244	250	260	266
C96008	168	186	217	229	238	246	261	260	277	279

000988

Appendix 3

Individual Body Weight Data (g)

PAGE: 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WEEK 1	WEEK 2	WEEK 3	WEEK 4	WEEK 5	WEEK 6	WEEK 7	WEEK 8	WEEK 9	WEEK 10
GROUP: FEMALE 2 - 3 PPM										
C96009	140	156	166	170	181	190	200	198	205	216
C96010	162	184	209	216	232	240	249	249	265	270
C96011	149	184	203	246	250					
C96012	149	167	193	213	223					
C96013	169	195	217	239	250					
C96014	167	193	214	245	260					
C96015	155	178	205	228	236	266	271	273	284	296
C96016	177	206	232	245	252	291	287	287	285	281
C96017	164	194	224	257	271	265	272	275	283	288
C96018	160	189	224	239	243	264	271	280	289	256
C96019	178	203	228	245	247	237	239	242	252	
C96020	154	184	193	222	224					
GROUP: FEMALE 3 - 30 PPM										
C96021	144	162	182	211	222	225	227	234	242	249
C96022	167	193	221	232	250	249	256	261	271	275
C96023	167	189	210	219	234	234	243	244	255	264
C96024	174	196	220	235	237	243	255	252	269	273
C96025	171	194	197	233	232	235	243	250	251	265
C96026	150	168	188	206	213	211	228	230	232	233
C96027	165	194	222	245	258	266	277	280	297	300
C96028	168	178	217	224	237	240	248	258	266	269
C96029	151	173	197	206	223	227	236	236	245	251
C96030	169	191	189	211	220	223	236	241	251	251
C96031	151	169	191	219	230					
C96032	159	182	191	220	230					
C96033	158	177	195	233	240					
C96034	160	190	210	246	262					
C96035	167	193	220	240	245					
C96036	169	203	227	240	231	238	249	254	261	266
C96036	164	185	206	226						

Appendix 3

Individual Body Weight Data (g)

PAGE: 6

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WEEK 1	WEEK 2	WEEK 3	WEEK 4	WEEK 5	WEEK 6	WEEK 7	WEEK 8	WEEK 9	WEEK 10
GROUP: FEMALE 3 - 30 PPM										
C96037	158	190	202	226	228	238	242	248	260	267
C96038	161	178	203	221	228	244	256	261	263	276
C96039	156	181	201	225	233	243	254	254	267	270
C96040	140	162	170	188	198	208	219	219	224	237
GROUP: FEMALE 4 - 100 PPM										
C96041	140	164	180	194	200	201	205	215	223	224
C96042	163	184	200	213	216	224	229	231	239	239
C96043	151	170	188	202	209	211	219	222	226	227
C96044	169	198	217	236	241	253	259	258	268	271
C96045	164	182	207	218	222	234	244	250	255	263
C96046	170	190	211	226	238	249	258	263	269	277
C96047	160	171	185	198	200	206	209	214	219	224
C96048	157	188	199	215	221	227	231	227	239	241
C96049	144	158	163	180	188	189	187	189	198	202
C96050	150	165	187	199	205	205	212	214	216	221
C96051	153	163	185	199	199					
C96052	179	192	208	223	228					
C96053	152	169	176	206	218					
C96054	156	174	180	198	209					
C96055	155	169	172	193	198					
C96056	155	174	190	194	206	215	222	214	222	227
C96057	166	196	218	222	237	248	256	258	262	264
C96058	169	193	224	228	230	241	246	251	257	257
C96059	141	164	176	174	193	199	210	206	218	222
C96060	165	188	213	221	233	245	243	247	258	262

000930

Appendix 3
Individual Body Weight Data (g)
13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 7

ANIMAL NUMBER	WEEK 11	WEEK 12	WEEK 13	WEEK 14
GROUP: MALE 1 - 0 PPM				
C95901	481	492	508	523
C95902	505	514	533	542
C95903	484	493	510	518
C95904	501	516	531	539
C95905	478	488	495	509
C95906	525	539	561	581
C95907	468	481	501	511
C95908	508	517	526	543
C95909	430	448	459	475
C95910	550	579	596	623
C95916	561	570	588	601
C95917	529	542	559	577
C95918	493	505	520	535
C95919	520	532	551	567
C95920	511	526	545	557
GROUP: MALE 2 - 3 PPM				
C95921	483	494	500	512
C95922	480	494	510	527
C95923	499	505	529	531
C95924	494	503	520	527
C95925	438	474	498	510
C95926	447	457	468	479
C95927	468	485	497	511
C95928	495	501	515	524
C95929	502	517	540	549
C95930	490	502	519	525
C95936	568	575	577	585
C95937	463	444	459	482
C95938	517	515	532	544
C95939	451	465	474	483

000391

Appendix 3

Individual Body Weight Data (g)

PAGE: 8

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WEEK 11	WEEK 12	WEEK 13	WEEK 14
GROUP: MALE 2 - 3 PPM				
C95940	487	504	518	524
GROUP: MALE 3 - 30 PPM				
C95941	419	426	435	448
C95942	421	427	449	454
C95943	501	509	529	534
C95944	444	452	466	475
C95945	486	501	513	524
C95946	476	480	492	502
C95947	457	458	463	471
C95948	478	493	504	515
C95949	534	542	517	413
C95950	473	482	495	502
C95956	518	534	543	558
C95957	479	487	499	516
C95958	471	485	500	502
C95959	448	458	467	476
C95960	460	466	477	481
GROUP: MALE 4 - 100 PPM				
C95961	497	509	513	522
C95962	434	443	457	468
C95963	478	488	501	511
C95964	466	474	485	499
C95965	422	430	443	453
C95966	451	461	468	482
C95967	449	460	472	480
C95968	458	468	485	494
C95969	475	481	485	500

000392

Appendix 3

Individual Body Weight Data (g)

PAGE: 9

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WEEK			
	11	12	13	14
GROUP: MALE 4 - 100 PPM				
C95970	460	462	468	473
C95976	393	404	408	418
C95977	458	469	479	490
C95978	465	475	485	490
C95979	441	452	459	469
C95980	442	450	458	465
GROUP: FEMALE 1 - 0 PPM				
C95981	292	313	307	331
C95982	262	267	271	275
C95983	240	241	241	246
C95984	280	285	284	299
C95985	293	297	296	312
C95986	281	285	290	297
C95987	251	256	264	272
C95988	264	261	276	283
C95989	284	299	311	310
C95990	330	332	349	341
C95996	278	281	292	291
C95997	256	266	271	277
C95998	257	262	266	267
C95999	362	372	388	394
C96000	286	292	290	298
GROUP: FEMALE 2 - 3 PPM				
C96001	245	253	253	263
C96002	236	237	243	251
C96003	234	231	232	244
C96004	267	268	279	298

000993

Appendix 3

Individual Body Weight Data (g)

PAGE: 10

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WEEK 11	WEEK 12	WEEK 13	WEEK 14
------------------	------------	------------	------------	------------

GROUP: FEMALE 2 - 3 PPM

C96005	257	261	266	269
C96006	251	258	266	274
C96007	268	271	280	292
C96008	289	296	300	305
C96009	216	218	226	231
C96010	274	275	283	292
C96017	303	301	318	303
C96018	299	292	307	306
C96019	300	298	313	321
C96020	257	261	260	267

GROUP: FEMALE 3 - 30 PPM

C96021	247	254	258	262
C96022	274	284	283	292
C96023	265	265	274	277
C96024	271	273	278	283
C96025	273	274	265	275
C96026	247	249	257	255
C96027	300	312	322	323
C96028	280	279	290	295
C96029	252	252	257	265
C96030	254	258	262	262
C96036	270	270	276	275
C96037	261	273	269	283
C96038	282	290	291	304
C96039	274	276	285	284
C96040	239	241	239	251

000994

Appendix 3

Individual Body Weight Data (g)

PAGE: 11

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WEEK 11	WEEK 12	WEEK 13	WEEK 14
------------------	------------	------------	------------	------------

GROUP: FEMALE 4 - 100 PPM

C96041	222	231	234	236
C96042	242	246	249	247
C96043	232	239	242	239
C96044	276	282	283	287
C96045	269	271	273	280
C96046	279	283	284	290
C96047	224	226	227	231
C96048	248	243	247	259
C96049	203	207	208	213
C96050	225	228	233	239
C96056	228	224	226	229
C96057	270	267	276	279
C96058	256	262	269	271
C96059	227	219	228	232
C96060	267	269	275	271

000995

APPENDIX 4

Individual Food Consumption Data (g)
Individual Test Material Consumption Data (mg/kg/day)

000396

Appendix 4

Individual Food Consumption Data (g)

PAGE: 1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WEEK 1	WEEK 2	WEEK 3	WEEK 4	WEEK 5	WEEK 6	WEEK 7	WEEK 8	WEEK 9	WEEK 10
GROUP: MALE 1 - 0 PPM										
C95901	165	168	171	183	159	179	181	181	176	170
C95902	172	174	179	190	178	197	187	187	192	190
C95903	160	173	176	187	162	190	182	181	182	177
C95904	176	182	191	202	177	203	202	208	213	215
C95905	164	165	175	182	166	182	SPILLED	SPILLED	214	184
C95906	161	166	182	199	168	199	195	188	183	192
C95907	156	158	169	174	157	180	177	167	166	161
C95908	177	183	192	202	172	197	199	192	190	192
C95909	152	151	154	SPILLED	147	170	164	165	160	153
C95910	175	179	191	198	176	194	200	193	208	SPILLED
C95911	172	165	182	172						
C95912	187	191	194	198						
C95913	181	179	169	174						
C95914	164	172	180	187						
C95915	136	SPILLED	163	142						
C95916	176	188	191	192	192	194	190	186	185	193
C95917	187	185	184	185	189	187	189	186	178	179
C95918	168	181	186	187	184	180	182	181	178	176
C95919	178	207	231	SPILLED	233	233	242	SPILLED	240	253
C95920	164	168	179	190	192	185	189	184	188	183
GROUP: MALE 2 - 3 PPM										
C95921	162	170	168	175	168	176	185	SPILLED	187	192
C95922	163	168	175	179	158	175	176	183	178	171
C95923	159	170	175	178	163	185	186	191	187	190
C95924	171	177	179	184	164	186	186	186	183	183
C95925	185	195	SPILLED	214	187	206	211	215	203	99
C95926	164	162	167	168	157	171	177	167	167	165
C95927	161	168	165	170	155	172	177	175	173	170
C95928	187	189	188	202	SPILLED	201	206	201	197	189

000937

Appendix 4

Individual Food Consumption Data (g)

PAGE: 2

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WEEK 1	WEEK 2	WEEK 3	WEEK 4	WEEK 5	WEEK 6	WEEK 7	WEEK 8	WEEK 9	WEEK 10
GROUP: MALE 2 - 3 PPM										
C95929	175	186	191	195	191	237	234	SPILLED	235	227
C95930	177	194	218	195	176	208	190	186	186	170
C95931	155	SPILLED	215	SPILLED						
C95932	160	165	166	173						
C95933	165	172	176	172						
C95934	156	161	164	169						
C95935	183	195	197	197	192	191	194	188	189	187
C95936	171	189	203	197	184	184	186	SPILLED	198	195
C95937	163	172	175	178	186	194	197	199	197	202
C95938	177	189	197	204	186	185	184	180	168	169
C95939	163	174	187	191	199	178	187	185	189	188
C95940	165	171	176	184	179					
GROUP: MALE 3 - 30 PPM										
C95941	154	159	158	166	152	162	159	153	145	145
C95942	156	162	161	166	152	170	145	153	158	166
C95943	178	184	190	197	162	186	189	183	185	178
C95944	174	179	183	186	164	189	180	178	169	168
C95945	168	184	191	193	179	198	197	187	187	183
C95946	169	168	180	189	173	186	184	189	178	175
C95947	151	161	166	171	152	168	162	155	157	149
C95948	166	179	200	188	176	185	183	185	182	181
C95949	172	186	197	201	181	207	209	204	200	199
C95950	168	180	189	188	168	200	191	189	190	180
C95951	181	185	193	192						
C95952	155	161	167	174						
C95953	178	186	192	187						
C95954	184	197	202	216						
C95955	194	197	200	210	SPILLED	259	246	SPILLED	236	223
C95956	208	SPILLED	214	228						

000000

Appendix 4

Individual Food Consumption Data (g)

PAGE: 3

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WEEK 1	WEEK 2	WEEK 3	WEEK 4	WEEK 5	WEEK 6	WEEK 7	WEEK 8	WEEK 9	WEEK 10
GROUP: MALE 3 - 30 PPM										
C95957	166	179	188	189	184	181	182	180	176	167
C95958	168	175	180	182	181	194	176	183	188	180
C95959	163	176	178	193	185	193	189	190	188	181
C95960	168	181	181	200	192	175	175	178	167	164
GROUP: MALE 4 - 100 PPM										
C95961	183	189	191	194	171	200	194	192	188	180
C95962	159	168	165	167	154	179	178	171	167	167
C95963	172	171	171	171	165	184	179	180	183	176
C95964	172	191	181	194	171	185	190	185	179	175
C95965	159	168	168	172	157	177	172	165	169	162
C95966	158	167	171	173	154	180	179	178	180	183
C95967	170	178	175	179	157	179	183	176	179	177
C95968	188	184	191	197	171	195	194	159	158	191
C95969	181	179	178	180	155	183	183	181	177	177
C95970	171	198	SPIILLED	187	170	195	187	187	174	173
C95971	159	166	170	152						
C95972	160	169	172	172						
C95973	165	170	164	167						
C95974	162	175	170	172						
C95975	164	168	169	173						
C95976	162	167	164	165	161	163	162	157	155	146
C95977	163	174	169	177	173	182	186	181	184	176
C95978	166	171	165	170	171	172	173	174	180	180
C95979	170	175	177	176	183	175	179	170	171	170
C95980	174	182	178	177	180	181	184	182	180	174

000939

Appendix 4

Individual Food Consumption Data (g)

PAGE: 4

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WEEK 1	WEEK 2	WEEK 3	WEEK 4	WEEK 5	WEEK 6	WEEK 7	WEEK 8	WEEK 9	WEEK 10
GROUP: FEMALE 1 - 0 PPM										
C95981	138	150	153	153	129	161	159	129	104	137
C95982	127	143	123	116	115	126	134	120	119	114
C95983	127	139	SPILLED	SPILLED	108	120	SPILLED	102	SPILLED	130
C95984	126	130	145	137	125	138	148	150	141	131
C95985	135	128	133	137	132	137	137	135	136	129
C95986	137	152	132	138	121	SPILLED	130	127	132	128
C95987	124	123	163	130	113	128	130	129	132	119
C95988	124	114	127	125	SPILLED	130	119	126	125	126
C95989	135	134	144	SPILLED	137	SPILLED	SPILLED	SPILLED	SPILLED	244
C95990	162	147	165	170	135	156	158	140	147	147
C95991	131	126	135	159						
C95992	111	101	116	116						
C95993	121	108	119	125						
C95994	129	133	138	146						
C95995	111	SPILLED	162	118	121	119	121	113	109	109
C95996	122	126	127	127	119	SPILLED	SPILLED	SPILLED	SPILLED	SPILLED
C95997	111	107	114	115	126	113	161	163	119	113
C95998	120	112	119	119	166	171	157	171	163	181
C95999	98	144	145	163	156	156	151		SPILLED	SPILLED
C96000	SPILLED	145	158	SPILLED						
GROUP: FEMALE 2 - 3 PPM										
C96001	SPILLED	SPILLED	SPILLED	SPILLED	SPILLED	121	176	SPILLED	SPILLED	107
C96002	103	106	104	121	106	109	111	119	113	106
C96003	106	113	110	107	105	106	106	SPILLED	109	109
C96004	169	45	127	SPILLED	125	138	134	140	144	166
C96005	125	120	122	125	119	125	127	161	129	120
C96006	SPILLED	SPILLED	SPILLED	SPILLED	112	SPILLED	SPILLED	123	SPILLED	SPILLED
C96007	119	122	133	137	SPILLED	132	149	137	133	136
C96008	148	SPILLED	144	148	113	145	137	146	135	136

001000

Appendix 4

Individual Food Consumption Data (g)

PAGE: 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WEEK 1	WEEK 2	WEEK 3	WEEK 4	WEEK 5	WEEK 6	WEEK 7	WEEK 8	WEEK 9	WEEK 10
GROUP: FEMALE 2 - 3 PPM										
C96009	115	SPIILLED	SPIILLED	108	99	147	171	125	119	104
C96010	117	122	127	137	124	141	SPIILLED	139	151	136
C96011	133	128	148	153						
C96012	113	143	SPIILLED	128						
C96013	SPIILLED	152	167	150						
C96014	131	135	144	150						
C96015	124	129	132	137						
C96016	122	131	130	127						
C96017	136	131	153	151	131	131	128	130	NOT TAKEN	138
C96018	140	144	158	SPIILLED	SPIILLED	SPIILLED	SPIILLED	SPIILLED	138	146
C96019	131	143	137	137	147	145	149	146	142	143
C96020	SPIILLED	120	126	126	126	120	128	125	117	114
GROUP: FEMALE 3 - 30 PPM										
C96021	SPIILLED	178	140	142	134	152	SPIILLED	127	SPIILLED	117
C96022	131	SPIILLED	149	150	SPIILLED	141	140	127	131	127
C96023	117	125	131	137	113	154	131	138	131	130
C96024	129	126	129	134	117	132	SPIILLED	131	121	121
C96025	133	123	135	135	120	137	129	134	139	127
C96026	SPIILLED	SPIILLED	SPIILLED	SPIILLED	SPIILLED	155	SPIILLED	SPIILLED	SPIILLED	171
C96027	208	SPIILLED	188	SPIILLED	SPIILLED	181	SPIILLED	SPIILLED	164	162
C96028	124	137	132	135	120	128	138	137	139	134
C96029	129	SPIILLED	138	135	115		SPIILLED	158	SPIILLED	220
C96030	131	130	139	139	130	139	146	130	133	130
C96031	135	SPIILLED	163	SPIILLED						
C96032	124	140	128	131						
C96033	124	129	133	133						
C96034	162	139	139	164						
C96035	149	151	136	142						
C96036	117	118	122	118	SPIILLED	159	SPIILLED	138	SPIILLED	SPIILLED

Appendix 4

Individual Food Consumption Data (g)

PAGE: 6

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL-
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WEEK 1	WEEK 2	WEEK 3	WEEK 4	WEEK 5	WEEK 6	WEEK 7	WEEK 8	WEEK 9	WEEK 10
GROUP: FEMALE 3 - 30 PPM										
C96037	130	127	178	SPILLED	125	130	129	136	127	122
C96038	120	161	SPILLED	SPILLED	SPILLED	194	SPILLED	SPILLED	SPILLED	149
C96039	185	SPILLED	SPILLED	SPILLED	SPILLED	SPILLED	SPILLED	SPILLED	225	SPILLED
C96040	SPILLED	SPILLED	SPILLED	SPILLED	SPILLED	SPILLED	SPILLED	SPILLED	SPILLED	SPILLED
GROUP: FEMALE 4 - 100 PPM										
C96041	135	110	109	120	106	125	117	113	111	110
C96042	100	92	96	99	99	107	106	109	104	96
C96043	107	104	103	105	102	109	105	102	104	106
C96044	136	127	137	142	132	142	129	141	135	128
C96045	118	128	124	145	124	148	142	140	146	140
C96046	SPILLED	128	127	130	128	136	SPILLED	151	149	125
C96047	108	107	105	99	102	106	104	109	103	100
C96048	128	111	102	111	100	110	102	111	114	115
C96049	101	SPILLED	99	102	94	105	98	106	108	102
C96050	133	SPILLED	124	120	104	117	114	109	111	108
C96051	101	107	101	106						
C96052	149	126	140	128						
C96053	113	108	118	121						
C96054	SPILLED	SPILLED	SPILLED	151						
C96055	105	100	104	108	112	111	103	107	101	102
C96056	109	108	110	116	149	137	132	133	124	135
C96057	175	133	130	159	127	125	129	124	121	127
C96058	124	133	124	121	121	124	125	SPILLED	SPILLED	SPILLED
C96059	118	143	107	121	121	124	125	121	123	115
C96060	129	123	116	120	122	115	125			

Appendix 4

Individual Food Consumption Data (g)

PAGE: 7

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WEEK 11	WEEK 12	WEEK 13
GROUP: MALE 1 - 0 PPM			
C95901	167	171	172
C95902	180	184	180
C95903	176	182	179
C95904	216	213	198
C95905	244	SPILLED	177
C95906	192	199	190
C95907	163	168	SPILLED
C95908	194	195	194
C95909	157	157	157
C95910	SPILLED	198	208
C95916	182	184	182
C95917	179	183	SPILLED
C95918	179	181	173
C95919	260	SPILLED	241
C95920	187	186	181

GROUP: MALE 2 - 3 PPM			
C95921	208	204	203
C95922	174	176	171
C95923	188	185	169
C95924	184	188	184
C95925	190	199	193
C95926	158	153	154
C95927	177	176	167
C95928	194	197	178
C95929	221	207	189
C95930	177	184	177
C95936	179	161	172
C95937	154	140	181
C95938	195	199	188
C95939	168	168	166

C01003

Appendix 4

Individual Food Consumption Data (g)

PAGE: 8

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WEEK 11	WEEK 12	WEEK 13
GROUP: MALE 2 - 3 PPM			
C95940	184	184	175
GROUP: MALE 3 - 30 PPM			
C95941	147	145	139
C95942	166	169	181
C95943	175	181	171
C95944	162	163	158
C95945	183	182	188
C95946	SPIILLED	174	167
C95947	143	141	138
C95948	182	178	182
C95949	192	175	7
C95950	184	190	188
C95956	233	224	207
C95957	164	167	165
C95958	184	182	173
C95959	183	185	178
C95960	161	166	156
GROUP: MALE 4 - 100 PPM			
C95961	177	178	178
C95962	159	164	157
C95963	182	183	175
C95964	177	178	178
C95965	163	169	168
C95966	175	172	164
C95967	177	177	166
C95968	197	199	175
C95969	169	170	170

001004

Appendix 4

Individual Food Consumption Data (g)

PAGE: 9

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WEEK 11	WEEK 12	WEEK 13
GROUP: MALE 4 - 100 PPM			
C95970	163	163	157
C95976	SPILLED	146	147
C95977	178	175	171
C95978	170	175	169
C95979	165	172	173
C95980	174	174	176
GROUP: FEMALE 1 - 0 PPM			
C95981	175	123	182
C95982	116	109	109
C95983	SPILLED	165	171
C95984	139	124	128
C95985	134	131	135
C95986	130	125	111
C95987	125	117	122
C95988	117	131	114
C95989	SPILLED	SPILLED	SPILLED
C95990	139	148	117
C95996	111	111	109
C95997	SPILLED	213	SPILLED
C95998	SPILLED	122	SPILLED
C95999	176	168	160
C96000	SPILLED	SPILLED	SPILLED
GROUP: FEMALE 2 - 3 PPM			
C96001	110	104	101
C96002	112	114	105
C96003	111	114	107
C96004	134	138	137

001005

Appendix 4

Individual Food Consumption Data (g)

PAGE: 10

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL-
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WEEK 11	WEEK 12	WEEK 13
GROUP: FEMALE 2 - 3 PPM			
C96005	120	116	113
C96006	SPILLED	SPILLED	SPILLED
C96007	141	149	136
C96008	132	130	126
C96009	SPILLED	129	SPILLED
C96010	135	150	130
C96017	118	127	97
C96018	123	147	SPILLED
C96019	139	143	150
C96020	110	119	113
GROUP: FEMALE 3 - 30 PPM			
C96021	195	127	125
C96022	132	121	126
C96023	125	121	121
C96024	123	123	118
C96025	125	107	115
C96026	SPILLED	250	220
C96027	193	183	148
C96028	116	120	124
C96029	SPILLED	177	SPILLED
C96030	122	123	125
C96036	SPILLED	129	212
C96037	128	128	134
C96038	139	134	130
C96039	197	172	162
C96040	SPILLED	174	SPILLED

001006

Appendix 4

Individual Food Consumption Data (g)

PAGE: 11

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WEEK 11	WEEK 12	WEEK 13
GROUP: FEMALE 4 - 100 PPM			
C96041	108	106	103
C96042	97	99	96
C96043	103	99	96
C96044	131	129	129
C96045	133	137	137
C96046	123	122	128
C96047	97	100	105
C96048	118	118	125
C96049	106	105	103
C96050	107	112	106
C96056	97	96	101
C96057	SPILLED	132	131
C96058	119	120	197
C96059	SPILLED	SPILLED	SPILLED
C96060	118	116	112

001007

Appendix 4

Individual Test Material Consumption Data (mg/kg/day)

PAGE: 1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WEEK 1	WEEK 2	WEEK 3	WEEK 4	WEEK 5	WEEK 6	WEEK 7	WEEK 8	WEEK 9	WEEK 10
GROUP: MALE 2 - 3 PPM										
C95921	0.3	0.3	0.2	0.2	0.2	0.2	0.2	SPILLED	0.2	0.2
C95922	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
C95923	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
C95924	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1
C95925	0.3	0.3	SPILLED	0.2	0.2	0.2	0.2	0.2	0.2	0.2
C95926	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
C95927	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
C95928	0.4	0.3	0.2	0.2	SPILLED	0.2	0.2	SPILLED	0.2	0.2
C95929	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
C95930	0.3	0.3	0.3	0.2	0.2	0.2	NOT TAKEN	0.2	0.2	0.2
C95931	0.3	SPILLED	0.3	SPILLED	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95932	0.3	0.3	0.2	0.2	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95933	0.3	0.3	0.2	0.2	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95934	0.3	0.3	0.2	0.2	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95935	0.3	0.3	0.2	0.2	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95936	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.1
C95937	0.3	0.2	0.2	0.2	0.2	0.2	0.2	SPILLED	0.2	0.2
C95938	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
C95939	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2
C95940	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
GROUP: MALE 3 - 30 PPM										
C95941	3.0	2.6	2.2	2.2	1.9	1.9	1.8	1.7	1.5	1.5
C95942	3.0	2.6	2.2	2.1	1.8	2.0	1.7	1.7	1.7	1.7
C95943	3.0	2.6	2.3	2.2	1.7	1.9	1.8	1.7	1.7	1.5
C95944	3.2	2.7	2.4	2.2	1.9	2.1	1.9	1.8	1.7	1.6
C95945	3.1	2.7	2.4	2.2	2.0	2.1	2.0	1.8	1.7	1.6
C95946	3.2	2.6	2.4	2.3	2.0	2.0	1.9	1.9	1.7	1.6
C95947	3.2	2.5	2.2	2.1	1.7	1.9	1.7	1.6	1.5	1.4
C95948	3.0	2.6	2.5	2.2	1.9	1.9	1.8	1.8	1.7	1.6

001008

Appendix 4

Individual Test Material Consumption Data (mg/kg/day)

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 2

ANIMAL NUMBER	WEEK 1	WEEK 2	WEEK 3	WEEK 4	WEEK 5	WEEK 6	WEEK 7	WEEK 8	WEEK 9	WEEK 10
GROUP: MALE 3 - 30 PPM										
C95949	3.0	2.6	2.4	2.2	1.8	2.0	1.9	1.8	1.7	1.6
C95950	3.0	2.6	2.4	2.1	1.8	2.0	1.9	1.8	1.8	1.6
C95951	2.9	2.4	2.2	2.0	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95952	2.9	2.5	2.3	2.2	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95953	3.3	2.8	2.5	2.3	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95954	3.1	2.8	2.5	2.4	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95955	3.6	2.9	2.6	2.5	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95956	3.8	SPILLED	2.7	2.6	SPILLED	2.5	2.3	SPILLED	1.9	1.9
C95957	2.8	2.5	2.3	2.1	2.0	1.9	1.8	1.7	1.6	1.5
C95958	3.1	2.6	2.4	2.2	2.0	2.1	1.8	1.8	1.8	1.7
C95959	3.2	2.8	2.5	2.4	2.2	2.2	2.0	2.0	1.9	1.8
C95960	3.0	2.7	2.3	2.3	2.1	1.8	1.8	1.7	1.6	1.5
GROUP: MALE 4 - 100 PPM										
C95961	10.2	8.7	7.7	7.2	6.0	6.7	6.2	5.9	5.6	5.2
C95962	9.9	8.8	7.7	7.2	6.3	6.9	6.5	6.0	5.9	5.6
C95963	10.2	8.5	7.5	6.9	6.2	6.6	6.1	5.9	5.7	5.3
C95964	10.1	9.4	7.7	7.6	6.3	6.5	6.4	6.0	5.7	5.4
C95965	10.3	9.1	8.1	7.6	6.6	7.1	6.5	6.1	6.0	5.6
C95966	10.2	9.0	8.1	7.4	6.2	6.9	6.5	6.1	5.9	5.6
C95967	10.6	9.1	7.9	7.3	6.1	6.6	6.5	6.1	5.9	5.7
C95968	11.6	9.3	8.3	7.8	6.4	6.9	6.5	5.3	5.2	6.0
C95969	10.4	8.6	7.6	7.1	5.8	6.5	6.2	5.9	5.6	5.4
C95970	9.5	9.3	SPILLED	7.2	6.2	6.8	6.2	6.0	5.5	5.4
C95971	10.0	8.6	7.7	6.4	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95972	10.1	8.7	7.6	6.9	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95973	11.3	9.6	8.3	7.9	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95974	10.4	9.3	7.9	7.4	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95975	9.8	8.5	7.5	7.2	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95976	10.4	9.1	7.9	7.3	6.7	6.6	6.4	6.0	5.8	5.3

001009

Appendix 4

Individual Test Material Consumption Data (mg/kg/day)

PAGE: 3

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WEEK 1	WEEK 2	WEEK 3	WEEK 4	WEEK 5	WEEK 6	WEEK 7	WEEK 8	WEEK 9	WEEK 10
GROUP: MALE 4 - 100 PPM										
C95977	10.0	9.0	7.7	7.4	6.7	6.7	6.5	6.1	6.0	5.6
C95978	9.9	8.4	7.2	6.8	6.4	6.1	5.9	5.8	5.8	5.6
C95979	10.6	9.0	8.0	7.3	7.1	6.4	6.4	5.9	5.9	5.7
C95980	10.2	8.9	7.8	7.1	6.8	6.6	6.5	6.2		
GROUP: FEMALE 2 - 3 PPM										
C96001	SPILLED	SPILLED	SPILLED	SPILLED	SPILLED	0.2	0.3	SPILLED	SPILLED	0.2
C96002	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2
C96003	0.3	0.3	0.2	0.2	0.2	0.2	0.2	SPILLED	0.2	0.2
C96004	0.4	0.1	0.3	SPILLED	0.2	0.2	0.2	0.2	0.2	0.3
C96005	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
C96006	SPILLED	SPILLED	SPILLED	SPILLED	0.2	SPILLED	SPILLED	0.2	SPILLED	SPILLED
C96007	0.3	0.3	0.3	0.3	0.2	0.2	0.3	0.2	0.2	0.2
C96008	0.4	SPILLED	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2
C96009	0.3	SPILLED	0.3	0.3	0.2	0.3	0.4	0.3	0.2	0.2
C96010	0.3	0.3	0.3	0.3	0.2	0.2	SPILLED	0.2	0.2	0.2
C96011	0.3	0.3	0.3	0.3	0.2	0.2	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96012	0.3	0.3	0.3	0.3	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96013	SPILLED	0.3	0.3	0.3	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96014	0.3	0.3	0.3	0.3	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96015	0.3	0.3	0.3	0.3	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96016	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
C96017	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2
C96018	0.3	0.3	0.3	SPILLED	SPILLED	SPILLED	SPILLED	SPILLED	0.2	0.2
C96019	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2
C96020	SPILLED	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2

001010

Appendix 4

Individual Test Material Consumption Data (mg/kg/day)

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 4

ANIMAL NUMBER	WEEK 1	WEEK 2	WEEK 3	WEEK 4	WEEK 5	WEEK 6	WEEK 7	WEEK 8	WEEK 9	WEEK 10
------------------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	------------

GROUP: FEMALE 3 - 30 PPM

C96021	SPILLED	4.4								
C96022	3.1	SPILLED	3.1	2.8	2.6	2.9	SPILLED	2.3	SPILLED	2.0
C96023	2.8	2.7	2.6	2.6	2.1	2.4	2.3	2.0	2.1	2.0
C96024	3.0	2.6	2.4	2.4	2.1	2.3	SPILLED	2.2	2.2	2.1
C96025	3.1	2.7	2.8	2.5	2.2	2.5	2.2	2.3	1.9	1.9
C96026	SPILLED		SPILLED	SPILLED	SPILLED	3.0	SPILLED	2.3	2.3	2.0
C96027	5.0	SPILLED	3.5	SPILLED	SPILLED	2.9	SPILLED	SPILLED	SPILLED	3.1
C96028	3.1	3.0	2.6	2.5	2.2	2.2	2.3	2.2	2.4	2.3
C96029	3.4	SPILLED	3.4	2.7	2.2	SPILLED	SPILLED	2.2	2.2	2.1
C96030	3.5	3.1	3.0	2.8	2.3	2.6	2.6	2.3	2.3	2.2
C96031	3.4	SPILLED	3.4	SPILLED	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96032	3.2	3.2	2.6	2.5	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96033	3.0	2.8	2.7	2.4	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96034	3.9	2.9	2.6	2.8	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96035	3.4	3.0	2.5	2.5	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96036	2.9	2.6	2.4	2.2	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96037	3.2	2.8	3.6	2.2	SPILLED	2.8	SPILLED	2.3	SPILLED	SPILLED
C96038	3.0	3.6	SPILLED	SPILLED	2.3	2.3	2.3	2.1	2.1	2.0
C96039	4.7	SPILLED	SPILLED	SPILLED	SPILLED	3.3	SPILLED	SPILLED	SPILLED	2.3
C96040	SPILLED	SPILLED	SPILLED	SPILLED	SPILLED	SPILLED	SPILLED	2.6	4.2	SPILLED

GROUP: FEMALE 4 - 100 PPM

C96041	12.7	9.1	8.3	8.7	7.6	8.8	8.0	7.4	7.1	7.0
C96042	8.2	6.8	6.6	6.6	6.4	6.7	6.6	6.6	6.2	5.7
C96043	9.5	8.3	7.5	7.3	6.9	7.2	6.8	6.5	6.6	6.6
C96044	10.6	8.7	8.6	8.5	7.6	7.9	7.1	7.7	7.2	6.7
C96045	9.7	9.4	8.3	9.4	7.6	8.8	8.2	7.9	8.1	7.5
C96046	SPILLED	SPILLED	8.3	8.0	7.5	7.7	SPILLED	8.1	7.8	6.4
C96047	9.3	8.6	7.8	7.1	7.2	7.3	7.0	7.2	6.6	6.4
C96048	10.6	8.2	7.0	7.3	6.4	6.9	6.4	6.8	6.8	6.7

001011

Appendix 4

Individual Test Material Consumption Data (mg/kg/day)

PAGE: 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WEEK 1	WEEK 2	WEEK 3	WEEK 4	WEEK 5	WEEK 6	WEEK 7	WEEK 8	WEEK 9	WEEK 10
GROUP: FEMALE 4 - 100 PPM										
C96049	9.6	SPILED	8.2	7.9	7.1	8.0	7.4	7.8	7.7	7.2
C96050	12.1	SPILED	9.2	8.5	7.2	8.0	7.6	7.2	7.3	6.9
C96051	9.1	8.8	7.5	7.6	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96052	11.5	9.0	9.3	8.1	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96053	10.1	8.9	8.8	8.2	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96054	SPILED	SPILED	SPILED	10.6	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96055	9.3	8.4	8.1	7.9	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96056	9.5	8.5	8.2	8.3	7.6	7.3	6.7	7.0	6.4	6.4
C96057	13.8	9.2	8.4	9.9	8.8	7.8	7.3	7.3	6.7	7.2
C96058	9.8	9.1	7.8	7.5	7.7	7.3	7.4	7.0	6.7	7.1
C96059	11.1	12.0	8.7	9.4	8.8	8.7	8.6	SPILED	SPILED	SPILED
C96060	10.4	8.8	7.6	7.6	7.3	6.7	7.3	6.8	6.8	6.2

001012

Appendix 4

Individual Test Material Consumption Data (mg/kg/day)

PAGE: 6

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WEEK 11	WEEK 12	WEEK 13
------------------	------------	------------	------------

GROUP: MALE 2 - 3 PPM

C95921	0.2	0.2	0.2
C95922	0.2	0.2	0.1
C95923	0.2	0.2	0.1
C95924	0.2	0.2	0.2
C95925	0.2	0.2	0.2
C95926	0.1	0.1	0.1
C95927	0.2	0.2	0.1
C95928	0.2	0.2	0.1
C95929	0.2	0.2	0.1
C95930	0.2	0.2	0.1
C95931	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95932	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95933	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95934	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95935	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95936	0.1	0.1	0.1
C95937	0.1	0.1	0.2
C95938	0.2	0.2	0.1
C95939	0.2	0.2	0.1
C95940	0.2	0.2	0.1

GROUP: MALE 3 - 30 PPM

C95941	1.5	1.4	1.3
C95942	1.7	1.7	1.7
C95943	1.5	1.5	1.4
C95944	1.5	1.5	1.4
C95945	1.6	1.5	1.6
C95946	SPILLED	1.5	1.4
C95947	1.3	1.3	1.3
C95948	1.6	1.5	1.5

001013

Appendix 4

Individual Test Material Consumption Data (mg/kg/day)

PAGE: 7

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WEEK 11	WEEK 12	WEEK 13
------------------	------------	------------	------------

GROUP: MALE 3 - 30 PPM

C95949	1.5	1.4	0.1
C95950	1.7	1.7	1.6
C95951	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95952	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95953	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95954	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95955	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95956	1.9	1.8	1.6
C95957	1.5	1.5	1.4
C95958	1.6	1.6	1.5
C95959	1.7	1.7	1.6
C95960	1.5	1.5	1.4

GROUP: MALE 4 - 100 PPM

C95961	5.0	5.0	4.9
C95962	5.2	5.2	4.8
C95963	5.4	5.3	4.9
C95964	5.4	5.3	5.2
C95965	5.5	5.5	5.4
C95966	5.5	5.3	4.9
C95967	5.6	5.4	5.0
C95968	6.1	6.0	5.1
C95969	5.1	5.0	4.9
C95970	5.1	5.0	4.8
C95971	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95972	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95973	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95974	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95975	NOT TAKEN	NOT TAKEN	NOT TAKEN
C95976	SPILLED	5.1	5.1

001014

Appendix 4

Individual Test Material Consumption Data (mg/kg/day)

PAGE: 8

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WEEK 11	WEEK 12	WEEK 13
GROUP: MALE 4 - 100 PPM			
C95977	5.5	5.3	5.0
C95978	5.2	5.2	5.0
C95979	5.3	5.4	5.3
C95980	5.6	5.5	5.4
GROUP: FEMALE 2 - 3 PPM			
C96001	0.2	0.2	0.2
C96002	0.2	0.2	0.2
C96003	0.2	0.2	0.2
C96004	0.2	0.2	0.2
C96005	0.2	0.2	0.2
C96006	SPILLED	SPILLED	SPILLED
C96007	0.2	0.2	0.2
C96008	0.2	0.2	0.2
C96009	SPILLED	0.2	SPILLED
C96010	0.2	0.2	0.2
C96011	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96012	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96013	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96014	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96015	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96016	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96017	0.2	0.2	0.1
C96018	0.2	0.2	SPILLED
C96019	0.2	0.2	0.2
C96020	0.2	0.2	0.2

001015

Appendix 4

Individual Test Material Consumption Data (mg/kg/day)

PAGE: 9

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL WEEK WEEK WEEK
NUMBER 11 12 13

GROUP: FEMALE 3 - 30 PPM

C96021	3.3	2.1	2.1
C96022	2.0	1.8	1.9
C96023	2.0	1.9	1.8
C96024	1.9	1.7	1.8
C96025	2.0	4.2	3.7
C96026	SPILLED	2.5	2.0
C96027	2.7	1.8	1.8
C96028	1.8	3.0	SPILLED
C96029	SPILLED	2.0	2.0
C96030	2.0	NOT TAKEN	NOT TAKEN
C96031	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96032	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96033	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96034	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96035	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96036	SPILLED	2.0	3.3
C96037	2.1	2.0	2.1
C96038	2.1	2.0	1.9
C96039	3.1	2.6	2.4
C96040	SPILLED	3.1	SPILLED

GROUP: FEMALE 4 - 100 PPM

C96041	6.8	6.5	6.3
C96042	5.7	5.7	5.5
C96043	6.2	5.9	5.7
C96044	6.7	6.5	6.5
C96045	7.0	7.2	7.1
C96046	6.3	6.1	6.4
C96047	6.2	6.3	6.6
C96048	6.9	6.9	7.1

001016

Appendix 4

Individual Test Material Consumption Data (mg/kg/day)

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 10

ANIMAL NUMBER	WEEK 11	WEEK 12	WEEK 13
------------------	------------	------------	------------

GROUP: FEMALE 4 - 100 PPM

C96049	7.4	7.2	7.0
C96050	6.7	6.9	6.4
C96051	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96052	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96053	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96054	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96055	NOT TAKEN	NOT TAKEN	NOT TAKEN
C96056	6.1	6.1	6.3
C96057	SPILLED	6.9	6.7
C96058	6.6	6.5	10.4
C96059	SPILLED	SPILLED	SPILLED
C96060	6.3	6.1	5.9

001017

APPENDIX 5

Individual Clinical Hematology Data

Individual Clinical Chemistry Data

Individual Clinical Urinalysis Data

C01018

Appendix 5

Individual Clinical Hematology Data

Males Week 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	RBC X10 ⁶ /μL	HGB G/DL	HCT %	MCV FL	MCH PG	MCHC %	PLT X10 ³ /μL
Group: 1 Dose Level: 0 Dosage Unit: ppm MeFOS							
C95901	8.89	17.3	50.1	56.4	19.4	34.5	1186
C95902	8.64	16.6	48.1	55.7	19.2	34.6	1223
C95903	8.23	16.4	47.5	57.6	19.9	34.6	1208
C95904	7.52	15.3	45.5	60.5	20.4	33.7	1227
C95905	7.92	15.3	42.7	54.0	19.3	35.7	1235
C95906	7.90	15.7	46.2	58.5	19.8	33.9	1221
C95907	8.18	16.3	46.7	57.1	19.9	34.9	1084
C95908	7.86	16.0	47.4	60.3	20.4	33.7	1086
C95909	8.63	17.0	49.1	56.9	19.7	34.7	1099
C95910	8.06	16.1	46.3	57.4	20.0	34.7	1162
MEAN	8.18	16.2	47.0	57.4	19.8	34.5	1173
S.D.	.424	.66	2.04	1.97	.42	.61	61.5
N	10	10	10	10	10	10	10
Group: 2 Dose Level: 3 Dosage Unit: ppm MeFOS							
C95921	8.38	16.8	49.2	58.7	20.0	34.1	1136
C95922	7.84	15.6	45.6	58.2	19.9	34.2	1297
C95923	8.57	17.0	48.6	56.8	19.9	35.0	1400
C95924	7.55	16.3	46.8	61.9	21.5	34.7	1177
C95925	8.09	16.2	46.7	57.7	20.1	34.8	1108
C95926	8.29	16.0	46.2	55.8	19.3	34.6	1172
C95927	8.07	16.2	47.9	59.3	20.0	33.8	1165
C95928	8.99	17.1	51.1	56.8	19.0	33.5	1143
C95929	7.76	16.1	46.0	59.3	20.8	35.0	917
C95930	7.89	16.4	47.1	59.7	20.7	34.7	993
MEAN	8.14	16.4	47.5	58.4	20.1	34.4	1151
S.D.	.427	.47	1.70	1.77	.73	.51	136.0
N	10	10	10	10	10	10	10

001019

Appendix 5

Individual Clinical Hematology Data

Males Week 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	RBC X10 ⁶ /μL	HGB G/DL	HCT %	MCV FL	MCH PG	MCHC %	PLT X10 ³ /μL
Group: 3	Dose Level: 30	Dosage Unit: ppm MeFOS					
C95941	8.09	16.1	45.9	56.7	19.9	35.1	1130
C95942	9.02	16.9	49.9	55.3	18.7	33.9	1133
C95943	8.19	15.9	45.9	56.1	19.4	34.6	1027
C95944	8.23	16.7	48.2	58.6	20.2	34.5	891
C95945	8.17	16.2	46.5	56.9	19.8	34.9	1074
C95946	7.68	16.2	46.0	59.9	21.1	35.2	893
C95947	8.10	16.0	45.3	55.9	19.8	35.4	1151
C95948	8.37	16.6	47.7	56.9	19.9	34.9	1027
C95949	9.01	16.7	49.4	54.8	18.6	33.9	958
C95950	7.56	15.8	45.8	60.6	20.9	34.4	1098
MEAN	8.24	16.3	47.1	57.2	19.8	34.7	1038
S.D.	.477	.38	1.64	1.93	.81	.52	96.9
N	10	10	10	10	10	10	10
Group: 4	Dose Level: 100	Dosage Unit: ppm MeFOS					
C95961	7.59	15.7	46.0	60.6	20.7	34.1	1141
C95962	8.16	16.0	46.4	56.9	19.6	34.5	1097
C95963	8.03	16.1	46.9	58.5	20.1	34.4	1177
C95964	7.64	16.1	46.1	60.3	21.1	34.9	986
C95965	8.33	16.9	48.7	58.5	20.2	34.6	1166
C95966	8.11	15.3	44.6	55.0	18.9	34.4	948
C95967	8.32	16.1	46.9	56.3	19.4	34.4	1001
C95968	7.57	14.9	43.2	57.1	19.7	34.4	1296
C95969	7.90	14.8	43.9	55.6	18.7	33.6	1032
C95970	8.29	16.8	47.7	57.5	20.3	35.3	970
MEAN	7.99	15.9	46.0	57.6	19.9	34.5	1081
S.D.	.303	.71	1.71	1.86	.76	.45	112.7
N	10	10	10	10	10	10	10

001020

Appendix 5

Individual Clinical Hematology Data

Males Week 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WBC $\times 10^3/\mu\text{L}$	N-SEG $\times 10^3/\mu\text{L}$	LYMPH $\times 10^3/\mu\text{L}$	MONO $\times 10^3/\mu\text{L}$	EOSIN $\times 10^3/\mu\text{L}$	BASO $\times 10^3/\mu\text{L}$	N-SEG% N-SEG% $\times 10^3/\mu\text{L}$	LYMPH% LYMPH% $\times 10^3/\mu\text{L}$	MONO% MONO% $\times 10^3/\mu\text{L}$	EOSIN% EOSIN% $\times 10^3/\mu\text{L}$	BASO% BASO% $\times 10^3/\mu\text{L}$
Group: 1	Dose Level: 0	Dosage Unit: ppm MeFOS									
C95901	8.6	.5	7.7	.4	.1	.0	6	89	4	1	0
C95902	9.7	.6	8.6	.4	.0	.0	6	89	5	0	0
C95903	6.7	.8	5.7	.2	.1	.0	11	85	3	1	0
C95904	11.9	.4	10.7	.3	.1	.0	3	90	6	1	0
C95905	8.4	.6	7.3	.3	.1	.0	8	87	4	1	0
C95906	6.9	.5	6.0	.3	.1	.0	7	87	4	1	0
C95907	8.1	.5	7.1	.4	.1	.0	6	88	5	1	1
C95908	12.7	.8	11.4	.4	.1	.0	6	90	3	1	0
C95909	8.1	.7	6.8	.5	.1	.0	9	84	6	1	0
C95910	9.5	.7	8.1	.6	.1	.0	7	85	7	1	0
MEAN	9.1	.6	7.9	.4	.1	.0	7	87	5	1	0
S.D.	1.96	.14	1.86	.15	.03	.00	2.1	2.2	1.3	.3	.3
N	10	10	10	10	10	10	10	10	10	10	10
Group: 2	Dose Level: 3	Dosage Unit: ppm MeFOS									
C95921	8.4	.5	7.5	.4	.1	.0	6	88	4	1	0
C95922	7.9	.6	6.6	.5	.2	.0	7	84	6	3	0
C95923	9.3	.6	8.2	.5	.1	.0	6	88	5	1	0
C95924	8.5	.6	7.2	.5	.1	.0	8	85	6	1	0
C95925	9.8	.5	8.9	.3	.1	.0	5	91	3	1	0
C95926	12.6	.7	10.9	.9	.1	.0	5	86	7	1	0
C95927	8.6	.7	7.2	.5	.1	.0	8	84	6	2	0
C95928	7.3	.4	6.8	.1	.0	.0	5	93	1	1	0
C95929	7.4	.7	6.2	.4	.1	.0	9	84	6	1	0
C95930	11.5	.5	10.6	.4	.1	.0	4	92	3	1	0
MEAN	9.1	.6	8.0	.4	.1	.0	6	88	5	1	0
S.D.	1.74	.10	1.64	.20	.05	.00	1.6	3.5	1.9	.7	.0
N	10	10	10	10	10	10	10	10	10	10	10

001021

Appendix 5

Individual Clinical Hematology Data

Males Week 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WBC $\times 10^3/\mu\text{L}$	N-SEG $\times 10^3/\mu\text{L}$	LYMPH $\times 10^3/\mu\text{L}$	MONO $\times 10^3/\mu\text{L}$	EOSIN $\times 10^3/\mu\text{L}$	BASO $\times 10^3/\mu\text{L}$	N-SEG%	LYMPH%	MONO%	EOSIN%	BASO%
Group: 3	Dose Level: 30	Dosage Unit: ppm MeFOS									
C95941	14.3	.8	12.8	.5	.2	.0	6	89	4	1	0
C95942	8.4	.4	7.6	.3	.0	.0	5	90	4	1	0
C95943	6.6	.4	5.8	.3	.1	.0	7	87	5	1	0
C95944	10.1	.5	9.2	.4	.1	.0	5	91	4	1	0
C95945	7.3	.6	6.4	.3	.1	.0	8	88	4	1	0
C95946	9.8	.3	9.1	.3	.1	.1	3	92	3	1	1
C95947	12.0	.7	10.7	.4	.1	.1	6	89	4	1	1
C95948	13.2	.6	12.0	.4	.1	.0	4	92	3	1	0
C95949	9.1	.5	8.2	.3	.0	.0	5	90	4	0	0
C95950	9.5	.5	8.3	.7	.1	.0	5	87	7	1	0
MEAN	10.0	.5	9.0	.4	.1	.0	5	90	4	1	0
S.D.	2.48	.15	2.27	.13	.06	.04	1.4	1.8	1.1	.3	.4
N	10	10	10	10	10	10	10	10	10	10	10
Group: 4	Dose Level: 100	Dosage Unit: ppm MeFOS									
C95961	14.4	.5	13.2	.6	.1	.1	3	92	4	0	1
C95962	10.2	1.0	8.4	.7	.1	.1	10	82	7	1	1
C95963	10.1	.5	8.9	.6	.1	.0	5	88	6	1	0
C95964	9.8	.8	8.5	.4	.1	.0	8	86	4	1	0
C95965	6.4	.7	5.2	.4	.1	.0	11	82	6	1	0
C95966	14.0	1.4	11.6	.8	.2	.0	10	83	6	1	0
C95967	4.2	.5	3.3	.3	.1	.0	12	79	8	1	0
C95968	14.7	1.1	12.7	.7	.1	.1	7	86	5	1	1
C95969	13.4	1.5	10.8	.9	.1	.0	12	80	7	1	0
C95970	9.8	1.4	7.8	.4	.1	.0	15	79	5	1	0
MEAN	10.7	.9	9.0	.6	.1	.0	9	84	6	1	0
S.D.	3.51	.40	3.17	.20	.03	.05	3.6	4.2	1.3	.3	.5
N	10	10	10	10	10	10	10	10	10	10	10

001022

Appendix 5

Individual Clinical Hematology Data

Males Week 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	ANISO	POLY	POIK	HYP0	TOXNEUT
-----	-----	-----	-----	-----	-----
Group: 1	Dose Level: 0			Dosage Unit: ppm MeFOS	
C95901	-	-	-	-	-
C95902	-	-	-	-	-
C95903	-	-	-	-	-
C95904	-	-	-	-	-
C95905	-	-	-	-	-
C95906	-	-	-	-	-
C95907	-	-	-	-	-
C95908	-	-	-	-	-
C95909	-	-	-	-	-
C95910	-	-	-	-	-
Group: 2	Dose Level: 3			Dosage Unit: ppm MeFOS	
C95921	-	-	-	-	-
C95922	-	-	-	-	-
C95923	-	-	-	-	-
C95924	-	-	-	-	-
C95925	-	-	-	-	-
C95926	-	-	-	-	-
C95927	-	-	-	-	-
C95928	-	-	-	-	-
C95929	-	-	-	-	-
C95930	-	-	-	-	-

001023

001024

Covance 6329-225
3M T-6314.1

Appendix 5
Individual Clinical Hematology Data

Males Week 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	ANISO	POLY	POIK	HYPO	TOXNEUT

Group: 3	Dose Level: 30		Dosage Unit: ppm MeFOS		
C95941	-	-	-	-	-
C95942	-	-	-	-	-
C95943	-	-	-	-	-
C95944	-	-	-	-	-
C95945	-	-	-	-	-
C95946	-	-	-	-	-
C95947	-	-	-	-	-
C95948	-	-	-	-	-
C95949	-	-	-	-	-
C95950	-	-	-	-	-
Group: 4	Dose Level: 100		Dosage Unit: ppm MeFOS		
C95961	-	-	-	-	-
C95962	-	-	-	-	-
C95963	-	-	-	-	-
C95964	-	-	-	-	-
C95965	-	-	-	-	-
C95966	-	-	-	-	-
C95967	-	-	-	-	-
C95968	-	-	-	-	-
C95969	-	-	-	-	-
C95970	-	-	-	-	-

Appendix 5

Individual Clinical Hematology Data

Females Week 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDE
ETHANOL (N-NEFOSE, T-6314) IN RATS

ANIMAL NUMBER	RBC X10 ⁶ /μL	HGB G/DL	HCT %	MCV FL	MCH PG	MCHC %	PLT X10 ³ /μL
Group: 1 Dose Level: 0 Dosage Unit: ppm MeFOS							
C95981	7.36	15.2	44.3	60.3	20.7	34.3	1230
C95982	7.74	15.7	45.1	58.3	20.3	34.8	1254
C95983	7.53	16.2	46.0	61.1	21.5	35.2	1094
C95984	7.56	14.5	42.6	56.4	19.1	33.9	1137
C95985	7.75	15.8	45.9	59.3	20.4	34.4	1012
C95986	8.17	15.9	47.0	57.5	19.5	33.9	1085
C95987	7.75	15.8	45.5	58.8	20.4	34.7	913
C95988	7.67	15.5	44.8	58.4	20.2	34.5	1190
C95989	7.63	15.7	46.1	60.4	20.6	34.1	1056
C95990	7.82	16.3	46.9	60.0	20.9	34.9	1171
MEAN	7.70	15.7	45.4	59.1	20.4	34.5	1114
S.D.	.214	.51	1.31	1.45	.68	.43	104.4
N	10	10	10	10	10	10	10
Group: 2 Dose Level: 3 Dosage Unit: ppm MeFOS							
C96001	8.35	15.9	45.1	54.0	19.1	35.3	1297
C96002	8.59	16.2	47.2	54.9	18.8	34.3	1181
C96003	8.09	16.3	47.7	58.9	20.2	34.2	1107
C96004	6.82	14.3	40.9	60.0	21.0	35.1	1267
C96005	8.04	16.3	47.4	59.0	20.3	34.5	1139
C96006	8.26	16.1	46.7	56.5	19.5	34.5	1166
C96007	7.89	16.3	46.5	59.0	20.7	35.1	1106
C96008	7.97	16.3	45.6	57.3	20.4	35.7	1303
C96009	7.86	16.1	46.1	58.7	20.5	34.9	1083
C96010	8.42	16.2	48.3	57.4	19.2	33.5	1129
MEAN	8.03	16.0	46.1	57.6	20.0	34.7	1178
S.D.	.488	.61	2.08	1.95	.76	.64	82.3
N	10	10	10	10	10	10	10

001025

Appendix 5
Individual Clinical Hematology Data
Females Week 5
13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	RBC X10 ⁶ /μL	HGB G/DL	HCT %	MCV FL	MCH PG	MCHC %	PLT X10 ³ /μL
Group: 3	Dose Level: 30	Dosage Unit: ppm MeFOS					
C96021	8.61	16.6	48.3	56.1	19.3	34.4	1199
C96022	7.97	16.0	45.7	57.4	20.1	35.0	1084
C96023	7.34	15.2	43.4	59.1	20.7	35.1	1037
C96024	8.01	16.2	46.3	57.8	20.2	35.0	1029
C96025	8.17	15.7	45.2	55.4	19.2	34.8	1082
C96026	7.78	16.0	45.2	58.2	20.6	35.5	1124
C96027	7.86	16.3	46.6	59.3	20.7	35.0	1057
C96028	7.66	15.3	43.5	56.8	20.0	35.3	1414
C96029	8.04	15.6	44.5	55.4	19.5	35.1	1199
C96030	8.36	16.9	49.3	58.9	20.2	34.3	1007
MEAN	7.98	16.0	45.8	57.4	20.0	35.0	1123
S.D.	.357	.55	1.91	1.48	.56	.37	121.8
N	10	10	10	10	10	10	10
Group: 4	Dose Level: 100	Dosage Unit: ppm MeFOS					
C96041	7.44	14.5	41.9	56.4	19.5	34.6	1007
C96042	7.66	15.1	43.1	56.2	19.7	35.1	1191
C96043	8.63	16.5	47.5	55.1	19.1	34.6	1235
C96044	7.63	15.3	44.9	58.8	20.1	34.2	1059
C96045	8.13	15.7	45.9	56.5	19.3	34.2	1091
C96046	7.84	15.7	46.0	58.7	20.0	34.0	535
C96047	7.91	15.8	43.4	54.9	19.9	36.3	1151
C96048	7.76	15.1	44.1	56.8	19.5	34.4	1333
C96049	7.95	15.2	44.9	56.4	19.2	34.0	942
C96050	7.95	15.7	44.8	56.3	19.8	35.2	1032
MEAN	7.89	15.5	44.7	56.6	19.6	34.7	1058
S.D.	.326	.54	1.62	1.28	.34	.71	217.2
N	10	10	10	10	10	10	10

001026

Appendix 5
Individual Clinical Hematology Data

Females Week 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WBC X10 ³ /μL	N-SEG X10 ³ /μL	LYMPH X10 ³ /μL	MONO X10 ³ /μL	EOSIN X10 ³ /μL	BASO X10 ³ /μL	N-SEG%	LYMPH%	MONO%	EOSIN%	BASO%
Group: 1	Dose Level: 0										
C95981	8.2	1.1	6.4	.6	.1	.0	14	77	8	1	0
C95982	7.6	.7	6.3	.5	.1	.0	9	83	6	1	1
C95983	5.3	.4	4.6	.2	.0	.0	7	88	5	1	0
C95984	7.3	.2	6.8	.3	.0	.0	3	93	4	1	0
C95985	7.6	.5	6.6	.4	.1	.0	7	87	5	1	0
C95986	6.8	.2	6.3	.2	.0	.0	4	92	3	1	0
C95987	3.3	.2	2.9	.1	.0	.0	7	89	3	1	0
C95988	6.9	.5	5.9	.3	.1	.0	7	87	5	2	0
C95989	5.2	.3	4.5	.3	.0	.0	7	87	5	1	0
C95990	4.6	.2	4.1	.3	.0	.0	4	89	6	1	0
MEAN	6.3	.4	5.4	.3	.0	.0	7	87	5	1	0
S.D.	1.59	.29	1.32	.15	.05	.00	3.1	4.5	1.5	.3	.3
N	10	10	10	10	10	10	10	10	10	10	10
Group: 2	Dose Level: 3										
C96001	5.7	.5	5.1	.1	.1	.0	9	88	1	1	0
C96002	3.5	.3	3.0	.1	.0	.0	9	86	4	1	0
C96003	7.6	.7	6.3	.4	.2	.0	10	83	5	2	0
C96004	7.2	.3	6.5	.4	.1	.0	4	90	5	1	0
C96005	11.1	.3	10.6	.2	.0	.0	2	96	1	0	0
C96006	5.8	.4	5.0	.3	.0	.0	6	88	5	1	0
C96007	10.4	.4	9.7	.3	.1	.0	4	93	3	1	0
C96008	7.0	.4	6.2	.3	.1	.0	5	89	5	1	0
C96009	7.9	.8	6.6	.3	.1	.0	10	84	4	2	0
C96010	7.4	.8	6.4	.1	.0	.0	11	86	2	0	0
MEAN	7.4	.5	6.5	.2	.1	.0	7	88	4	1	0
S.D.	2.21	.20	2.20	.12	.07	.00	3.2	4.0	1.6	.7	.0
N	10	10	10	10	10	10	10	10	10	10	10

001027

Individual Clinical Hematology Data

Females
Week 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

[illegible]

Appendix 5

Individual Clinical Hematology Data

Females Week 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	ANISO	POLY	POIK	HYPO	TOXNEUT
-----	-----	-----	-----	-----	-----
Group: 1	Dose Level: 0		Dosage Unit: ppm MeFOS		
C95981	-	-	-	-	-
C95982	-	-	-	-	-
C95983	-	-	-	-	-
C95984	-	-	-	-	-
C95985	-	-	-	-	-
C95986	-	-	-	-	-
C95987	-	-	-	-	-
C95988	-	-	-	-	-
C95989	-	-	-	-	-
C95990	-	-	-	-	-
Group: 2	Dose Level: 3		Dosage Unit: ppm MeFOS		
C96001	-	-	-	-	-
C96002	-	-	-	-	-
C96003	-	-	-	-	-
C96004	-	-	-	-	-
C96005	-	-	-	-	-
C96006	-	-	-	-	-
C96007	-	-	-	-	-
C96008	-	-	-	-	-
C96009	-	-	-	-	-
C96010	-	-	-	-	-

001029

Appendix 5
Individual Clinical Hematology Data
Females Week 5
13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	ANISO	POLY	POIK	HYPO	TOXNEUT

Group: 3	Dose Level: 30		Dosage Unit: ppm MeFOS		
C96021	-	-	-	-	-
C96022	-	-	-	-	-
C96023	-	-	-	-	-
C96024	-	-	-	-	-
C96025	-	-	-	-	-
C96026	-	-	-	-	-
C96027	-	-	-	-	-
C96028	-	-	-	-	-
C96029	-	-	-	-	-
C96030	-	-	-	-	-
Group: 4	Dose Level: 100		Dosage Unit: ppm MeFOS		
C96041	-	-	-	-	-
C96042	-	-	-	-	-
C96043	-	-	-	-	-
C96044	-	-	-	-	-
C96045	-	-	-	-	-
C96046	-	-	-	-	-
C96047	-	-	-	-	-
C96048	-	-	-	-	-
C96049	-	-	-	-	-
C96050	-	-	-	-	-

001030

Appendix 5

Individual Clinical Hematology Data

Males Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDE
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	RBC X10 ⁶ /μL	HGB G/DL	HCT %	MCV FL	MCH PG	MCHC %	PLT X10 ³ /μL
Group: 1 Dose Level: 0 Dosage Unit: ppm MeFOS							
C95901	9.37	16.2	49.4	52.7	17.3	32.9	1131
C95902	8.66	15.1	46.4	53.5	17.5	32.6	1142
C95903	9.11	16.1	49.9	54.7	17.6	32.2	1001
C95904	7.96	14.8	45.9	57.6	18.5	32.2	1130
C95905	8.19	14.3	43.7	53.3	17.4	32.7	1174
C95906	8.16	14.4	45.5	55.7	17.7	31.7	1148
C95909	8.63	15.7	48.0	55.6	18.2	32.7	935
C95918	8.25	14.8	45.7	55.4	17.9	32.3	1072
C95919	8.36	15.3	46.3	55.4	18.3	33.0	1095
C95920	8.96	16.1	49.0	54.7	17.9	32.8	1390
MEAN	8.56	15.3	47.0	54.9	17.8	32.5	1122
S.D.	.463	.71	2.00	1.43	.40	.40	119.5
N	10	10	10	10	10	10	10
Group: 2 Dose Level: 3 Dosage Unit: ppm MeFOS							
C95921	8.71	15.4	47.4	54.4	17.7	32.5	1019
C95922	8.33	15.3	46.0	55.2	18.3	33.2	1225
C95924	7.14	14.2	43.1	60.4	19.8	32.9	1042
C95925	8.57	15.4	47.6	55.6	17.9	32.3	1038
C95926	8.91	15.4	47.1	52.9	17.3	32.7	1115
C95927	9.13	16.5	50.0	54.8	18.1	33.0	1124
C95928	9.29	16.0	49.2	53.0	17.2	32.5	1028
C95936	8.05	15.0	45.8	57.0	18.7	32.8	988
C95937	8.34	14.9	45.9	55.0	17.9	32.5	1003
C95939	7.72	14.5	44.9	58.1	18.7	32.2	1019
MEAN	8.42	15.3	46.7	55.6	18.2	32.7	1060
S.D.	.657	.67	2.02	2.31	.77	.32	73.0
N	10	10	10	10	10	10	10

001031

Appendix 5

Individual Clinical Hematology Data

Males Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	RBC X10 ⁶ /μL	HGB G/DL	HCT %	MCV FL	MCH PG	MCHC %	PLT X10 ³ /μL
Group: 3	Dose Level: 30	Dosage Unit: ppm MeFOS					
C95941	8.33	14.7	44.1	53.0	17.7	33.3	1015
C95942	9.62	16.0	50.5	52.5	16.6	31.6	1065
C95943	8.89	15.4	47.2	53.2	17.4	32.7	1109
C95944	8.53	15.5	47.5	55.7	18.2	32.7	932
C95947	9.15	15.2	47.1	51.5	16.6	32.3	1107
C95948	9.24	16.1	49.0	53.0	17.4	32.9	1006
C95949	8.68	14.6	46.3	53.4	16.8	31.6	636
C95950	8.26	15.4	45.6	55.2	18.7	33.9	986
C95958	8.07	14.4	44.2	54.8	17.9	32.6	1100
C95960	8.03	15.2	46.1	57.5	18.9	33.0	1045
MEAN	8.68	15.2	46.8	54.0	17.6	32.7	1000
S.D.	.535	.56	1.99	1.79	.82	.71	140.3
N	10	10	10	10	10	10	10
Group: 4	Dose Level: 100	Dosage Unit: ppm MeFOS					
C95961	8.07	14.7	44.6	55.3	18.3	33.0	957
C95962	8.58	14.9	45.4	52.9	17.4	32.8	992
C95963	8.15	14.6	44.6	54.8	17.9	32.7	1153
C95964	7.96	14.4	43.4	54.6	18.1	33.3	1014
C95965	8.54	15.8	47.0	55.1	18.5	33.6	1104
C95967	8.41	14.9	45.2	53.7	17.7	33.0	907
C95968	8.44	14.3	43.6	51.7	17.0	32.8	1306
C95969	7.75	13.0	39.7	51.2	16.7	32.7	902
C95977	8.14	14.6	43.8	53.8	17.9	33.3	878
C95980	8.42	15.0	45.3	53.9	17.8	33.1	897
MEAN	8.25	14.6	44.3	53.7	17.7	33.0	1011
S.D.	.273	.71	1.92	1.39	.56	.30	138.4
N	10	10	10	10	10	10	10

001032

Appendix 5
Individual Clinical Hematology Data

Males Week 14
13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WBC X10 ³ /μL	N-SEG X10 ³ /μL	LYMPH X10 ³ /μL	MONO X10 ³ /μL	EOSIN X10 ³ /μL	BASO X10 ³ /μL	N-SEG% X10 ³	LYMPH% X10 ³	MONO% X10 ³	EOSIN% X10 ³	BASO% X10 ³
Group: 1	Dose Level: 0										
	Dosage Unit: ppm MeFOS										
C95901	7.5	.8	6.0	.6	.1	.0	11	80	7	1	0
C95902	6.0	.9	4.8	.2	.0	.0	15	81	4	1	0
C95903	6.8	1.2	4.9	.6	.1	.0	17	72	9	2	0
C95904	6.0	.7	4.8	.5	.1	.0	11	79	8	2	0
C95905	6.4	.6	5.3	.3	.1	.0	10	82	5	2	0
C95906	6.0	1.0	4.4	.5	.1	.0	16	74	8	2	1
C95909	6.9	1.0	5.2	.6	.1	.0	15	75	9	1	0
C95918	7.0	.6	5.7	.6	.1	.0	8	82	8	1	1
C95919	8.8	.8	7.4	.4	.1	.0	10	84	5	2	0
C95920	7.4	1.5	5.4	.4	.1	.0	21	73	5	1	0
MEAN	6.9	.9	5.4	.5	.1	.0	13	78	7	2	0
S.D.	.88	.28	.85	.14	.03	.00	4.0	4.3	1.9	.5	.4
N	10	10	10	10	10	10	10	10	10	10	10
Group: 2	Dose Level: 3										
	Dosage Unit: ppm MeFOS										
C95921	4.8	.8	3.6	.3	.1	.0	16	76	6	2	0
C95922	7.0	1.1	5.0	.6	.2	.0	16	72	9	3	0
C95924	7.7	.9	6.4	.4	.1	.0	12	83	5	1	0
C95925	6.1	1.4	4.2	.4	.1	.0	23	69	6	1	0
C95926	8.0	1.3	5.9	.6	.2	.0	16	73	8	2	0
C95927	6.4	.8	5.1	.3	.2	.0	12	80	5	2	0
C95928	6.0	1.0	4.0	.8	.1	.0	17	66	14	2	1
C95936	7.2	.9	5.7	.4	.2	.0	13	80	5	2	0
C95937	4.3	.7	3.2	.2	.1	.0	17	75	5	2	0
C95939	9.2	1.1	7.5	.5	.1	.0	12	81	6	1	0
MEAN	6.7	1.0	5.1	.4	.1	.0	15	76	7	2	0
S.D.	1.48	.23	1.35	.18	.05	.00	3.4	5.6	2.8	.6	.3
N	10	10	10	10	10	10	10	10	10	10	10

001033

Appendix 5

Individual Clinical Hematology Data

Males Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WBC X10 ³ /μL	N-SEG X10 ³ /μL	LYMPH X10 ³ /μL	MONO X10 ³ /μL	EOSIN X10 ³ /μL	BASO X10 ³ /μL	N-SEG%	LYMPH%	MONO%	EOSIN%	BASO%
Group: 3	Dose Level: 30		Dosage Unit: ppm MeFOS								
C95941	10.8	1.2	9.0	.4	.1	.0	11	84	3	1	0
C95942	3.7	.7	2.8	.1	.0	.0	20	76	3	1	0
C95943	7.9	.8	6.4	.5	.1	.0	10	82	7	1	0
C95944	6.1	.7	4.9	.4	.1	.0	11	81	6	2	0
C95947	12.0	.9	10.6	.3	.1	.0	8	88	3	1	0
C95948	10.8	.7	9.5	.5	.1	.0	6	88	4	1	0
C95949	7.4	1.3	5.3	.6	.1	.0	18	72	9	1	0
C95950	5.9	.9	4.5	.4	.1	.0	15	76	7	2	0
C95958	6.2	1.1	4.6	.3	.1	.0	18	74	5	3	0
C95960	6.0	.6	4.9	.4	.1	.0	10	82	6	1	0
MEAN	7.7	.9	6.2	.4	.1	.0	13	80	5	1	0
S.D.	2.68	.24	2.57	.14	.05	.00	4.7	5.6	2.1	.7	.0
N	10	10	10	10	10	10	10	10	10	10	10
Group: 4	Dose Level: 100		Dosage Unit: ppm MeFOS								
C95961	5.0	.8	3.9	.2	.1	.0	17	77	4	1	0
C95962	5.9	1.0	4.6	.3	.1	.0	16	77	5	1	0
C95963	8.8	1.0	7.4	.3	.1	.0	11	84	3	2	0
C95964	6.7	1.2	5.0	.4	.1	.0	18	74	6	1	0
C95965	5.4	1.1	4.0	.2	.1	.0	20	74	4	1	0
C95967	4.4	.7	3.3	.4	.0	.0	15	75	8	1	0
C95968	12.7	1.6	10.0	1.0	.2	.1	12	78	8	1	1
C95969	10.3	1.8	7.8	.4	.1	.0	18	76	4	1	0
C95977	5.9	.8	4.6	.3	.1	.0	14	79	6	1	0
C95980	6.6	1.1	5.0	.4	.1	.0	16	76	6	2	0
MEAN	7.2	1.1	5.6	.4	.1	.0	16	77	5	1	0
S.D.	2.63	.35	2.13	.23	.05	.03	2.8	2.9	1.7	.4	.3
N	10	10	10	10	10	10	10	10	10	10	10

001034

Appendix 5

Individual Clinical Hematology Data

Males Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	ANISO	POLY	POIK	HYPO	TOXNEUT

Group: 1	Dose Level: 0		Dosage Unit: ppm MeFOS		
C95901	-	-	-	-	-
C95902	-	-	-	-	-
C95903	-	-	-	-	-
C95904	-	-	-	-	-
C95905	-	-	-	-	-
C95906	-	-	-	-	-
C95909	-	-	-	-	-
C95918	-	-	-	-	-
C95919	-	-	-	-	-
C95920	-	-	-	-	-
Group: 2	Dose Level: 3		Dosage Unit: ppm MeFOS		
C95921	-	-	-	-	-
C95922	-	-	-	-	-
C95924	-	-	-	-	-
C95925	-	-	-	-	-
C95926	-	-	-	-	-
C95927	-	-	-	-	-
C95928	-	-	-	-	-
C95936	-	-	-	-	-
C95937	-	-	-	-	-
C95939	-	-	-	-	-

001035

Appendix 5
Individual Clinical Hematology Data
Males Week 14
13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	ANISO	POLY	POIK	HYP0	TOXNEUT
-----	-----	-----	-----	-----	-----
Group: 3	Dose Level: 30		Dosage Unit: ppm MeFOS		
C95941	-	-	-	-	-
C95942	-	-	-	-	-
C95943	-	-	-	-	-
C95944	-	-	-	-	-
C95947	-	-	-	-	-
C95948	-	-	-	-	-
C95949	-	-	-	-	-
C95950	-	-	-	-	-
C95958	-	-	-	-	-
C95960	-	-	-	-	-
Group: 4	Dose Level: 100		Dosage Unit: ppm MeFOS		
C95961	-	-	-	-	-
C95962	-	-	-	-	-
C95963	-	-	-	-	-
C95964	-	-	-	-	-
C95965	-	-	-	-	-
C95967	-	-	-	-	-
C95968	-	-	-	-	-
C95969	-	-	-	-	-
C95977	-	-	-	-	-
C95980	-	-	-	-	-

001036

Appendix 5
Individual Clinical Hematology Data

Females Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	RBC x10 ⁶ /μL	HGB G/DL	HCT %	MCV FL	MCH PG	MCHC %	PLT x10 ³ /μL
Group: 1	Dose Level: 0	Dosage Unit: ppm MeFOS					
C95983	7.70	15.1	46.1	59.9	19.7	32.8	940
C95984	8.09	14.8	45.7	56.5	18.2	32.3	1026
C95985	7.55	14.6	44.9	59.4	19.3	32.5	894
C95986	7.54	14.2	43.4	57.6	18.8	32.6	1107
C95987	7.59	14.6	44.9	59.1	19.2	32.5	848
C95988	7.77	14.7	44.7	57.6	18.9	32.9	1087
C95990	7.85	14.6	45.0	57.4	18.6	32.5	1098
C95996	7.80	14.3	44.5	57.0	18.3	32.2	1163
C95997	7.78	14.8	45.0	57.8	19.0	32.9	951
C96000	7.96	15.8	46.9	58.9	19.8	33.6	1049
MEAN	7.76	14.8	45.1	58.1	19.0	32.7	1016
S.D.	.178	.45	.95	1.13	.54	.40	103.3
N	10	10	10	10	10	10	10
Group: 2	Dose Level: 3	Dosage Unit: ppm MeFOS					
C96001	8.33	14.6	45.4	54.5	17.5	32.1	1176
C96002	8.06	14.9	44.6	55.4	18.5	33.4	1045
C96003	7.54	14.4	44.8	59.4	19.1	32.1	938
C96006	7.77	14.5	44.3	57.0	18.7	32.8	965
C96007	8.34	16.6	49.8	59.7	19.9	33.3	1108
C96008	7.73	14.3	43.9	56.8	18.5	32.6	1055
C96010	8.55	15.8	48.0	56.1	18.5	33.0	1142
C96019	8.13	15.1	46.8	57.6	18.5	32.2	1263
C96020	8.18	15.3	46.9	57.4	18.7	32.5	969
MEAN	8.07	15.1	46.1	57.1	18.7	32.7	1073
S.D.	.330	.75	1.97	1.70	.63	.49	108.7
N	9	9	9	9	9	9	9

001037

Appendix 5

Individual Clinical Hematology Data

Females Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDE
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	RBC X10 ⁶ /μL	HGB G/DL	HCT %	MCV FL	MCH PG	MCHC %	PLT X10 ³ /μL
Group: 3	Dose Level: 30	Dosage Unit: ppm MeFOS					
C96021	8.56	15.4	47.5	55.5	18.0	32.4	931
C96022	8.45	15.9	48.2	57.1	18.9	33.0	1017
C96023	7.78	15.0	45.5	58.5	19.3	32.9	1047
C96024	7.76	14.5	44.2	57.0	18.6	32.7	915
C96025	8.49	15.3	46.7	55.1	18.1	32.8	1079
C96026	7.62	14.6	44.9	58.9	19.1	32.5	918
C96027	8.33	15.7	48.4	58.2	18.8	32.4	1059
C96030	8.05	15.6	47.8	59.4	19.4	32.7	809
C96038	7.86	14.2	43.3	55.1	18.0	32.7	905
C96040	7.99	15.1	45.7	57.2	18.9	33.0	1117
MEAN	8.09	15.1	46.2	57.2	18.7	32.7	980
S.D.	.343	.56	1.77	1.57	.52	.22	97.8
N	10	10	10	10	10	10	10
Group: 4	Dose Level: 100	Dosage Unit: ppm MeFOS					
C96042	7.74	13.9	42.7	55.1	18.0	32.6	1203
C96044	7.55	14.0	43.3	57.4	18.6	32.4	945
C96045	8.20	15.1	45.7	55.7	18.5	33.1	965
C96046	7.75	15.0	44.7	57.7	19.4	33.6	1049
C96047	7.33	13.0	39.9	54.4	17.7	32.6	1088
C96048	7.47	14.0	42.3	56.6	18.7	33.1	1189
C96049	8.10	14.7	44.6	55.1	18.1	32.9	988
C96050	7.41	13.7	41.4	55.9	18.6	33.2	925
C96056	7.52	13.8	41.5	55.2	18.4	33.3	1210
C96060	7.80	14.8	45.3	58.1	19.0	32.8	995
MEAN	7.69	14.2	43.1	56.1	18.5	33.0	1056
S.D.	.289	.67	1.91	1.26	.49	.37	110.7
N	10	10	10	10	10	10	10

001038

Appendix 5
Individual Clinical Hematology Data

Females Week 14
13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WBC X10 ³ /μL	N-SEG X10 ³ /μL	LYMPH X10 ³ /μL	MONO X10 ³ /μL	EOSIN X10 ³ /μL	BASO X10 ³ /μL	N-SEG X10 ³ /μL	LYMPH X10 ³ /μL	MONO X10 ³ /μL	EOSIN X10 ³ /μL	BASO X10 ³ /μL
Group: 1	Dose Level: 0										
	Dosage Unit: ppm MeFOS										
C95983	4.2	1.0	3.1	.0	.1	.0	.0	24	74	1	2
C95984	2.3	.3	1.9	.1	.0	.0	.0	12	82	4	0
C95985	4.9	1.9	2.6	.3	.1	.0	.0	39	53	7	0
C95986	4.8	.7	4.0	.0	.1	.0	.0	15	82	0	0
C95987	3.6	.4	2.9	.2	.1	.0	.0	11	81	6	0
C95988	5.6	1.1	4.2	.2	.1	.0	.0	19	75	4	0
C95990	3.1	.5	2.4	.1	.1	.0	.0	15	78	4	0
C95996	4.1	.9	2.7	.3	.1	.0	.0	22	67	7	1
C95997	3.6	.4	2.8	.3	.1	.0	.0	11	77	9	1
C96000	4.2	.8	3.0	.2	.1	.0	.0	20	72	5	0
MEAN	4.0	.8	3.0	.2	.1	.0	.0	19	74	5	0
S.D.	.95	.47	.69	.12	.03	.00	.00	8.5	8.8	2.8	.6
N	10	10	10	10	10	10	10	10	10	10	10
Group: 2	Dose Level: 3										
	Dosage Unit: ppm MeFOS										
C96001	3.6	.4	2.9	.3	.1	.0	.0	10	80	7	0
C96002	3.1	.6	2.1	.2	.1	.0	.0	19	70	8	0
C96003	4.0	.7	2.9	.2	.1	.0	.0	19	72	6	0
C96006	4.0	1.0	2.7	.3	.1	.0	.0	24	67	7	0
C96007	6.7	.5	5.8	.3	.1	.0	.0	8	86	4	0
C96008	6.6	.8	5.4	.3	.1	.0	.0	12	82	5	0
C96010	5.0	.7	4.0	.3	.1	.0	.0	13	80	5	0
C96019	4.3	.8	3.3	.2	.1	.0	.0	18	75	5	0
C96020	5.3	.6	4.4	.2	.1	.0	.0	11	82	5	0
MEAN	4.7	.7	3.7	.3	.1	.0	.0	15	77	6	0
S.D.	1.27	.18	1.27	.05	.00	.00	.00	5.3	6.4	1.3	.8
N	9	9	9	9	9	9	9	9	9	9	9

Covance 6329-225
3M T-6314.1

Appendix 5

Individual Clinical Hematology Data

Females Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	WBC $\times 10^3/\mu\text{L}$	N-SEG $\times 10^3/\mu\text{L}$	LYMPH $\times 10^3/\mu\text{L}$	MONO $\times 10^3/\mu\text{L}$	EOSIN $\times 10^3/\mu\text{L}$	BASO $\times 10^3/\mu\text{L}$	N-SEG%	LYMPH%	MONO%	EOSIN%	BASO%
Group: 3	Dose Level: 30	Dosage Unit: ppm MeFOS									
C96021	6.9	.6	6.1	.1	.1	.0	8	89	2	1	0
C96022	5.1	.8	3.9	.3	.1	.0	15	77	6	1	0
C96023	5.1	.8	3.8	.4	.1	.0	16	74	8	2	0
C96024	4.9	.9	3.5	.3	.1	.0	19	72	6	2	0
C96025	4.5	.5	3.8	.2	.1	.0	10	83	4	1	0
C96026	6.6	1.9	4.2	.3	.1	.0	29	65	5	1	0
C96027	5.4	.9	4.1	.4	.1	.0	16	75	7	1	0
C96030	7.8	.6	6.7	.4	.1	.1	7	86	5	2	1
C96038	6.0	1.8	3.9	.2	.1	.0	29	65	4	1	0
C96040	4.8	.8	3.6	.3	.1	.0	17	75	6	1	0
MEAN	5.7	1.0	4.4	.3	.1	.0	17	76	5	1	0
S.D.	1.08	.49	1.10	.10	.00	.03	7.6	8.0	1.7	.5	.3
N	10	10	10	10	10	10	10	10	10	10	10
Group: 4	Dose Level: 100	Dosage Unit: ppm MeFOS									
C96042	2.9	.4	2.3	.1	.0	.0	14	81	4	1	0
C96044	3.6	.6	2.8	.2	.1	.0	17	76	4	2	0
C96045	9.7	1.0	8.4	.2	.1	.0	10	86	2	1	0
C96046	6.4	.6	5.5	.3	.1	.0	9	85	4	1	1
C96047	3.9	.3	3.4	.0	.1	.0	8	89	0	1	0
C96048	4.0	1.1	2.6	.2	.1	.0	28	65	5	2	0
C96049	4.7	.4	4.1	.2	.0	.0	7	88	4	1	0
C96050	3.8	.9	2.6	.3	.1	.0	24	67	7	3	0
C96056	3.0	.4	2.5	.1	.1	.0	13	82	3	2	0
C96060	4.8	.5	4.0	.2	.1	.0	11	84	4	2	0
MEAN	4.7	.6	3.8	.2	.1	.0	14	80	4	2	0
S.D.	2.03	.28	1.89	.09	.04	.00	7.0	8.4	1.8	.7	.3
N	10	10	10	10	10	10	10	10	10	10	10

001040

Appendix 5

Individual Clinical Hematology Data

Females Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	ANISO	POLY	POIK	HYPO	TOXNEUT

Group: 1	Dose Level: 0		Dosage Unit: ppm MeFOS		
C95983	-	-	-	-	-
C95984	-	-	-	-	-
C95985	-	-	-	-	-
C95986	-	-	-	-	-
C95987	-	-	-	-	-
C95988	-	-	-	-	-
C95990	-	-	-	-	-
C95996	-	-	-	-	-
C95997	-	-	-	-	-
C96000	-	-	-	-	-
Group: 2	Dose Level: 3		Dosage Unit: ppm MeFOS		
C96001	-	-	-	-	-
C96002	-	-	-	-	-
C96003	-	-	-	-	-
C96006	-	-	-	-	-
C96007	-	-	-	-	-
C96008	-	-	-	-	-
C96010	-	-	-	-	-
C96019	-	-	-	-	-
C96020	-	-	-	-	-

001041

Appendix 5

Individual Clinical Hematology Data

Females Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	ANISO	POLY	POIK	HYP0	TOXNEUT
Group: 3	Dose Level: 30 Dosage Unit: ppm MeFOS				
C96021	-	-	-	-	-
C96022	-	-	-	-	-
C96023	-	-	-	-	-
C96024	-	-	-	-	-
C96025	-	-	-	-	-
C96026	-	-	-	-	-
C96027	-	-	-	-	-
C96030	-	-	-	-	-
C96038	-	-	-	-	-
C96040	-	-	-	-	-
Group: 4	Dose Level: 100 Dosage Unit: ppm MeFOS				
C96042	-	-	-	-	-
C96044	-	-	-	-	-
C96045	-	-	-	-	-
C96046	-	-	-	-	-
C96047	-	-	-	-	-
C96048	-	-	-	-	-
C96049	-	-	-	-	-
C96050	-	-	-	-	-
C96056	-	-	-	-	-
C96060	-	-	-	-	-

001042

Appendix 5
Individual Clinical Chemistry Data

Males Week 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	GLU MG/DL	UN MG/DL	CREAT MG/DL	T PRO G/DL	ALB G/DL	GLOB G/DL	T BILLI MG/DL	CHOL MG/DL	TRIG MG/DL
Group: 1 Dose Level: 0 Dosage Unit: ppm MeFOS									
C95901	101	9	.7	7.3	4.8	2.5	.2	60	47
C95902	98	11	.7	7.3	4.8	2.5	.2	57	31
C95903	109	14	.6	7.0	4.5	2.5	.2	76	33
C95904	107	12	.7	6.5	4.4	2.1	.2	50	40
C95905	96	11	.6	7.0	4.8	2.2	.2	56	42
C95906	100	9	.6	7.4	5.0	2.4	.2	83	94
C95907	107	9	.5	6.3	4.3	2.0	.2	32	37
C95908	110	13	.7	7.1	4.9	2.2	.2	76	33
C95909	94	12	.6	6.5	4.7	1.8	.2	54	49
C95910	105	9	.7	6.7	4.8	1.9	.2	47	46
MEAN	103	11	.6	6.9	4.7	2.2	.2	59	45
S.D.	5.7	1.9	.07	.39	.23	.26	.00	15.4	18.3
N	10	10	10	10	10	10	10	10	10
Group: 2 Dose Level: 3 Dosage Unit: ppm MeFOS									
C95921	116	10	.7	7.1	5.1	2.0	.2	50	44
C95922	109	11	.6	6.5	4.5	2.0	.2	51	51
C95923	117	10	.6	7.0	4.8	2.2	.2	54	30
C95924	107	14	.7	6.6	4.5	2.1	.2	88	44
C95925	148	9	.7	7.1	4.7	2.4	.2	62	50
C95926	108	12	.6	7.0	4.7	2.3	.2	50	53
C95927	104	12	.6	6.9	4.7	2.2	.2	59	75
C95928	103	10	.6	6.9	4.9	2.0	.2	45	92
C95929	103	10	.6	6.6	4.6	2.0	.2	60	69
C95930	127	11	.7	7.1	4.9	2.2	.2	63	38
MEAN	114	11	.6	6.9	4.7	2.1	.2	58	55
S.D.	14.1	1.4	.05	.23	.19	.14	.00	12.1	18.8
N	10	10	10	10	10	10	10	10	10

001043

Appendix 5

Individual Clinical Chemistry Data

Males Week 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	GLU MG/DL	UN MG/DL	CREAT MG/DL	T PRO G/DL	ALB G/DL	GLOB G/DL	T BILI MG/DL	CHOL MG/DL	TRIG MG/DL
Group: 3	Dose Level: 30 Dosage Unit: ppm MeFOS								
C95941	93	12	.6	6.5	4.8	1.7	.2	51	41
C95942	114	10	.6	7.1	4.8	2.3	.2	48	63
C95943	129	11	.6	6.5	4.6	1.9	.2	46	26
C95944	91	10	.6	6.8	4.7	2.1	.2	67	40
C95945	111	10	.6	6.5	4.5	2.0	.2	42	30
C95946	104	15	.6	6.9	4.8	2.1	.2	58	33
C95947	116	14	.6	6.4	4.6	1.8	.1	73	39
C95948	124	11	.6	6.6	4.9	1.7	.2	57	39
C95949	84	11	.6	6.7	4.6	2.1	.2	54	49
C95950	90	11	.5	6.5	4.5	2.0	.2	40	36
MEAN	106	12	.6	6.6	4.7	2.0	.2	54	40
S.D.	15.6	1.7	.03	.22	.14	.19	.03	10.6	10.4
N	10	10	10	10	10	10	10	10	10
Group: 4	Dose Level: 100 Dosage Unit: ppm MeFOS								
C95961	152	17	.7	6.4	4.4	2.0	.2	26	12
C95962	90	11	.6	6.6	4.7	1.9	.2	25	27
C95963	97	12	.7	6.8	4.8	2.0	.2	60	45
C95964	102	13	.6	6.4	4.8	1.6	.2	64	29
C95965	94	12	.6	7.0	4.9	2.1	.2	41	25
C95966	110	13	.5	6.3	4.4	1.9	.2	34	16
C95967	99	13	.6	6.9	4.8	2.1	.2	55	24
C95968	111	12	.6	7.1	4.8	2.3	.2	54	35
C95969	105	13	.5	6.5	4.5	2.0	.2	35	17
C95970	98	13	.6	7.0	5.0	2.0	.2	51	18
MEAN	106	13	.6	6.7	4.7	2.0	.2	44	25
S.D.	17.5	1.6	.07	.29	.21	.18	.00	14.1	9.9
N	10	10	10	10	10	10	10	10	10

001044

Appendix 5

Individual Clinical Chemistry Data

Males Week 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	AST/SGOT IU/L	ALT/SGPT IU/L	GGT IU/L	CA MG/DL	I PHOS MG/DL	NA MMOL/L	K MMOL/L	CL MMOL/L
Group: 1	Dose Level: 0	Dosage Unit: ppm MeFOS						
C95901	98	34	1	11.7	10.2	149	6.6	107
C95902	86	35	0	11.3	9.9	149	6.8	107
C95903	71	25	1	11.0	9.0	145	6.1	102
C95904	99	31	0	10.4	10.5	146	5.7	103
C95905	97	34	0	10.4	8.5	148	6.1	107
C95906	143	35	0	11.6	9.6	146	6.8	101
C95907	114	26	1	10.5	8.9	147	6.4	106
C95908	100	33	0	11.0	9.7	147	6.1	104
C95909	111	33	0	10.7	9.3	147	6.0	103
C95910	109	30	0	10.8	9.3	146	5.8	106
MEAN	103	32	0	10.9	9.5	147	6.2	105
S.D.	18.9	3.6	0.5	0.47	0.61	1.3	0.39	2.3
N	10	10	10	10	10	10	10	10
Group: 2	Dose Level: 3	Dosage Unit: ppm MeFOS						
C95921	92	33	0	11.1	11.1	150	7.0	108
C95922	115	35	1	10.6	9.0	145	6.0	102
C95923	95	39	0	10.7	10.3	147	7.0	105
C95924	86	26	1	11.0	10.3	147	6.1	106
C95925	79	29	0	10.7	9.0	145	5.9	105
C95926	108	41	0	10.9	9.5	147	6.1	104
C95927	74	35	0	11.1	9.6	146	5.9	104
C95928	123	32	0	10.9	10.5	149	6.2	105
C95929	83	30	0	10.7	9.7	147	5.9	103
C95930	103	30	1	10.6	9.1	148	6.3	105
MEAN	96	33	0	10.8	9.8	147	6.2	105
S.D.	16.1	4.6	0.5	0.19	0.71	1.6	0.42	1.6
N	10	10	10	10	10	10	10	10

001045

Appendix 5

Individual Clinical Chemistry Data

Males Week 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	AST/SGOT IU/L	ALT/SGPT IU/L	GGT IU/L	CA MG/DL	I PHOS MG/DL	NA MMOL/L	K MMOL/L	CL MMOL/L
Group: 3	Dose Level: 30	Dosage Unit: ppm MeFOS						
C95941	131	33	0	10.5	9.0	147	6.6	106
C95942	109	36	0	10.8	9.0	150	6.5	107
C95943	95	32	2	10.1	8.7	147	6.3	105
C95944	113	32	0	10.9	9.9	147	6.6	105
C95945	76	27	0	10.6	9.0	146	5.4	104
C95946	82	31	0	10.3	7.7	147	5.2	108
C95947	90	36	0	10.1	8.2	147	5.6	107
C95948	88	34	1	10.6	10.4	148	6.0	107
C95949	125	38	1	10.4	8.9	149	6.3	105
C95950	105	26	0	10.9	9.5	147	6.5	106
MEAN	101	32	0	10.5	9.0	148	6.1	106
S.D.	18.3	3.8	.7	.30	.78	1.2	.52	1.2
N	10	10	10	10	10	10	10	10
Group: 4	Dose Level: 100	Dosage Unit: ppm MeFOS						
C95961	86	36	0	10.2	11.4	146	5.8	103
C95962	113	41	0	10.2	8.9	148	6.0	107
C95963	93	35	0	10.9	9.6	148	6.5	106
C95964	122	41	1	10.8	9.0	149	6.5	109
C95965	103	41	0	10.6	9.3	148	6.5	104
C95966	91	41	0	10.0	7.9	143	5.1	104
C95967	94	50	0	11.2	9.2	150	6.1	108
C95968	88	40	0	10.9	9.1	146	6.5	103
C95969	90	41	0	10.0	7.8	142	6.1	102
C95970	86	34	0	10.7	8.3	146	5.7	102
MEAN	97	40	0	10.6	9.0	147	6.1	105
S.D.	12.2	4.5	.3	.42	1.02	2.5	.46	2.5
N	10	10	10	10	10	10	10	10

001046

Appendix 5

Individual Clinical Chemistry Data

Females Week 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	GLU MG/DL	UN MG/DL	CREAT MG/DL	T PRO G/DL	ALB G/DL	GLOB G/DL	T BILI MG/DL	CHOL MG/DL	TRIG MG/DL
Group: 1	Dose Level: 0 Dosage Unit: ppm MeFOS								
C95981	101	13	.7	8.1	5.6	2.5	.2	80	34
C95982	119	11	.7	7.3	5.5	1.8	.2	76	26
C95983	117	13	.7	7.3	4.9	2.4	.2	67	30
C95984	111	13	.7	7.7	5.4	2.3	.3	58	30
C95985	110	14	.7	7.4	5.5	1.9	.3	93	39
C95986	92	12	.7	7.3	5.3	2.0	.2	64	16
C95987	105	11	.7	7.2	5.3	1.9	.2	61	35
C95988	99	13	.7	7.3	5.2	2.1	.2	58	20
C95989	120	12	.7	7.0	5.1	1.9	.2	55	26
C95990	111	10	.6	7.6	5.4	2.2	.3	66	42
MEAN	108	12	.7	7.4	5.3	2.1	.2	68	30
S.D.	9.2	1.2	.03	.31	.21	.24	.05	11.9	8.1
N	10	10	10	10	10	10	10	10	10
Group: 2	Dose Level: 3 Dosage Unit: ppm MeFOS								
C96001	99	9	.6	7.3	5.4	1.9	.2	53	20
C96002	99	11	.7	7.6	5.8	1.8	.2	54	19
C96003	107	10	.7	7.0	4.9	2.1	.2	87	40
C96004	95	9	.6	6.9	5.0	1.9	.2	70	43
C96005	115	14	.7	7.4	5.0	2.4	.2	81	25
C96006	116	14	.7	6.9	4.8	2.1	.2	68	24
C96007	103	10	.7	7.5	5.3	2.2	.2	54	19
C96008	116	11	.7	8.5	6.0	2.5	.2	75	36
C96009	105	14	.7	7.2	5.2	2.0	.2	78	22
C96010	121	13	.7	7.2	5.4	1.8	.2	69	28
MEAN	108	12	.7	7.4	5.3	2.1	.2	69	28
S.D.	8.9	2.1	.04	.47	.39	.24	.00	12.0	8.9
N	10	10	10	10	10	10	10	10	10

001047

Appendix 5
Individual Clinical Chemistry Data

Females Week 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	GLU MG/DL	UN MG/DL	CREAT MG/DL	T PRO G/DL	ALB G/DL	GLOB G/DL	T BILI MG/DL	CHOL MG/DL	TRIG MG/DL
Group: 3	Dose Level: 30 Dosage Unit: ppm MeFOS								
C96021	102	10	.7	7.6	5.5	2.1	.2	62	28
C96022	116	13	.7	7.1	5.3	1.8	.2	52	30
C96023	113	12	.7	7.9	5.9	2.0	.2	75	25
C96024	95	12	.6	7.1	5.3	1.8	.2	73	37
C96025	111	13	.7	7.7	5.4	2.3	.2	69	21
C96026	94	12	.7	6.9	4.9	2.0	.2	53	25
C96027	105	10	.7	7.1	5.3	1.8	.2	76	43
C96028	123	11	.7	7.0	4.9	2.1	.2	53	26
C96029	117	13	.7	7.0	5.3	1.7	.2	58	28
C96030	99	15	.7	7.2	5.4	1.8	.1	71	21
MEAN	108	12	.7	7.3	5.3	1.9	.2	64	28
S.D.	10.0	1.5	.03	.34	.29	.19	.03	9.7	6.9
N	10	10	10	10	10	10	10	10	10
Group: 4	Dose Level: 100 Dosage Unit: ppm MeFOS								
C96041	89	12	.6	7.1	5.4	1.7	.2	46	24
C96042	98	15	.6	7.5	5.5	2.0	.2	100	48
C96043	75	13	.6	7.5	5.4	2.1	.2	43	17
C96044	96	12	.6	7.5	5.4	2.1	.2	86	34
C96045	105	11	.6	6.8	5.0	1.8	.2	50	26
C96046	90	15	.6	6.6	4.7	1.9	.2	38	24
C96047	84	14	.8	7.1	5.4	1.7	.2	48	20
C96048	117	15	.7	8.0	5.7	2.3	.2	37	21
C96049	89	15	.6	7.0	5.4	1.6	.2	39	17
C96050	104	13	.7	7.3	5.2	2.1	.2	45	26
MEAN	95	14	.6	7.2	5.3	1.9	.2	53	26
S.D.	12.0	1.5	.07	.41	.28	.23	.00	21.7	9.3
N	10	10	10	10	10	10	10	10	10

C91048

Appendix 5
Individual Clinical Chemistry Data

Females Week 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	AST/SGOT IU/L	ALT/SGPT IU/L	GGT IU/L	CA MG/DL	I PHOS MG/DL	NA MMOL/L	K MMOL/L	CL MMOL/L
Group: 1	Dose Level: 0	Dosage Unit: ppm MeFOS						
C95981	71	31	1	12.0	8.5	147	6.0	105
C95982	91	24	0	10.8	8.0	148	6.5	107
C95983	85	23	1	10.6	7.0	148	6.2	105
C95984	120	34	0	10.7	7.8	147	6.0	106
C95985	91	25	0	11.2	8.1	144	6.3	103
C95986	99	25	1	10.4	8.1	146	5.8	104
C95987	99	36	0	10.3	7.1	145	5.5	105
C95988	101	24	0	10.5	8.0	146	5.9	104
C95989	111	28	1	10.5	7.4	146	5.7	104
C95990	119	33	0	11.2	9.2	146	6.3	104
MEAN	99	28	0	10.8	7.9	146	6.0	105
S.D.	15.3	4.8		.52	.65	1.3	.31	1.2
N	10	10	10	10	10	10	10	10
Group: 2	Dose Level: 3	Dosage Unit: ppm MeFOS						
C96001	91	30	0	10.4	7.9	144	5.4	105
C96002	99	25	0	10.9	8.7	149	5.9	104
C96003	81	22	1	10.9	8.2	146	6.4	106
C96004	95	25	0	10.6	8.8	145	5.5	105
C96005	105	36	0	11.3	9.8	149	6.1	105
C96006	83	23	1	10.3	7.9	147	5.5	107
C96007	94	30	0	10.8	8.3	147	5.8	105
C96008	97	21	0	11.3	8.3	147	6.5	102
C96009	89	23	2	10.2	7.3	146	5.5	106
C96010	100	31	0	10.4	7.1	147	5.7	106
MEAN	93	27	0	10.7	8.2	147	5.8	105
S.D.	7.5	4.9		.7	.77	1.6	.39	1.4
N	10	10	10	10	10	10	10	10

001049

Appendix 5

Individual Clinical Chemistry Data

Females Week 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	AST/SGOT IU/L	ALT/SGPT IU/L	GGT IU/L	CA MG/DL	I PHOS MG/DL	NA MMOL/L	K MMOL/L	CL MMOL/L
Group: 3	Dose Level: 30	Dosage Unit: ppm MeFOS						
C96021	91	30	0	11.4	9.4	150	7.2	107
C96022	99	29	1	10.6	8.0	146	6.0	102
C96023	86	25	3	10.7	6.2	145	5.1	103
C96024	118	25	0	10.9	8.8	144	6.4	103
C96025	83	28	0	11.0	6.9	145	5.8	104
C96026	80	21	2	10.5	8.3	146	5.6	107
C96027	97	31	0	11.1	8.6	146	6.4	104
C96028	73	31	1	10.7	7.2	145	5.7	104
C96029	112	26	1	10.0	6.4	145	5.4	103
C96030	93	32	2	10.8	7.4	149	6.0	107
MEAN	93	28	1	10.8	7.7	146	6.0	104
S.D.	14.0	3.5	1.1	.38	1.07	1.9	.60	1.9
N	10	10	10	10	10	10	10	10
Group: 4	Dose Level: 100	Dosage Unit: ppm MeFOS						
C96041	96	26	1	10.4	8.2	149	5.6	107
C96042	119	23	0	11.2	8.4	143	6.3	101
C96043	106	31	1	10.7	8.2	149	5.7	108
C96044	82	34	0	11.2	9.6	146	6.7	103
C96045	67	28	1	10.7	8.6	148	5.9	107
C96046	88	36	0	10.6	8.9	146	5.9	107
C96047	113	39	0	10.4	9.2	147	5.8	104
C96048	86	26	2	10.3	6.8	147	5.0	109
C96049	91	34	2	10.5	8.5	150	5.8	109
C96050	80	22	1	10.5	6.1	147	5.0	106
MEAN	93	30	1	10.6	8.2	147	5.8	106
S.D.	16.0	5.8	.8	.32	1.06	2.0	.52	2.6
N	10	10	10	10	10	10	10	10

001050

Appendix 5

Individual Clinical Chemistry Data

Males Week 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	PCOAO IU/G	

Group: 1	Dose Level: 0	Dosage Unit: ppm MeFOS
C95911	11	
C95912	9	
C95913	13	
C95914	10	
C95915	18	
MEAN	12	
S.D.	3.6	
N	5	
Group: 2	Dose Level: 3	Dosage Unit: ppm MeFOS
C95931	12	
C95932	15	
C95933	13	
C95934	11	
C95935	16	
MEAN	13	
S.D.	2.1	
N	5	
Group: 3	Dose Level: 30	Dosage Unit: ppm MeFOS
C95951	14	
C95952	7	
C95953	2	
C95954	8	
C95955	9	
MEAN	8	
S.D.	4.3	
N	5	

001051

Appendix 5

Individual Clinical Chemistry Data

Males Week 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	PCOAO IU/G	Group: 4	Dose Level: 100	Dosage Unit: ppm MeFOS
C95971			7	
C95972			8	
C95973			9	
C95974			12	
C95975			9	
MEAN			9	
S.D.			1.9	
N			5	

001052

Appendix 5

Individual Clinical Chemistry Data

Females Week 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	PCOAO IU/G	

Group: 1	Dose Level: 0	Dosage Unit: ppm MeFOS
C95991	6	
C95992	9	
C95993	3	
C95994	7	
C95995	9	
MEAN	7	
S.D.	2.5	
N	5	
Group: 2	Dose Level: 3	Dosage Unit: ppm MeFOS
C96011	6	
C96012	3	
C96013	9	
C96014	3	
C96015	2	
MEAN	5	
S.D.	2.9	
N	5	
Group: 3	Dose Level: 30	Dosage Unit: ppm MeFOS
C96031	7	
C96032	6	
C96033	5	
C96034	5	
C96035	13	
MEAN	7	
S.D.	3.3	
N	5	

001053

Appendix 5

Individual Clinical Chemistry Data

Females Week 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER -----	PCAO IU/G -----	Group: 4	Dose Level: 100	Dosage Unit: ppm MeFOS
C96051			8	
C96052			9	
C96053			11	
C96054			10	
C96055			15	
MEAN			11	
S.D.			2.7	
N			5	

001054

Appendix 5
Individual Clinical Chemistry Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS											
Males Week 14											
ANIMAL NUMBER	GLU MG/DL	UN MG/DL	CREAT MG/DL	T PRO G/DL	ALB G/DL	GLOB G/DL	T BILI MG/DL	CHOL MG/DL	TRIG MG/DL		
Group: 1	Dose Level: 0 Dosage Unit: ppm MeFOS										
C95901	125	12	.8	7.5	4.8	2.7	.2	54	76		
C95902	123	13	.8	7.5	4.6	2.9	.1	71	34		
C95903	107	14	.8	7.3	4.7	2.6	.2	69	61		
C95904	120	17	.8	6.8	4.5	2.3	.2	80	61		
C95905	106	11	.7	6.8	4.6	2.2	.1	53	64		
C95906	120	11	.7	7.6	5.1	2.5	.2	100	111		
C95909	106	14	.8	6.7	4.4	2.3	.1	54	53		
C95918	115	13	.8	7.3	4.9	2.4	.1	82	50		
C95919	126	10	.7	7.5	5.1	2.4	.1	64	43		
C95920	125	11	.7	7.2	4.7	2.5	.2	60	39		
MEAN	117	13	.8	7.2	4.7	2.5	.2	69	59		
S.D.	8.2	2.1	.05	.34	.24	.21	.05	15.2	22.2		
N	10	10	10	10	10	10	10	10	10		
Group: 2	Dose Level: 3 Dosage Unit: ppm MeFOS										
C95921	115	10	.7	6.9	4.8	2.1	.2	37	44		
C95922	129	15	.8	6.7	4.6	2.1	.1	44	83		
C95924	121	15	.7	6.5	4.4	2.1	.1	87	42		
C95925	126	10	.7	7.1	4.6	2.5	.1	60	41		
C95926	118	13	.8	7.1	4.7	2.4	.2	41	43		
C95927	125	14	.8	7.3	4.9	2.4	.2	56	84		
C95928	115	12	.8	7.1	5.0	2.1	.1	38	75		
C95936	105	10	.7	6.8	4.5	2.3	.1	61	89		
C95937	109	10	.7	7.6	4.9	2.7	.1	69	28		
C95939	126	13	.7	6.6	4.5	2.1	.1	79	36		
MEAN	119	12	.7	7.0	4.7	2.3	.1	57	56		
S.D.	7.9	2.1	.05	.34	.20	.21	.05	17.4	23.3		
N	10	10	10	10	10	10	10	10	10		

001055

Appendix 5

Individual Clinical Chemistry Data

Males Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	GLU MG/DL	UN MG/DL	CREAT MG/DL	T PRO G/DL	ALB G/DL	GLOB G/DL	T BILI MG/DL	CHOL MG/DL	TRIG MG/DL
Group: 3	Dose Level: 30	Dosage Unit: ppm MeFOS							
C95941	122	14	.7	6.6	4.8	1.8	.1	39	31
C95942	119	12	.8	7.6	5.1	2.5	.2	45	33
C95943	124	11	.7	6.7	4.7	2.0	.2	33	58
C95944	100	12	.7	6.7	4.6	2.1	.2	59	38
C95947	115	16	.8	6.6	4.5	2.1	.1	71	35
C95948	108	12	.9	7.0	4.7	2.3	.2	57	47
C95949	94	15	.7	5.4	3.8	1.6	.2	36	25
C95950	119	15	.7	6.7	4.8	1.9	.1	31	23
C95958	117	13	.8	7.0	4.5	2.5	.1	72	39
C95960	114	17	.8	7.2	4.9	2.3	.2	50	32
MEAN	113	14	.8	6.8	4.6	2.1	.2	49	36
S.D.	9.7	2.0	.07	.57	.35	.30	.05	15.1	10.3
N	10	10	10	10	10	10	10	10	10
Group: 4	Dose Level: 100	Dosage Unit: ppm MeFOS							
C95961	131	16	.7	7.3	5.1	2.2	.2	13	15
C95962	121	13	.7	6.5	4.9	1.6	.3	11	12
C95963	153	16	.8	7.5	5.6	1.9	.2	39	27
C95964	140	16	.8	7.4	5.6	1.8	.2	37	21
C95965	120	14	.7	7.3	5.3	2.0	.2	25	18
C95967	123	15	.7	7.5	5.3	2.2	.2	46	21
C95968	120	13	.8	7.7	5.2	2.5	.2	31	16
C95969	123	17	.7	6.8	4.7	2.1	.2	15	9
C95977	116	14	.7	6.7	4.9	1.8	.1	19	19
C95980	131	15	.7	7.1	5.1	2.0	.2	20	14
MEAN	128	15	.7	7.2	5.2	2.0	.2	26	17
S.D.	11.3	1.4	.05	.39	.29	.26	.05	12.1	5.2
N	10	10	10	10	10	10	10	10	10

001056

Appendix 5

Individual Clinical Chemistry Data

Males Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	AST/SGOT IU/L	ALT/SGPT IU/L	GGT IU/L	CA MG/DL	I PHOS MG/DL	NA MMOL/L	K MMOL/L	CL MMOL/L
Group: 1	Dose Level: 0 Dosage Unit: ppm MeFOS							
C95901	101	44	3	11.3	7.8	154	6.2	113
C95902	95	38	3	10.6	6.0	148	6.4	107
C95903	87	35	1	10.8	7.6	152	6.5	110
C95904	107	42	0	10.3	7.0	149	6.1	109
C95905	110	39	3	9.9	6.2	147	6.1	106
C95906	87	38	3	11.3	6.2	146	5.7	102
C95909	109	47	3	10.2	6.2	148	5.9	107
C95918	59	29	2	11.2	6.2	150	5.9	109
C95919	68	35	0	11.0	6.4	151	6.3	111
C95920	70	36	1	10.6	6.2	151	5.5	113
MEAN	89	38	2	10.7	6.6	150	6.1	109
S.D.	18.4	5.1	1.3	.49	.65	2.5	.31	3.4
N	10	10	10	10	10	10	10	10
Group: 2	Dose Level: 3 Dosage Unit: ppm MeFOS							
C95921	86	49	2	10.3	6.7	150	5.9	111
C95922	80	44	2	10.8	6.7	151	6.2	110
C95924	83	36	2	10.3	6.7	146	6.1	105
C95925	95	43	3	10.2	6.2	148	6.1	106
C95926	94	50	3	10.6	6.1	152	6.3	111
C95927	69	39	2	10.9	6.9	148	6.1	105
C95928	91	44	3	10.8	6.5	150	6.4	110
C95936	100	30	3	10.3	6.1	147	5.5	105
C95937	75	34	3	10.7	6.4	152	5.8	112
C95939	92	39	1	10.7	6.7	152	6.4	112
MEAN	86	41	2	10.6	6.5	150	6.1	109
S.D.	9.7	6.4	.7	.26	.29	2.2	.28	3.1
N	10	10	10	10	10	10	10	10

001057

Appendix 5

Individual Clinical Chemistry Data

Males Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	AST/SGOT IU/L	ALT/SGPT IU/L	GGT IU/L	CA MG/DL	I PHOS MG/DL	NA MMOL/L	K MMOL/L	CL MMOL/L
Group: 3	Dose Level: 30 Dosage Unit: ppm MeFOS							
C95941	96	44	3	10.3	6.3	150	6.5	112
C95942	111	45	1	10.3	6.2	151	6.4	111
C95943	82	31	3	10.0	6.8	149	6.1	109
C95944	74	38	3	10.6	7.5	150	6.5	111
C95947	75	36	1	10.1	6.4	145	5.4	104
C95948	84	45	3	10.6	7.2	150	6.3	105
C95949	126	49	0	9.2	5.5	145	5.8	104
C95950	94	34	2	10.3	6.8	150	6.1	108
C95958	88	28	3	10.4	6.0	151	6.2	110
C95960	78	34	4	10.4	6.6	150	6.5	110
MEAN	91	38	2	10.2	6.5	149	6.2	108
S.D.	16.7	7.0	1.3	.40	.58	2.2	.36	3.0
N	10	10	10	10	10	10	10	10
Group: 4	Dose Level: 100 Dosage Unit: ppm MeFOS							
C95961	73	46	3	10.6	6.6	152	6.9	112
C95962	83	48	2	10.0	6.8	150	6.5	108
C95963	63	33	2	10.9	7.1	154	6.4	112
C95964	100	64	1	10.6	7.3	152	7.0	112
C95965	99	65	3	10.6	7.3	153	6.7	109
C95967	95	57	2	10.6	6.7	147	5.6	102
C95968	108	70	3	10.5	7.3	147	6.6	103
C95969	86	36	2	10.0	6.2	147	6.3	108
C95977	99	70	1	10.1	6.4	150	5.6	111
C95980	102	108	2	10.5	7.0	151	6.5	112
MEAN	91	60	2	10.4	6.9	150	6.4	109
S.D.	14.3	21.6	.7	.30	.39	2.6	.48	3.8
N	10	10	10	10	10	10	10	10

001058

Appendix 5

Individual Clinical Chemistry Data

Females Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDE
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	GLU MG/DL	UN MG/DL	CREAT MG/DL	T PRO G/DL	ALB G/DL	GLOB G/DL	T BILI MG/DL	CHOL MG/DL	TRIG MG/DL
Group: 1	Dose Level: 0 Dosage Unit: ppm MeFOS								
C95983	123	14	.9	8.3	5.9	2.4	.2	69	55
C95984	137	11	.9	8.5	5.9	2.6	.2	68	40
C95985	125	15	.9	8.1	6.2	1.9	.2	101	37
C95986	133	9	.8	7.5	5.7	1.8	.2	60	31
C95987	115	11	.8	7.9	6.1	1.8	.2	66	35
C95988	114	15	.9	8.0	5.7	2.3	.2	68	28
C95990	107	12	.9	8.7	6.4	2.3	.3	90	86
C95996	120	13	.7	7.2	5.4	1.8	.2	70	41
C95997	114	11	.8	7.5	5.6	1.9	.2	71	35
C96000	104	13	.9	7.4	5.0	2.4	.2	66	39
MEAN	119	12	.8	7.9	5.8	2.1	.2	73	43
S.D.	10.6	2.0	.07	.50	.41	.31	.03	12.6	16.8
N	10	10	10	10	10	10	10	10	10
Group: 2	Dose Level: 3 Dosage Unit: ppm MeFOS								
C96001	120	10	.7	7.6	5.6	2.0	.1	52	19
C96002	123	11	.8	7.6	5.9	1.7	.2	57	24
C96003	124	13	.8	7.1	5.0	2.1	.2	78	32
C96006	113	17	.7	7.0	5.0	2.0	.1	69	34
C96007	138	13	.9	8.6	6.0	2.6	.1	70	34
C96008	125	11	.7	8.3	6.1	2.2	.2	67	92
C96010	111	14	.8	7.3	5.2	2.1	.2	73	28
C96019	116	14	.8	7.9	5.7	2.2	.2	76	45
C96020	123	12	.8	7.7	5.7	2.0	.1	80	34
MEAN	121	13	.8	7.7	5.6	2.1	.2	69	38
S.D.	8.0	2.1	.07	.53	.42	.24	.05	9.4	21.5
N	9	9	9	9	9	9	9	9	9

001059

Appendix 5

Individual Clinical Chemistry Data

Females Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDE
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	GLU MG/DL	UN MG/DL	CREAT MG/DL	T PRO G/DL	ALB G/DL	GLOB G/DL	T BILI MG/DL	CHOL MG/DL	TRIG MG/DL
Group: 3	Dose Level: 30 Dosage Unit: ppm MeFOS								
C96021	109	13	.8	7.7	5.6	2.1	.2	57	35
C96022	111	16	.9	7.8	5.9	1.9	.2	58	30
C96023	97	13	.8	8.3	6.1	2.2	.2	74	29
C96024	105	12	.7	7.8	5.6	2.2	.2	66	36
C96025	116	13	.9	8.8	6.0	2.8	.2	75	26
C96026	122	12	.7	7.6	5.5	2.1	.2	59	32
C96027	103	13	.8	7.6	5.7	1.9	.2	78	69
C96030	113	17	.8	7.7	5.7	2.0	.2	78	26
C96038	113	16	.9	8.2	5.7	2.5	.2	74	44
C96040	111	13	.8	8.5	6.2	2.3	.1	86	37
MEAN	110	14	.8	8.0	5.8	2.2	.2	70	36
S.D.	7.0	1.8	.07	.42	.24	.28	.03	9.9	12.7
N	10	10	10	10	10	10	10	10	10
Group: 4	Dose Level: 100 Dosage Unit: ppm MeFOS								
C96042	100	15	.8	9.0	6.6	2.4	.2	84	32
C96044	116	14	.9	8.6	6.4	2.2	.2	65	29
C96045	125	15	.8	7.6	5.6	2.0	.2	45	26
C96046	116	18	.9	7.8	5.6	2.2	.2	56	23
C96047	109	17	.8	8.1	6.2	1.9	.2	50	18
C96048	120	18	.9	8.6	6.1	2.5	.2	30	21
C96049	111	18	.8	7.7	5.9	1.8	.2	35	25
C96050	104	20	.7	7.9	5.7	2.2	.2	34	22
C96056	95	19	.8	8.7	6.4	2.3	.2	42	22
C96060	109	14	.8	7.8	5.9	1.9	.2	71	28
MEAN	110	17	.8	8.2	6.0	2.1	.2	51	25
S.D.	9.2	2.1	.06	.50	.36	.23	.00	17.7	4.2
N	10	10	10	10	10	10	10	10	10

001060

Appendix 5

Individual Clinical Chemistry Data

Females Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	AST/SGOT IU/L	ALT/SGPT IU/L	GGT IU/L	CA MG/DL	I MG/DL	PHOS MG/DL	NA MMOL/L	K MMOL/L	CL MMOL/L
Group: 1	Dose Level: 0 Dosage Unit: ppm MeFOS								
C95983	74	27	4	11.0	4.2	152	6.1	113	113
C95984	92	34	3	10.9	4.2	153	5.7	113	113
C95985	79	31	2	11.0	4.6	149	5.4	110	110
C95986	77	31	3	10.7	4.4	151	5.3	114	114
C95987	79	35	1	11.0	5.0	149	5.8	110	110
C95988	101	35	3	11.1	5.6	148	5.5	107	107
C95990	88	48	1	11.7	5.4	152	6.2	112	112
C95996	78	37	3	11.2	5.0	150	5.6	113	113
C95997	75	30	1	10.9	5.5	149	5.6	111	111
C96000	99	33	4	10.5	4.4	147	5.9	106	106
MEAN	84	34	2	11.0	4.8	150	5.7	111	111
S.D.	10.1	5.7	1.2	32	54	1.9	29	2.7	10
N	10	10	10	10	10	10	10	10	10
Group: 2	Dose Level: 3 Dosage Unit: ppm MeFOS								
C96001	91	36	1	10.7	5.3	150	6.0	115	115
C96002	100	31	1	10.6	4.3	149	5.8	108	108
C96003	190	96	3	10.6	4.3	150	5.9	111	111
C96006	89	30	3	10.4	5.0	147	5.8	109	109
C96007	101	44	3	11.7	7.3	159	7.7	114	114
C96008	81	30	4	10.7	4.3	147	5.1	105	105
C96010	101	42	3	10.5	5.0	149	6.2	110	110
C96019	94	35	1	11.1	5.3	150	6.1	114	114
C96020	114	40	4	11.1	5.4	151	6.4	113	113
MEAN	107	43	3	10.8	5.1	150	6.1	111	111
S.D.	32.6	20.7	1.2	41	93	3.6	70	3.3	9
N	9	9	9	9	9	9	9	9	9

001061

Appendix 5

Individual Clinical Chemistry Data

Females Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	AST/SGOT IU/L	ALT/SGPT IU/L	GGT IU/L	CA MG/DL	I PHOS MG/DL	NA MMOL/L	K MMOL/L	CL MMOL/L
Group: 3	Dose Level: 30 Dosage Unit: ppm MeFOS							
C96021	84	30	6	11.2	6.0	153	6.9	112
C96022	81	29	4	11.2	6.2	156	6.7	113
C96023	88	30	1	11.0	4.3	147	5.9	105
C96024	65	22	1	11.0	5.6	150	5.6	111
C96025	87	35	2	11.6	6.5	132	6.2	109
C96026	76	29	2	10.6	4.7	148	5.3	109
C96027	106	34	4	10.9	5.2	149	6.4	106
C96030	78	33	2	10.9	6.5	151	6.3	110
C96038	130	34	3	10.9	5.3	149	5.8	107
C96040	485	364	5	11.1	5.9	151	6.5	111
MEAN	128	64	3	11.0	5.6	151	6.2	109
S.D.	126.7	105.5	1.7	.26	.74	2.6	.50	2.6
N	10	10	10	10	10	10	10	10
Group: 4	Dose Level: 100 Dosage Unit: ppm MeFOS							
C96042	95	30	3	11.5	5.3	150	6.0	110
C96044	75	31	3	11.3	5.6	154	6.0	111
C96045	70	33	5	10.6	5.9	147	5.5	104
C96046	78	41	3	11.0	5.8	151	6.2	110
C96047	79	32	1	10.9	6.8	149	5.8	108
C96048	79	31	2	10.7	4.8	151	5.1	110
C96049	91	48	3	10.9	6.3	155	6.5	112
C96050	66	31	2	10.8	5.4	149	5.4	110
C96056	90	38	3	10.8	3.6	147	5.4	105
C96060	82	41	3	10.9	4.7	152	5.5	114
MEAN	80	36	3	10.9	5.4	150	5.7	109
S.D.	9.3	6.1	1.0	.27	.90	2.7	.43	3.0
N	10	10	10	10	10	10	10	10

001062

Appendix 5

Individual Clinical Chemistry Data

Males Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	PCOAO IU/G	Group: 1	Dose Level: 0	Dosage Unit: ppm MeFOS
C95901			4	
C95902			11	
C95903			11	
C95904			10	
C95905			11	
C95906			12	
C95907			8	
C95908			10	
C95909			10	
C95910			8	
C95916			11	
C95917			11	
C95918			13	
C95919			8	
C95920			7	
MEAN			10	
S.D.			2.3	
N			15	

001063

Appendix 5
Individual Clinical Chemistry Data
Males Week 14
13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	PCOAO IU/G	Group: 2	Dose Level: 3	Dosage Unit: ppm MeFOS
C95921			10	
C95922			10	
C95923			7	
C95924			11	
C95925			10	
C95926			11	
C95927			12	
C95928			12	
C95929			12	
C95930			12	
C95936			10	
C95937			4	
C95938			9	
C95939			8	
C95940			4	
MEAN			9	
S.D.			2.7	
N			15	

001064

Appendix 5

Individual Clinical Chemistry Data

Males Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	PCOAO IU/G	Group: 3	Dose Level: 30	Dosage Unit: ppm MeFOS
C95941	10			
C95942	9			
C95943	8			
C95944	10			
C95945	7			
C95946	9			
C95947	9			
C95948	12			
C95949	6			
C95950	12			
C95956	5			
C95957	11			
C95958	10			
C95959	9			
C95960	9			
MEAN	9			
S.D.	2.0			
N	15			

001065

Appendix 5
Individual Clinical Chemistry Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

Males Week 14

ANIMAL NUMBER	PCOAO IU/G	Group: 4	Dose Level: 100	Dosage Unit: ppm MeFOS
C95961	17			
C95962	11			
C95963	12			
C95964	11			
C95965	5			
C95966	12			
C95967	38			
C95968	13			
C95969	19			
C95970	22			
C95976	22			
C95977	5			
C95978	12			
C95979	15			
C95980	16			
MEAN	15			
S.D.	8.1			
N	15			

001066

Appendix 5

Individual Clinical Chemistry Data

Females Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	PCOAO IU/G	Group: 1	Dose Level: 0	Dosage Unit: ppm MeFOS
C95981	10			
C95982	11			
C95983	5			
C95984	5			
C95985	9			
C95986	4			
C95987	4			
C95988	5			
C95989	7			
C95990	8			
C95996	3			
C95997	8			
C95998	10			
C95999	7			
C96000	7			
MEAN	7			
S.D.	2.5			
N	15			

001967

Appendix 5
Individual Clinical Chemistry Data
Females Week 14
13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	PCOAG IU/G	Group: 2	Dose Level: 3	Dosage Unit: ppm MeFOS
C96001	6			
C96002	4			
C96003	10			
C96004	5			
C96005	8			
C96006	3			
C96007	4			
C96008	5			
C96009	10			
C96010	10			
C96017	4			
C96018	8			
C96019	11			
C96020	10			
MEAN	7			
S.D.	2.9			
N	14			

001068

Appendix 5

Individual Clinical Chemistry Data

Females Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	PCOAO IU/G	Group: 3	Dose Level: 30	Dosage Unit: ppm MeFOS
-----	-----			
C96021	4			
C96022	11			
C96023	14			
C96024	5			
C96025	8			
C96026	11			
C96027	7			
C96028	9			
C96029	11			
C96030	12			
C96036	9			
C96037	10			
C96038	10			
C96039	10			
C96040	5			
MEAN	9			
S.D.	2.8			
N	15			

001069

Appendix 5

Individual Clinical Chemistry Data

Females Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	PCAO IU/G	Group: 4	Dose Level: 100	Dosage Unit: ppm MeFOS
C96041	27			
C96042	22			
C96043	16			
C96044	20			
C96045	17			
C96046	12			
C96047	36			
C96048	11			
C96049	20			
C96050	23			
C96056	19			
C96057	22			
C96058	21			
C96059	19			
C96060	8			
MEAN	20			
S.D.	6.7			
N	15			

001070

Appendix 5
Individual Clinical Urinalysis Data
Males Week 14
13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	U VOL ML	SP GR	U PH	

Group: 1	Dose Level: 0	Dosage Unit: ppm MeFOS		
C95901	18.0	1.018	7.0	
C95902	7.6	1.034	7.0	
C95903	12.5	1.026	7.0	
C95904	6.4	1.044	6.5	
C95905	4.0	1.039	7.0	
C95906	7.7	1.030	7.0	
C95909	.8	1.074	6.5	
C95918	12.0	1.025	7.0	
C95919	24.7	1.013	7.5	
C95920	5.0	1.052	6.5	
MEAN	9.9	1.036	6.9	
S.D.	7.15	.0179	.32	
N	10	10	10	
Group: 2	Dose Level: 3	Dosage Unit: ppm MeFOS		
C95921	10.2	1.023	7.5	
C95922	6.4	1.026	7.0	
C95924	10.6	1.030	7.0	
C95925	19.6	1.016	7.0	
C95926	2.6	1.040	6.5	
C95927	4.7	1.043	6.5	
C95928	17.5	1.016	7.0	
C95936	8.0	1.030	6.5	
C95937	15.3	1.020	7.5	
C95939	11.2	1.021	6.5	
MEAN	10.6	1.026	6.9	
S.D.	5.52	.0093	.39	
N	10	10	10	

001071

Appendix 5

Individual Clinical Urinalysis Data

Males Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	U VOL ML	SP GR	U PH	
Group: 3	Dose Level: 30			Dosage Unit: ppm MeFOS
C95941	9.1	1.022		7.5
C95942	21.1	1.013		6.5
C95943	17.1	1.021		7.0
C95944	3.5	1.037		7.0
C95947	10.4	1.024		7.0
C95948	13.6	1.014		7.0
C95949	47.5 a	1.002		6.0
C95950	6.0	1.037		6.5
C95958	1.3	1.060		6.5
C95960	8.0	1.030		7.0
MEAN	13.8	1.026		6.8
S.D.	13.29	.0162		.42
N	10	10		10
Group: 4	Dose Level: 100			Dosage Unit: ppm MeFOS
C95961	38.5	1.010		7.0
C95962	6.3	1.042		6.5
C95963	14.4	1.024		6.5
C95964	5.6	1.047		6.5
C95965	10.0	1.031		6.5
C95967	19.0	1.019		7.0
C95968	48.2 a	1.007		6.5
C95969	45.2 a	1.004		7.0
C95977	14.5	1.023		6.5
C95980	8.3	1.037		6.5
MEAN	21.0	1.024		6.6
S.D.	16.52	.0148		.24
N	10	10		10

a Sample was overflowing at the time of collection.

001072

Appendix 5

Individual Clinical Urinalysis Data

Males Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	U PRO MG/DL	U GLU	U KET	U BILI	U BLOOD	UROBILI

Group: 1	Dose Level: 0		Dosage Unit: ppm MeFOS			
C95901	30	-	-	-	-	-
C95902	30	-	+	-	-	-
C95903	30	-	+	-	-	-
C95904	300	-	+++	-	-	-
C95905	30	-	++	-	-	-
C95906	30	-	-	-	-	-
C95909	100	-	+++	-	-	-
C95918	30	-	+	-	++	-
C95919	30	-	+	-	-	-
C95920	30	-	+	-	-	-
Group: 2	Dose Level: 3		Dosage Unit: ppm MeFOS			
C95921	30	-	-	-	-	-
C95922	30	-	+	-	-	-
C95924	30	-	-	-	-	-
C95925	30	-	-	-	-	-
C95926	30	-	-	-	-	-
C95927	30	-	++	-	-	-
C95928	30	-	++	-	-	-
C95936	30	-	++	-	-	-
C95937	30	-	-	-	-	-
C95939	10	-	-	-	-	-

001073

Appendix 5
Individual Clinical Urinalysis Data
Males Week 14
13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	U PRO MG/DL	U GLU	U KET	U BILI	U BLOOD	UROBILI
Group: 3	Dose Level: 30					
C95941	30	-	+	-	-	-
C95942	10	-	-	-	-	-
C95943	10	-	+	-	-	-
C95944	30	-	-	-	-	-
C95947	30	-	-	-	-	-
C95948	10	-	+	-	-	-
C95949	0	-	-	-	-	-
C95950	30	-	++	-	-	-
C95958	30	-	+	-	-	-
C95960	30	-	+	-	-	-
Group: 4	Dose Level: 100					
C95961	10	-	-	-	-	-
C95962	30	-	++	-	-	-
C95963	10	-	-	-	-	-
C95964	30	-	++	-	-	-
C95965	30	-	+	-	-	-
C95967	30	-	+	-	-	-
C95968	10	-	-	-	-	-
C95969	10	-	-	-	+++	-
C95977	10	-	+++	-	-	-
C95980	10	-	+	-	-	-

091074

Appendix 5

Individual Clinical Urinalysis Data

Males Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	RBC PER HPF	WBC PER HPF	EPITH PER HPF	BACT PER HPF	CASTS PER LPF	CRYSTALS PER LPF1	URINE APP1	COMMENTS
Group: 1 Dose Level: 0 Dosage Unit: ppm MeFOS								
C95901	0	0	0	3	0	1B4D	BJ	P
C95902	1	1	1	3	0	4D	BJ	PQ
C95903	0	0	1	3	0	4D	BJ	PQ
C95904	0	0	1	2	0	2B4D	BK	PQ
C95905	0	0	1	3	0	0	BJ	P
C95906	0	0	1	2	0	4D	BJ	PQ
C95909	0	0	0	1	0	3D	CJ	-
C95918	4	1	1	3	0	4D	BJ	PQ
C95919	0	1	1	2	0	2B1D	BJ	QR
C95920	0	0	1	2	0	4D	CJ	Q
Group: 2 Dose Level: 3 Dosage Unit: ppm MeFOS								
C95921	0	1	1	2	0	4D	BKM	PQ
C95922	0	0	1	2	0	3D	BJ	Q
C95924	0	0	1	2	0	3D	BJ	PQ
C95925	0	0	1	2	0	4D	BJ	P
C95926	0	1	0	2	0	3D	BJ	P
C95927	0	0	1	3	0	4D	BJ	Q
C95928	0	1	1	2	0	1B4D	BJ	Q
C95936	0	0	1	3	0	4D	BJ	PQ
C95937	0	0	1	3	0	4D	BJ	PQ
C95939	0	1	1	2	0	2D	BJ	Q

001075

Appendix 5
Individual Clinical Urinalysis Data

Males Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	RBC PER HPF	WBC PER HPF	EPITH PER HPF	BACT PER HPF	CASTS PER LPF	CRYSTALS PER LPF1	URINE APPL	COMMENTS
Group: 3	Dose Level: 30				Dosage Unit: ppm MeFOS			
C95941	0	0	0	3	0	2B3D	BJ	QR
C95942	0	1	1	1	0	4D	BJ	PQ
C95943	0	1	0	2	0	3D	BJ	Q
C95944	0	0	0	2	0	4D	BJ	PQ
C95947	0	0	1	2	0	4D	BJ	PQ
C95948	0	0	1	2	0	4D	BJ	PQ
C95949	0	1	0	1	0	0	AJ	-
C95950	0	1	1	3	0	4D	BJ	PQ
C95958	0	0	1	1	0	4D	CJ	PQ
C95960	0	0	1	2	0	3D	BJ	PQ
Group: 4	Dose Level: 100				Dosage Unit: ppm MeFOS			
C95961	0	1	0	1	0	0	AJ	Q
C95962	0	0	1	2	0	4D	BJ	Q
C95963	0	0	1	2	0	2D	BJ	Q
C95964	0	0	1	1	0	4D	BJ	PQ
C95965	0	0	1	3	0	4D	BJ	PQ
C95967	2	0	1	2	0	4D	BJ	PQ
C95968	0	0	1	3	0	4D	AJ	Q
C95969	4	0	0	0	0	0	AJ	-
C95977	0	0	1	2	0	4D	BJ	PQ
C95980	0	1	1	2	0	3D	BJ	Q

001076

Appendix 5
Individual Clinical Urinalysis Data
Females Week 14
13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	U VOL ML	SP GR	U PH	
-----	-----	-----	-----	-----
Group: 1	Dose Level: 0		Dosage Unit: ppm MeFOS	
C95983	4.3	1.027	6.5	
C95984	17.5	1.010	7.0	
C95985	25.6	1.006	7.0	
C95986	36.0	1.005	6.5	
C95987	4.3	1.027	6.5	
C95988	41.0	1.002	6.5	
C95990	2.6	1.034	6.5	
C95996	24.3	1.008	7.0	
C95997	5.8	1.023	6.5	
C96000	8.2	1.017	7.0	
MEAN	17.0	1.016	6.7	
S.D.	14.13	.0112	.26	
N	10	10	10	
Group: 2	Dose Level: 3		Dosage Unit: ppm MeFOS	
C96001	3.1	1.027	6.5	
C96002	8.2	1.022	6.5	
C96003	2.3	1.036	6.5	
C96006	7.6	1.022	7.5	
C96007	30.4	1.006	6.5	
C96008	26.1	1.006	7.5	
C96010	12.2	1.015	6.5	
C96019	3.8	1.028	6.5	
C96020	9.1	1.014	7.0	
MEAN	11.4	1.020	6.8	
S.D.	10.11	.0102	.44	
N	9	9	9	

001077

Covance 6329-225
3M T-6314.1

Appendix 5
Individual Clinical Urinalysis Data
Females Week 14
13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	U VOL ML	SP GR	U PH	

Group: 3	Dose Level: 30	Dosage Unit: ppm MeFOS		
C96021	9.0	1.021	7.0	
C96022	9.9	1.019	6.5	
C96023	22.1	1.009	6.5	
C96024	41.5	1.004	7.0	
C96025	2.4	1.030	6.5	
C96026	43.8	1.006	6.5	
C96027	6.3	1.025	7.0	
C96030	7.6	1.020	6.5	
C96038	11.3	1.016	7.0	
C96040	12.8	1.013	7.0	
MEAN	16.7	1.016	6.8	
S.D.	14.62	.0084	.26	
N	10	10	10	
Group: 4	Dose Level: 100	Dosage Unit: ppm MeFOS		
C96042	7.0	1.029	6.5	
C96044	2.4	1.039	6.5	
C96045	19.0	1.014	6.5	
C96046	7.2	1.026	6.5	
C96047	18.7	1.010	6.5	
C96048	18.0	1.012	6.5	
C96049	5.3	1.024	6.5	
C96050	9.1	1.026	6.5	
C96056	40.5	1.004	7.0	
C96060	29.5	1.007	7.0	
MEAN	15.7	1.019	6.6	
S.D.	12.05	.0113	.21	
N	10	10	10	

001078

Appendix 5
Individual Clinical Urinalysis Data

Females Week 14
13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	U PRO MG/DL	U GLU	U KET	U BILI	U BLOOD	UROBILI
Group: 1	Dose Level: 0					
	Dosage Unit: ppm MeFOS					
C95983	30	-	-	-	-	-
C95984	0	-	-	-	-	-
C95985	0	-	-	-	-	-
C95986	0	-	-	-	-	-
C95987	10	-	-	-	-	-
C95988	0	-	-	-	-	-
C95990	30	-	-	-	-	-
C95996	0	-	-	-	-	-
C95997	10	-	-	-	-	-
C96000	10	-	-	-	-	-
Group: 2	Dose Level: 3					
	Dosage Unit: ppm MeFOS					
C96001	10	-	-	-	-	-
C96002	10	-	-	-	-	-
C96003	10	-	-	-	-	-
C96006	10	-	-	-	-	-
C96007	0	-	-	-	-	-
C96008	10	-	-	-	-	-
C96010	10	-	-	-	-	-
C96019	10	-	-	-	-	-
C96020	0	-	-	-	-	-

001079

Appendix 5

Individual Clinical Urinalysis Data

Females Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	U PRO MG/DL	U GLU	U KET	U BILI	U BLOOD	UROBILI

Group: 3	Dose Level: 30	Dosage Unit: ppm MeFOS				
C96021	10	-	-	-	-	-
C96022	0	-	-	-	-	-
C96023	10	-	-	-	-	-
C96024	0	-	-	-	-	-
C96025	10	-	-	-	-	-
C96026	10	-	-	-	-	-
C96027	10	-	-	-	-	-
C96030	10	-	-	-	-	-
C96038	10	-	-	-	-	-
C96040	0	-	-	-	-	-
Group: 4	Dose Level: 100	Dosage Unit: ppm MeFOS				
C96042	10	-	-	-	-	-
C96044	30	-	-	-	-	-
C96045	10	-	-	-	-	-
C96046	10	-	-	-	-	-
C96047	0	-	-	-	-	-
C96048	10	-	-	-	-	-
C96049	10	-	-	-	-	-
C96050	10	-	-	-	-	-
C96056	0	-	-	-	-	-
C96060	0	-	-	-	-	-

001030

Appendix 5
Individual Clinical Urinalysis Data

Females Week 14													
13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS													
ANIMAL NUMBER	RBC PER HPF	WBC PER HPF	EPITH PER HPF	BACT PER HPF	CASTS PER LPF	CRYSTALS PER LPF1	URINE APPL	COMMENTS					
Group: 1	Dose Level: 0			Dosage Unit: ppm MeFOS									
C95983	0	1	1	2	0	2D	BJ						
C95984	0	0	1	1	0	1D	AJ						
C95985	0	0	0	1	0	0	AJ						
C95986	0	1	1	1	0	0	AJ						
C95987	0	1	1	2	0	4D	BJ						
C95988	0	0	1	3	0	4D	AJ						
C95990	0	0	1	1	0	2D	BJ						
C95996	0	0	0	1	0	2D	AJ						
C95997	0	0	1	3	0	4D	BJ						
C96000	0	0	1	1	0	4D	BJ						
Group: 2	Dose Level: 3			Dosage Unit: ppm MeFOS									
C96001	0	0	1	1	0	4D	BJ						
C96002	0	0	1	2	0	4D	BJ						
C96003	0	0	1	3	0	4D	BJ						
C96006	0	1	1	2	0	4D	BJ						
C96007	0	0	0	1	0	0	AJ						
C96008	0	0	1	3	0	1D	AJ						
C96010	0	1	1	2	0	0	BJ						
C96019	0	0	1	1	0	4D	BJ						
C96020	0	0	0	1	0	2D	BJ						

001031

Appendix 5
Individual Clinical Urinalysis Data

Females Week 14

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO
ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER	RBC PER HPF	WBC PER HPF	EPITH PER HPF	BACT PER HPF	CASTS PER LPF	CRYSTALS PER LPF1	URINE APPL	COMMENTS
Group: 3	Dose Level: 30				Dosage Unit: ppm MeFOS			
C96021	0	1	1	2	0	4D	BJ	P
C96022	0	0	0	1	0	3D	BJ	-
C96023	0	0	1	1	0	0	AJ	-
C96024	1	0	1	1	0	0	AJ	-
C96025	0	0	1	1	0	0	BJ	-
C96026	0	0	1	1	0	1B	AJ	-
C96027	0	1	1	2	0	4D	BJ	P
C96030	0	1	1	1	0	0	BJ	-
C96038	0	0	1	2	0	0	BJ	-
C96040	0	0	1	1	0	0	BJ	P
Group: 4	Dose Level: 100				Dosage Unit: ppm MeFOS			
C96042	0	0	1	1	0	2D	BJ	-
C96044	0	1	0	2	0	1D	BJ	-
C96045	0	0	1	2	0	0	AJ	-
C96046	0	1	1	2	0	4D	BJ	P
C96047	0	1	1	1	0	1B	AJ	-
C96048	0	0	1	1	0	0	BJ	P
C96049	0	0	1	2	0	4D	BJ	P
C96050	0	1	1	2	0	4D	BJ	-
C96056	0	0	0	2	0	0	AJ	-
C96060	0	0	1	1	0	0	AJ	-

001082

APPENDIX 6

Individual Anatomic Pathology Data

001033

APPENDIX 6

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 1ANIMAL NUMBER: C95901
SEX: MALE
DOSE GROUP: 1
SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98
STUDY WEEK OF DEATH: 14
TERMINAL BODY WEIGHT: 477.9 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 12:16
PROSECTOR: LAURIE J. SCHULLER
RECORDER: TERRI L. WILLIAMS
POST-FIX WEIGHTER: NOT AVAILABLE
PATHOLOGIST: JOHN EIGHMY
WEIGHER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0507	.0106 %	.0234	WEIGHT TAKEN
BRAIN (BR)	2.1637	.4528 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	3.6235	.7582 %	1.6747	WEIGHT TAKEN
LIVER (LI)	13.2353	2.7695 %	6.1170	WEIGHT TAKEN
SPLEEN (SP)	.6399	.1339 %	.2957	WEIGHT TAKEN
TESTIS (TE)	3.0353	.6351 %	1.4028	WEIGHT TAKEN
THYMUS (TH)	.3522	.0737 %	.1628	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0353	.0074 %	.0163	WEIGHT TAKEN

O B S E R V A T I O N S

NECROPSY

HISTOPATHOLOGY

LUNG (LU) :
-MINERALIZATION, VASCULAR, -MINIMAL
LIVER (LI) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
HEART (HT) :
-DEGENERATION/NECROSIS, -MINIMAL
THYROID (TY) :
-CYST, ULTIMOBRANCHIAL, -PRESENT
PANCREAS (PA) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
PROSTATE (PR) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

C01084

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 2

ANIMAL NUMBER: C95901 SEX: MALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 477.9 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 12:16 PROSECTOR: LAURIE J. SCHULLER RECORDER: TERRI L. WILLIAMS
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN),
SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA),
AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL),
CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG),
AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP)

001085

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 3

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95902 SEX: MALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 500.6 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:38 PROSECTOR: JODI THIEREN REORDER: TERRI L. WILLIAMS
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0754	.0151 %	.0353	WEIGHT TAKEN
BRAIN (BR)	2.1335	.4262 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	4.1240	.8238 %	1.9330	WEIGHT TAKEN
LIVER (LI)	15.9820	3.1926 %	7.4910	WEIGHT TAKEN
SPLEEN (SP)	.8310	.1660 %	.3895	WEIGHT TAKEN
TESTIS (TE)	3.5597	.7111 %	1.6685	WEIGHT TAKEN
THYMUS (TH)	.2540	.0507 %	.1191	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0198	.0040 %	.0093	WEIGHT TAKEN

NECROPSY OBSERVATIONS HISTOPATHOLOGY

THYMUS (TH) :
-MOTTLED; BILATERAL; RED AND TAN
LN, MANDIBULAR (MN) :
-DIFFUSELY RED; BILATERAL; DARK RED
GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
LUNG (LU) :
-MINERALIZATION, VASCULAR, -MINIMAL
KIDNEY (KD) :
-MINERALIZATION, TUBULAR, -MINIMAL
THYMUS (TH) :
-HEMORRHAGE, -MINIMAL
LN, MANDIBULAR (MN) :
-HEMORRHAGE, -SLIGHT
PROSTATE (PR) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -SLIGHT

001036

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 4

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95902 SEX: MALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 500.6 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:38 PROSECTOR: JODI THIEREN RECORDER: TERRI L. WILLIAMS
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC),
ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG),
HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV),
TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN),
SPLEEN (SP), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA),
AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE),
ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS),
SKIN (SK), URINARY BLADDER (UB), SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP)

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 5

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95903 SEX: MALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 482.8 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:07 PROSECTOR: KIMBERLY DURLAND REORDER: TERRI L. WILLIAMS
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0585	.0121 %	.0273	WEIGHT TAKEN
BRAIN (BR)	2.1393	.4431 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	3.4896	.7228 %	1.6312	WEIGHT TAKEN
LIVER (LI)	13.3362	2.7623 %	6.2339	WEIGHT TAKEN
SPLEEN (SP)	.8041	.1665 %	.3759	WEIGHT TAKEN
TESTIS (TE)	3.5494	.7352 %	1.6591	WEIGHT TAKEN
THYMUS (TH)	.2730	.0565 %	.1276	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0280	.0058 %	.0131	WEIGHT TAKEN

NECROPSY OBSERVATIONS HISTOPATHOLOGY

LUNG (LU) :
-HEMORRHAGE, -MINIMAL
-INFILTRATE, MACROPHAGE, ALVEOLAR, -MINIMAL
LIVER (LI) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL
THYROID (TY) :
-ECTOPIC THYMUS, -PRESENT
LN, MANDIBULAR (MN) :
-HEMORRHAGE, -MINIMAL
SKIN (SK) :
-EXUDATE, EPIDERMAL SURFACE, -MINIMAL
-INFLAMMATION, SUBACUTE, -MINIMAL

LN, MANDIBULAR (MN) :
-MOTTLED; BILATERAL; RED AND TAN

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN

001038

APPENDIX 6

Covance 6329-225
3M T-6314.1

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 6

ANIMAL NUMBER: C95903 SEX: MALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 482.8 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:07 PROSECTOR: KIMBERLY DURLAND REORDER: TERRI L. WILLIAMS
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LONG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TV), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG),
HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV),
TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN),
SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA),
AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE),
ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS),
URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP)

APPENDIX 6
Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 7

ANIMAL NUMBER: C95904 SEX: MALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 502.8 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 12:07 PROSECTOR: JERALD W. JOHNSON REORDER: WENDY CHERRY
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0551	.0110 %	.0249	WEIGHT TAKEN
BRAIN (BR)	2.2087	.4393 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	4.1957	.8345 %	1.8996	WEIGHT TAKEN
LIVER (LI)	14.9025	2.9639 %	6.7472	WEIGHT TAKEN
SPLEEN (SP)	.8734	.1737 %	.3954	WEIGHT TAKEN
TESTIS (TE)	3.4791	.6919 %	1.5752	WEIGHT TAKEN
THYMUS (TH)	.3279	.0652 %	.1485	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0119	.0024 %	.0054	WEIGHT TAKEN

OBSERVATIONS

NECROPSY

HISTOPATHOLOGY

LIVER (LI) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL
KIDNEY (KD) :
-BASOPHILIC TUBULES, CORTEX, -MINIMAL
-THICKENING, BASEMENT MEMBRANE, -MINIMAL
HEART (HT) :
-DEGENERATION/NECROSIS, -MINIMAL
THYMUS (TH) :
-HEMORRHAGE, -MINIMAL
THYROID (TY) :
-ONE EXAMINED, -PRESENT
PARATHYROID (PT) :
-ONE EXAMINED, -PRESENT
PROSTATE (PR) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL

THYROID (TY) :
>NOTE:>TRIM COMMENT: ONE MISSING AT TIME OF TRIMMING
PARATHYROID (PT) :
>NOTE:>TRIM COMMENT: ONE MISSING AT TIME OF TRIMMING

001090

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 8

ANIMAL NUMBER: C95904
DATE OF DEATH: 12/02/98
DATE AND TIME OF NECROPSY: 12/02/98 12:07
POST-FIX WEIGHER: NOT AVAILABLE
SEX: MALE
STUDY DAY OF DEATH: 93
DOSE GROUP: 1
SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
STUDY WEEK OF DEATH: 14
TERMINAL BODY WEIGHT: 502.8 GRAMS
PROSECTOR: JERALD W. JOHNSON
RECORDED: WENDY CHERRY
PATHOLOGIST: JOHN EIGHY
WEIGHER: LYNN HORNER

OBSERVATIONS (CONTINUED)

NECROPSY

HISTOPATHOLOGY

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN), SPLEEN (SP),
LN, MESENTERIC (MS), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PT), STOMACH, GL (ST),
STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE),
LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK),
URINARY BLADDER (UB), SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP)

001091

APPENDIX 6
Individual Anatomic Pathology Data

PAGE: 9

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95905 SEX: MALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 477.8 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:03 PROSECTOR: NANCY DIEDRICH REORDER: TERRI L. WILLIAMS
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATISTICS
ADRENAL (AD)	.0413	.0086 %	.0188	WEIGHT TAKEN
BRAIN (BR)	2.1962	.4596 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	3.2242	.6748 %	1.4681	WEIGHT TAKEN
LIVER (LI)	13.4331	2.8114 %	6.1165	WEIGHT TAKEN
SPLEEN (SP)	.8531	.1785 %	.3884	WEIGHT TAKEN
TESTIS (TE)	3.1958	.6689 %	1.4551	WEIGHT TAKEN
THYMUS (TH)	.3319	.0695 %	.1511	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0209	.0044 %	.0095	WEIGHT TAKEN

OBSERVATIONS
NECROPSY
HISTOPATHOLOGY

LIVER (LI) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL
THYMUS (TH) :
-HEMORRHAGE, -MINIMAL
PROSTATE (PR) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -SLIGHT

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

001092

APPENDIX 6

Covance 6329-225
3M T-6314.1

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 10

ANIMAL NUMBER: C95905 SEX: MALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL, SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 477.8 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:03 PROSECTOR: NANCY DIEDRICH RECORDED: TERRI L. WILLIAMS
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC),
ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP)

601093

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 11

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95906
SEX: MALE
DATE OF DEATH: 12/02/98
STUDY DAY OF DEATH: 93
DATE AND TIME OF NECROPSY: 12/02/98 10:06
POST-FIX WEIGHTER: NOT AVAILABLE
DOSE GROUP: 1
SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
TERMINAL BODY WEIGHT: 532.3 GRAMS
PROSECTOR: JOHN S. HALFORD
RECORDED: WENDY CHERRY
PATHOLOGIST: JOHN EIGHMY
WEIGHER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS	WEIGHT TAKEN
ADRENAL (AD)	.0596	.0112 %	.0270		WEIGHT TAKEN
BRAIN (BR)	2.2070	.4146 %	1.0000		WEIGHT TAKEN
KIDNEY (KD)	3.6715	.6897 %	1.6636		WEIGHT TAKEN
LIVER (LI)	17.6850	3.3224 %	8.0131		WEIGHT TAKEN
SPLEEN (SP)	.9226	.1733 %	.4180		WEIGHT TAKEN
TESTIS (TE)	3.5489	.6667 %	1.6080		WEIGHT TAKEN
THYMUS (TH)	.2941	.0553 %	.1333		WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0273	.0051 %	.0124		WEIGHT TAKEN

HISTOPATHOLOGY

OBSERVATIONS

NECROPSY

LUNG (LU) :
-MINERALIZATION, VASCULAR, -MINIMAL
THYROID (TY) :
-CYST, ULTIMOBRANCHIAL, -PRESENT
PANCREAS (PA) :
-PIGMENT, BROWN, -MINIMAL

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 12

ANIMAL NUMBER: C95906
DATE OF DEATH: 12/02/98
DATE AND TIME OF NECROPSY: 12/02/98 10:06
POST-FIX WEIGHTER: NOT AVAILABLE

SEX: MALE
STUDY DAY OF DEATH: 93
STUDY WEEK OF DEATH: 14
PROSECTOR: JOHN S. HALFORD
PATHOLOGIST: JOHN EIGHMY

DOSE GROUP: 1
SACRIFICE STATUS: SCHEDULED, TERMINAL, SACRIFICE
TERMINAL BODY WEIGHT: 532.3 GRAMS
REORDER: WENDY CHERRY
WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), PARATHYROID (PT), ADRENAL, CORTEX (AC),
ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), DUODENUM (DU),
JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG),
HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV),
TESTIS (TE), EPIDIDYMIDES (EP)

001095

APPENDIX 6
Individual Anatomic Pathology Data

PAGE: 13

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95907	SEX: MALE	DOSE GROUP: 1	SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE	
DATE OF DEATH: 12/02/98	STUDY DAY OF DEATH: 93	STUDY WEEK OF DEATH: 14	TERMINAL BODY WEIGHT: 469.2 GRAMS	
DATE AND TIME OF NECROPSY: 12/02/98 7:56	PROSECTOR: TODD ACKER		RECORDER: WENDY CHERRY	
POST-FIX WEIGHER: NOT AVAILABLE	PATHOLOGIST: JOHN EIGHMY		WEIGHER: LYNN HORNER	
ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0456	.0097 %	.0216	WEIGHT TAKEN
BRAIN (BR)	2.1146	.4507 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	3.3900	.7225 %	1.6031	WEIGHT TAKEN
LIVER (LI)	12.7160	2.7101 %	6.0134	WEIGHT TAKEN
SPLEEN (SP)	.6803	.1450 %	.3217	WEIGHT TAKEN
TESTIS (TE)	3.1705	.6757 %	1.4993	WEIGHT TAKEN
THYMUS (TH)	.3000	.0639 %	.1419	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0298	.0064 %	.0141	WEIGHT TAKEN

NECROPSY	OBSERVATIONS	HISTOPATHOLOGY
	THYROID (TY) :	
	-CYST, ULTIMOBRANCHIAL, -PRESENT	
	PANCREAS (PA) :	
	-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL	
	-PIGMENT, BROWN, -MINIMAL	
	RECTUM (RE) :	
	-PARASITISM, -PRESENT	
	PROSTATE (PR) :	
	-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL	

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

001096

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 14

ANIMAL NUMBER: C95907 SEX: MALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 469.2 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 7:56 PROSECTOR: TODD ACKER REORDER: WENDY CHERRY
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), PARATHYROID (PT), ADRENAL, CORTEX (AC),
ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), DUODENUM (DU),
JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG),
AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP)

001037

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 15

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95908		SEX: MALE		DOSE GROUP: 1		SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE	
DATE OF DEATH: 12/02/98		STUDY DAY OF DEATH: 93		STUDY WEEK OF DEATH: 14		TERMINAL BODY WEIGHT: 505.0 GRAMS	
DATE AND TIME OF NECROPSY: 12/02/98 7:36		PROSECTOR: JOHN S. HALFORD		RECORDER: WENDY CHERRY			
POST-FIX WEIGHT: NOT AVAILABLE		PATHOLOGIST: JOHN EIGHMY		WEIGHER: LYNN HORNER			

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0571	.0113 %	.0269	WEIGHT TAKEN
BRAIN (BR)	2.1256	.4209 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	3.4921	.6915 %	1.6429	WEIGHT TAKEN
LIVER (LI)	13.0677	2.5877 %	6.1478	WEIGHT TAKEN
SPLEEN (SP)	.8254	.1634 %	.3883	WEIGHT TAKEN
TESTIS (TE)	3.3149	.6564 %	1.5595	WEIGHT TAKEN
THYMUS (TH)	.5078	.1006 %	.2389	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0193	.0038 %	.0091	WEIGHT TAKEN

NECROPSY	OBSERVATIONS	HISTOPATHOLOGY
LN, MANDIBULAR (MN) :	THYROID (TY) :	
-DIFFUSELY RED; BILATERAL	-CYST, ULTIMOBRANCHIAL, -PRESENT	
	LN, MANDIBULAR (MN) :	
	-HEMORRHAGE, -MINIMAL	
	PROSTATE (PR) :	
	-INFILTRATE, LYMPHOHISTIOCYTIC, -SLIGHT	
GENERAL COMMENT (GC) :		
-BONE MARROW SMEAR TAKEN		

001038

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 16

ANIMAL NUMBER: C95908 SEX: MALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 505.0 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 7:36 PROSECTOR: JOHN S. HALFORD REORDER: WENDY CHERRY
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG),
HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV),
TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), PARATHYROID (PT), ADRENAL, CORTEX (AC),
ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG),
HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), SEMINAL VESICLES (SV), TESTIS (TE),
EPIDIDYMIDES (EP)

001039

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 17

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95909	SEX: MALE	DOSE GROUP: 1	SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98	STUDY DAY OF DEATH: 93	STUDY WEEK OF DEATH: 14	TERMINAL BODY WEIGHT: 434.0 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:29	PROSECTOR: JOHN S. HALFORD	REORDER: WENDY CHERRY	
POST-FIX WEIGHTER: NOT AVAILABLE	PATHOLOGIST: JOHN EIGHMY	WEIGHER: LYNN HORNER	
	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO
ORGAN NAME			ORGAN STATUS
ADRENAL (AD)	.0629	.0145 %	.0287 WEIGHT TAKEN
BRAIN (BR)	2.1932	.5053 %	1.0000 WEIGHT TAKEN
KIDNEY (KD)	3.1195	.7188 %	1.4224 WEIGHT TAKEN
LIVER (LI)	11.3174	2.6077 %	5.1602 WEIGHT TAKEN
SPLEEN (SP)	.6744	.1554 %	.3075 WEIGHT TAKEN
TESTIS (TE)	3.6084	.8314 %	1.6453 WEIGHT TAKEN
THYMUS (TH)	.3922	.0904 %	.1788 WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0256	.0059 %	.0117 WEIGHT TAKEN

NECROPSY	OBSERVATIONS	HISTOPATHOLOGY
	LUNG (LU) :	-INFILTRATE, MACROPHAGE, ALVEOLAR, -MINIMAL
	LIVER (LI) :	-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
	THYMUS (TH) :	-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL
	THYROID (TY) :	-HEMORRHAGE, -MINIMAL
	HARDERIAN GLAND (HG) :	-CYST, ULTIMOBRANCHIAL, -PRESENT
	PROSTATE (PR) :	-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
		-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-MALOCCLUSION; BILATERAL, UPPER INCISORS

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 18

ANIMAL NUMBER: C95909 SEX: MALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 434.0 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:29 PROSECTOR: JOHN S. HALFORD RECORDER: WENDY CHERRY
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN),
SPLEEN (SP), LN, MESENTERIC (MS), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO),
PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL),
CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), AUDITORY SEB GL (AS), SKIN (SK),
URINARY BLADDER (UB), SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP)

001101

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 19

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95910 SEX: MALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 573.8 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 7:54 PROSECTOR: NANCY DIEDRICH RECORDED: STEVE VAN ADESTINE
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0412	.0072 %	.0183	WEIGHT TAKEN
BRAIN (BR)	2.2483	.3918 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	4.4547	.7764 %	1.9814	WEIGHT TAKEN
LIVER (LI)	16.2577	2.8333 %	7.2311	WEIGHT TAKEN
SPLEEN (SP)	.8328	.1451 %	.3704	WEIGHT TAKEN
TESTIS (TE)	3.6513	.6363 %	1.6240	WEIGHT TAKEN
THYMUS (TH)	.5244	.0914 %	.2332	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0345	.0060 %	.0153	WEIGHT TAKEN

NECROPSY	OBSERVATIONS	HISTOPATHOLOGY
	LUNG (LU) :	-INFILTRATE, MACROPHAGE, ALVEOLAR, -MINIMAL
		-MINERALIZATION, VASCULAR, -MINIMAL
	LIVER (LI) :	-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
	KIDNEY (KD) :	-BASOPHILIC TUBULES, CORTEX, -MINIMAL
	THYROID (TY) :	-CYST, ULTIMOBRANCHIAL, -PRESENT
		-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
	PROSTATE (PR) :	-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
GENERAL COMMENT (GC) :		
-BONE MARROW SMEAR TAKEN		
-NO MACROSCOPIC LESIONS		

APPENDIX 6

Covance 6329-225
3M T-6314.1

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 20

ANIMAL NUMBER: C95910 SEX: MALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 573.8 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 7:54 PROSECTOR: NANCY DIEDRICH
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY REORDER: STEVE VAN ADESTINE
WEIGHER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), TRACHEA (TR), ESOPHAGUS (ES), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN),
SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA),
AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE),
ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG),
AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP)

001103

APPENDIX 6

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 21ANIMAL NUMBER: C95911
DATE OF DEATH: 10/01/98
DATE AND TIME OF NECROPSY: 10/01/98 11:59
POST-FIX WEIGHER: NOT AVAILABLE
SEX: MALE
STUDY DAY OF DEATH: 31
DOSE GROUP: 1
SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
TERMINAL BODY WEIGHT: 324.5 GRAMS
REORDER: KEVIN BILLINGS
WEIGHER: MICHELE ZIMA
PROSECTOR: TONY HANFELD
PATHOLOGIST: JOHN EIGHMY

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	---	---	---	NOT TAKEN
BRAIN (BR)	---	---	---	NOT TAKEN
KIDNEY (KD)	---	---	---	NOT TAKEN
LIVER (LI)	---	---	---	WEIGHT TAKEN
SPLEEN (SP)	9.9529	3.0671 %	---	NOT TAKEN
TESTIS (TE)	---	---	---	NOT TAKEN
THYMUS (TH)	---	---	---	NOT TAKEN
THYROID/PARATHYR (TT)	---	---	---	NOT TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI), GENERAL COMMENT (GC)

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 22

ANIMAL NUMBER: C95912 SEX: MALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
DATE OF DEATH: 10/01/98 STUDY DAY OF DEATH: 31 STUDY WEEK OF DEATH: 5 TERMINAL BODY WEIGHT: 376.3 GRAMS
DATE AND TIME OF NECROPSY: 10/01/98 11:31 PROSECTOR: MATTHEW AXTMAN RECORDER: KEVIN BILLINGS
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: MICHELE ZIMA

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	---	---	---	NOT TAKEN
BRAIN (BR)	---	---	---	NOT TAKEN
KIDNEY (KD)	---	---	---	NOT TAKEN
LIVER (LI)	---	---	---	WEIGHT TAKEN
SPLEEN (SP)	12.7772	3.3955 %	---	NOT TAKEN
TESTIS (TE)	---	---	---	NOT TAKEN
THYMUS (TH)	---	---	---	NOT TAKEN
THYROID/PARATHYR (TT)	---	---	---	NOT TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI), GENERAL COMMENT (GC)

001105

APPENDIX 6

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 23

ANIMAL NUMBER: C95913 SEX: MALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
DATE OF DEATH: 10/01/98 STUDY DAY OF DEATH: 31 STUDY WEEK OF DEATH: 5 TERMINAL BODY WEIGHT: 367.1 GRAMS
DATE AND TIME OF NECROPSY: 10/01/98 11:45 PROSECTOR: WENDY CHERRY REORDER: KEVIN BILLINGS
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: MICHELE ZIMA

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	---	---	---	NOT TAKEN
BRAIN (BR)	---	---	---	NOT TAKEN
KIDNEY (KD)	---	---	---	NOT TAKEN
LIVER (LI)	11.6383	3.1703 %	---	WEIGHT TAKEN
SPLEEN (SP)	---	---	---	NOT TAKEN
TESTIS (TE)	---	---	---	NOT TAKEN
THYMUS (TH)	---	---	---	NOT TAKEN
THYROID/PARATHYR (TT)	---	---	---	NOT TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI), GENERAL COMMENT (GC)

001106

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 24

ANIMAL NUMBER: C95914
SEX: MALE
DOSE GROUP: 1
SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
DATE OF DEATH: 10/01/98
STUDY DAY OF DEATH: 31
STUDY WEEK OF DEATH: 5
TERMINAL BODY WEIGHT: 346.2 GRAMS
DATE AND TIME OF NECROPSY: 10/01/98 11:36
PROSECTOR: MATTHEW AXTMAN
REORDER: KEVIN BILLINGS
POST-FIX WEIGHER: NOT AVAILABLE
PATHOLOGIST: JOHN EIGHMY
WEIGHER: MICHELE ZIMA

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	---	---	---	NOT TAKEN
BRAIN (BR)	---	---	---	NOT TAKEN
KIDNEY (KD)	---	---	---	NOT TAKEN
LIVER (LI)	---	---	---	WEIGHT TAKEN
SPLEEN (SP)	11.8655	3.4274 %	---	NOT TAKEN
TESTIS (TE)	---	---	---	NOT TAKEN
THYMUS (TH)	---	---	---	NOT TAKEN
THYROID/PARATHYR (TT)	---	---	---	NOT TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EV), BRAIN (BR), SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI), GENERAL COMMENT (GC)

001107

APPENDIX 6

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 25ANIMAL NUMBER: C95915
SEX: MALE
DOSE GROUP: 1
SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
DATE OF DEATH: 10/01/98
STUDY DAY OF DEATH: 31
STUDY WEEK OF DEATH: 5
TERMINAL BODY WEIGHT: 262.0 GRAMS
DATE AND TIME OF NECROPSY: 10/01/98 11:42
PROSECTOR: WENDY CHERRY
REORDER: KEVIN BILLINGS
POST-FIX WEIGHT: NOT AVAILABLE
PATHOLOGIST: JOHN EIGHMY
WEIGHER: MICHELE ZIMA

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	---	---	---	NOT TAKEN
BRAIN (BR)	---	---	---	NOT TAKEN
KIDNEY (KD)	---	---	---	NOT TAKEN
LIVER (LI)	---	---	---	WEIGHT TAKEN
SPLEEN (SP)	7.5432	2.8791 %	---	NOT TAKEN
TESTIS (TE)	---	---	---	NOT TAKEN
THYMUS (TH)	---	---	---	NOT TAKEN
THYROID/PARATHYR (TT)	---	---	---	NOT TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI), GENERAL COMMENT (GC)

001108

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.113-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 26

ANIMAL NUMBER: C95916 SEX: MALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
 DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 557.9 GRAMS
 DATE AND TIME OF NECROPSY: 12/02/98 8:32 PROSECTOR: NANCY DIEDRICH REORDER: STEVE VAN ADESTINE
 POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0570	.0102 %	.0246	WEIGHT TAKEN
BRAIN (BR)	2.3125	.4145 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	4.2219	.7567 %	1.8257	WEIGHT TAKEN
LIVER (LI)	15.3513	2.7516 %	6.6384	WEIGHT TAKEN
SPLEEN (SP)	.8545	.1532 %	1.3695	WEIGHT TAKEN
TESTIS (TE)	3.7162	.6661 %	1.6070	WEIGHT TAKEN
THYMUS (TH)	.3622	.0649 %	.1566	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0252	.0045 %	.0109	WEIGHT TAKEN

O B S E R V A T I O N S

NECROPSY

HISTOPATHOLOGY

KIDNEY (KD) :
 -LARGE PELVIS(ES); RIGHT
 THYMUS (TH) :
 -RED FOCUS(I)/AREA(S); DIFFUSE: MULTIPLE DARK RED AREAS, UP TO 3 X 2 MM
 STOMACH, GL (ST) :
 >NOTE:>TRIMMING COMMENT: STOMACH NOT FOUND IN WET TISSUE
 STOMACH, NONGL (SU) :
 >NOTE:>TRIMMING COMMENT: STOMACH NOT FOUND IN WET TISSUE
 DUODENUM (DU) :
 >NOTE:>TRIMMING COMMENT: DUODENUM NOT FOUND IN WET TISSUE

LUNG (LU) :
 -INFLAMMATION, SUBACUTE, -MINIMAL
 KIDNEY (KD) :
 -DILATATION, PELVIC, -MINIMAL
 THYMUS (TH) :
 -HEMORRHAGE, -MINIMAL
 STOMACH, GL (ST) :
 >TISSUE MISSING
 STOMACH, NONGL (SU) :
 >TISSUE MISSING
 PANCREAS (PA) :
 -INFLAMMATION, CHRONIC-ACTIVE, -SLIGHT
 DUODENUM (DU) :
 >TISSUE MISSING
 PROSTATE (PR) :
 -INFILTRATE, LYMPHOHISTIOCYTIC, -SLIGHT

001109

APPENDIX 6
Individual Anatomic Pathology Data

PAGE: 27

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95916 SEX: MALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 557.9 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:32 PROSECTOR: NANCY DIEDRICH REORDER: STEVE VAN ADESTINE
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHT: TONY HANFELD

O B S E R V A T I O N S (CONTINUED)

HISTOPATHOLOGY

NECROPSY

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC),
ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN),
SPLEEN (SP), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA),
AORTA (AO), PITUITARY (PI), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP)

001110

APPENDIX 6

Covance 6329-225
3M T-6314.1

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 28ANIMAL NUMBER: C95917
SEX: MALE
DOSE GROUP: 1
SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98
STUDY DAY OF DEATH: 93
STUDY WEEK OF DEATH: 14
TERMINAL BODY WEIGHT: 531.9 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 7:37
PROSECTOR: JERALD W. JOHNSON
REORDER: WENDY CHERRY
POST-FIX WEIGHER: NOT AVAILABLE
PATHOLOGIST: JOHN EIGHMY
WEIGHER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0545	.0102 %	.0261	WEIGHT TAKEN
BRAIN (BR)	2.0848	.3920 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	4.3299	.8140 %	2.0769	WEIGHT TAKEN
LIVER (LI)	13.8740	2.6084 %	6.6548	WEIGHT TAKEN
SPLEEN (SP)	.7099	.1335 %	.3405	WEIGHT TAKEN
TESTIS (TE)	3.4976	.6576 %	1.6777	WEIGHT TAKEN
THYMUS (TH)	.3380	.0635 %	.1621	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0278	.0052 %	.0133	WEIGHT TAKEN

NECROPSY

OBSERVATIONS

HISTOPATHOLOGY

LUNG (LU) :
-MINERALIZATION, VASCULAR, -MINIMAL
LIVER (LI) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL
THYROID (TY) :
-CYST, ULTIMOBRANCHIAL, -PRESENT
RECTUM (RE) :
-PARASITISM, -PRESENT
PROSTATE (PR) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -SLIGHT

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

001111

APPENDIX 6

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 29

ANIMAL NUMBER: C95917 SEX: MALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 531.9 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 7:37 PROSECTOR: JERALD W. JOHNSON RECORDED: WENDY CHERRY
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHTMY WEIGHTER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN),
SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA),
AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE),
ILEUM (IL), CECUM (CE), COLON (CO), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG),
AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP)

001112

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 30

ANIMAL NUMBER: C95918
SEX: MALE
DOSE GROUP: 1
SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98
STUDY DAY OF DEATH: 93
STUDY WEEK OF DEATH: 14
TERMINAL BODY WEIGHT: 494.7 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:58
PROSECTOR: LAURIE J. SCHULLER
REORDER: TERRI L. WILLIAMS
POST-FIX WEIGHT: NOT AVAILABLE
PATHOLOGIST: JOHN EIGHNY
WEIGHER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0547	.0111 %	.0261	WEIGHT TAKEN
BRAIN (BR)	2.0931	.4231 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	3.6402	.7358 %	1.7391	WEIGHT TAKEN
LIVER (LI)	14.9341	3.0188 %	7.1349	WEIGHT TAKEN
SPLEEN (SP)	.7976	.1612 %	.3811	WEIGHT TAKEN
TESTIS (TE)	3.2582	.6586 %	1.5566	WEIGHT TAKEN
THYMUS (TH)	.2806	.0567 %	.1341	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0265	.0054 %	.0127	WEIGHT TAKEN

HISTOPATHOLOGY

OBSERVATIONS

NECROPSY

LUNG (LU) :
-INFILTRATE, MACROPHAGE, ALVEOLAR, -MINIMAL
LIVER (LI) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
THYMUS (TH) :
-HEMORRHAGE, -MINIMAL

THYMUS (TH) :
-RED FOCUS (I)/AREA (S) :
TO 3 X 2 MM
GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN

APPENDIX 6

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 31

ANIMAL NUMBER: C95918
DATE OF DEATH: 12/02/98
DATE AND TIME OF NECROPSY: 12/02/98 10:58
POST-FIX WEIGHTER: NOT AVAILABLE
SEX: MALE
STUDY DAY OF DEATH: 93
STUDY WEEK OF DEATH: 14
PROSECTOR: LAURIE J. SCHULLER
PATHOLOGIST: JOHN EIGHMY
DOSE GROUP: 1
SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
TERMINAL BODY WEIGHT: 494.7 GRAMS
RECORDED: TERRI L. WILLIAMS
WEIGHED: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC),
ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN),
SPLEEN (SP), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA),
AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE),
ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG),
AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV), TESTIS (TE),
EPIDIDYMIDES (EP)

001114

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 32

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95919 SEX: MALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 523.9 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:30 PROSECTOR: NANCY DIEDRICH RECORDER: TEREI L. WILLIAMS
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0603	.0115 %	.0267	WEIGHT TAKEN
BRAIN (BR)	2.2555	.4305 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	4.1905	.7999 %	1.8579	WEIGHT TAKEN
LIVER (LI)	16.3018	3.1116 %	7.2276	WEIGHT TAKEN
SPLEEN (SP)	.9470	.1808 %	.4199	WEIGHT TAKEN
TESTIS (TE)	3.1638	.6039 %	1.4027	WEIGHT TAKEN
THYMUS (TH)	.2340	.0447 %	.1037	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0308	.0059 %	.0137	WEIGHT TAKEN

HISTOPATHOLOGY

LIVER (LI) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL
HEART (HT) :
-DEGENERATION/NECROSIS, -MINIMAL
THYROID (TY) :
-CYST, ULTIMOBRANCHIAL, -PRESENT
PARATHYROID (PT) :
-CYST, -PRESENT
PITUITARY (PI) :
-CYST, -PRESENT
LN, MANDIBULAR (MN) :
-HEMORRHAGE, -MINIMAL
PROSTATE (PR) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL

NECROPSY

LN, MANDIBULAR (MN) :
-DIFFUSELY RED; RIGHT: DARK RED

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 33

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95919 SEX: MALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 523.9 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:30 PROSECTOR: NANCY DIEDRICH RECORDED: TERRI L. WILLIAMS
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG),
HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV),
TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN),
SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO),
STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE),
COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK),
URINARY BLADDER (UB), SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP)

001116

APPENDIX 6

Covance 6329-225
3M T-6314.1

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS PAGE: 34

ANIMAL NUMBER: C95920 SEX: MALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
 DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 514.2 GRAMS
 DATE AND TIME OF NECROPSY: 12/02/98 11:35 PROSECTOR: JOHN S. HALFORD RECORDED: WENDY CHERRY
 POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUSES
ADRENAL (AD)	.0685	.0133 %	.0314	WEIGHT TAKEN
BRAIN (BR)	2.1808	.4241 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	4.1319	.8036 %	1.8947	WEIGHT TAKEN
LIVER (LI)	15.1321	2.9428 %	6.9388	WEIGHT TAKEN
SPLEEN (SP)	.8162	.1587 %	.3743	WEIGHT TAKEN
TESTIS (TE)	3.1121	.6052 %	1.4270	WEIGHT TAKEN
THYMUS (TH)	.4094	.0796 %	.1877	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0193	.0038 %	.0088	WEIGHT TAKEN

NECROPSY

OBSERVATIONS

HISTOPATHOLOGY

LIVER (LI) :
 -INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
 THYROID (TY) :
 -CYST, ULTIMOBRANCHIAL, -PRESENT
 PROSTATE (PR) :
 -INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL

GENERAL COMMENT (GC) :
 -BONE MARROW SMEAR TAKEN
 -NO MACROSCOPIC LESIONS

APPENDIX 6

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 35

ANIMAL NUMBER: C95920
SEX: MALE
DOSE GROUP: 1
SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98
STUDY DAY OF DEATH: 93
STUDY WEEK OF DEATH: 14
TERMINAL BODY WEIGHT: 514.2 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:35
PROSECTOR: JOHN S. HALFORD
RECORDED: WENDY CHERRY
POST-FIX WEIGHT: NOT AVAILABLE
PATHOLOGIST: JOHN EIGHMY
WEIGHED: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), PARATHYROID (PT), ADRENAL, CORTEX (AC),
ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP)

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.113-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 36

ANIMAL NUMBER: C95921 SEX: MALE DOSE GROUP: 2 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
 DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 478.4 GRAMS
 DATE AND TIME OF NECROPSY: 12/02/98 11:11 PROSECTOR: JOHN S. HALFORD RECORDED: WENDY CHERRY
 POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUSES
ADRENAL (AD)	.0513	.0107 %	.0226	WEIGHT TAKEN
BRAIN (BR)	2.2655	.4736 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	3.4063	.7120 %	1.5036	WEIGHT TAKEN
LIVER (LI)	13.6319	2.8495 %	6.0172	WEIGHT TAKEN
SPLEEN (SP)	.7113	.1487 %	.3140	WEIGHT TAKEN
TESTIS (TE)	3.1045	.6489 %	1.3703	WEIGHT TAKEN
THYMUS (TH)	.2703	.0565 %	.1193	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0315	.0066 %	.0139	WEIGHT TAKEN

NECROPSY

THYMUS (TH) :
 -DIFFUSELY RED; LEFT LOBE: DARK RED
 GENERAL COMMENT (GC) :
 -BONE MARROW SMEAR TAKEN

OBSERVATIONS

HISTOPATHOLOGY

LIVER (LI) :
 -INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
 THYMUS (TH) :
 -HEMORRHAGE, -SLIGHT

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
 SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
 NERVE, SCIATIC (SN), SPLEEN (SP), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC),
 ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA),
 DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
 SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
 SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

001119

APPENDIX 6

Covance 6329-225
3M T-6314.1

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 37ANIMAL NUMBER: C95922
SEX: MALE
DOSE GROUP: 2
SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98
STUDY DAY OF DEATH: 93
STUDY WEEK OF DEATH: 14
TERMINAL BODY WEIGHT: 486.8 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:30
PROSECTOR: LAURIE J. SCHULLER
REORDER: TERRI L. WILLIAMS
POST-FIX WEIGHTER: NOT AVAILABLE
PATHOLOGIST: JOHN EIGHY
WEIGHER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0448	.0092 %	.0206	WEIGHT TAKEN
BRAIN (BR)	2.1727	.4463 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	3.4077	.7000 %	1.5684	WEIGHT TAKEN
LIVER (LI)	13.7456	2.8237 %	6.3265	WEIGHT TAKEN
SPLEEN (SP)	.7358	.1512 %	.3387	WEIGHT TAKEN
TESTIS (TE)	3.6580	.7514 %	1.6836	WEIGHT TAKEN
THYMUS (TH)	.2339	.0480 %	.1077	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0446	.0092 %	.0205	WEIGHT TAKEN

NECROPSY

O B S E R V A T I O N S

HISTOPATHOLOGY

LIVER (LI):
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMALGENERAL COMMENT (GC):
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR), SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

001120

APPENDIX 6

Covance 6329-225
3M T-6314.1

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 38ANIMAL NUMBER: C95923
SEX: MALE
DOSE GROUP: 2
SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98
STUDY DAY OF DEATH: 93
STUDY WEEK OF DEATH: 14
TERMINAL BODY WEIGHT: 491.7 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:47
PROSECTOR: JERALD W. JOHNSON
RECORDED: WENDY CHERRY
POST-FIX WEIGHTER: NOT AVAILABLE
PATHOLOGIST: JOHN EIGHMY
WEIGHER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	O R G A N	
				S T A T U S	W E I G H T
ADRENAL (AD)	.0499	.0101 %	.0232	WEIGHT TAKEN	
BRAIN (BR)	2.1488	.4370 %	1.0000	WEIGHT TAKEN	
KIDNEY (KD)	4.0171	.8170 %	1.8695	WEIGHT TAKEN	
LIVER (LI)	13.8730	2.8214 %	6.4562	WEIGHT TAKEN	
SPLEEN (SP)	.6287	.1279 %	.2926	WEIGHT TAKEN	
TESTIS (TE)	3.6620	.7448 %	1.7042	WEIGHT TAKEN	
THYMUS (TH)	.3346	.0680 %	.1557	WEIGHT TAKEN	
THYROID/PARATHYR (TT)	.0122	.0025 %	.0057	WEIGHT TAKEN	

NECROPSY

O B S E R V A T I O N S

HISTOPATHOLOGY

LIVER (LI) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL

GENERAL COMMENT (GC) :

-BONE MARROW SMEAR TAKEN

-NO MACROSCOPIC LESIONS

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TV), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 39

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95924 SEX: MALE DOSE GROUP: 2 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 498.6 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 9:45 PROSECTOR: GREGORY CODDINGTON RECORDER: TERRI L. WILLIAMS
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN S T A T U S
ADRENAL (AD)	.0561	.0113 %	.0258	WEIGHT TAKEN
BRAIN (BR)	2.1774	.4367 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	3.7379	.7497 %	1.7167	WEIGHT TAKEN
LIVER (LI)	13.9536	2.7986 %	6.4084	WEIGHT TAKEN
SPLEEN (SP)	.9245	.1854 %	.4246	WEIGHT TAKEN
TESTIS (TE)	3.5483	.7117 %	1.6296	WEIGHT TAKEN
THYMUS (TH)	.3028	.0607 %	.1391	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0248	.0050 %	.0114	WEIGHT TAKEN

O B S E R V A T I O N S
NECROPSY
HISTOPATHOLOGY

LIVER (LI) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:
BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TV), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

APPENDIX 6
Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 40

ANIMAL NUMBER: C95925 SEX: MALE DOSE GROUP: 2 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 468.3 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:10 PROSECTOR: KIMBERLY DURLAND REORDER: TERRI L. WILLIAMS
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATISTICS
ADRENAL (AD)	.0628	.0134 %	.0283	WEIGHT TAKEN
BRAIN (BR)	2.2194	.4739 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	3.9763	.8491 %	1.7916	WEIGHT TAKEN
LIVER (LI)	14.6970	3.1384 %	6.6221	WEIGHT TAKEN
SPLEEN (SP)	.6523	.1393 %	.2939	WEIGHT TAKEN
TESTIS (TE)	3.1815	.6794 %	1.4335	WEIGHT TAKEN
THYMUS (TH)	.2829	.0604 %	.1275	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0261	.0056 %	.0118	WEIGHT TAKEN

NECROPSY OBSERVATIONS HISTOPATHOLOGY

STOMACH, GL (ST) :
-LIGHT FOCUS(I)/AREA(S); MUCOSA, NEAR ESOPHAGEAL GROVE:
SINGLE WHITE, SLIGHTLY RAISED AREA, 1 MM IN DIAMETER
LN, MANDIBULAR (MN) :
-DIFFUSELY RED; BILATERAL: DARK RED
GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-MALOCCLUSION; BILATERAL, UPPER INCISORS

LIVER (LI) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL
STOMACH, GL (ST) :
>UNREMARKABLE
LN, MANDIBULAR (MN) :
-HEMORRHAGE, -SLIGHT

001123

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 41

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95925 SEX: MALE DOSE GROUP: 2 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 468.3 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:10 PROSECTOR: KIMBERLY DURLAND RECORDER: TERRI L. WILLIAMS
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG),
HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV),
TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

001124

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.113-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 42ANIMAL NUMBER: C95926 SEX: MALE DOSE GROUP: 2 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 438.7 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:19 PROSECTOR: KEVIN BILLINGS
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY RECORDER: WENDY CHERRY
WEIGHER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS	WEIGHT TAKEN
ADRENAL (AD)	.0518	.0118 %	.0261		WEIGHT TAKEN
BRAIN (BR)	1.9876	.4531 %	1.0000		WEIGHT TAKEN
KIDNEY (KD)	2.8500	.6496 %	1.4339		WEIGHT TAKEN
LIVER (LI)	11.7726	2.6835 %	5.9230		WEIGHT TAKEN
SPLEEN (SP)	.6868	.1566 %	.3455		WEIGHT TAKEN
TESTIS (TE)	2.7873	.6354 %	1.4023		WEIGHT TAKEN
THYMUS (TH)	.3553	.0810 %	.1788		WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0163	.0037 %	.0082		WEIGHT TAKEN

NECROPSY

OBSERVATIONS

HISTOPATHOLOGY

LIVER (LI):

-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMALGENERAL COMMENT (GC):
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

APPENDIX 6

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 43ANIMAL NUMBER: C95927 SEX: MALE DOSE GROUP: 2 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 469.4 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 9:50 PROSECTOR: JODI THIEREN REORDER: TERRI L. WILLIAMS
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0504	.0107 %	.0239	WEIGHT TAKEN
BRAIN (BR)	2.1110	.4497 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	4.0122	.8548 %	1.9006	WEIGHT TAKEN
LIVER (LI)	14.9670	3.1885 %	7.0900	WEIGHT TAKEN
SPLEEN (SP)	.6945	.1480 %	.3290	WEIGHT TAKEN
TESTIS (TE)	3.0365	.6469 %	1.4384	WEIGHT TAKEN
THYMUS (TH)	.3438	.0732 %	.1629	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0230	.0049 %	.0109	WEIGHT TAKEN

O B S E R V A T I O N S

NECROPSY

HISTOPATHOLOGY

LIVER (LI) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
STOMACH, GL (ST) :
-EROSION, -SLIGHTSTOMACH, GL (ST) :
-DARK FOCUS(I)/AREA(S) : MUCOSA: SINGLE DARK BROWN LINEAR
AREA, 3 X 1 MMGENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

APPENDIX 6

Covance 6329-225
3M T-6314.1

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 44

ANIMAL NUMBER: C95928
SEX: MALE
DOSE GROUP: 2
SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98
STUDY DAY OF DEATH: 93
WEEK OF DEATH: 14
TERMINAL BODY WEIGHT: 486.1 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:23
PROSECTOR: MICHELE ZIMA
RECORDER: WENDY CHERRY
POST-FIX WEIGHT: NOT AVAILABLE
PATHOLOGIST: JOHN EIGHMY
WEIGHER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0537	.0110 %	.0259	WEIGHT TAKEN
BRAIN (BR)	2.0720	.4262 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	3.6794	.7569 %	1.7758	WEIGHT TAKEN
LIVER (LI)	14.1057	2.9018 %	6.8078	WEIGHT TAKEN
SPLEEN (SP)	.5995	.1233 %	.2893	WEIGHT TAKEN
TESTIS (TE)	3.4858	.7171 %	1.6823	WEIGHT TAKEN
THYMUS (TH)	.2143	.0441 %	.1034	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0300	.0062 %	.0145	WEIGHT TAKEN

NECROPSY

OBSERVATIONS

HISTOPATHOLOGY

LIVER (LI) :

-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 45

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95929 SEX: MALE DOSE GROUP: 2 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 505.5 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:20 PROSECTOR: JOHN S. HALFORD RECORDED: WENDY CHERRY
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	O R G A N	
				S T A T U S	W E I G H T
ADRENAL (AD)	.0687	.0136 %	.0317	WEIGHT TAKEN	
BRAIN (BR)	2.1644	.4282 %	1.0000	WEIGHT TAKEN	
KIDNEY (KD)	3.1590	.6249 %	1.4595	WEIGHT TAKEN	
LIVER (LI)	13.3939	2.6496 %	6.1883	WEIGHT TAKEN	
SPLEEN (SP)	.8459	.1673 %	.3908	WEIGHT TAKEN	
TESTIS (TE)	3.7218	.7363 %	1.7196	WEIGHT TAKEN	
THYMUS (TH)	.2340	.0463 %	.1081	WEIGHT TAKEN	
THYROID/PARATHYR (TT)	.0298	.0059 %	.0138	WEIGHT TAKEN	

O B S E R V A T I O N S
NECROPSY
HISTOPATHOLOGY

LIVER (LI) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -SLIGHT
-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL
THYMUS (TH) :
-HEMORRHAGE, -SLIGHT

THYMUS (TH) :
-RED FOCUS(1)/AREA(S); RIGHT LOBE: SINGLE DARK RED AREA, 6 X
3 MM

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC),
ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

001123

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 46

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95930	SEX: MALE	DOSE GROUP: 2	SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98	STUDY DAY OF DEATH: 93	STUDY WEEK OF DEATH: 14	TERMINAL BODY WEIGHT: 484.0 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:59	PROSECTOR: KEVIN BILLINGS	RECORDER: WENDY CHERRY	
POST-FIX WEIGHTER: NOT AVAILABLE	PATHOLOGIST: JOHN EIGHMY	WEIGHER: LYNN HORNER	
ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO
ADRENAL (AD)	.0711	.0147 %	.0350
BRAIN (BR)	2.0340	.4202 %	1.0000
KIDNEY (KD)	3.8914	.8040 %	1.9132
LIVER (LI)	15.3859	3.1789 %	7.5644
SPLEEN (SP)	.8442	.1744 %	.4150
TESTIS (TE)	3.4684	.7166 %	1.7052
THYMUS (TH)	.4143	.0856 %	.2037
THYROID/PARATHYR (TT)	.0169	.0035 %	.0083

NECROPSY	OBSERVATIONS	HISTOPATHOLOGY
KIDNEY (KD) :		LIVER (LI) :
-LARGE PELVIS(ES); LEFT		-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-GRANULAR MATERIAL; LEFT, WITHIN PELVIS: MULTIPLE TAN,		-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL
PINPOINT GRANULES		KIDNEY (KD) :
GENERAL COMMENT (GC) :		-DILATATION, PELVIC, -SLIGHT
-BONE MARROW SMEAR TAKEN		

001129

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 47

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95930 SEX: MALE DOSE GROUP: 2 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 484.0 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:59 PROSECTOR: KEVIN BILLINGS REORDER: WENDY CHERRY
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

001130

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 48

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95931	SEX: MALE	DOSE GROUP: 2	SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE	
DATE OF DEATH: 10/01/98	STUDY DAY OF DEATH: 31	STUDY WEEK OF DEATH: 5	TERMINAL BODY WEIGHT: 362.7 GRAMS	
DATE AND TIME OF NECROPSY: 10/01/98 11:47	PROSECTOR: MATTHEW AXTMAN		RECORDER: KEVIN BILLINGS	
POST-FIX WEIGHTER: NOT AVAILABLE	PATHOLOGIST: JOHN EIGHMY		WEIGHER: MICHELE ZIMA	
ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	---	---	---	NOT TAKEN
BRAIN (BR)	---	---	---	NOT TAKEN
KIDNEY (KD)	---	---	---	NOT TAKEN
LIVER (LI)	11.8205	3.2590 %	---	WEIGHT TAKEN
SPLEEN (SP)	---	---	---	NOT TAKEN
TESTIS (TE)	---	---	---	NOT TAKEN
THYMUS (TH)	---	---	---	NOT TAKEN
THYROID/PARATHYR (TT)	---	---	---	NOT TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR), SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI), GENERAL COMMENT (GC)

001131

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 49

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95932 SEX: MALE DOSE GROUP: 2 SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
DATE OF DEATH: 10/01/98 STUDY DAY OF DEATH: 31 STUDY WEEK OF DEATH: 5 TERMINAL BODY WEIGHT: 329.1 GRAMS
DATE AND TIME OF NECROPSY: 10/01/98 11:42 PROSECTOR: MATTHEW AXTMAN REORDER: KEVIN BILLINGS
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: MICHELE ZIMA

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	---	---	---	NOT TAKEN
BRAIN (BR)	---	---	---	NOT TAKEN
KIDNEY (KD)	---	---	---	NOT TAKEN
LIVER (LI)	9.9205	3.0144 %	---	WEIGHT TAKEN
SPLEEN (SP)	---	---	---	NOT TAKEN
TESTIS (TE)	---	---	---	NOT TAKEN
THYMUS (TH)	---	---	---	NOT TAKEN
THYROID/PARATHYR (TT)	---	---	---	NOT TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TV), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI), GENERAL COMMENT (GC)

001132

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 50

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95933	SEX: MALE	DOSE GROUP: 2	SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
DATE OF DEATH: 10/01/98	STUDY DAY OF DEATH: 31	STUDY WEEK OF DEATH: 5	TERMINAL BODY WEIGHT: 341.3 GRAMS
DATE AND TIME OF NECROPSY: 10/01/98 11:44	PROSECTOR: TONY HANFELD		RECORDER: KEVIN BILLINGS
POST-FIX WEIGHTER: NOT AVAILABLE	PATHOLOGIST: JOHN EIGHMY		WEIGHER: MICHELE ZIMA
</			

601133

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 51

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95934 SEX: MALE DOSE GROUP: 2 SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
DATE OF DEATH: 10/01/98 STUDY DAY OF DEATH: 31 STUDY WEEK OF DEATH: 5 TERMINAL BODY WEIGHT: 317.0 GRAMS
DATE AND TIME OF NECROPSY: 10/01/98 11:58 PROSECTOR: WENDY CHERRY REORDER: KEVIN BILLINGS
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: MICHELE ZIMA

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	---	---	---	NOT TAKEN
BRAIN (BR)	---	---	---	NOT TAKEN
KIDNEY (KD)	---	---	---	WEIGHT TAKEN
LIVER (LI)	9.9286	3.1321 %	---	NOT TAKEN
SPLEEN (SP)	---	---	---	NOT TAKEN
TESTIS (TE)	---	---	---	NOT TAKEN
THYMUS (TH)	---	---	---	NOT TAKEN
THYROID/PARATHYR (TT)	---	---	---	---

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI), GENERAL COMMENT (GC)

C91134

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 52

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95935	SEX: MALE	DOSE GROUP: 2	SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
DATE OF DEATH: 10/01/98	STUDY DAY OF DEATH: 31	STUDY WEEK OF DEATH: 5	TERMINAL BODY WEIGHT: 354.6 GRAMS
DATE AND TIME OF NECROPSY: 10/01/98 12:07	PROSECTOR: TONY HANFELD		RECORDER: KEVIN BILLINGS
POST-FIX WEIGHTER: NOT AVAILABLE	PATHOLOGIST: JOHN EIGHMY		WEIGHER: MICHELE ZIMA
	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO
ORGAN NAME			S T A T U S
ADRENAL (AD)	----	----	NOT TAKEN
BRAIN (BR)	----	----	NOT TAKEN
KIDNEY (KD)	----	----	NOT TAKEN
LIVER (LI)	11.8361	3.3379 %	WEIGHT TAKEN
SPLEEN (SP)	----	----	NOT TAKEN
TESTIS (TE)	----	----	NOT TAKEN
THYMUS (TH)	----	----	NOT TAKEN
THYROID/PARATHYR (TT)	----	----	NOT TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR), SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI), GENERAL COMMENT (GC)

001125

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 53

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95936	SEX: MALE	DOSE GROUP: 2	SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98	STUDY DAY OF DEATH: 93	STUDY WEEK OF DEATH: 14	TERMINAL BODY WEIGHT: 550.9 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 9:51	PROSECTOR: JERALD W. JOHNSON	REORDER: WENDY CHERRY	
POST-FIX WEIGHT: NOT AVAILABLE	PATHOLOGIST: JOHN EIGHMY	WEIGHER: LYNN HORNER	
ABSOLUTE ORGAN WEIGHT (GRAMS)		ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	WEIGHT TAKEN
ADRENAL (AD)	.0484	.0088 %	.0234
BRAIN (BR)	2.0694	.3756 %	1.0000
KIDNEY (KD)	4.1177	.7474 %	1.9898
LIVER (LI)	15.6498	2.8408 %	7.5625
SPLEEN (SP)	.7959	.1445 %	.3846
TESTIS (TE)	3.1581	.5733 %	1.5261
THYMUS (TH)	.3262	.0592 %	.1576
THYROID/PARATHYR (TT)	.0188	.0034 %	.0091

HISTOPATHOLOGY

OBSERVATIONS

NECROPSY

LIVER (LI) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-MALOCCLUSION; LEFT, UPPER INCISOR

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

001136

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 54

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95937	SEX: MALE	DOSE GROUP: 2	SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98	STUDY DAY OF DEATH: 93	STUDY WEEK OF DEATH: 14	TERMINAL BODY WEIGHT: 444.5 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:10	PROSECTOR: TODD ACKER		RECORDER: WENDY CHERRY
POST-FIX WEIGHTER: NOT AVAILABLE	PATHOLOGIST: JOHN EIGHMY		WEIGHER: LYNN HORNER
			</

OBSERVATIONS

NECROPSY

HISTOPATHOLOGY

LIVER (LI):
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL

GENERAL COMMENT (GC):
-BONE MARROW SMEAR TAKEN
-MALOCCLUSION; BILATERAL, UPPER INCISORS

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EV), BRAIN (BR), SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

001137

APPENDIX 6
Individual Anatomic Pathology Data

PAGE: 55

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95938		SEX: MALE		DOSE GROUP: 2		SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE	
DATE OF DEATH: 12/02/98		STUDY DAY OF DEATH: 93		STUDY WEEK OF DEATH: 14		TERMINAL BODY WEIGHT: 497.6 GRAMS	
DATE AND TIME OF NECROPSY: 12/02/98 7:59		PROSECTOR: MICHELE ZIMA		PATHOLOGIST: JOHN EIGHMY		RECORDER: WENDY CHERRY	
POST-FIX WEIGHT: NOT AVAILABLE						WEIGHER: LYNN HORNER	

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 56

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95939 SEX: MALE DOSE GROUP: 2 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 447.8 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 12:23 PROSECTOR: JOHN S. HALFORD
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY RECORDER: WENDY CHERRY
WEIGHER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0634	.0142 %	.0273	WEIGHT TAKEN
BRAIN (BR)	2.3237	.5189 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	3.6350	.8117 %	1.5643	WEIGHT TAKEN
LIVER (LI)	12.6033	2.8145 %	5.4238	WEIGHT TAKEN
SPLEEN (SP)	1.1088	.2476 %	.4772	WEIGHT TAKEN
TESTIS (TE)	3.2485	.7254 %	1.3980	WEIGHT TAKEN
THYMUS (TH)	.4355	.0973 %	.1874	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0289	.0065 %	.0124	WEIGHT TAKEN

OBSERVATIONS

NECROPSY

HISTOPATHOLOGY

LIVER (LI) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 57

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95940 SEX: MALE DOSE GROUP: 2 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 483.1 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:10 PROSECTOR: KIMBERLY DURLAND REORDER: STEVE VAN ADESTINE
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	O R G A N	
				S T A T U S	WEIGHT TAKEN
ADRENAL (AD)	.0659	.0136 %	.0308		WEIGHT TAKEN
BRAIN (BR)	2.1363	.4422 %	1.0000		WEIGHT TAKEN
KIDNEY (KD)	4.1869	.8667 %	1.9599		WEIGHT TAKEN
LIVER (LI)	14.1874	2.9367 %	6.6411		WEIGHT TAKEN
SPLEEN (SP)	.8620	.1784 %	.4035		WEIGHT TAKEN
TESTIS (TE)	3.8586	.7987 %	1.8062		WEIGHT TAKEN
THYMUS (TH)	.2849	.0590 %	.1334		WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0224	.0046 %	.0105		WEIGHT TAKEN

O B S E R V A T I O N S

HISTOPATHOLOGY

NECROPSY

LIVER (LI) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL
LN, MANDIBULAR (MN) :
-HEMORRHAGE, -SLIGHT

LN, MANDIBULAR (MN) :
-DIFFUSELY RED; BILATERAL: DARK RED
GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-MALOCCLUSION; BILATERAL UPPER INCISORS

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG),
HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV),
TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

C01140

APPENDIX 6
Individual Anatomic Pathology Data

PAGE: 58

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL-
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95941		DOSE GROUP: 3		SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE	
SEX: MALE		STUDY WEEK OF DEATH: 14		TERMINAL BODY WEIGHT: 411.5 GRAMS	
DATE OF DEATH: 12/02/98		PROSECTOR: KEVIN BILLINGS		REORDER: WENDY CHERRY	
DATE AND TIME OF NECROPSY: 12/02/98 11:42		PATHOLOGIST: JOHN EIGHMY		WEIGHER: LYNN HORNER	
POST-FIX WEIGHER: NOT AVAILABLE					

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS	WEIGHT TAKEN
ADRENAL (AD)	.0457	.0111 %	.0216		WEIGHT TAKEN
BRAIN (BR)	2.1186	.5148 %	1.0000		WEIGHT TAKEN
KIDNEY (KD)	3.5249	.8566 %	1.6638		WEIGHT TAKEN
LIVER (LI)	13.6671	3.3213 %	6.4510		WEIGHT TAKEN
SPLEEN (SP)	.8387	.2038 %	.3959		WEIGHT TAKEN
TESTIS (TE)	3.2827	.7977 %	1.5495		WEIGHT TAKEN
THYMUS (TH)	.3528	.0857 %	.1665		WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0182	.0044 %	.0086		WEIGHT TAKEN

NECROPSY	OBSERVATIONS	HISTOPATHOLOGY
GENERAL COMMENT (GC) :	LIVER (LI) :	
-BONE MARROW SMEAR TAKEN	-FIBROSIS, CAPSULAR, -MINIMAL	
-NO MACROSCOPIC LESIONS	-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MINIMAL	
	-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL	

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 59

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95941 SEX: MALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 411.5 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:42 PROSECTOR: KEVIN BILLINGS REORDER: WENDY CHERRY
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EPI), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

STOMACH, GL (ST), STOMACH, NONGL (SU)

C01142

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 60

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95942		SEX: MALE		DOSE GROUP: 3		SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE	
DATE OF DEATH: 12/02/98		STUDY DAY OF DEATH: 93		STUDY WEEK OF DEATH: 14		TERMINAL BODY WEIGHT: 418.7 GRAMS	
DATE AND TIME OF NECROPSY: 12/02/98 10:58		PROSECTOR: JERALD W. JOHNSON		RECORDER: WENDY CHERRY		WEIGHER: LYNN HORNER	
POST-FIX WEIGHER: NOT AVAILABLE		PATHOLOGIST: JOHN EIGHMY					

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0694	.0166 %	.0335	WEIGHT TAKEN
BRAIN (BR)	2.0705	.4945 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	4.1159	.9830 %	1.9879	WEIGHT TAKEN
LIVER (LI)	14.8588	3.5488 %	7.1764	WEIGHT TAKEN
SPLEEN (SP)	.6815	.1628 %	.3291	WEIGHT TAKEN
TESTIS (TE)	3.5997	.8597 %	1.7386	WEIGHT TAKEN
THYMUS (TH)	.2568	.0613 %	.1240	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0174	.0042 %	.0084	WEIGHT TAKEN

NECROPSY	OBSERVATIONS	HISTOPATHOLOGY
LN, MANDIBULAR (MN) :		
-DIFFUSELY RED; BILATERAL: DARK RED		
GENERAL COMMENT (GC) :		
-BONE MARROW SMEAR TAKEN		
-MALOCCLUSION; BILATERAL, UPPER INCISORS		
LIVER (LI) :		
-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MINIMAL		
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL		
-VACUOLATION, HEPATOCELLULAR, PERIportal, -MINIMAL		
LN, MANDIBULAR (MN) :		
-HEMORRHAGE, -SLIGHT		

001143

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 61

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95942 SEX: MALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 418.7 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:58 PROSECTOR: JERALD W. JOHNSON RECORDER: WENDY CHERRY
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG),
HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV),
TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

STOMACH, GL (ST), STOMACH, NONGL (SU)

0911144

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 62

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95943
DATE OF DEATH: 12/02/98
DATE AND TIME OF NECROPSY: 12/02/98 11:21
POST-FIX WEIGHT: NOT AVAILABLE

SEX: MALE
STUDY DAY OF DEATH: 93
PROSECTOR: JERALD W. JOHNSON
PATHOLOGIST: JOHN EIGHMY

DOSE GROUP: 3
SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
STUDY WEEK OF DEATH: 14
TERMINAL BODY WEIGHT: 500.1 GRAMS
RECORDER: WENDY CHERRY
WEIGHER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0531	.0106 %	.0251	WEIGHT TAKEN
BRAIN (BR)	2.1128	.4225 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	4.1504	.8299 %	1.9644	WEIGHT TAKEN
LIVER (LI)	19.4681	3.8928 %	9.2144	WEIGHT TAKEN
SPLEEN (SP)	.6715	.1343 %	.3178	WEIGHT TAKEN
TESTIS (TE)	3.5995	.7198 %	1.7037	WEIGHT TAKEN
THYMUS (TH)	.3527	.0705 %	.1669	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0275	.0055 %	.0130	WEIGHT TAKEN

OBSERVATIONS

HISTOPATHOLOGY

NECROPSY

LIVER (LI) :
-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -SLIGHT
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-VACUOLATION, HEPATOCELLULAR, DIFFUSE, -MINIMAL

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

601145

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 63

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95943 SEX: MALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 500.1 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:21 PROSECTOR: JERALD W. JOHNSON REORDER: WENDY CHERRY
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

STOMACH, GL (ST), STOMACH, NONGL (SU)

001146

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 64

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95944		SEX: MALE		DOSE GROUP: 3		SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE	
DATE OF DEATH: 12/02/98		STUDY DAY OF DEATH: 93		STUDY WEEK OF DEATH: 14		TERMINAL BODY WEIGHT: 433.2 GRAMS	
DATE AND TIME OF NECROPSY: 12/02/98 11:31		PROSECTOR: TODD ACKER		PATHOLOGIST: JOHN EIGHMY		RECORDER: WENDY CHERRY	
POST-FIX WEIGHER: NOT AVAILABLE		ABSOLUTE ORGAN WEIGHT (GRAMS)		ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)		ORGAN TO BRAIN WEIGHT RATIO	
		ORGAN NAME				O R G A N S T A T U S	
		ADRENAL (AD)		.0511		WEIGHT TAKEN	
		BRAIN (BR)		2.1737		WEIGHT TAKEN	
		KIDNEY (KD)		3.9789		WEIGHT TAKEN	
		LIVER (LI)		15.6049		WEIGHT TAKEN	
		SPLEEN (SP)		.6847		WEIGHT TAKEN	
		TESTIS (TE)		3.4797		WEIGHT TAKEN	
		THYMUS (TH)		.2276		WEIGHT TAKEN	
		THYROID/PARATHYR (TT)		.0265		WEIGHT TAKEN	

NECROPSY		O B S E R V A T I O N S		HISTOPATHOLOGY	
LN, MANDIBULAR (MN) :		LIVER (LI) :		LIVER (LI) :	
-DIFFUSELY RED; BILATERAL: DARK RED		-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MINIMAL		LN, MANDIBULAR (MN) :	
GENERAL COMMENT (GC) :		-HEMORRHAGE, -SLIGHT			
-BONE MARROW SMEAR TAKEN					

001147

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 65

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95944
SEX: MALE
DATE OF DEATH: 12/02/98
STUDY DAY OF DEATH: 93
DATE AND TIME OF NECROPSY: 12/02/98 11:31
PROSECTOR: TODD ACKER
POST-FIX WEIGHER: NOT AVAILABLE
DOSE GROUP: 3
SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
TERMINAL BODY WEIGHT: 433.2 GRAMS
REORDER: WENDY CHERRY
WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG),
HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV),
TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

STOMACH, GL (ST), STOMACH, NONGL (SU)

001148

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 66

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95945	SEX: MALE	DOSE GROUP: 3	SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98	STUDY DAY OF DEATH: 93	STUDY WEEK OF DEATH: 14	TERMINAL BODY WEIGHT: 491.3 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:17	PROSECTOR: KEVIN BILLINGS	RECORDED: WENDY CHERRY	
POST-FIX WEIGHTER: NOT AVAILABLE	PATHOLOGIST: JOHN EIGHMY	WEIGHER: LYNN HORNER	
	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO
ORGAN NAME			S T A T U S
ADRENAL (AD)	.0502	.0102 %	.0232
BRAIN (BR)	2.1630	.4403 %	1.0000
KIDNEY (KD)	4.1701	.8488 %	1.9279
LIVER (LI)	17.2249	3.5060 %	7.9634
SPLEEN (SP)	.6713	.1366 %	.3104
TESTIS (TE)	3.8878	.7913 %	1.7974
THYMUS (TH)	.2839	.0578 %	.1313
THYROID/PARATHYR (TT)	.0216	.0044 %	.0100

OBSERVATIONS

NECROPSY

HISTOPATHOLOGY

LIVER (LI) :
-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MINIMAL
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL
-VACUOLATION, HEPATOCELLULAR, DIFFUSE, -MINIMAL

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 67

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95945 SEX: MALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 491.3 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:17 PROSECTOR: KEVIN BILLINGS REORDER: WENDY CHERRY
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

STOMACH, GL (ST), STOMACH, NONGL (SU)

001150

APPENDIX 6
Individual Anatomic Pathology Data

PAGE: 68

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95946 SEX: MALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 469.0 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 7:59 PROSECTOR: JOHN S. HALFORD RECORDER: WENDY CHERRY
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	O R G A N	
				S T A T U S	WEIGHT TAKEN
ADRENAL (AD)	.0648	.0138 %	.0273		WEIGHT TAKEN
BRAIN (BR)	2.3771	.5068 %	1.0000		WEIGHT TAKEN
KIDNEY (KD)	3.3029	.7042 %	1.3895		WEIGHT TAKEN
LIVER (LI)	15.7438	3.3569 %	6.6231		WEIGHT TAKEN
SPLEEN (SP)	.7863	.1677 %	.3308		WEIGHT TAKEN
TESTIS (TE)	3.5288	.7524 %	1.4845		WEIGHT TAKEN
THYMUS (TH)	.1989	.0424 %	.0837		WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0365	.0078 %	.0154		WEIGHT TAKEN

O B S E R V A T I O N S
HISTOPATHOLOGY
LIVER (LI) :
-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MINIMAL
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

001151

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 69

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95946 SEX: MALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 469.0 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 7:59 PROSECTOR: JOHN S. HALFORD RECORDER: WENDY CHERRY
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TV), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

STOMACH, GL (ST), STOMACH, NONGL (SU)

001152

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 70

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL-
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95947 SEX: MALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 449.9 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 9:45 PROSECTOR: NANCY DIEDRICH RECORDED: TERRI L. WILLIAMS
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN S T A T U S	WEIGHT TAKEN
ADRENAL (AD)	.0507	.0113 %	.0231		WEIGHT TAKEN
BRAIN (BR)	2.1942	.4877 %	1.0000		WEIGHT TAKEN
KIDNEY (KD)	3.2317	.7183 %	1.4728		WEIGHT TAKEN
LIVER (LI)	15.1188	3.3605 %	6.8903		WEIGHT TAKEN
SPLEEN (SP)	.8297	.1844 %	.3781		WEIGHT TAKEN
TESTIS (TE)	3.2832	.7298 %	1.4963		WEIGHT TAKEN
THYMUS (TH)	.3773	.0839 %	.1720		WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0232	.0052 %	.0106		WEIGHT TAKEN

O B S E R V A T I O N S

HISTOPATHOLOGY

NECROPSY

LIVER (LI) :
-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MINIMAL
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL
THYMUS (TH) :
-HEMORRHAGE, -MINIMAL
LN, MANDIBULAR (MN) :
-HEMORRHAGE, -MINIMAL

THYMUS (TH) :
-RED FOCUS(I)/AREA(S); DIFFUSE: MULTIPLE DARK RED PINPOINT FOCI

LN, MANDIBULAR (MN) :
-MOTTLED; BILATERAL: DARK RED AND TAN

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN

601153

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 71

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95947 SEX: MALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 449.9 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 9:45 PROSECTOR: NANCY DIEDRICH RECORDED: TERRI L. WILLIAMS
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), LN, MESENTERIC (MS), THYROID (TV), PARATHYROID (PT), ADRENAL, CORTEX (AC),
ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG),
HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV),
TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

STOMACH, GL (ST), STOMACH, NONGL (SU)

001154

APPENDIX 6

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95948	SEX: MALE	DOSE GROUP: 3	SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98	STUDY DAY OF DEATH: 93	STUDY WEEK OF DEATH: 14	TERMINAL BODY WEIGHT: 476.3 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 9:44	PROSECTOR: JOHN S. HALFORD		REORDER: WENDY CHERRY
POST-FIX WEIGHTER: NOT AVAILABLE	PATHOLOGIST: JOHN EIGHMY		WEIGHER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0458	.0096 %	.0214	WEIGHT TAKEN
BRAIN (BR)	2.1430	.4499 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	3.6102	.7580 %	1.6846	WEIGHT TAKEN
LIVER (LI)	13.4487	2.8236 %	6.2756	WEIGHT TAKEN
SPLEEN (SP)	.8221	.1726 %	.3836	WEIGHT TAKEN
TESTIS (TE)	3.2511	.6826 %	1.5171	WEIGHT TAKEN
THYMUS (TH)	.3464	.0727 %	.1616	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0202	.0042 %	.0094	WEIGHT TAKEN

NECROPSY	OBSERVATIONS	HISTOPATHOLOGY
THYMUS (TH) : -DIFFUSELY RED; DARK RED LN, MANDIBULAR (MN) : -DIFFUSELY RED; BILATERAL: DARK RED GENERAL COMMENT (GC) : -BONE MARROW SMEAR TAKEN GENERAL INFORMATION (XX) : >NOTE:>ANIMAL IDENTIFICATION TRANSPONDER DAMAGED AT NECROPSY	LIVER (LI) : -HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -SLIGHT -INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL THYMUS (TH) : -HEMORRHAGE, -SLIGHT LN, MANDIBULAR (MN) : -HEMORRHAGE, -SLIGHT	

601135

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 73

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL-
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
TERMINAL BODY WEIGHT: 476.3 GRAMS
RECORDED: WENDY CHERRY
WEIGHED: LYNN HORNER

DOSE GROUP: 3
STUDY WEEK OF DEATH: 14
PROSECTOR: JOHN S. HALFORD
PATHOLOGIST: JOHN EIGHMY

ANIMAL NUMBER: C95948
SEX: MALE
STUDY DAY OF DEATH: 93
DATE OF DEATH: 12/02/98
DATE AND TIME OF NECROPSY: 12/02/98 9:44
POST-FIX WEIGHED: NOT AVAILABLE

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC),
ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG), SEMINAL VESICLES (SV),
HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV),
TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

STOMACH, GL (ST), STOMACH, NONGL (SU)

001156

APPENDIX 6

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95949		SEX: MALE	DOSE GROUP: 3	SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE	
DATE OF DEATH: 12/02/98		STUDY DAY OF DEATH: 93	STUDY WEEK OF DEATH: 14	TERMINAL BODY WEIGHT: 421.1 GRAMS	
DATE AND TIME OF NECROPSY: 12/02/98 11:44		PROSECTOR: JILL PAUS		REORDER: WENDY CHERRY	
POST-FIX WEIGHTER: NOT AVAILABLE		PATHOLOGIST: JOHN EIGHMY		WEIGHER: LYNN HORNER	
		ABSOLUTE ORGAN WEIGHT (GRAMS)		ORGAN TO BRAIN WEIGHT RATIO	
ORGAN NAME		ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)		O R G A N S T A T U S	
ADRENAL (AD)		.0133 %		WEIGHT TAKEN	
BRAIN (BR)		.4879 %		WEIGHT TAKEN	
KIDNEY (KD)		.9355 %		WEIGHT TAKEN	
LIVER (LI)		3.5517 %		WEIGHT TAKEN	
SPLEEN (SP)		.1758 %		WEIGHT TAKEN	
TESTIS (TE)		.8505 %		WEIGHT TAKEN	
THYMUS (TH)		.0376 %		WEIGHT TAKEN	
THYROID/PARATHYR (TT)		.0052 %		WEIGHT TAKEN	

O B S E R V A T I O N S

HISTOPATHOLOGY

NECROPSY

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-MALOCCLUSION: BILATERAL, UPPER INCISORS
-STAINS-PERINEUM/PERIANAL; YELLOW AND MOIST

001157

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 75

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95949 SEX: MALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 421.1 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:44 PROSECTOR: JILL PAUS REORDER: WENDY CHERRY
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

LIVER (LI), STOMACH, GL (ST), STOMACH, NONGL (SU)

001158

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 76

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95950 SEX: MALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 467.6 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:29 PROSECTOR: GREGORY CODDINGTON RECORDED BY: TERRI L. WILLIAMS
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0593	.0127 %	.0267	WEIGHT TAKEN
BRAIN (BR)	2.2213	.4750 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	3.8346	.8201 %	1.7263	WEIGHT TAKEN
LIVER (LI)	15.5750	3.3308 %	7.0117	WEIGHT TAKEN
SPLEEN (SP)	.6608	.1413 %	.2975	WEIGHT TAKEN
TESTIS (TE)	3.4323	.7340 %	1.5452	WEIGHT TAKEN
THYMUS (TH)	.2306	.0493 %	.1038	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0259	.0055 %	.0117	WEIGHT TAKEN

OBSERVATIONS

HISTOPATHOLOGY

NECROPSY

LIVER (LI) :
-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -SLIGHT
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL
THYMUS (TH) :
-HEMORRHAGE, -SLIGHT
-INVOLUTION, -MINIMAL

THYMUS (TH) :
-RED FOCUS(I)/AREA(S); BOTH LOBES: MULTIPLE DARK RED AREAS,
UP TO 5 MM IN DIAMETER
GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN

001159

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 77

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95950 SEX: MALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 467.6 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:29 PROSECTOR: GREGORY CODDINGTON REORDER: TERRI L. WILLIAMS
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EICHMY WEIGHT: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), LN, MESENTERIC (MS), THYROID (TV), PARATHYROID (PT), ADRENAL, CORTEX (AC),
ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

STOMACH, GL (ST), STOMACH, NONGL (SU)

C01160

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 78

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95951	SEX: MALE	DOSE GROUP: 3	SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
DATE OF DEATH: 10/01/98	STUDY DAY OF DEATH: 31	STUDY WEEK OF DEATH: 5	TERMINAL BODY WEIGHT: 381.8 GRAMS
DATE AND TIME OF NECROPSY: 10/01/98 12:06	PROSECTOR: WENDY CHERRY		RECORDER: KEVIN BILLINGS
POST-FIX WEIGHER: NOT AVAILABLE	PATHOLOGIST: JOHN EIGHMY		WEIGHER: MICHELE ZIMA

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO
ADRENAL (AD)	-----	-----	-----
BRAIN (BR)	-----	-----	NOT TAKEN
KIDNEY (KD)	-----	-----	NOT TAKEN
LIVER (LI)	-----	-----	WEIGHT TAKEN
	13.6836	3.5840 %	NOT TAKEN
SPLEEN (SP)	-----	-----	NOT TAKEN
TESTIS (TE)	-----	-----	NOT TAKEN
THYMUS (TH)	-----	-----	NOT TAKEN
THYROID/PARATHYR (TT)	-----	-----	-----

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:			
BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),			
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),			
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),			
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),			
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),			
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),			
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI), GENERAL COMMENT (GC)			

001161

Individual Anatomic Pathology Data

PAGE: 79

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95952
DATE OF DEATH: 10/01/98
SEX: MALE
STUDY DAY OF DEATH: 31
DOSE GROUP: 3
STUDY WEEK OF DEATH: 5
SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
TERMINAL BODY WEIGHT: 316.5 GRAMS
RECORDER: KEVIN BILLINGS
WEIGHER: MICHELE ZIMA
PROSECTOR: WENDY CHERRY
10/01/98 11:25

DATE AND TIME OF NECROPSY		PATHOLOGIST: JOHN E. HARRIS		ORGAN TO BRAIN		ORGAN TO STOMACH	
POST-FIX WEIGHT: NOT AVAILABLE		ABSOLUTE ORGAN WEIGHT (GRAMS)		ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)		WEIGHT RATIO	
ORGAN NAME							
ADRENAL (AD)						NOT TAKEN	
BRAIN (BR)						NOT TAKEN	
KIDNEY (KD)				3.6961 %		WEIGHT TAKEN	
LIVER (LI)		11.6980				NOT TAKEN	
SPLEEN (SP)						NOT TAKEN	
TESTIS (TE)						NOT TAKEN	
THYMUS (TH)						NOT TAKEN	
THYROID/PARATHYR (TT)							

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:
 BONE, MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
 BONE, FEMUR (FE), MARROW, FEMUR (FM), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
 SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), THYROID (TY), PARATHYROID (PT),
 SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
 NERVE, SCIATIC (SN), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
 ADRENAL, CORTX (AC), ADRENAL, MEDULLA (MA), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
 ADRENAL, CORTX (AC), ADRENAL, MEDULLA (MA), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
 PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
 SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
 SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI), GENERAL COMMENT (GC)

001162

APPENDIX 6
Individual Anatomic Pathology Data13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 80ANIMAL NUMBER: C95953
SEX: MALE
DOSE GROUP: 3
SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
DATE OF DEATH: 10/01/98
STUDY DAY OF DEATH: 31
STUDY WEEK OF DEATH: 5
TERMINAL BODY WEIGHT: 326.4 GRAMS
DATE AND TIME OF NECROPSY: 10/01/98 11:13
PROSECTOR: TONY HANFELD
REORDER: KEVIN BILLINGS
POST-FIX WEIGHT: NOT AVAILABLE
PATHOLOGIST: JOHN EIGHMY
WEIGHER: MICHELE ZIMA

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	---	---	---	NOT TAKEN
BRAIN (BR)	---	---	---	NOT TAKEN
KIDNEY (KD)	---	---	---	NOT TAKEN
LIVER (LI)	11.6552	3.5708 %	---	WEIGHT TAKEN
SPLEEN (SP)	---	---	---	NOT TAKEN
TESTIS (TE)	---	---	---	NOT TAKEN
THYMUS (TH)	---	---	---	NOT TAKEN
THYROID/PARATHYR (TT)	---	---	---	NOT TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI), GENERAL COMMENT (GC)

C01103

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 81

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
TERMINAL BODY WEIGHT: 365.4 GRAMS
REORDER: KEVIN BILLINGS
WEIGHER: MICHELE ZIMA

DOSE GROUP: 3
STUDY WEEK OF DEATH: 5
PROSECTOR: TONY HANFELD
PATHOLOGIST: JOHN EIGHMY

ANIMAL NUMBER: C95954

SEX: MALE

STUDY DAY OF DEATH: 31

DATE OF DEATH: 10/01/98

DATE AND TIME OF NECROPSY: 10/01/98 11:38

POST-FIX WEIGHER: NOT AVAILABLE

ABSOLUTE ORGAN WEIGHT
(GRAMS)

ORGAN NAME

ADRENAL (AD)
BRAIN (BR)
KIDNEY (KD)
LIVER (LI)
SPLEEN (SP)
TESTIS (TE)
THYMUS (TH)

THYROID/PARATHYR (TT)

ORGAN WEIGHT RELATIVE
TO BODY WEIGHT (%)

3.6968 %

ORGAN TO BRAIN
WEIGHT RATIO

ORGAN

STATUS

NOT TAKEN
NOT TAKEN
NOT TAKEN
NOT TAKEN
NOT TAKEN
NOT TAKEN
NOT TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI), GENERAL COMMENT (GC)

001164

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 82

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95955		SEX: MALE	DOSE GROUP: 3	SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
DATE OF DEATH: 10/01/98		STUDY DAY OF DEATH: 31	STUDY WEEK OF DEATH: 5	TERMINAL BODY WEIGHT: 346.9 GRAMS
DATE AND TIME OF NECROPSY: 10/01/98 11:55		PROSECTOR: TONY HANFELD		REORDER: KEVIN BILLINGS
POST-FIX WEIGHT: NOT AVAILABLE		PATHOLOGIST: JOHN EIGHMY		WEIGHER: MICHELE ZIMA
ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	---	---	---	NOT TAKEN
BRAIN (BR)	---	---	---	NOT TAKEN
KIDNEY (KD)	---	---	---	NOT TAKEN
LIVER (LI)	15.0803	4.3472 %	---	WEIGHT TAKEN
SPLEEN (SP)	---	---	---	NOT TAKEN
TESTIS (TE)	---	---	---	NOT TAKEN
THYMUS (TH)	---	---	---	NOT TAKEN
THYROID/PARATHYR (TT)	---	---	---	---

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR), SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), NONGL (SU), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI), GENERAL COMMENT (GC)

001165

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 83

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95956		SEX: MALE	DOSE GROUP: 3	SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98		STUDY DAY OF DEATH: 93	STUDY WEEK OF DEATH: 14	TERMINAL BODY WEIGHT: 523.3 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:27		PROSECTOR: MICHELE ZIMA		RECORDED: WENDY CHERRY
POST-FIX WEIGHTER: NOT AVAILABLE		PATHOLOGIST: JOHN EIGHMY		WEIGHER: LYNN HORNER
		ABSOLUTE ORGAN WEIGHT (GRAMS)		ORGAN TO BRAIN WEIGHT RATIO
				STATUS
ORGAN NAME		ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)		WEIGHT TAKEN
ADRENAL (AD)		.0101 %		WEIGHT TAKEN
BRAIN (BR)		.4093 %		WEIGHT TAKEN
KIDNEY (KD)		.7374 %		WEIGHT TAKEN
LIVER (LI)		3.3374 %		WEIGHT TAKEN
SPLEEN (SP)		.1526 %		WEIGHT TAKEN
TESTIS (TE)		.5321 %		WEIGHT TAKEN
THYMUS (TH)		.0598 %		WEIGHT TAKEN
THYROID/PARATHYR (TT)		.0051 %		WEIGHT TAKEN
		.0268		.0246
				1.0000
				1.8014
				8.1534
				.3728
				1.3000
				.1462
				.0125

HISTOPATHOLOGY

NECROPSY

LIVER (LI) :
-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MINIMAL
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

031106

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 84

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95956 SEX: MALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 523.3 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:27 PROSECTOR: MICHELE ZIMA RECORDER: WENDY CHERRY
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:
BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:
STOMACH, GL (ST), STOMACH, NONGL (SU)

694167

APPENDIX 6
Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 85

ANIMAL NUMBER: C95957 SEX: MALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 474.7 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:54 PROSECTOR: JILL PAUS REORDER: WENDY CHERRY
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	O R G A N	
				S T A T U S	W E I G H T T A K E N
ADRENAL (AD)	.0530	.0112 %	.0243		WEIGHT TAKEN
BRAIN (BR)	2.1797	.4592 %	1.0000		WEIGHT TAKEN
KIDNEY (KD)	4.2081	.8865 %	1.9306		WEIGHT TAKEN
LIVER (LI)	16.5473	3.4858 %	7.5915		WEIGHT TAKEN
SPLEEN (SP)	.8376	.1764 %	.3843		WEIGHT TAKEN
TESTIS (TE)	3.8841	.8182 %	1.7819		WEIGHT TAKEN
THYMUS (TH)	.2384	.0502 %	.1094		WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0268	.0056 %	.0123		WEIGHT TAKEN

NECROPSY
O B S E R V A T I O N S
HISTOPATHOLOGY
LIVER (LI) :
- HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MINIMAL
- INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
- VACUOLATION, HEPATOCELLULAR, CENTRILOBULAR TO MIDZONAL, -
MINIMAL

GENERAL COMMENT (GC) :
- BONE MARROW SMEAR TAKEN
- NO MACROSCOPIC LESIONS

001168

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

PAGE: 86

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95957 SEX: MALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 474.7 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:54 PROSECTOR: JILL PAUS RECORDER: WENDY CHERRY
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

STOMACH, GL (ST), STOMACH, NONGL (SU)

001169

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 87

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95958 SEX: MALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 471.3 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:40 PROSECTOR: KEVIN BILLINGS RECORDED: WENDY CHERRY
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0509	.0108 %	.0238	WEIGHT TAKEN
BRAIN (BR)	2.1388	.4538 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	3.6164	.7673 %	1.6909	WEIGHT TAKEN
LIVER (LI)	16.8594	3.5772 %	7.8826	WEIGHT TAKEN
SPLEEN (SP)	.7451	.1581 %	.3484	WEIGHT TAKEN
TESTIS (TE)	3.0781	.6531 %	1.4392	WEIGHT TAKEN
THYMUS (TH)	.2885	.0612 %	.1349	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0212	.0045 %	.0099	WEIGHT TAKEN

NECROPSY	OBSERVATIONS	HISTOPATHOLOGY
	LIVER (LI) :	
	-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MINIMAL	
	-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL	
	-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL	

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

001170

APPENDIX 6

Covance 6329-225
3M T-6314.1

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 88

ANIMAL NUMBER: C95958 SEX: MALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 471.3 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:40 PROSECTOR: KEVIN BILLINGS RECORDER: WENDY CHERRY
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

STOMACH, GL (ST), STOMACH, NONGL (SU)

APPENDIX 6
Individual Anatomic Pathology Data

PAGE: 89

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95959 SEX: MALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 TERMINAL BODY WEIGHT: 443.4 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 7:39 PROSECTOR: KIMBERLY DURLAND
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	O R G A N	
				S T A T U S	WEIGHT TAKEN
ADRENAL (AD)	.0479	.0108 %	.0214	WEIGHT TAKEN	WEIGHT TAKEN
BRAIN (BR)	2.2347	.5040 %	1.0000	WEIGHT TAKEN	WEIGHT TAKEN
KIDNEY (KD)	3.6999	.8344 %	1.6557	WEIGHT TAKEN	WEIGHT TAKEN
LIVER (LI)	15.7371	3.5492 %	7.0422	WEIGHT TAKEN	WEIGHT TAKEN
SPLEEN (SP)	.6008	.1355 %	.2689	WEIGHT TAKEN	WEIGHT TAKEN
TESTIS (TE)	3.2245	.7272 %	1.4429	WEIGHT TAKEN	WEIGHT TAKEN
THYMUS (TH)	.2848	.0642 %	.1274	WEIGHT TAKEN	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0286	.0065 %	.0128	WEIGHT TAKEN	WEIGHT TAKEN

O B S E R V A T I O N S

NECROPSY

HISTOPATHOLOGY

LIVER (LI) :
- HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MINIMAL
- INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
LN, MANDIBULAR (MN) :
- HEMORRHAGE, -MINIMAL

LN, MANDIBULAR (MN) :
- DIFFUSELY RED; BILATERAL: DARK RED
GENERAL COMMENT (GC) :
- BONE MARROW SMEAR TAKEN

001172

APPENDIX 6

Covance 6329-225
3M T-6314.1

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 90

ANIMAL NUMBER: C95959 SEX: MALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 443.4 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 7:39 PROSECTOR: KIMBERLY DURLAND RECORDED: STEVE VAN ADESTINE
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG),
HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV),
TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

STOMACH, GL (ST), STOMACH, NONGL (SU)

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 91

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95960 SEX: MALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 449.6 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:43 PROSECTOR: MICHELE ZIMA RECORDED: WENDY CHERRY
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0760	.0169 %	.0346	WEIGHT TAKEN
BRAIN (BR)	2.1953	.4883 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	3.8855	.8642 %	1.7699	WEIGHT TAKEN
LIVER (LI)	16.2467	3.6136 %	7.4007	WEIGHT TAKEN
SPLEEN (SP)	.6429	.1430 %	.2929	WEIGHT TAKEN
TESTIS (TE)	3.8243	.8506 %	1.7420	WEIGHT TAKEN
THYMUS (TH)	.1769	.0393 %	.0806	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0243	.0054 %	.0111	WEIGHT TAKEN

NECROPSY	OBSERVATIONS	HISTOPATHOLOGY
THYMUS (TH) :		
-RED FOCUS(I)/AREA(S) :		
PINPOINT FOCI		
GENERAL COMMENT (GC) :		
-BONE MARROW SMEAR TAKEN		
LIVER (LI) :		
-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MINIMAL		
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL		
-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL		
-VACUOLATION, HEPATOCELLULAR, CENTRILOBULAR TO MIDZONAL, -		
MINIMAL		
THYMUS (TH) :		
-HEMORRHAGE, -SLIGHT		

001171

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 92

ANIMAL NUMBER: C95960 SEX: MALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 449.6 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:43 PROSECTOR: MICHELE ZIMA RECORDED: WENDY CHERRY
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC),
ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

STOMACH, GL (ST), STOMACH, NONGL (SU)

APPENDIX 6
Individual Anatomic Pathology Data13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 93ANIMAL NUMBER: C95961
SEX: MALE
DOSE GROUP: 4
SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98
STUDY WEEK OF DEATH: 14
TERMINAL BODY WEIGHT: 483.1 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:53
PROSECTOR: LAURIE J. SCHULLER
RECORDED: TERRI L. WILLIAMS
POST-FIX WEIGHT: NOT AVAILABLE
PATHOLOGIST: JOHN EIGHMY
WEIGHED: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0572	.0118 %	.0248	WEIGHT TAKEN
BRAIN (BR)	2.3075	.4776 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	4.4633	.9239 %	1.9343	WEIGHT TAKEN
LIVER (LI)	27.5733	5.7076 %	11.9494	WEIGHT TAKEN
SPLEEN (SP)	.7101	.1470 %	.3077	WEIGHT TAKEN
TESTIS (TE)	3.9981	.8276 %	1.7327	WEIGHT TAKEN
THYMUS (TH)	.3288	.0681 %	.1425	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0241	.0050 %	.0104	WEIGHT TAKEN

O B S E R V A T I O N S

NECROPSY

HISTOPATHOLOGY

LIVER (LI) :
-LARGE; ALL LOBES, MARGINS ROUNDED

LUNG (LU) :
-INFILTRATE, MACROPHAGE, ALVEOLAR, -MINIMAL

LIVER (LI) :
-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MODERATE
-VACUOLATION, HEPATOCELLULAR, CENTRILOBULAR TO MIDZONAL, -SLIGHT

THYMUS (TH) :
-HEMORRHAGE, -MINIMAL

THYROID (TY) :
-CYST, ULTIMOBRANCHIAL, -PRESENT

LN, MANDIBULAR (MN) :
-HEMORRHAGE, -MINIMAL

LN, MANDIBULAR (MN) :
-DIFFUSELY RED; BILATERAL; DARK RED

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 94

ANIMAL NUMBER: C95961 SEX: MALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 483.1 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:53 PROSECTOR: LAURIE J. SCHULLER REORDER: TERRI L. WILLIAMS
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG),
HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV),
TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN),
SPLEEN (SP), LN, MESENTERIC (MS), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO),
PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL),
CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK),
URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP)

C01177

APPENDIX 6
Individual Anatomic Pathology Data

PAGE: 95

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95962	SEX: MALE	DOSE GROUP: 4	SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE	
DATE OF DEATH: 12/02/98	STUDY DAY OF DEATH: 93	STUDY WEEK OF DEATH: 14	TERMINAL BODY WEIGHT: 428.1 GRAMS	
DATE AND TIME OF NECROPSY: 12/02/98 10:31	PROSECTOR: TODD ACKER		RECORDED: WENDY CHERRY	
POST-FIX WEIGHT: NOT AVAILABLE	PATHOLOGIST: JOHN EIGHMY		WEIGHER: LYNN HORNER	
ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0573	.0134 %	.0263	WEIGHT TAKEN
BRAIN (BR)	2.1762	.5083 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	3.9386	.9200 %	1.8099	WEIGHT TAKEN
LIVER (LI)	21.3603	4.9896 %	9.8154	WEIGHT TAKEN
SPLEEN (SP)	.6749	.1577 %	.3101	WEIGHT TAKEN
TESTIS (TE)	3.5204	.8223 %	1.6177	WEIGHT TAKEN
THYMUS (TH)	.2975	.0695 %	.1367	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0265	.0062 %	.0122	WEIGHT TAKEN

NECROPSY	OBSERVATIONS	HISTOPATHOLOGY
	LIVER (LI) :	
	-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MODERATE	
	-VACUOLATION, HEPATOCELLULAR, CENTRILOBULAR TO MIDZONAL, -	
	MINIMAL	
	LN, MANDIBULAR (MN) :	
	-HEMORRHAGE, -MINIMAL	
	HARDERIAN GLAND (HG) :	
	-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL	

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 96

ANIMAL NUMBER: C95962 SEX: MALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 428.1 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:31 PROSECTOR: TODD ACKER
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY RECORDER: WENDY CHERRY
WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG),
AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV), TESTIS (TE),
EPIDIDYMIDES (EP)

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 97

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95963 SEX: MALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 473.2 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 12:30 PROSECTOR: NANCY DIEDRICH RECORDER: TERRI L. WILLIAMS
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0519	.0110 %	.0250	WEIGHT TAKEN
BRAIN (BR)	2.0730	.4381 %	1.0000	WEIGHT TAKEN
KIDNEY (KO)	3.9593	.8367 %	1.9099	WEIGHT TAKEN
LIVER (LI)	26.0579	5.5067 %	12.5701	WEIGHT TAKEN
SPLEEN (SP)	.6285	.1328 %	.3032	WEIGHT TAKEN
TESTIS (TE)	3.4019	.7189 %	1.6411	WEIGHT TAKEN
THYMUS (TH)	.2799	.0592 %	.1350	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0246	.0052 %	.0119	WEIGHT TAKEN

OBSERVATIONS

NECROPSY

HISTOPATHOLOGY

LIVER (LI) :
-LARGE: ALL LOBES THICKENED, MARGINS ROUNDED
-DIFFUSELY DARK; ALL LOBES: DARK BROWN
STOMACH, GL (ST) :
-DARK FOCUS(I)/AREA(S); MUCOSA: SINGLE DARK BROWN FOCUS, 1
MM IN DIAMETER
LUNG (LU) :
-MINERALIZATION, VASCULAR, -MINIMAL
LIVER (LI) :
-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MODERATE
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
STOMACH, GL (ST) :
-EROSION, -SLIGHT
PANCREAS (PA) :
-ATROPHY, ACINAR, -MINIMAL
-INFLAMMATION, CHRONIC-ACTIVE, -SLIGHT
-PIGMENT, BROWN, -MINIMAL
HARDERIAN GLAND (HG) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
PROSTATE (PR) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -SLIGHT

001130

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 98

ANIMAL NUMBER: C95963 SEX: MALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 473.2 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 12:30 PROSECTOR: NANCY DIEDRICH
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY RECORDER: TERRI L. WILLIAMS
WEIGHER: TONY HANFELD

O B S E R V A T I O N S (CONTINUED)

HISTOPATHOLOGY

NECROPSY

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LONG (LU), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN),
SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC),
ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, NONGL (SU), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL),
CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), AUDITORY SEB GL (AS), SKIN (SK),
URINARY BLADDER (UB), SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP)

001181

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 99

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95964 SEX: MALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 462.9 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 12:07 PROSECTOR: KEVIN BILLINGS RECORDED: WENDY CHERRY
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0554	.0120 %	.0243	WEIGHT TAKEN
BRAIN (BR)	2.2840	.4934 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	4.5315	.9789 %	1.9840	WEIGHT TAKEN
LIVER (LI)	30.5553	6.6008 %	13.3780	WEIGHT TAKEN
SPLEEN (SP)	.7488	.1618 %	.3278	WEIGHT TAKEN
TESTIS (TE)	3.8137	.8239 %	1.6697	WEIGHT TAKEN
THYMUS (TH)	.2714	.0586 %	.1188	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0220	.0048 %	.0096	WEIGHT TAKEN

OBSERVATIONS

NECROPSY

HISTOPATHOLOGY

LUNG (LU) :
-INFILTRATE, MACROPHAGE, ALVEOLAR, -MINIMAL
LIVER (LI) :
-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MODERATE
-NECROSIS, COAGULATIVE, -MINIMAL
-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL
THYMUS (TH) :
-HEMORRHAGE, -MINIMAL
LN, MESENTERIC (MS) :
-HEMORRHAGE, -MINIMAL
LN, MANDIBULAR (MN) :
-HEMORRHAGE, -MINIMAL
URINARY BLADDER (UB) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
PROSTATE (PR) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -SLIGHT

001132

APPENDIX 6

Covance 6329-225
3M T-6314.1

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 100

ANIMAL NUMBER: C95964
SEX: MALE
DOSE GROUP: 4
SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98
STUDY DAY OF DEATH: 93
STUDY WEEK OF DEATH: 14
TERMINAL BODY WEIGHT: 462.9 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 12:07
PROSECTOR: KEVIN BILLINGS
RECORDED: WENDY CHERRY
POST-FIX WEIGHT: NOT AVAILABLE
PATHOLOGIST: JOHN EIGHMY
WEIGHER: LYNN HORNER

O B S E R V A T I O N S (CONTINUED)

NECROPSY

HISTOPATHOLOGY

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN),
SPLEEN (SP), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO),
PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL),
CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP)

001183

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 101

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95965 SEX: MALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 421.9 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:04 PROSECTOR: NANCY DIEDRICH RECORDER: TERRI L. WILLIAMS
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN S T A T U S
ADRENAL (AD)	.0569	.0135 %	.0271	WEIGHT TAKEN
BRAIN (BR)	2.1026	.4984 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	4.2312	1.0029 %	2.0124	WEIGHT TAKEN
LIVER (LI)	24.8958	5.9009 %	11.8405	WEIGHT TAKEN
SPLEEN (SP)	.5544	.1314 %	.2637	WEIGHT TAKEN
TESTIS (TE)	3.5478	.8409 %	1.6873	WEIGHT TAKEN
THYMUS (TH)	.2759	.0654 %	.1312	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0214	.0051 %	.0102	WEIGHT TAKEN

O B S E R V A T I O N S

NECROPSY

LIVER (LI) :
-LARGE; ALL LOBES: THICKENED
-DIFFUSELY DARK; ALL LOBES: DARK BROWN

HISTOPATHOLOGY

LIVER (LI) :
-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MODERATE
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-VACUOLATION, HEPATOCELLULAR, CENTRILOBULAR TO MIDZONAL, -
MINIMAL
KIDNEY (KD) :
-MINERALIZATION, TUBULAR, -MINIMAL
HEART (HT) :
-DEGENERATION/NECROSIS, -MINIMAL
THYROID (TY) :
-CYST, ULTIMOBRANCHIAL, -PRESENT
JEJUNUM (JE) :
>UNREMARKABLE
>NOTE:>ON SLIDE 13.

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN

001134

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 102

ANIMAL NUMBER: C95965 SEX: MALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 421.9 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:04 PROSECTOR: NANCY DIEDRICH
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY
WEIGHER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN), SPLEEN (SP),
THYMUS (TH), LN, MESENTERIC (MS), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO),
PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), ILEUM (IL), CECUM (CE),
COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS),
SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP)

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 103

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95966		SEX: MALE		DOSE GROUP: 4		SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE	
DATE OF DEATH: 12/02/98		STUDY DAY OF DEATH: 93		STUDY WEEK OF DEATH: 14		TERMINAL BODY WEIGHT: 444.3 GRAMS	
DATE AND TIME OF NECROPSY: 12/02/98 7:53		PROSECTOR: KEVIN BILLINGS		PATHOLOGIST: JOHN EIGHMY		RECORDER: WENDY CHERRY	
POST-FIX WEIGHTER: NOT AVAILABLE						WEIGHER: LYNN HORNER	
				ABSOLUTE ORGAN WEIGHT (GRAMS)		ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	
ORGAN NAME						ORGAN TO BRAIN WEIGHT RATIO	
						O R G A N S T A T U S	
ADRENAL (AD)				.0524		.0118 %	WEIGHT TAKEN
BRAIN (BR)				2.0426		.4597 %	WEIGHT TAKEN
KIDNEY (KD)				4.8321		1.0876 %	WEIGHT TAKEN
LIVER (LI)				20.9821		4.7225 %	WEIGHT TAKEN
SPLEEN (SP)				.7381		.1661 %	WEIGHT TAKEN
TESTIS (TE)				3.0674		.6904 %	WEIGHT TAKEN
THYMUS (TH)				.3532		.0795 %	WEIGHT TAKEN
THYROID/PARATHYR (TT)				.0227		.0051 %	WEIGHT TAKEN

HISTOPATHOLOGY

NECROPSY

LUNG (LU) :
-MINERALIZATION, VASCULAR, -MINIMAL
LIVER (LI) :
-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -SLIGHT
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL
THYMUS (TH) :
-HEMORRHAGE, -MINIMAL
LN, MESENTERIC (MS) :
-HEMORRHAGE, -MINIMAL
THYROID (TY) :
-CYST, ULTIMOBRANCHIAL, -PRESENT
URINARY BLADDER (UB) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

APPENDIX 6
Individual Anatomic Pathology Data

PAGE: 104

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95966 SEX: MALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 444.3 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 7:53 PROSECTOR: KEVIN BILLINGS RECORDED: WENDY CHERRY
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:
BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:
BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN),
SPLEEN (SP), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI),
STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE),
COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS),
SKIN (SK), PROSTATE (PR), SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP)

001137

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 105

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95967
DATE OF DEATH: 12/02/98
DATE AND TIME OF NECROPSY: 12/02/98 9:54
POST-FIX WEIGHTER: NOT AVAILABLE

SEX: MALE
STUDY DAY OF DEATH: 93
STUDY WEEK OF DEATH: 14
PROSECTOR: MICHELE ZIMA
PATHOLOGIST: JOHN EIGHMY

DOSE GROUP: 4
SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
TERMINAL BODY WEIGHT: 447.8 GRAMS
RECORDER: WENDY CHERRY
WEIGHER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN S T A T U S
ADRENAL (AD)	.0473	.0106 %	.0221	WEIGHT TAKEN
BRAIN (BR)	2.1398	.4778 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	4.1495	.9266 %	1.9392	WEIGHT TAKEN
LIVER (LI)	25.0600	5.5962 %	11.7114	WEIGHT TAKEN
SPLEEN (SP)	.4205	.0939 %	.1965	WEIGHT TAKEN
TESTIS (TE)	3.2710	.7305 %	1.5286	WEIGHT TAKEN
THYMUS (TH)	.1537	.0343 %	.0718	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0278	.0062 %	.0130	WEIGHT TAKEN

O B S E R V A T I O N S

NECROPSY

LUNG (LU) :
-INFILTRATE, MACROPHAGE, ALVEOLAR, -MINIMAL

LIVER (LI) :
-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MODERATE
-PIGMENT, HEPATOCELLULAR, -MINIMAL
-VACUOLATION, HEPATOCELLULAR, DIFFUSE, -SLIGHT
>NOTE:>PIGMENT IS CONCENTRATED IN THE CYTOPLASM OF CENTRILOBULAR HEPATOCYTES NEAR BILE CANALICULI.

KIDNEY (KD) :
-DILATATION, PELVIC, -SLIGHT

THYMUS (TH) :
-HEMORRHAGE, -MINIMAL
-INVOLUTION, -MINIMAL

THYROID (TY) :
-CYST, ULTIMOBRANCHIAL, -PRESENT

PITUITARY (PI) :
-CYST, -PRESENT

KIDNEY (KD) :
-LARGE PELVIS(ES); LEFT
THYMUS (TH) :
-DIFFUSELY RED; RIGHT LOBE

001138

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 106

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95967 SEX: MALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 TERMINAL BODY WEIGHT: 447.8 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 9:54 PROSECTOR: MICHELE ZIMA RECORDED: WENDY CHERRY
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EICHMY WEIGHER: LYNN HORNER

O B S E R V A T I O N S (CONTINUED) HISTOPATHOLOGY

NECROPSY

LN, MANDIBULAR (MN) :
-DIFFUSELY RED; BILATERAL: DARK RED

LN, MANDIBULAR (MN) :
-HEMORRHAGE, -SLIGHT
AUDITORY SEB GL (AS) :
-ONE EXAMINED, -PRESENT
PROSTATE (PR) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC),
ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG),
HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV),
TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), TRACHEA (TR), ESOPHAGUS (ES), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN),
SPLEEN (SP), LN, MESENTERIC (MS), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO),
STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE),
COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), SKIN (SK), URINARY BLADDER (UB),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP)

001189

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 107

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95968		SEX: MALE		DOSE GROUP: 4		SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE	
DATE OF DEATH: 12/02/98		STUDY DAY OF DEATH: 93		STUDY WEEK OF DEATH: 14		TERMINAL BODY WEIGHT: 459.5 GRAMS	
DATE AND TIME OF NECROPSY: 12/02/98 9:58		PROSECTOR: KEVIN BILLINGS		REORDER: WENDY CHERRY			
POST-FIX WEIGHT: NOT AVAILABLE		PATHOLOGIST: JOHN EIGHMY		WEIGHER: LYNN HORNER			
		ABSOLUTE ORGAN WEIGHT (GRAMS)		ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)		ORGAN TO BRAIN WEIGHT RATIO	
ORGAN NAME						O R G A N S T A T U S	
ADRENAL (AD)		.0730		.0159 %		WEIGHT TAKEN	
BRAIN (BR)		2.1136		.4600 %		WEIGHT TAKEN	
KIDNEY (KD)		4.2887		.9333 %		WEIGHT TAKEN	
LIVER (LI)		24.0192		5.2272 %		WEIGHT TAKEN	
SPLEEN (SP)		.9478		.2063 %		WEIGHT TAKEN	
TESTIS (TE)		2.8087		.6113 %		WEIGHT TAKEN	
THYMUS (TH)		.2901		.0631 %		WEIGHT TAKEN	
THYROID/PARATHYR (TT)		.0236		.0051 %		WEIGHT TAKEN	

NECROPSY	OBSERVATIONS	HISTOPATHOLOGY
	LUNG (LU) :	
	-INFILTRATE, MACROPHAGE, ALVEOLAR, -MINIMAL	
	LIVER (LI) :	
	-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MODERATE	
	-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL	
	-VACUOLATION, HEPATOCELLULAR, CENTRILOBULAR TO MIDZONAL, -MINIMAL	
	ADRENAL, CORTEX (AC) :	
	-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL	
	PROSTATE (PR) :	
	-INFILTRATE, LYMPHOHISTIOCYTIC, -SLIGHT	

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-MALOCCLUSION; BILATERAL, UPPER INCISORS

001190

APPENDIX 6
Individual Anatomic Pathology Data

PAGE: 108

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95968 SEX: MALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 459.5 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 9:58 PROSECTOR: KEVIN BILLINGS RECORDER: WENDY CHERRY
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:
BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), STOMACH, NONGL (SU),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), RECTUM (RE), LN, MANDIBULAR (MN),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:
BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN),
SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, MEDULLA (MA), AORTA (AO),
PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL),
CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG),
AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP)

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 109

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95969	SEX: MALE	DOSE GROUP: 4	SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98	STUDY DAY OF DEATH: 93	STUDY WEEK OF DEATH: 14	TERMINAL BODY WEIGHT: 465.8 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:41	PROSECTOR: JERALD W. JOHNSON	RECORDED: WENDY CHERRY	
POST-FIX WEIGHTER: NOT AVAILABLE	PATHOLOGIST: JOHN EIGHMY	WEIGHER: LYNN HORNER	
ABSOLUTE ORGAN WEIGHT (GRAMS)		ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	
ORGAN NAME		ORGAN TO BRAIN WEIGHT RATIO	
ADRENAL (AD)		.0334	
BRAIN (BR)		1.0000	
KIDNEY (KD)		2.4366	
LIVER (LI)		10.8901	
SPLEEN (SP)		.3664	
TESTIS (TE)		1.6395	
THYMUS (TH)		.1204	
THYROID/PARATHYR (TT)		.0122	
OBSERVATIONS		HISTOPATHOLOGY	
NECROPSY		LIVER (LI) : -HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MODERATE -INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL -NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL -VACUOLATION, HEPATOCELLULAR, CENTRILOBULAR TO MIDZONAL, -MINIMAL	
KIDNEY (KD) : -LARGE PELVIS(ES); RIGHT		KIDNEY (KD) : -DILATATION, PELVIC, -MODERATE HEART (HT) : -DEGENERATION/NECROSIS, -MINIMAL THYMUS (TH) : -HEMORRHAGE, -MINIMAL STOMACH, GL (ST) : -EROSION, -MINIMAL	
STOMACH, GL (ST) : -DARK FOCUS(I)/AREA(S); UP TO 2 X 1 MM		PROSTATE (PR) : -INFILTRATE, LYMPHOHISTIOCYTIC, -SLIGHT	
MUCOSA: MULTIPLE DARK BROWN AREAS,			

001192

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 110

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL-
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95969 SEX: MALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 465.8 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:41 PROSECTOR: JERALD W. JOHNSON RECORDED: WENDY CHERRY
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: LYNN HORNER

O B S E R V A T I O N S (CONTINUED)

HISTOPATHOLOGY

NECROPSY

PROSTATE (PR) :
-INFLAMMATION, CHRONIC-ACTIVE, -MINIMAL

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN), SPLEEN (SP),
LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO),
PITUITARY (PI), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE),
COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS),
SKIN (SK), URINARY BLADDER (UB), SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP)

001193

APPENDIX 6
Individual Anatomic Pathology Data

PAGE: 111

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95970
SEX: MALE
DATE OF DEATH: 12/02/98
STUDY DAY OF DEATH: 93
DATE AND TIME OF NECROPSY: 12/02/98 8:47
POST-FIX WEIGHER: NOT AVAILABLE

DOSE GROUP: 4
SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
TERMINAL BODY WEIGHT: 439.7 GRAMS
RECORDED: STEVE VAN ADESTINE
WEIGHER: TONY HANFELD

PROSECTOR: GREGORY CODDINGTON
PATHOLOGIST: JOHN EIGHMY

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS	WEIGHT TAKEN
ADRENAL (AD)	.0560	.0127 %	.0249		WEIGHT TAKEN
BRAIN (BR)	2.2490	.5115 %	1.0000		WEIGHT TAKEN
KIDNEY (KD)	3.7296	.8482 %	1.6583		WEIGHT TAKEN
LIVER (LI)	23.2250	5.2820 %	10.3268		WEIGHT TAKEN
SPLEEN (SP)	.6436	.1464 %	.2862		WEIGHT TAKEN
TESTIS (TE)	3.5189	.8003 %	1.5647		WEIGHT TAKEN
THYMUS (TH)	.2718	.0618 %	.1209		WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0230	.0052 %	.0102		WEIGHT TAKEN

NECROPSY

THYROID/PARATHYR (TT)

THYROID (TH) :
-HEMORRHAGE, -MINIMAL

PANCREAS (PA) :
-INFLAMMATION, CHRONIC-ACTIVE, -MINIMAL
-PIGMENT, BROWN, -MINIMAL
LN, MANDIBULAR (MN) :
-HEMORRHAGE, -MINIMAL

LIVER (LI) :
-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MODERATE
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-NECROSIS, COAGULATIVE, -MINIMAL
-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL
-PIGMENT, HEPATOCELLULAR, -MINIMAL

THYMUS (TH) :
-HEMORRHAGE, -MINIMAL

PANCREAS (PA) :
-INFLAMMATION, CHRONIC-ACTIVE, -MINIMAL
-PIGMENT, BROWN, -MINIMAL
LN, MANDIBULAR (MN) :
-HEMORRHAGE, -MINIMAL

LN, MANDIBULAR (MN) :
-MOTTLED; BILATERAL; DARK RED AND TAN

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-MALOCCLUSION; BILATERAL UPPER INCISORS

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 112

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95970 SEX: MALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 439.7 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:47 PROSECTOR: GREGORY CODDINGTON RECORDER: STEVE VAN ADESTINE
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TV), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG),
HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV),
TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), LN, MESENTERIC (MS), THYROID (TV), PARATHYROID (PT), ADRENAL, CORTEX (AC),
ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), DUODENUM (DU),
JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG),
AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV), TESTIS (TE),
EPIDIDYMIDES (EP)

001195

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 113

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95971 SEX: MALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
DATE OF DEATH: 10/01/98 STUDY DAY OF DEATH: 31 STUDY WEEK OF DEATH: 5 TERMINAL BODY WEIGHT: 309.1 GRAMS
DATE AND TIME OF NECROPSY: 10/01/98 11:57 PROSECTOR: MATTHEW AXTMAN RECORDED: KEVIN BILLINGS
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHED: MICHELE ZIMA

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	O R G A N	
				S T A T U S	
ADRENAL (AD)	---	---	---	NOT TAKEN	
BRAIN (BR)	---	---	---	NOT TAKEN	
KIDNEY (KD)	---	---	---	NOT TAKEN	
LIVER (LI)	---	---	---	WEIGHT TAKEN	
SPLEEN (SP)	14.5443	4.7054 %	---	NOT TAKEN	
TESTIS (TE)	---	---	---	NOT TAKEN	
THYMUS (TH)	---	---	---	NOT TAKEN	
THYROID/PARATHYR (TT)	---	---	---	NOT TAKEN	

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:
BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI), GENERAL COMMENT (GC)

001196

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 114

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL-
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95972 SEX: MALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
DATE OF DEATH: 10/01/98 STUDY DAY OF DEATH: 31 STUDY WEEK OF DEATH: 5 TERMINAL BODY WEIGHT: 333.3 GRAMS
DATE AND TIME OF NECROPSY: 10/01/98 12:06 PROSECTOR: MATTHEW AXTMAN RECORDER: KEVIN BILLINGS
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: MICHELE ZIMA

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	---	---	---	NOT TAKEN
BRAIN (BR)	---	---	---	NOT TAKEN
KIDNEY (KD)	---	---	---	NOT TAKEN
LIVER (LI)	14.9723	4.4921 %	---	WEIGHT TAKEN
SPLEEN (SP)	---	---	---	NOT TAKEN
TESTIS (TE)	---	---	---	NOT TAKEN
THYMUS (TH)	---	---	---	NOT TAKEN
THYROID/PARATHYR (TT)	---	---	---	NOT TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI), GENERAL COMMENT (GC)

C01197

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 115

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95973	SEX: MALE	DOSE GROUP: 4	SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
DATE OF DEATH: 10/01/98	STUDY DAY OF DEATH: 31	STUDY WEEK OF DEATH: 5	TERMINAL BODY WEIGHT: 276.7 GRAMS
DATE AND TIME OF NECROPSY: 10/01/98 11:48	PROSECTOR: WENDY CHERRY		REORDER: KEVIN BILLINGS
POST-FIX WEIGHTER: NOT AVAILABLE	PATHOLOGIST: JOHN EIGHMY		WEIGHER: MICHELE ZIMA
	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO
ORGAN NAME			S T A T U S
ADRENAL (AD)			NOT TAKEN
BRAIN (BR)			NOT TAKEN
KIDNEY (KD)			NOT TAKEN
LIVER (LI)		4.7703 %	WEIGHT TAKEN
SPLEEN (SP)	13.1995		NOT TAKEN
TESTIS (TE)			NOT TAKEN
THYMUS (TH)			NOT TAKEN
THYROID/PARATHYR (TT)			
THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:			
BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),			
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),			
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),			
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),			
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),			
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),			
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI), GENERAL COMMENT (GC)			

001198

APPENDIX 6
Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 116

ANIMAL NUMBER: C95974	SEX: MALE	DOSE GROUP: 4	SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
DATE OF DEATH: 10/01/98	STUDY DAY OF DEATH: 31	STUDY WEEK OF DEATH: 5	TERMINAL BODY WEIGHT: 311.4 GRAMS
DATE AND TIME OF NECROPSY: 10/01/98 11:22	PROSECTOR: TONY HANFELD		REORDER: KEVIN BILLINGS
POST-FIX WEIGHT: NOT AVAILABLE	PATHOLOGIST: JOHN EIGHMY		WEIGHER: MICHELE ZIMA
ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO
ADRENAL (AD)	-----	-----	-----
BRAIN (BR)	-----	-----	NOT TAKEN
KIDNEY (KD)	-----	-----	NOT TAKEN
LIVER (LI)	-----	-----	NOT TAKEN
SPLEEN (SP)	14.4142	4.6288 %	WEIGHT TAKEN
TESTIS (TE)	-----	-----	NOT TAKEN
THYROID (TH)	-----	-----	NOT TAKEN
THYROID/PARATHYR (TT)	-----	-----	NOT TAKEN
THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:			
BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),			
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),			
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),			
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),			
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),			
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),			
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI), GENERAL COMMENT (GC)			

001199

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 117

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95975	SEX: MALE	DOSE GROUP: 4	SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
DATE OF DEATH: 10/01/98	STUDY DAY OF DEATH: 31	STUDY WEEK OF DEATH: 5	TERMINAL BODY WEIGHT: 327.4 GRAMS
DATE AND TIME OF NECROPSY: 10/01/98 11:53	PROSECTOR: MATTHEW AXTMAN		REORDER: KEVIN BILLINGS
POST-FIX WEIGHT: NOT AVAILABLE	PATHOLOGIST: JOHN EIGHMY		WEIGHER: MICHELE ZIMA
	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO
ORGAN NAME			S T A T U S
ADRENAL (AD)	---	---	NOT TAKEN
BRAIN (BR)	---	---	NOT TAKEN
KIDNEY (KD)	---	---	NOT TAKEN
LIVER (LI)	---	5.0104 %	WEIGHT TAKEN
SPLEEN (SP)	16.4042	---	NOT TAKEN
TESTIS (TE)	---	---	NOT TAKEN
THYMUS (TH)	---	---	NOT TAKEN
THYROID/PARATHYR (TT)	---	---	
THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:			
BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR), SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI), GENERAL COMMENT (GC)			

001200

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 118

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95976	SEX: MALE	DOSE GROUP: 4	SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE	TERMINAL BODY WEIGHT: 382.8 GRAMS
DATE OF DEATH: 12/02/98	STUDY DAY OF DEATH: 93	STUDY WEEK OF DEATH: 14	REORDER: WENDY CHERRY	
DATE AND TIME OF NECROPSY: 12/02/98 8:23	PROSECTOR: JILL PAUS		WEIGHER: LYNN HORNER	
POST-FIX WEIGHER: NOT AVAILABLE	PATHOLOGIST: JOHN EIGHMY			
ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN S T A T U S
ADRENAL (AD)	.0447	.0117 %	.0203	WEIGHT TAKEN
BRAIN (BR)	2.1994	.5746 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	3.5516	.9278 %	1.6148	WEIGHT TAKEN
LIVER (LI)	21.3634	5.5808 %	9.7133	WEIGHT TAKEN
SPLEEN (SP)	.4872	.1273 %	.2215	WEIGHT TAKEN
TESTIS (TE)	3.4775	.9084 %	1.5811	WEIGHT TAKEN
THYMUS (TH)	.2380	.0622 %	.1082	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0300	.0078 %	.0136	WEIGHT TAKEN

NECROPSY	O B S E R V A T I O N S	HISTOPATHOLOGY
	LIVER (LI) :	
	-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MODERATE	
	-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL	
	-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL	
	-VACUOLATION, HEPATOCELLULAR, CENTRILOBULAR TO MIDZONAL, -MINIMAL	
	THYMUS (TH) :	
	-HEMORRHAGE, -MINIMAL	
	THYROID (TY) :	
	-CYST, ULTIMOBRANCHIAL, -PRESENT	
	LN, MANDIBULAR (MN) :	
	-HEMORRHAGE, -SLIGHT	

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

001201

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 119

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95976 SEX: MALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 382.8 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:23 PROSECTOR: JILL PAUS RECORDED: WENDY CHERRY
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:
BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), NONGL (SU),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:
BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), LN, MESENTERIC (MS), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA),
AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE),
ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS),
SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP)

001202

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 120

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95977 SEX: MALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL, SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 454.6 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:58 PROSECTOR: MICHELE ZIMA REORDER: WENDY CHERRY
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS	WEIGHT TAKEN
ADRENAL (AD)	.0587	.0129 %	.0280		WEIGHT TAKEN
BRAIN (BR)	2.0929	.4604 %	1.0000		WEIGHT TAKEN
KIDNEY (KD)	4.2673	.9387 %	2.0389		WEIGHT TAKEN
LIVER (LI)	25.0541	5.5112 %	11.9710		WEIGHT TAKEN
SPLEEN (SP)	.7871	.1731 %	.3761		WEIGHT TAKEN
TESTIS (TE)	3.4440	.7576 %	1.6456		WEIGHT TAKEN
THYMUS (TH)	.2202	.0484 %	.1052		WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0339	.0075 %	.0162		WEIGHT TAKEN

OBSERVATIONS

HISTOPATHOLOGY

NECROPSY

LUNG (LU) :
-INFILTRATE, MACROPHAGE, ALVEOLAR, -MINIMAL
LIVER (LI) :
-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MODERATE
-VACUOLATION, HEPATOCELLULAR, DIFFUSE, -SLIGHT
HEART (HT) :
-DEGENERATION/NECROSIS, -MINIMAL
THYMUS (TH) :
-HEMORRHAGE, -MINIMAL
LN, MANDIBULAR (MN) :
-HEMORRHAGE, -MINIMAL

THYMUS (TH) :
-RED FOCUS (I)/AREA (S); BOTH LOBES: MULTIPLE DARK RED,
PINPOINT FOCI
LN, MANDIBULAR (MN) :
-DIFFUSELY RED; BILATERAL
GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN

C01203

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 121

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
TERMINAL BODY WEIGHT: 454.6 GRAMS
RECORDED: WENDY CHERRY
WEIGHED: LYNN HORNER

ANIMAL NUMBER: C95977 SEX: MALE DOSE GROUP: 4
STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14
DATE OF DEATH: 12/02/98 PROSECTOR: MICHELE ZIMA
DATE AND TIME OF NECROPSY: 12/02/98 10:58 PATHOLOGIST: JOHN EIGHMY
POST-FIX WEIGHT: NOT AVAILABLE

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC),
ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG),
HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV),
TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN),
SPLEEN (SP), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA),
AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE),
ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS),
SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP)

001204

APPENDIX 6

Covance 6329-225
3M T-6314.1

Individual Anatomic Pathology Data

PAGE: 122

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95978 SEX: MALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
 DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 454.2 GRAMS
 DATE AND TIME OF NECROPSY: 12/02/98 8:40 PROSECTOR: KEVIN BILLINGS REORDER: WENDY CHERRY
 POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0568	.0125 %	.0266	WEIGHT TAKEN
BRAIN (BR)	2.1377	.4707 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	3.6874	.8118 %	1.7249	WEIGHT TAKEN
LIVER (LI)	28.4115	6.2553 %	13.2907	WEIGHT TAKEN
SPLEEN (SP)	.6243	.1375 %	.2920	WEIGHT TAKEN
TESTIS (TE)	3.8868	.8537 %	1.8182	WEIGHT TAKEN
THYROID (TH)	.1836	.0404 %	.0859	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0245	.0054 %	.0115	WEIGHT TAKEN

O B S E R V A T I O N S

HISTOPATHOLOGY

NECROPSY

LIVER (LI) :
 -LIGHT FOCUS(I)/AREA(S) : DIAPHRAGMATIC SURFACE OF LEFT
 LATERAL LOBE: MULTIPLE TAN AREAS, PINPOINT AND UP TO 1 MM IN
 DIAMETER; COLLECTED ON ROUTINE SECTION

LUNG (LU) :
 -MINERALIZATION, VASCULAR, -MINIMAL

LIVER (LI) :
 -HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MODERATE
 -INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
 -NECROSIS, COAGULATIVE, -SLIGHT
 -VACUOLATION, HEPATOCELLULAR, CENTRILLOBULAR TO MIDZONAL, -
 SLIGHT

THYMUS (TH) :
 -HEMORRHAGE, -MINIMAL

THYROID (TY) :
 -CYST, ULTIMOBRANCHIAL, -PRESENT

PROSTATE (PR) :
 -INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL

GENERAL COMMENT (GC) :
 -BONE MARROW SMEAR TAKEN

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 123

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

SCHEDULED, TERMINAL SACRIFICE
TERMINAL BODY WEIGHT: 454.2 GRAMS

RECORDED: WENDY CHERRY
WEIGHED: LYNN HORNER

DOSE GROUP: 4 SACRIFICE STATUS: 14

STUDY WEEK OF DEATH: 93

SEX: MALE
STUDY DAY OF DEATH: 12/02/98 8:40

DATE OF DEATH: 12/02/98
POST-FIX WEIGHT: NOT AVAILABLE

PROSECTOR: KEVIN BILLINGS
PATHOLOGIST: JOHN EIGHMY

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR), (SM),
SPINAL CORD (SC), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SK),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), GL (ST), STOMACH, NONGL (SU),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), LN, MANDIBULAR (MN),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN),
SPLEEN (SP), LN, MESENTERIC (MS), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO),
PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL),
CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG),
AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP)

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 124

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
TERMINAL BODY WEIGHT: 435.0 GRAMS
REORDER: STEVE VAN ADESTINE
WEIGHER: TONY HANFELD

DOSE GROUP: 4
STUDY WEEK OF DEATH: 14
PROSECTOR: LAURIE J. SCHULLER
PATHOLOGIST: JOHN EIGHMY

ANIMAL NUMBER: C95979
SEX: MALE
STUDY DAY OF DEATH: 93
DATE OF DEATH: 12/02/98 8:16
DATE AND TIME OF NECROPSY: 12/02/98 8:16
POST-FIX WEIGHT: NOT AVAILABLE

ORGAN
STATISTICS

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	WEIGHT TAKEN
ADRENAL (AD)	.0499	.0115 %	.0221	WEIGHT TAKEN
BRAIN (BR)	2.2608	.5197 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	3.8911	.8945 %	1.7211	WEIGHT TAKEN
LIVER (LI)	23.2324	5.3408 %	10.2762	WEIGHT TAKEN
SPLEEN (SP)	.6649	.1529 %	.2941	WEIGHT TAKEN
TESTIS (TE)	3.4924	.8029 %	1.5448	WEIGHT TAKEN
THYROID (TH)	.2278	.0524 %	.1008	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0253	.0058 %	.0112	WEIGHT TAKEN

OBSERVATIONS HISTOPATHOLOGY

NECROPSY

LIVER (LI) :
-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MODERATE
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-NECROSIS, INDIVIDUAL HEPATOCYTE, -SLIGHT
THYROID (TY) :
-CYST, ULTIMOBRANCHIAL, -PRESENT
AUDITORY SEB GL (AS) :
-ONE EXAMINED, -PRESENT

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 125

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
TERMINAL BODY WEIGHT: 435.0 GRAMS
REORDER: STEVE VAN ADESTINE
WEIGHER: TONY HANFELD

DOSE GROUP: 4
STUDY WEEK OF DEATH: 14
PROSECTOR: LAURIE J. SCHULLER
PATHOLOGIST: JOHN EIGHMY

SEX: MALE
STUDY DAY OF DEATH: 12/02/98 8:16
DATE OF NECROPSY: 12/02/98
POST-FIX WEIGHER: NOT AVAILABLE

ANIMAL NUMBER: C95979

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EV), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EV), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
ADRENAL, MEDULLA (MA), AORTA (AO), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG),
TESTIS (TE), EPIDIDYMIDES (EP)

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 126

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL-
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95980 SEX: MALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 TERMINAL BODY WEIGHT: 434.0 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 12:24 PROSECTOR: KEVIN BILLINGS RECORDER: WENDY CHERRY
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	O R G A N	
				S T A T U S	WEIGHT TAKEN
ADRENAL (AD)	.0539	.0124 %	.0237		WEIGHT TAKEN
BRAIN (BR)	2.2717	.5234 %	1.0000		WEIGHT TAKEN
KIDNEY (KD)	4.2176	.9718 %	1.8566		WEIGHT TAKEN
LIVER (LI)	24.1794	5.5713 %	10.6437		WEIGHT TAKEN
SPLEEN (SP)	.6736	.1552 %	.2965		WEIGHT TAKEN
TESTIS (TE)	3.5469	.8173 %	1.5613		WEIGHT TAKEN
THYMUS (TH)	.2741	.0632 %	.1207		WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0237	.0055 %	.0104		WEIGHT TAKEN

O B S E R V A T I O N S

NECROPSY

HISTOPATHOLOGY

LIVER (LI) :
-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MODERATE
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
LN, MANDIBULAR (MN) :
-HEMORRHAGE, -MINIMAL
AUDITORY SEB GL (AS) :
-ONE EXAMINED, -PRESENT

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

001209

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 127

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95980 SEX: MALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 434.0 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 12:24 PROSECTOR: KEVIN BILLINGS RECORDED: WENDY CHERRY
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TV), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR),
SEMINAL VESICLES (SV), TESTIS (TE), EPIDIDYMIDES (EP), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TV), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG),
HARDERIAN GLAND (HG), SKIN (SK), URINARY BLADDER (UB), PROSTATE (PR), SEMINAL VESICLES (SV), TESTIS (TE),
EPIDIDYMIDES (EP)

6329210

APPENDIX 6

Covance 6329-225
3M T-6314.1

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 128

ANIMAL NUMBER: C95981
 DATE OF DEATH: 12/02/98
 DATE AND TIME OF NECROPSY: 12/02/98 8:39
 POST-FIX WEIGHT: NOT AVAILABLE

SEX: FEMALE
 STUDY DAY OF DEATH: 93
 PROSECTOR: JOHN S. HALFORD
 PATHOLOGIST: JOHN EIGHMY

DOSE GROUP: 1
 STUDY WEEK OF DEATH: 14
 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
 TERMINAL BODY WEIGHT: 296.6 GRAMS
 RECORDER: WENDY CHERRY
 WEIGHER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0884	.0298 %	.0444	WEIGHT TAKEN
BRAIN (BR)	1.9910	.6713 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.5928	.8742 %	1.3023	WEIGHT TAKEN
LIVER (LI)	11.3754	3.8353 %	5.7134	WEIGHT TAKEN
OVARY (OV)	.0971	.0327 %	.0488	WEIGHT TAKEN
SPLEEN (SP)	.9631	.3247 %	.4837	WEIGHT TAKEN
THYMUS (TH)	.2919	.0984 %	.1466	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0246	.0083 %	.0124	WEIGHT TAKEN

NECROPSY

SPLEEN (SP) :
 -ADHESION(S); ENTIRE SPLEEN ADHERED TO PANCREAS AND SURROUNDING ADIPOSE TISSUE; MULTIPLE GRAY AND TAN FIBROUS ADHESIONS; COLLECTED INTACT

PITUITARY (PI) :
 -LARGE; 6 X 5 X 3 MM; APPEARS SWOLLEN

HISTOPATHOLOGY

LIVER (LI) :
 -INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL

SPLEEN (SP) :
 -INFLAMMATION, CHRONIC, CAPSULAR, -MODERATE
 >NOTE:>THERE ARE NUMEROUS PIGMENT-CONTAINING MACROPHAGES IN THE AREAS OF CHRONIC INFLAMMATION.

THYROID (TY) :
 -CYST, ULTIMOBRANCHIAL, -PRESENT

PITUITARY (PI) :
 -HYPERPLASIA, -MODERATE

PANCREAS (PA) :
 -ATROPHY, ACINAR, -MINIMAL

RECTUM (RE) :
 -INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL

LN, MANDIBULAR (MN) :
 -PARASITISM, -PRESENT
 -HYPERPLASIA, PLASMACYTIC, -MODERATE

001211

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 129

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95981 SEX: FEMALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 296.6 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:39 PROSECTOR: JOHN S. HALFORD RECORDED: WENDY CHERRY
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

O B S E R V A T I O N S (CONTINUED)

NECROPSY

HISTOPATHOLOGY

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC),
ADRENAL, MEDULLA (MA), AORTA (AO), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU),
JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG),
HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF), URINARY BLADDER (UB), UTERUS (UT),
CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), THYMUS (TH), LN, MESENTERIC (MS), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA),
AORTA (AO), STOMACH, GL (ST), STOMACH, NONGL (SU), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA)

001212

APPENDIX 6

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 130

ANIMAL NUMBER: C95982	SEX: FEMALE	DOSE GROUP: 1	SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98	STUDY DAY OF DEATH: 93	STUDY WEEK OF DEATH: 14	TERMINAL BODY WEIGHT: 251.3 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:38	PROSECTOR: TODD ACKER		REORDER: WENDY CHERRY
POST-FIX WEIGHER: NOT AVAILABLE	PATHOLOGIST: JOHN EIGHMY		WEIGHER: LYNN HORNER
ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO
ADRENAL (AD)	.0501	.0199 %	
BRAIN (BR)	1.7308	.6887 %	WEIGHT TAKEN
KIDNEY (KD)	1.8839	.7497 %	WEIGHT TAKEN
LIVER (LI)	7.3455	2.9230 %	WEIGHT TAKEN
OVARY (OV)	.1041	.0414 %	WEIGHT TAKEN
SPLEEN (SP)	.4764	.1896 %	WEIGHT TAKEN
THYMUS (TH)	.3310	.1317 %	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0135	.0054 %	WEIGHT TAKEN

NECROPSY		OBSERVATIONS	HISTOPATHOLOGY
GENERAL COMMENT (GC) :			
-BONE MARROW SMEAR TAKEN			
-NO MACROSCOPIC LESIONS			

001223

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 131

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95982 SEX: FEMALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 251.3 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:38 PROSECTOR: TODD ACKER RECORDER: WENDY CHERRY
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA)

001214

APPENDIX 6

Covance 6329-225
3M T-6314.1

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 132

ANIMAL NUMBER: C95983
 DATE OF DEATH: 12/02/98
 DATE AND TIME OF NECROPSY: 12/02/98 12:35
 POST-FIX WEIGHT: NOT AVAILABLE

SEX: FEMALE
 STUDY DAY OF DEATH: 93
 PROSECTOR: TODD ACKER
 PATHOLOGIST: JOHN EIGHMY

DOSE GROUP: 1
 STUDY WEEK OF DEATH: 14
 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
 TERMINAL BODY WEIGHT: 223.7 GRAMS
 RECORDER: WENDY CHERRY
 WEIGHTER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0677	.0303 %	.0369	WEIGHT TAKEN
BRAIN (BR)	1.8357	.8206 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.1211	.9482 %	1.1555	WEIGHT TAKEN
LIVER (LI)	7.7858	3.4805 %	4.2413	WEIGHT TAKEN
OVARY (OV)	.1169	.0523 %	.0637	WEIGHT TAKEN
SPLEEN (SP)	.4244	.1897 %	.2312	WEIGHT TAKEN
THYMUS (TH)	.2297	.1027 %	.1251	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0160	.0072 %	.0087	WEIGHT TAKEN

NECROPSY

OBSERVATIONS

HISTOPATHOLOGY

LIVER (LI) :
 -FIBROSIS, CAPSULAR, -MINIMAL
 -INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
 -VACUOLATION, HEPATOCELLULAR, PERIPORTAL, -MINIMAL
 THYMUS (TH) :
 -HEMORRHAGE, -MINIMAL
 -INVOLUTION, -MINIMAL
 THYROID (TY) :
 -CYST, ULTIMOBRANCHIAL, -PRESENT

GENERAL COMMENT (GC) :
 -BONE MARROW SMEAR TAKEN
 -NO MACROSCOPIC LESIONS

001245

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 133

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95983 SEX: FEMALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 223.7 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 12:35 PROSECTOR: TODD ACKER REORDER: WENDY CHERRY
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TV), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), LN, MESENTERIC (MS), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA),
AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE),
ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG),
AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF), URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV),
VAGINA (VA)

001216

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 134

ANIMAL NUMBER: C95984
DATE OF DEATH: 12/02/98
DATE AND TIME OF NECROPSY: 12/02/98 12:26
POST-FIX WEIGHER: NOT AVAILABLE
SEX: FEMALE
STUDY DAY OF DEATH: 93
STUDY WEEK OF DEATH: 14
PROSECTOR: JERALD W. JOHNSON
PATHOLOGIST: JOHN EIGHMY
DOSE GROUP: 1
SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
TERMINAL BODY WEIGHT: 266.8 GRAMS
RECORDER: WENDY CHERRY
WEIGHER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN S T A T U S
ADRENAL (AD)	.0689	.0258 %	.0327	WEIGHT TAKEN
BRAIN (BR)	2.1061	.7894 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.2832	.8558 %	1.0841	WEIGHT TAKEN
LIVER (LI)	8.6507	3.2424 %	4.1074	WEIGHT TAKEN
OVARY (OV)	.1184	.0444 %	.0562	WEIGHT TAKEN
SPLEEN (SP)	.5182	.1942 %	.2460	WEIGHT TAKEN
THYMUS (TH)	.2544	.0954 %	.1208	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0185	.0069 %	.0088	WEIGHT TAKEN

O B S E R V A T I O N S

NECROPSY

HISTOPATHOLOGY

UTERUS (UT) :
-LUMEN FILLED WITH FLUID; BILATERAL: CLEAR FLUID
GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN

THYROID (TY) :
-CYST, ULTIMOBRANCHIAL, -PRESENT
UTERUS (UT) :
>UNREMARKABLE

001217

APPENDIX 6

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 135

ANIMAL NUMBER: C95984 SEX: FEMALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 266.8 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 12:26 PROSECTOR: JERALD W. JOHNSON REORDER: WENDY CHERRY
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), PARATHYROID (PT), ADRENAL, CORTEX (AC),
ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), CERVIX (CV), OVARY (OV), VAGINA (VA)

001218

APPENDIX 6

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 136

ANIMAL NUMBER: C95985		SEX: FEMALE	DOSE GROUP: 1	SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98		STUDY DAY OF DEATH: 93	STUDY WEEK OF DEATH: 14	TERMINAL BODY WEIGHT: 283.0 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:20		PROSECTOR: JILL PAUS		REORDER: WENDY CHERRY
POST-FIX WEIGHTER: NOT AVAILABLE		PATHOLOGIST: JOHN EIGHMY		WEIGHER: LYNN HORNER
ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0785	.0277 %	.0387	WEIGHT TAKEN
BRAIN (BR)	2.0288	.7169 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.1159	.7477 %	1.0429	WEIGHT TAKEN
LIVER (LI)	8.9903	3.1768 %	4.4313	WEIGHT TAKEN
OVARY (OV)	.1524	.0539 %	.0751	WEIGHT TAKEN
SPLEEN (SP)	.5420	.1915 %	.2672	WEIGHT TAKEN
THYMUS (TH)	.2466	.0871 %	.1215	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0227	.0080 %	.0112	WEIGHT TAKEN

NECROPSY	OBSERVATIONS	HISTOPATHOLOGY
	LIVER (LI) :	
	-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL	
	LN, MANDIBULAR (MN) :	
	-HEMORRHAGE, -MINIMAL	
	AUDITORY SEB GL (AS) :	
	-ONE EXAMINED, -PRESENT	
GENERAL COMMENT (GC) :		
-BONE MARROW SMEAR TAKEN		
-NO MACROSCOPIC LESIONS		

001219

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 137

ANIMAL NUMBER: C95985 SEX: FEMALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 283.0 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:20 PROSECTOR: JILL PAUS RECORDER: WENDY CHERRY
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG),
HARDERIAN GLAND (HG), SKIN (SK), MAMMARY, FEMALE (MF), URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV),
VAGINA (VA)

APPENDIX 6
Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 138

ANIMAL NUMBER: C95986 SEX: FEMALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 264.3 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 12:34 PROSECUTOR: GREGORY CODDINGTON REORDER: TERRI L. WILLIAMS
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0824	.0312 %	.0414	WEIGHT TAKEN
BRAIN (BR)	1.9890	.7526 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.1350	.8078 %	1.0734	WEIGHT TAKEN
LIVER (LI)	7.8859	2.9837 %	3.9648	WEIGHT TAKEN
OVARY (OV)	.1421	.0538 %	.0714	WEIGHT TAKEN
SPLEEN (SP)	.5597	.2118 %	.2814	WEIGHT TAKEN
THYMUS (TH)	.3403	.1288 %	.1711	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0156	.0059 %	.0078	WEIGHT TAKEN

NECROPSY OBSERVATIONS HISTOPATHOLOGY

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

001231

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 139

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95986 SEX: FEMALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 264.3 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 12:34 PROSECTOR: GREGORY CODDINGTON RECORDED: TERRI L. WILLIAMS
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TV), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TV), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA)

001222

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.113-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 140

ANIMAL NUMBER: C95987
 DATE OF DEATH: 12/02/98
 DATE AND TIME OF NECROPSY: 12/02/98 10:51
 POST-FIX WEIGHT: NOT AVAILABLE

SEX: FEMALE
 STUDY DAY OF DEATH: 93
 PROSECTOR: JOHN S. HALFORD
 PATHOLOGIST: JOHN EIGHMY

DOSE GROUP: 1
 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
 STUDY WEEK OF DEATH: 14
 TERMINAL BODY WEIGHT: 244.1 GRAMS
 RECORDER: WENDY CHERRY
 WEIGHER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0980	.0401 %	.0519	WEIGHT TAKEN
BRAIN (BR)	1.8897	.7741 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	1.9180	.7857 %	1.0150	WEIGHT TAKEN
LIVER (LI)	7.2576	2.9732 %	3.8406	WEIGHT TAKEN
OVARY (OV)	.1316	.0539 %	.0696	WEIGHT TAKEN
SPLEEN (SP)	.4780	.1958 %	.2530	WEIGHT TAKEN
THYMUS (TH)	.3293	.1349 %	.1743	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0233	.0095 %	.0123	WEIGHT TAKEN

NECROPSY

O B S E R V A T I O N S

HISTOPATHOLOGY

LUNG (LU) :
 -INFILTRATE, MACROPHAGE, ALVEOLAR, -MINIMAL
 LIVER (LI) :
 -INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
 THYROID (TY) :
 -CYST, ULTIMOBRANCHIAL, -PRESENT
 LN, MANDIBULAR (MN) :
 -HEMORRHAGE, -MINIMAL
 UTERUS (UT) :
 -DILATATION, -MINIMAL

GENERAL COMMENT (GC) :
 -BONE MARROW SMEAR TAKEN
 -NO MACROSCOPIC LESIONS

001223

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 141

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95987 SEX: FEMALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 244.1 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:51 PROSECTOR: JOHN S. HALFORD REORDER: WENDY CHERRY
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN),
SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA),
AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE),
ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS),
SKIN (SK), MAMMARY, FEMALE (MF), URINARY BLADDER (UB), CERVIX (CV), OVARY (OV), VAGINA (VA)

001234

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 142

ANIMAL NUMBER: C95988 SEX: FEMALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 252.6 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:22 PROSECTOR: KEVIN BILLINGS REORDER: WENDY CHERRY
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0716	.0283 %	.0387	WEIGHT TAKEN
BRAIN (BR)	1.8513	.7329 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	1.8853	.7464 %	1.0184	WEIGHT TAKEN
LIVER (LI)	7.0591	2.7946 %	3.8131	WEIGHT TAKEN
OVARY (OV)	.1193	.0472 %	.0644	WEIGHT TAKEN
SPLEEN (SP)	.4753	.1882 %	.2567	WEIGHT TAKEN
THYMUS (TH)	.3660	.1449 %	.1977	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0136	.0054 %	.0073	WEIGHT TAKEN

NECROPSY

OBSERVATIONS

HISTOPATHOLOGY

LIVER (LI) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
KIDNEY (KD) :
-MINERALIZATION, TUBULAR, -MINIMAL
THYMUS (TH) :
-HEMORRHAGE, -MINIMAL
THYROID (TY) :
-CYST, ULTIMOBRANCHIAL, -PRESENT

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

001225

APPENDIX 6

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 143

ANIMAL NUMBER: C95988
DATE OF DEATH: 12/02/98
DATE AND TIME OF NECROPSY: 12/02/98 10:22
POST-FIX WEIGHER: NOT AVAILABLE
SEX: FEMALE
STUDY DAY OF DEATH: 93
DOSE GROUP: 1
SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
TERMINAL BODY WEIGHT: 252.6 GRAMS
PROSECTOR: KEVIN BILLINGS
RECORDED: WENDY CHERRY
PATHOLOGIST: JOHN EIGHTY
WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN),
SPLEEN (SP), LN, MESENTERIC (MS), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO),
PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL),
CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG),
AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF), URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV),
VAGINA (VA)

001226

APPENDIX 6
Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 144

ANIMAL NUMBER: C95989 SEX: FEMALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 277.0 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:40 PROSECTOR: LAURIE J. SCHULLER REORDER: STEVE VAN ADESTINE
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUSES
ADRENAL (AD)	.0673	.0243 %	.0353	WEIGHT TAKEN
BRAIN (BR)	1.9052	.6878 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	1.9871	.7174 %	1.0430	WEIGHT TAKEN
LIVER (LI)	7.3952	2.6697 %	3.8816	WEIGHT TAKEN
OVARY (OV)	.1033	.0373 %	.0542	WEIGHT TAKEN
SPLEEN (SP)	.5285	.1908 %	.2774	WEIGHT TAKEN
THYMUS (TH)	.2674	.0965 %	.1404	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0121	.0044 %	.0064	WEIGHT TAKEN

OBSERVATIONS

NECROPSY

HISTOPATHOLOGY

LUNG (LU) :
-MINERALIZATION, VASCULAR, -MINIMAL
LIVER (LI) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL
THYROID (TY) :
-CYST, ULTIMOBRANCHIAL, -PRESENT
AUDITORY SEB GL (AS) :
-ONE EXAMINED, -PRESENT

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 145

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95989 SEX: FEMALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 TERMINAL BODY WEIGHT: 277.0 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:40 PROSECTOR: LAURIE J. SCHULLER RECORDER: STEVE VAN ADESTINE
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN),
SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA),
AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE),
ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG),
SKIN (SK), MAMMARY, FEMALE (MF), URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA)

001238

APPENDIX 6

Covance 6329-225
3M T-6314.1

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 146ANIMAL NUMBER: C95990
SEX: FEMALE
DOSE GROUP: 1
SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98
STUDY DAY OF DEATH: 93
STUDY WEEK OF DEATH: 14
TERMINAL BODY WEIGHT: 313.0 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 12:30
PROSECTOR: KIMBERLY DURLAND
RECORDER: TERRI L. WILLIAMS
POST-FIX WEIGHTER: NOT AVAILABLE
PATHOLOGIST: JOHN EIGHMY
WEIGHER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0642	.0205 %	.0338	WEIGHT TAKEN
BRAIN (BR)	1.8999	.6070 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.3886	.7631 %	1.2572	WEIGHT TAKEN
LIVER (LI)	8.8486	2.8270 %	4.6574	WEIGHT TAKEN
OVARY (OV)	.1396	.0446 %	.0735	WEIGHT TAKEN
SPLEEN (SP)	.4933	.1576 %	.2596	WEIGHT TAKEN
THYMUS (TH)	.2034	.0650 %	.1071	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0280	.0089 %	.0147	WEIGHT TAKEN

OBSERVATIONS

NECROPSY

HISTOPATHOLOGY

LUNG (LU) :
-MINERALIZATION, VASCULAR, -MINIMAL
LIVER (LI) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 147

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95990 SEX: FEMALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 313.0 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 12:30 PROSECTOR: KIMBERLY DURLAND RECORDED: TERRI L. WILLIAMS
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TV), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN),
SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC),
ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA)

001230

ANIMAL NUMBER: C95991
SEX: FEMALE
DATE OF DEATH: 10/01/98
STUDY DAY OF DEATH: 31
SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
TERMINAL BODY WEIGHT: 205.1 GRAMS
DATE AND TIME OF NECROPSY: 10/01/98 11:32
PROSECTOR: TONY HANFELD
REORDER: KEVIN BILLINGS
POST-FIX WEIGHER: NOT AVAILABLE
PATHOLOGIST: JOHN EIGHMY
WEIGHER: MICHELE ZIMA

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)				NOT TAKEN
BRAIN (BR)				NOT TAKEN
KIDNEY (KD)				NOT TAKEN
LIVER (LI)	6.8351	3.3326 %		WEIGHT TAKEN
OVARY (OV)				NOT TAKEN
SPLEEN (SP)				NOT TAKEN
THYMUS (TH)				NOT TAKEN
THYROID/PARATHYR (TT)				NOT TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI), GENERAL COMMENT (GC)

001231

APPENDIX 6

Individual Anatomical Pathology Data

PAGE: 149

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95992
DATE OF DEATH: 10/01/98
DATE AND TIME OF NECROPSY: 10/01/98 11:51
POST-FIX WEIGHTER: NOT AVAILABLE

SEX: FEMALE
STUDY DAY OF DEATH: 31
STUDY WEEK OF DEATH: 5
PROSECTOR: WENDY CHERRY
PATHOLOGIST: JOHN EIGHMY

DOSE GROUP: 1
SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
TERMINAL BODY WEIGHT: 180.6 GRAMS
RECORDED: KEVIN BILLINGS
WEIGHER: MICHELE ZIMA

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	---	---	---	NOT TAKEN
BRAIN (BR)	---	---	---	NOT TAKEN
KIDNEY (KD)	---	---	---	NOT TAKEN
LIVER (LI)	---	---	---	WEIGHT TAKEN
OVARY (OV)	5.8602	3.2449 %	---	NOT TAKEN
SPLEEN (SP)	---	---	---	NOT TAKEN
THYMUS (TH)	---	---	---	NOT TAKEN
THYROID/PARATHYR (TT)	---	---	---	NOT TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI), GENERAL COMMENT (GC)

001232

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL-
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 150

ANIMAL NUMBER: C95993	SEX: FEMALE	DOSE GROUP: 1	SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE	
DATE OF DEATH: 10/01/98	STUDY DAY OF DEATH: 31	STUDY WEEK OF DEATH: 5	TERMINAL BODY WEIGHT: 201.3 GRAMS	
DATE AND TIME OF NECROPSY: 10/01/98 11:38	PROSECTOR: WENDY CHERRY		REORDER: KEVIN BILLINGS	
POST-FIX WEIGHT: NOT AVAILABLE	PATHOLOGIST: JOHN EIGHMY		WEIGHER: MICHELE ZIMA	
ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	---	---	---	NOT TAKEN
BRAIN (BR)	---	---	---	NOT TAKEN
KIDNEY (KD)	---	---	---	NOT TAKEN
LIVER (LI)	---	---	---	WEIGHT TAKEN
OVARY (OV)	6.0180	2.9896 %	---	NOT TAKEN
SPLEEN (SP)	---	---	---	NOT TAKEN
THYMUS (TH)	---	---	---	NOT TAKEN
THYROID/PARATHYR (TT)	---	---	---	NOT TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR), SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF), URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI), GENERAL COMMENT (GC)

001233

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 151

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95994		SEX: FEMALE		DOSE GROUP: 1		SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE	
DATE OF DEATH: 10/01/98		STUDY DAY OF DEATH: 31		STUDY WEEK OF DEATH: 5		TERMINAL BODY WEIGHT: 215.5 GRAMS	
DATE AND TIME OF NECROPSY: 10/01/98 11:25		PROSECTOR: MATTHEW AXTMAN		PATHOLOGIST: JOHN EIGHMY		REORDER: KEVIN BILLINGS	
POST-FIX WEIGHTER: NOT AVAILABLE						WEIGHER: MICHELE ZIMA	
		ABSOLUTE ORGAN WEIGHT (GRAMS)		ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)		ORGAN TO BRAIN WEIGHT RATIO	
ORGAN NAME						S T A T U S	
ADRENAL (AD)		---		---		NOT TAKEN	
BRAIN (BR)		---		---		NOT TAKEN	
KIDNEY (KD)		---		---		NOT TAKEN	
LIVER (LI)		7.0305		3.2624 %		WEIGHT TAKEN	
OVARY (OV)		---		---		NOT TAKEN	
SPLEEN (SP)		---		---		NOT TAKEN	
THYMUS (TH)		---		---		NOT TAKEN	
THYROID/PARATHYR (TT)		---		---		NOT TAKEN	

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI), GENERAL COMMENT (GC)

001234

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.113-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 152ANIMAL NUMBER: C95995
SEX: FEMALE
DOSE GROUP: 1
SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
DATE OF DEATH: 10/01/98
STUDY DAY OF DEATH: 31
STUDY WEEK OF DEATH: 5
TERMINAL BODY WEIGHT: 188.5 GRAMS
DATE AND TIME OF NECROPSY: 10/01/98 12:01
PROSECTOR: WENDY CHERRY
REORDER: KEVIN BILLINGS
POST-FIX WEIGHTER: NOT AVAILABLE
PATHOLOGIST: JOHN EIGHMY
WEIGHER: MICHELE ZIMA

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	---	---	---	NOT TAKEN
BRAIN (BR)	---	---	---	NOT TAKEN
KIDNEY (KD)	---	---	---	NOT TAKEN
LIVER (LI)	---	---	---	WEIGHT TAKEN
OVARY (OV)	5.7678	3.0598 %	---	NOT TAKEN
SPLEEN (SP)	---	---	---	NOT TAKEN
THYMUS (TH)	---	---	---	NOT TAKEN
THYROID/PARATHYR (TT)	---	---	---	NOT TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR), SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF), URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI), GENERAL COMMENT (GC)

001235

APPENDIX 6
Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 153

ANIMAL NUMBER: C95996
SEX: FEMALE
DATE OF DEATH: 12/02/98
STUDY DAY OF DEATH: 93
SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
TERMINAL BODY WEIGHT: 268.9 GRAMS
REORDER: WENDY CHERRY
DATE AND TIME OF NECROPSY: 12/02/98 12:35
PROSECTOR: JILL PAUS
WEIGHER: LYNN HORNER
POST-FIX WEIGHER: NOT AVAILABLE
PATHOLOGIST: JOHN EIGHMY

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0538	.0200 %	.0269	WEIGHT TAKEN
BRAIN (BR)	2.0029	.7448 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.2491	.8364 %	1.1229	WEIGHT TAKEN
LIVER (LI)	7.1747	2.6682 %	3.5822	WEIGHT TAKEN
OVARY (OV)	.1609	.0598 %	.0803	WEIGHT TAKEN
SPLEEN (SP)	.4536	.1687 %	.2265	WEIGHT TAKEN
THYMUS (TH)	.2401	.0893 %	.1199	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0251	.0093 %	.0125	WEIGHT TAKEN

NECROPSY
OBSERVATIONS
HISTOPATHOLOGY
LIVER (LI):
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL
-VACUOLATION, HEPATOCELLULAR, PERIportal, -MINIMAL
THYMUS (TH):
-HEMORRHAGE, -MINIMAL

GENERAL COMMENT (GC):
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

001236

APPENDIX 6

Covance 6329-225
3M T-6314.1

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL-
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 154

ANIMAL NUMBER: C95996 SEX: FEMALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 PROSECTOR: JILL PAUS TERMINAL BODY WEIGHT: 268.9 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 12:35 PATHOLOGIST: JOHN EIGHMY RECORDER: WENDY CHERRY
POST-FIX WEIGHT: NOT AVAILABLE

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC),
ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA)

001237

APPENDIX 6
Individual Anatomic Pathology Data

PAGE: 155

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95997 SEX: FEMALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 247.6 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:02 PROSECTOR: KEVIN BILLINGS RECORDED: WENDY CHERRY
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATISTICS
ADRENAL (AD)	.0693	.0280 %	.0377	WEIGHT TAKEN
BRAIN (BR)	1.8393	.7429 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.0840	.8417 %	1.1330	WEIGHT TAKEN
LIVER (LI)	6.8198	2.7544 %	3.7078	WEIGHT TAKEN
OVARY (OV)	.1272	.0514 %	.0692	WEIGHT TAKEN
SPLEEN (SP)	.4192	.1693 %	.2279	WEIGHT TAKEN
THYMUS (TH)	.2108	.0851 %	.1146	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0177	.0071 %	.0096	WEIGHT TAKEN

NECROPSY OBSERVATIONS HISTOPATHOLOGY

THYMUS (TH) :
-HEMORRHAGE, -MINIMAL
THYROID (TY) :
-CYST, ULTIMOBRANCHIAL, -PRESENT

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

001238

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 156

ANIMAL NUMBER: C95997 SEX: FEMALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 247.6 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:02 PROSECTOR: KEVIN BILLINGS REORDER: WENDY CHERRY
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), LN, MESENTERIC (MS), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA),
AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE),
ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG),
AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF), URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV),
VAGINA (VA)

001239

APPENDIX 6
Individual Anatomic Pathology Data

PAGE: 157

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95998 SEX: FEMALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 240.2 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:40 PROSECTOR: KIMBERLY DURLAND REORDER: STEVE VAN ADESTINE
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0641	.0267 %	.0316	WEIGHT TAKEN
BRAIN (BR)	2.0289	.8447 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	1.8478	.7693 %	.9107	WEIGHT TAKEN
LIVER (LI)	6.0936	2.5369 %	3.0034	WEIGHT TAKEN
OVARY (OV)	.1647	.0686 %	.0812	WEIGHT TAKEN
SPLEEN (SP)	.4325	.1801 %	.2132	WEIGHT TAKEN
THYMUS (TH)	.3476	.1447 %	.1713	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0341	.0142 %	.0168	WEIGHT TAKEN

HISTOPATHOLOGY

OBSERVATIONS

NECROPSY

THYROID (TY) :
-CYST, ULTIMOBRANCHIAL, -PRESENT
AUDITORY SEB GL (AS) :
-ONE EXAMINED, -PRESENT

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

001240

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 158

ANIMAL NUMBER: C95998 SEX: FEMALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 240.2 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:40 PROSECTOR: KIMBERLY DURLAND REORDER: STEVE VAN ADESTINE
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), PARATHYROID (PT), ADRENAL, CORTEX (AC),
ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), SKIN (SK), MAMMARY, FEMALE (MF), URINARY BLADDER (UB), UTERUS (UT),
CERVIX (CV), OVARY (OV), VAGINA (VA)

001241

APPENDIX 6
Individual Anatomic Pathology Data

PAGE: 159

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C95999 SEX: FEMALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 358.4 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:25 PROSECTOR: JERALD W. JOHNSON REORDER: WENDY CHERRY
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	O R G A N	
				S T A T U S	WEIGHT TAKEN
ADRENAL (AD)	.0752	.0210 %	.0359		WEIGHT TAKEN
BRAIN (BR)	2.0931	.5840 %	1.0000		WEIGHT TAKEN
KIDNEY (KD)	2.4475	.6829 %	1.1693		WEIGHT TAKEN
LIVER (LI)	9.9107	2.7653 %	4.7349		WEIGHT TAKEN
OVARY (OV)	.1232	.0344 %	.0589		WEIGHT TAKEN
SPLEEN (SP)	.5403	.1508 %	.2581		WEIGHT TAKEN
THYMUS (TH)	.3750	.1046 %	.1792		WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0228	.0064 %	.0109		WEIGHT TAKEN

NECROPSY OBSERVATIONS HISTOPATHOLOGY

LIVER (LI) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
THYROID (TY) :
-CYST, ULTIMOBRANCHIAL, -PRESENT

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

001242

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 160

ANIMAL NUMBER: C95999 SEX: FEMALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 358.4 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:25 PROSECTOR: JERALD W. JOHNSON REORDER: WENDY CHERRY
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), PARATHYROID (PT), ADRENAL, CORTEX (AC),
ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA)

001243

APPENDIX 6

Individual Anatomical Pathology Data

PAGE: 161

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96000 SEX: FEMALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 269.7 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:11 PROSECTOR: TODD ACKER RECORDED: WENDY CHERRY
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0647	.0240 %	.0330	WEIGHT TAKEN
BRAIN (BR)	1.9596	.7266 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	1.9224	.7128 %	.9810	WEIGHT TAKEN
LIVER (LI)	8.1750	3.0311 %	4.1718	WEIGHT TAKEN
OVARY (OV)	.1191	.0442 %	.0608	WEIGHT TAKEN
SPLEEN (SP)	.4243	.1573 %	.2165	WEIGHT TAKEN
THYMUS (TH)	.1879	.0697 %	.0959	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0169	.0063 %	.0086	WEIGHT TAKEN

NECROPSY OBSERVATIONS HISTOPATHOLOGY

LIVER (LI) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
AUDITORY SEE GL (AS) :
-ONE EXAMINED, -PRESENT

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

001244

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 162

ANIMAL NUMBER: C96000 SEX: FEMALE DOSE GROUP: 1 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 269.7 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:11 PROSECTOR: TODD ACKER RECORDED: WENDY CHERRY
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), SKIN (SK), MAMMARY, FEMALE (MF), URINARY BLADDER (UB), UTERUS (UT),
CERVIX (CV), OVARY (OV), VAGINA (VA)

0012A5

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 163

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96001 SEX: FEMALE DOSE GROUP: 2 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 PROSECTOR: JILL PAUS TERMINAL BODY WEIGHT: 233.4 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 12:11 PATHOLOGIST: JOHN EIGHMY REORDER: WENDY CHERRY
POST-FIX WEIGHER: NOT AVAILABLE

WEIGHER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0574	.0246 %	.0314	WEIGHT TAKEN
BRAIN (BR)	1.8284	.7834 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.1472	.9200 %	1.1744	WEIGHT TAKEN
LIVER (LI)	7.2356	3.1001 %	3.9573	WEIGHT TAKEN
OVARY (OV)	.1433	.0614 %	.0784	WEIGHT TAKEN
SPLEEN (SP)	.5303	.2272 %	.2900	WEIGHT TAKEN
THYMUS (TH)	.3077	.1318 %	.1683	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0305	.0131 %	.0167	WEIGHT TAKEN

O B S E R V A T I O N S

NECROPSY

HISTOPATHOLOGY

LIVER (LI) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-VACUOLATION, HEPATOCELLULAR, PERIportal, -MINIMAL

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:
BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

001246

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.113-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 164

ANIMAL NUMBER: C96002
 DATE OF DEATH: 12/02/98
 DATE AND TIME OF NECROPSY: 12/02/98 10:25
 POST-FIX WEIGHTER: NOT AVAILABLE

SEX: FEMALE
 STUDY DAY OF DEATH: 93
 PROSECTOR: NANCY DIEDRICH
 PATHOLOGIST: JOHN EIGHMY

DOSE GROUP: 2
 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
 STUDY WEEK OF DEATH: 14
 TERMINAL BODY WEIGHT: 228.4 GRAMS
 RECORDER: TERRI L. WILLIAMS
 WEIGHTER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATISTICS
ADRENAL (AD)	.0630	.0276 %	.0311	WEIGHT TAKEN
BRAIN (BR)	2.0244	.8863 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	1.9083	.8355 %	.9426	WEIGHT TAKEN
LIVER (LI)	7.1598	3.1348 %	3.5368	WEIGHT TAKEN
OVARY (OV)	.1099	.0481 %	.0543	WEIGHT TAKEN
SPLEEN (SP)	.4890	.2141 %	.2416	WEIGHT TAKEN
THYMUS (TH)	.2793	.1223 %	.1380	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0179	.0078 %	.0088	WEIGHT TAKEN

O B S E R V A T I O N S

NECROPSY

HISTOPATHOLOGY

UTERUS (UT) :
 -LUMEN FILLED WITH FLUID; BILATERAL; CLEAR FLUID
 GENERAL COMMENT (GC) :
 -BONE MARROW SMEAR TAKEN

LIVER (LI) :
 -INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
 -NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL
 UTERUS (UT) :
 -DILATATION, -SLIGHT

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
 SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
 NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
 ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
 PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
 SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
 URINARY BLADDER (UB), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

APPENDIX 6
Individual Anatomic Pathology Data

PAGE: 165

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96003 SEX: FEMALE DOSE GROUP: 2 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 217.5 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:50 PROSECTOR: TODD ACKER REORDER: WENDY CHERRY
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0598	.0275 %	.0301	WEIGHT TAKEN
BRAIN (BR)	1.9851	.9127 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	1.7313	.7960 %	.8721	WEIGHT TAKEN
LIVER (LI)	6.5350	3.0046 %	3.2920	WEIGHT TAKEN
OVARY (OV)	.1142	.0525 %	.0575	WEIGHT TAKEN
SPLEEN (SP)	.5506	.2531 %	.2774	WEIGHT TAKEN
THYMUS (TH)	.2760	.1269 %	.1390	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0194	.0089 %	.0098	WEIGHT TAKEN

HISTOPATHOLOGY

NECROPSY

UTERUS (UT) :
-LUMEN FILLED WITH FLUID; BILATERAL: CLEAR FLUID
GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN

001248

APPENDIX 6
Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 166

ANIMAL NUMBER: C96003
SEX: FEMALE
DATE OF DEATH: 12/02/98
STUDY DAY OF DEATH: 93
STUDY WEEK OF DEATH: 14
SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE AND TIME OF NECROPSY: 12/02/98 10:50
PROSECTOR: TODD ACKER
TERMINAL BODY WEIGHT: 217.5 GRAMS
POST-FIX WEIGHT: NOT AVAILABLE
PATHOLOGIST: JOHN EIGHMY
REORDER: WENDY CHERRY
WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:
LIVER (LI)

001249

APPENDIX 6
Individual Anatomic Pathology Data

PAGE: 167

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
TERMINAL BODY WEIGHT: 263.6 GRAMS
REORDER: WENDY CHERRY
WEIGHER: LYNN HORNER

DOSE GROUP: 2
STUDY WEEK OF DEATH: 14
PROSECTOR: TODD ACKER
PATHOLOGIST: JOHN EIGHMY

ANIMAL NUMBER: C96004
SEX: FEMALE
STUDY DAY OF DEATH: 93
DATE OF DEATH: 12/02/98
DATE AND TIME OF NECROPSY: 12/02/98 7:34
POST-FIX WEIGHER: NOT AVAILABLE

ORGAN
STATUSES

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	WEIGHT TAKEN
ADRENAL (AD)	.0516	.0196 %	.0265	WEIGHT TAKEN
BRAIN (BR)	1.9505	.7399 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.2579	.8566 %	1.1576	WEIGHT TAKEN
LIVER (LI)	8.3001	3.1487 %	4.2554	WEIGHT TAKEN
OVARY (OV)	.1230	.0467 %	.0631	WEIGHT TAKEN
SPLEEN (SP)	.5005	.1899 %	.2566	WEIGHT TAKEN
THYMUS (TH)	.2900	.1100 %	.1487	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0143	.0054 %	.0073	WEIGHT TAKEN

HISTOPATHOLOGY

NECROPSY

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

001250

455

APPENDIX 6
Individual Anatomic Pathology Data

PAGE: 168

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96004 SEX: FEMALE DOSE GROUP: 2 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 TERMINAL BODY WEIGHT: 263.6 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 7:34 PROSECTOR: TODD ACKER REORDER: WENDY CHERRY
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:
BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:
LIVER (LI)

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 169

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96005		SEX: FEMALE		DOSE GROUP: 2		SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE	
DATE OF DEATH: 12/02/98		STUDY DAY OF DEATH: 93		STUDY WEEK OF DEATH: 14		TERMINAL BODY WEIGHT: 243.7 GRAMS	
DATE AND TIME OF NECROPSY: 12/02/98 8:05		PROSECTOR: JERALD W. JOHNSON		RECORDED: WENDY CHERRY		WEIGHED: LYNN HORNER	
POST-FIX WEIGHTER: NOT AVAILABLE		PATHOLOGIST: JOHN EIGHMY		ORGAN TO BRAIN		O R G A N	
		ABSOLUTE ORGAN WEIGHT (GRAMS)		ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)		S T A T U S	
ORGAN NAME						WEIGHT TAKEN	
ADRENAL (AD)		.0702		.0288 %		.0370	
BRAIN (BR)		1.8993		.7794 %		1.0000	
KIDNEY (KD)		2.2248		.9129 %		1.1714	
LIVER (LI)		8.2448		3.3832 %		4.3410	
OVARY (OV)		.1103		.0453 %		.0581	
SPLEEN (SP)		.4497		.1845 %		.2368	
THYMUS (TH)		.3365		.1381 %		.1772	
THYROID/PARATHYR (TT)		.0162		.0066 %		.0085	

HISTOPATHOLOGY

NECROPSY

LIVER (LI) :
-INFILTRATE, LYMPHOHISTIOCYTIC, MINIMAL
-NECROSIS, INDIVIDUAL HEPATOCYTE, MINIMAL

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), VAGINA (VA), TAIL (TI)

001252

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 170

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96006 SEX: FEMALE DOSE GROUP: 2 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 249.2 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:36 PROSECTOR: MICHELE ZIMA REORDER: WENDY CHERRY
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS	WEIGHT TAKEN
ADRENAL (AD)	.0512	.0205 %	.0259		WEIGHT TAKEN
BRAIN (BR)	1.9754	.7927 %	1.0000		WEIGHT TAKEN
KIDNEY (KD)	1.8567	.7451 %	.9399		WEIGHT TAKEN
LIVER (LI)	7.3435	2.9468 %	3.7175		WEIGHT TAKEN
OVARY (OV)	.1079	.0433 %	.0546		WEIGHT TAKEN
SPLEEN (SP)	.4943	.1984 %	.2502		WEIGHT TAKEN
THYMUS (TH)	.2233	.0896 %	.1130		WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0032	.0013 %	.0016		WEIGHT TAKEN

HISTOPATHOLOGY

NECROPSY

OBSERVATIONS

LIVER (LI): Lymphohistiocytic, minimal
-INFILTRATE,GENERAL COMMENT (GC):
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:
BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 171

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96007 SEX: FEMALE DOSE GROUP: 2 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 TERMINAL BODY WEIGHT: 252.5 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 12:06 PROSECTOR: MICHELE ZIMA RECORDED: WENDY CHERRY
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	O R G A N	
				S T A T U S	WEIGHT TAKEN
ADRENAL (AD)	.0765	.0303 %	.0379		WEIGHT TAKEN
BRAIN (BR)	2.0179	.7992 %	1.0000		WEIGHT TAKEN
KIDNEY (KD)	2.2417	.8878 %	1.1109		WEIGHT TAKEN
LIVER (LI)	8.5804	3.3982 %	4.2521		WEIGHT TAKEN
OVARY (OV)	.1210	.0479 %	.0600		WEIGHT TAKEN
SPLEEN (SP)	.6280	.2487 %	.3112		WEIGHT TAKEN
THYMUS (TH)	.3639	.1441 %	.1803		WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0251	.0099 %	.0124		WEIGHT TAKEN

O B S E R V A T I O N S

NECROPSY

HISTOPATHOLOGY

LIVER (LI) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-VACUOLATION, HEPATOCELLULAR, PERIportal, -MINIMAL

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:
BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TV), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

001254

APPENDIX 6
Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 172

ANIMAL NUMBER: C96008 SEX: FEMALE DOSE GROUP: 2 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 TERMINAL BODY WEIGHT: 277.6 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:19 PROSECTOR: MICHELE ZIMA RECORDED: WENDY CHERRY
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0821	.0296 %	.0412	WEIGHT TAKEN
BRAIN (BR)	1.9908	.7171 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.4080	.8674 %	1.2096	WEIGHT TAKEN
LIVER (LI)	9.2167	3.3201 %	4.6296	WEIGHT TAKEN
OVARY (OV)	.1217	.0438 %	.0611	WEIGHT TAKEN
SPLEEN (SP)	.6382	.2299 %	.3206	WEIGHT TAKEN
THYMUS (TH)	.3024	.1089 %	.1519	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0224	.0081 %	.0113	WEIGHT TAKEN

NECROPSY
OBSERVATIONS
HISTOPATHOLOGY
LIVER (LI) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS
GENERAL INFORMATION (XX) :
>NOTE:>ANIMAL IDENTIFICATION TRANSPONDER DAMAGED AT NECROPSY

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:
BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 173

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96009 SEX: FEMALE DOSE GROUP: 2 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 208.7 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:20 PROSECTOR: TODD ACKER RECORDER: WENDY CHERRY
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0564	.0270 %	.0293	WEIGHT TAKEN
BRAIN (BR)	1.9241	.9219 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	1.8706	.8963 %	.9722	WEIGHT TAKEN
LIVER (LI)	5.9471	2.8496 %	3.0908	WEIGHT TAKEN
OVARY (OV)	.1006	.0482 %	.0523	WEIGHT TAKEN
SPLEEN (SP)	.4206	.2015 %	.2186	WEIGHT TAKEN
THYMUS (TH)	.1966	.0942 %	.1022	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0090	.0043 %	.0047	WEIGHT TAKEN

NECROPSY OBSERVATIONS HISTOPATHOLOGY

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

001256

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 174

ANIMAL NUMBER: C96009 SEX: FEMALE DOSE GROUP: 2 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 208.7 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:20 PROSECTOR: TODD ACKER RECORDED: WENDY CHERRY
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EICHMY WEIGHT: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:
LIVER (LI)

001257

APPENDIX 6

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 175ANIMAL NUMBER: C96010 SEX: FEMALE DOSE GROUP: 2 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 263.0 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:35 PROSECTOR: LAURIE J. SCHULLER RECORDED: TERRI L. WILLIAMS
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0658	.0250 %	.0317	WEIGHT TAKEN
BRAIN (BR)	2.0743	.7887 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.0370	.7745 %	.9820	WEIGHT TAKEN
LIVER (LI)	8.2431	3.1343 %	3.9739	WEIGHT TAKEN
OVARY (OV)	.0965	.0367 %	.0465	WEIGHT TAKEN
SPLEEN (SP)	.5876	.2234 %	.2833	WEIGHT TAKEN
THYMUS (TH)	.3850	.1464 %	.1856	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0181	.0069 %	.0087	WEIGHT TAKEN

NECROPSY	OBSERVATIONS	HISTOPATHOLOGY
	LIVER (LI) :	
	-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL	
	-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL	
	-VACUOLATION, HEPATOCELLULAR, PERIPORTAL, -MINIMAL	

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TV), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

APPENDIX 6

Covance 6329-225
3M T-6314.1

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 176

ANIMAL NUMBER: C96011 SEX: FEMALE DOSE GROUP: 2 SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
DATE OF DEATH: 10/01/98 STUDY DAY OF DEATH: 31 STUDY WEEK OF DEATH: 5 TERMINAL BODY WEIGHT: 226.8 GRAMS
DATE AND TIME OF NECROPSY: 10/01/98 12:09 PROSECTOR: WENDY CHERRY RECORDED: KEVIN BILLINGS
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: MICHELE ZIMA

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	---	---	---	NOT TAKEN
BRAIN (BR)	---	---	---	NOT TAKEN
KIDNEY (KD)	---	---	---	NOT TAKEN
LIVER (LI)	---	---	---	WEIGHT TAKEN
OVARY (OV)	7.5294	3.3198 %	---	NOT TAKEN
SPLEEN (SP)	---	---	---	NOT TAKEN
THYMUS (TH)	---	---	---	NOT TAKEN
THYROID/PARATHYR (TT)	---	---	---	NOT TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TV), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI), GENERAL COMMENT (GC)

C01259

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 177

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96012 SEX: FEMALE DOSE GROUP: 2 SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
DATE OF DEATH: 10/01/98 STUDY DAY OF DEATH: 31 STUDY WEEK OF DEATH: 5 TERMINAL BODY WEIGHT: 191.9 GRAMS
DATE AND TIME OF NECROPSY: 10/01/98 11:05 PROSECTOR: TONY HANFELD RECORDER: KEVIN BILLINGS
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: MICHELE ZIMA

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	---	---	---	NOT TAKEN
BRAIN (BR)	---	---	---	NOT TAKEN
KIDNEY (KD)	---	---	---	NOT TAKEN
LIVER (LI)	6.5011	3.3878 %	---	WEIGHT TAKEN
OVARY (OV)	---	---	---	NOT TAKEN
SPLEEN (SP)	---	---	---	NOT TAKEN
THYMUS (TH)	---	---	---	NOT TAKEN
THYROID/PARATHYR (TT)	---	---	---	NOT TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI), GENERAL COMMENT (GC)

001260

APPENDIX 6

Covance 6329-225
3M T-6314.1

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 178ANIMAL NUMBER: C96013
SEX: FEMALE
DATE OF DEATH: 10/01/98
STUDY DAY OF DEATH: 31
SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
DATE AND TIME OF NECROPSY: 10/01/98 11:31
PROSECTOR: WENDY CHERRY
TERMINAL BODY WEIGHT: 223.1 GRAMS
POST-FIX WEIGHT: NOT AVAILABLE
PATHOLOGIST: JOHN EIGHMY
RECORDED: KEVIN BILLINGS
WEIGHER: MICHELE ZIMA

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	---	---	---	NOT TAKEN
BRAIN (BR)	---	---	---	NOT TAKEN
KIDNEY (KD)	---	---	---	NOT TAKEN
LIVER (LI)	---	---	---	WEIGHT TAKEN
OVARY (OV)	7.7373	3.4681 %	---	NOT TAKEN
SPLEEN (SP)	---	---	---	NOT TAKEN
THYMUS (TH)	---	---	---	NOT TAKEN
THYROID/PARATHYR (TT)	---	---	---	NOT TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TV), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI), GENERAL COMMENT (GC)

APPENDIX 6

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 179ANIMAL NUMBER: C96014
SEX: FEMALE
DOSE GROUP: 2
SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
DATE OF DEATH: 10/01/98
STUDY DAY OF DEATH: 31
STUDY WEEK OF DEATH: 5
TERMINAL BODY WEIGHT: 227.7 GRAMS
DATE AND TIME OF NECROPSY: 10/01/98 11:07
PROSECTOR: MATTHEW AXTMAN
RECORDED: KEVIN BILLINGS
POST-FIX WEIGHTER: NOT AVAILABLE
PATHOLOGIST: JOHN EIGHMY
WEIGHER: MICHELE ZIMA

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	---	---	---	NOT TAKEN
BRAIN (BR)	---	---	---	NOT TAKEN
KIDNEY (KD)	---	---	---	NOT TAKEN
LIVER (LI)	7.4920	3.2903 %	---	WEIGHT TAKEN
OVARY (OV)	---	---	---	NOT TAKEN
SPLEEN (SP)	---	---	---	NOT TAKEN
THYMUS (TH)	---	---	---	NOT TAKEN
THYROID/PARATHYR (TT)	---	---	---	NOT TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), STOMACH, NONGL (SU),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), RECTUM (RE), LN, MANDIBULAR (MN),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI), GENERAL COMMENT (GC)

001262

APPENDIX 6

Covance 6329-225
3M T-6314.1

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 180

ANIMAL NUMBER: C96015 SEX: FEMALE DOSE GROUP: 2 SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
DATE OF DEATH: 10/01/98 STUDY DAY OF DEATH: 31 STUDY WEEK OF DEATH: 5 TERMINAL BODY WEIGHT: 210.4 GRAMS
DATE AND TIME OF NECROPSY: 10/01/98 11:14 PROSECTOR: MATTHEW AXTMAN REORDER: KEVIN BILLINGS
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: MICHELE ZIMA

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	---	---	---	NOT TAKEN
BRAIN (BR)	---	---	---	NOT TAKEN
KIDNEY (KD)	---	---	---	NOT TAKEN
LIVER (LI)	7.0360	3.3441 %	---	WEIGHT TAKEN
OVARY (OV)	---	---	---	NOT TAKEN
SPLEEN (SP)	---	---	---	NOT TAKEN
THYMUS (TH)	---	---	---	NOT TAKEN
THYROID/PARATHYR (TT)	---	---	---	NOT TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI), GENERAL COMMENT (GC)

001263

APPENDIX 6
Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 181

ANIMAL NUMBER: C96016 SEX: FEMALE DOSE GROUP: 2 SACRIFICE STATUS: UNSCHEDULED (M)
DATE OF DEATH: 10/29/98 STUDY DAY OF DEATH: 59 STUDY WEEK OF DEATH: 9 TERMINAL BODY WEIGHT: 272.9 GRAMS
DATE AND TIME OF NECROPSY: 10/29/98 15:09 PROSECTOR: JAMES T. CALLEN RECORDER: JAMES T. CALLEN
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: JAMES T. CALLEN

*** ORGAN WEIGHTS WERE NOT RECORDED ***

O B S E R V A T I O N S		HISTOPATHOLOGY
NECROPSY		
LIVER (LI) :		
-INFILTRATE, LYMPHOHISTIOCYTIC,-MINIMAL		
-NECROSIS, INDIVIDUAL HEPATOCYTE,-MINIMAL		
TAIL (TI) :		
-HEMORRHAGE,-SLIGHT		
-ULCERATION,-MINIMAL		
TAIL (TI) :		
-LACERATION(S); 20 MM FROM BASE OF TAIL: SINGLE LACERATION,		
20 X 10 MM; SURROUNDING TISSUE APPEARS SWOLLEN AND EXUDES A		
DARK RED SEMIFLUID; COLLECTED		

001264

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 182

ANIMAL NUMBER: C96016 SEX: FEMALE DOSE GROUP: 2 SACRIFICE STATUS: UNSCHEDULED (M)
DATE OF DEATH: 10/29/98 STUDY DAY OF DEATH: 59 STUDY WEEK OF DEATH: 9 TERMINAL BODY WEIGHT: 272.9 GRAMS
DATE AND TIME OF NECROPSY: 10/29/98 15:09 PROSECTOR: JAMES T. CALLEN REORDER: JAMES T. CALLEN
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHNY WEIGHER: JAMES T. CALLEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), GENERAL COMMENT (GC)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA)

001265

APPENDIX 6

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 183ANIMAL NUMBER: C96017 SEX: FEMALE DOSE GROUP: 2 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 277.4 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 7:41 PROSECTOR: JODI THIEREN REORDER: STEVE VAN ADESTINE
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0693	.0250 %	.0342	WEIGHT TAKEN
BRAIN (BR)	2.0282	.7311 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.2400	.8075 %	1.1044	WEIGHT TAKEN
LIVER (LI)	8.0588	2.9051 %	3.9734	WEIGHT TAKEN
OVARY (OV)	.1886	.0680 %	.0930	WEIGHT TAKEN
SPLEEN (SP)	.5476	.1974 %	.2700	WEIGHT TAKEN
THYMUS (TH)	.3084	.1112 %	.1521	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0200	.0072 %	.0099	WEIGHT TAKEN

O B S E R V A T I O N S

NECROPSY

HISTOPATHOLOGY

LIVER (LI) :

-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-MALOCCLUSION; BILATERAL UPPER INCISORS

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

001266

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.113-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 184ANIMAL NUMBER: C96018
SEX: FEMALE
DOSE GROUP: 2
SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98
STUDY DAY OF DEATH: 93
STUDY WEEK OF DEATH: 14
TERMINAL BODY WEIGHT: 279.7 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:50
PROSECTOR: MICHELE ZIMA
REORDER: WENDY CHERRY
POST-FIX WEIGHTER: NOT AVAILABLE
PATHOLOGIST: JOHN EIGHMY
WEIGHER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0805	.0288 %	.0423	WEIGHT TAKEN
BRAIN (BR)	1.9028	.6803 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.0616	.7371 %	1.0835	WEIGHT TAKEN
LIVER (LI)	8.0095	2.8636 %	4.2093	WEIGHT TAKEN
OVARY (OV)	.1174	.0420 %	.0617	WEIGHT TAKEN
SPLEEN (SP)	.5094	.1821 %	.2677	WEIGHT TAKEN
THYMUS (TH)	.2793	.0999 %	.1468	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0175	.0063 %	.0092	WEIGHT TAKEN

NECROPSY

OBSERVATIONS

HISTOPATHOLOGY

THYMUS (TH) :
-DIFFUSELY RED; DARK RED
GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN

LIVER (LI) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL
THYMUS (TH) :
-HEMORRHAGE, -SLIGHT
-INVOLUTION, -MINIMAL

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR), SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN), SPLEEN (SP), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF), URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

001267

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 185

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96019		SEX: FEMALE	DOSE GROUP: 2	SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98		STUDY DAY OF DEATH: 93	STUDY WEEK OF DEATH: 14	TERMINAL BODY WEIGHT: 283.2 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 12:12		PROSECTOR: NANCY DIEDRICH	REORDER: TERRI L. WILLIAMS	
POST-FIX WEIGHTER: NOT AVAILABLE		PATHOLOGIST: JOHN EIGHMY	WEIGHER: TONY HANFELD	
		ABSOLUTE ORGAN WEIGHT (GRAMS)		ORGAN TO BRAIN WEIGHT RATIO
ORGAN NAME				S T A T U S
ADRENAL (AD)	.0860	.0304 %	.0420	WEIGHT TAKEN
BRAIN (BR)	2.0462	.7225 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.6887	.9494 %	1.3140	WEIGHT TAKEN
LIVER (LI)	8.9464	3.1590 %	4.3722	WEIGHT TAKEN
OVARY (OV)	.1290	.0456 %	.0630	WEIGHT TAKEN
SPLEEN (SP)	.5329	.1882 %	.2604	WEIGHT TAKEN
THYMUS (TH)	.3098	.1094 %	.1514	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0209	.0074 %	.0102	WEIGHT TAKEN

HISTOPATHOLOGY

O B S E R V A T I O N S

NECROPSY

UTERUS (UT) :
-DILATATION, -SLIGHT

UTERUS (UT) :
-LUMEN FILLED WITH FLUID; BILATERAL: CLEAR FLUID

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN

001268

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 186

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96019 SEX: FEMALE DOSE GROUP: 2 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 283.2 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 12:12 PROSECTOR: NANCY DIEDRICH REORDER: FERRI L. WILLIAMS
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

LIVER (LI)

001269

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 187

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96020		SEX: FEMALE	DOSE GROUP: 2	SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE	
DATE OF DEATH: 12/02/98		STUDY DAY OF DEATH: 93	STUDY WEEK OF DEATH: 14	TERMINAL BODY WEIGHT: 239.9 GRAMS	
DATE AND TIME OF NECROPSY: 12/02/98 12:06		PROSECTOR: JOHN S. HALFORD		RECORDER: WENDY CHERRY	
POST-FIX WEIGHTER: NOT AVAILABLE		PATHOLOGIST: JOHN EIGHMY		WEIGHER: LYNN HORNER	
ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)		ORGAN TO BRAIN WEIGHT RATIO	O R G A N S T A T U S

ADRENAL (AD)	.0746	.0311 %		.0392	WEIGHT TAKEN
BRAIN (BR)	1.9018	.7927 %		1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.0893	.8709 %		1.0986	WEIGHT TAKEN
LIVER (LI)	8.3210	3.4685 %		4.3753	WEIGHT TAKEN
OVARY (OV)	.1320	.0550 %		.0694	WEIGHT TAKEN
SPLEEN (SP)	.4508	.1879 %		.2370	WEIGHT TAKEN
THYMUS (TH)	.2735	.1140 %		.1438	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0172	.0072 %		.0090	WEIGHT TAKEN
O B S E R V A T I O N S					

O B S E R V A T I O N S

NECROPSY

LIVER (LI) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LONG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

001270

APPENDIX 6
Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL-
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEPOSE, T-6314) IN RATS

PAGE: 188

ANIMAL NUMBER: C96021
DATE OF DEATH: 12/02/98
SEX: FEMALE
STUDY DAY OF DEATH: 93
DOSE GROUP: 3
SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE AND TIME OF NECROPSY: 12/02/98 10:36
PROSECTOR: JERALD W. JOHNSON
TERMINAL BODY WEIGHT: 239.7 GRAMS
POST-FIX WEIGHTER: NOT AVAILABLE
PATHOLOGIST: JOHN EIGHMY
REORDER: WENDY CHERRY
WEIGHER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUSES
ADRENAL (AD)	.0599	.0250 %	.0302	WEIGHT TAKEN
BRAIN (BR)	1.9833	.8274 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.1418	.8935 %	1.0799	WEIGHT TAKEN
LIVER (LI)	7.9743	3.3268 %	4.0207	WEIGHT TAKEN
OVARY (OV)	.1125	.0469 %	.0567	WEIGHT TAKEN
SPLEEN (SP)	.5182	.2162 %	.2613	WEIGHT TAKEN
THYMUS (TH)	.2110	.0880 %	.1064	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0159	.0066 %	.0080	WEIGHT TAKEN

NECROPSY
OBSERVATIONS
HISTOPATHOLOGY
LIVER (LI) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

C01271

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 189

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96021 SEX: FEMALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 239.7 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:36 PROSECTOR: JERALD W. JOHNSON RECORDED: WENDY CHERRY
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

STOMACH, GL (ST), STOMACH, NONGL (SU)

001272

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 190

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96022		SEX: FEMALE		DOSE GROUP: 3		SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE	
DATE OF DEATH: 12/02/98		STUDY DAY OF DEATH: 93		STUDY WEEK OF DEATH: 14		TERMINAL BODY WEIGHT: 264.4 GRAMS	
DATE AND TIME OF NECROPSY: 12/02/98 12:09		PROSECTOR: GREGORY CODDINGTON		PATHOLOGIST: JOHN EIGHMY		RECORDER: TERRI L. WILLIAMS	
POST-FIX WEIGHTER: NOT AVAILABLE						WEIGHER: TONY HANFELD	
ORGAN NAME		ABSOLUTE ORGAN WEIGHT (GRAMS)		ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)		ORGAN TO BRAIN WEIGHT RATIO	
						O R G A N S T A T U S	
ADRENAL (AD)		.0765		.0289 %		.0367	
BRAIN (BR)		2.0837		.7881 %		1.0000	
KIDNEY (KD)		2.1627		.8180 %		1.0379	
LIVER (LI)		8.7642		3.3147 %		4.2061	
OVARY (OV)		.1119		.0423 %		.0537	
SPLEEN (SP)		.5033		.1904 %		.2415	
THYMUS (TH)		.2463		.0932 %		.1182	
THYROID/PARATHYR (TT)		.0279		.0106 %		.0134	

NECROPSY		O B S E R V A T I O N S		HISTOPATHOLOGY	
				LIVER (LI) :	
				-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MINIMAL	
				-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL	

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 191

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96022 SEX: FEMALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 264.4 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 12:09 PROSECTOR: GREGORY CODDINGTON RECORDER: TERRI L. WILLIAMS
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:
BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:
STOMACH, GL (ST), STOMACH, NONGL (SU)

C01274

APPENDIX 6
Individual Anatomic Pathology Data

PAGE: 192

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96023	SEX: FEMALE	DOSE GROUP: 3	SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98	STUDY DAY OF DEATH: 93	STUDY WEEK OF DEATH: 14	TERMINAL BODY WEIGHT: 253.0 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:06	PROSECTOR: GREGORY CODDINGTON	RECORDER: TERRI L. WILLIAMS	
POST-FIX WEIGHTER: NOT AVAILABLE	PATHOLOGIST: JOHN EIGHMY	WEIGHER: TONY HANFELD	
ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO
ADRENAL (AD)	.0695	.0275 %	.0337
BRAIN (BR)	2.0599	.8142 %	1.0000
KIDNEY (KD)	2.5238	.9975 %	1.2252
LIVER (LI)	10.5886	4.1852 %	5.1403
OVARY (OV)	.1184	.0468 %	.0575
SPLEEN (SP)	.5279	.2087 %	.2563
THYMUS (TH)	.3996	.1579 %	.1940
THYROID/PARATHYR (TT)	.0150	.0059 %	.0073
O B S E R V A T I O N S		HISTOPATHOLOGY	
NECROPSY		LIVER (LI) :	
		-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MINIMAL	
		-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL	
		-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL	
GENERAL COMMENT (GC) :			
-BONE MARROW SMEAR TAKEN			
-NO MACROSCOPIC LESIONS			

001275

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 193

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96023 SEX: FEMALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 253.0 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:06 PROSECTOR: GREGORY CODDINGTON RECORDER: TERRI L. WILLIAMS
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:
BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), STOMACH, NONGL (SU),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:
STOMACH, GL (ST), STOMACH, NONGL (SU)

001276

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 194

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL-
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96024		SEX: FEMALE		DOSE GROUP: 3		SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE	
DATE OF DEATH: 12/02/98		STUDY DAY OF DEATH: 93		STUDY WEEK OF DEATH: 14		TERMINAL BODY WEIGHT: 257.7 GRAMS	
DATE AND TIME OF NECROPSY: 12/02/98 12:12		PROSECTOR: JODI THIENEN		REORDER: TERRI L. WILLIAMS			
POST-FIX WEIGHTER: NOT AVAILABLE		PATHOLOGIST: JOHN EIGHMY		WEIGHER: TONY HANFELD			

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS	WEIGHT TAKEN
ADRENAL (AD)	.0753	.0292 %	.0350		WEIGHT TAKEN
BRAIN (BR)	2.1505	.8345 %	1.0000		WEIGHT TAKEN
KIDNEY (KD)	2.5361	.9841 %	1.1793		WEIGHT TAKEN
LIVER (LI)	8.6565	3.3591 %	4.0253		WEIGHT TAKEN
OVARY (OV)	.1361	.0528 %	.0633		WEIGHT TAKEN
SPLEEN (SP)	.4379	.1699 %	.2036		WEIGHT TAKEN
THYMUS (TH)	.2907	.1128 %	.1352		WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0183	.0071 %	.0085		WEIGHT TAKEN

NECROPSY	OBSERVATIONS	HISTOPATHOLOGY
	LIVER (LI) :	
	-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MINIMAL	
	-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL	

GENERAL COMMENT (GC) :

-BONE MARROW SMEAR TAKEN

-NO MACROSCOPIC LESIONS

C01277

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 195

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96024 SEX: FEMALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 257.7 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 12:12 PROSECTOR: JODI THIENEN RECORDED: TERRI L. WILLIAMS
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), NONGL (SU),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

STOMACH, GL (ST), STOMACH, NONGL (SU)

001278

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 196

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96025 SEX: FEMALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 253.9 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:33 PROSECTOR: KIMBERLY DURLAND REORDER: TERRI L. WILLIAMS
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN S T A T U S
ADRENAL (AD)	.0788	.0310 %	.0395	WEIGHT TAKEN
BRAIN (BR)	1.9934	.7851 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.3465	.9242 %	1.1771	WEIGHT TAKEN
LIVER (LI)	9.8976	3.8982 %	4.9652	WEIGHT TAKEN
OVARY (OV)	.1407	.0554 %	.0706	WEIGHT TAKEN
SPLEEN (SP)	.5995	.2361 %	.3007	WEIGHT TAKEN
THYMUS (TH)	.2171	.0855 %	.1089	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0268	.0106 %	.0134	WEIGHT TAKEN

O B S E R V A T I O N S

NECROPSY

HISTOPATHOLOGY

STOMACH, GL (ST) :
-LIGHT FOCUS(I)/AREA(S); MUCOSA, NEAR ESOPHAGEAL GROOVE:
SINGLE WHITE PINPOINT SLIGHTLY RAISED FOCUS

LIVER (LI) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
STOMACH, GL (ST) :
>UNREMARKABLE
>NOTE:>THERE IS A FOCAL AREA OF NORMAL-APPEARING SKELETAL MUSCLE IN THE TUNICA MUSCULARIS IN THE REGION OF THE LIMITING RIDGE.

LN, MANDIBULAR (MN) :
-MOTTLED; BILATERAL: RED AND TAN
OVARY (OV) :
-DIFFUSELY RED; BILATERAL
GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN

LN, MANDIBULAR (MN) :
-HEMORRHAGE, -MINIMAL
OVARY (OV) :
>UNREMARKABLE

001279

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 197

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96025 SEX: FEMALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 TERMINAL BODY WEIGHT: 253.9 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:33 PROSECTOR: KIMBERLY DURLAND RECORDED: TERRI L. WILLIAMS
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG),
HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF), URINARY BLADDER (UB), UTERUS (UT),
CERVIX (CV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

STOMACH, NONGL (SU)

C961230

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 198

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96026 SEX: FEMALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 232.5 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:08 PROSECTOR: GREGORY CODDINGTON RECORDED: TERRI L. WILLIAMS
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATISTICS
ADRENAL (AD)	.0700	.0301 %	.0353	WEIGHT TAKEN
BRAIN (BR)	1.9824	.8526 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.0111	.8650 %	1.0145	WEIGHT TAKEN
LIVER (LI)	8.3034	3.5714 %	4.1886	WEIGHT TAKEN
OVARY (OV)	.1165	.0501 %	.0588	WEIGHT TAKEN
SPLEEN (SP)	.4610	.1983 %	.2325	WEIGHT TAKEN
THYMUS (TH)	.2781	.1196 %	.1403	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0139	.0060 %	.0070	WEIGHT TAKEN

OBSERVATIONS

NECROPSY

HISTOPATHOLOGY

LIVER (LI) :
-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MINIMAL
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

001281

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 199

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96026 SEX: FEMALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 232.5 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:08 PROSECTOR: GREGORY CODDINGTON REORDER: TERRI L. WILLIAMS
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

STOMACH, GL (ST), STOMACH, NONGL (SU)

091282

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 200

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96027	SEX: FEMALE	DOSE GROUP: 3	SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98	STUDY DAY OF DEATH: 93	STUDY WEEK OF DEATH: 14	TERMINAL BODY WEIGHT: 296.8 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:10	PROSECTOR: LAURIE J. SCHULLER	RECORDER: TERRI L. WILLIAMS	
POST-FIX WEIGHT: NOT AVAILABLE	PATHOLOGIST: JOHN EIGHMY	WEIGHER: TONY HANFELD	
	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO
ORGAN NAME			S T A T U S
ADRENAL (AD)	.0659	.0222 %	WEIGHT TAKEN
BRAIN (BR)	1.9373	.6527 %	WEIGHT TAKEN
KIDNEY (KD)	2.3111	.7787 %	WEIGHT TAKEN
LIVER (LI)	9.8850	3.3305 %	WEIGHT TAKEN
OVARY (OV)	.1046	.0352 %	WEIGHT TAKEN
SPLEEN (SP)	.5319	.1792 %	WEIGHT TAKEN
THYMUS (TH)	.3779	.1273 %	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0177	.0060 %	WEIGHT TAKEN

NECROPSY	OBSERVATIONS	HISTOPATHOLOGY
GENERAL COMMENT (GC) :	LIVER (LI) :	
-BONE MARROW SMEAR TAKEN	-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL	
-NO MACROSCOPIC LESIONS	-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL	

001283

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 201

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96027 SEX: FEMALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 TERMINAL BODY WEIGHT: 296.8 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:10 PROSECTOR: LAURIE J. SCHULLER REORDER: TERRI L. WILLIAMS
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

STOMACH, GL (ST), STOMACH, NONGL (SU)

601284

APPENDIX 6

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96028	SEX: FEMALE	DOSE GROUP: 3	SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98	STUDY DAY OF DEATH: 93	STUDY WEEK OF DEATH: 14	TERMINAL BODY WEIGHT: 259.6 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 9:00	PROSECTOR: LAURIE J. SCHULLER		RECORDER: STEVE VAN ADESTINE
POST-FIX WEIGHTER: NOT AVAILABLE	PATHOLOGIST: JOHN EIGHMY		WEIGHER: TONY HANFELD
ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO
ADRENAL (AD)	.0628	.0242 %	.0315
BRAIN (BR)	1.9952	.7686 %	1.0000
KIDNEY (KD)	2.0909	.8054 %	1.0480
LIVER (LI)	8.7646	3.3762 %	4.3928
OVARY (OV)	.1106	.0426 %	.0554
SPLEEN (SP)	.5702	.2196 %	.2858
THYMUS (TH)	.3955	.1523 %	.1982
THYROID/PARATHYR (TT)	.0193	.0074 %	.0097
O R G A N S T A T U S			
WEIGHT TAKEN			
WEIGHT TAKEN			
WEIGHT TAKEN			
WEIGHT TAKEN			
WEIGHT TAKEN			
WEIGHT TAKEN			

O B S E R V A T I O N S		HISTOPATHOLOGY
NECROPSY		
LIVER (LI) :		
-INFILTRATE, LYMPHOHISTIOCYTIC,-MINIMAL		

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-MALOCCLUSION; BILATERAL UPPER INCISORS

601285

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 203

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96028 SEX: FEMALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 259.6 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 9:00 PROSECTOR: LAURIE J. SCHULLER REORDER: STEVE VAN ADESTINE
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

STOMACH, GL (ST), STOMACH, NONGL (SU)

001286

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 204

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

SEX: FEMALE		DOSE GROUP: 3		SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE	
ANIMAL NUMBER: C96029		STUDY DAY OF DEATH: 93		TERMINAL BODY WEIGHT: 239.5 GRAMS	
DATE OF DEATH: 12/02/98		STUDY WEEK OF DEATH: 14		RECORDED: STEVE VAN ADESTINE	
DATE AND TIME OF NECROPSY: 12/02/98 7:41		PROSECTOR: NANCY DIEDRICH		WEIGHER: TONY HANFELD	
POST-FIX WEIGHER: NOT AVAILABLE		PATHOLOGIST: JOHN EIGHMY			
ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	S T A T U S	
ADRENAL (AD)	.0817	.0341 %	.0412	WEIGHT TAKEN	
BRAIN (BR)	1.9819	.8275 %	1.0000	WEIGHT TAKEN	
KIDNEY (KD)	2.1671	.9048 %	1.0934	WEIGHT TAKEN	
LIVER (LI)	8.7181	3.8401 %	4.3989	WEIGHT TAKEN	
OVARY (OV)	.1243	.0519 %	.0627	WEIGHT TAKEN	
SPLEEN (SP)	.5235	.2186 %	.2641	WEIGHT TAKEN	
THYMUS (TH)	.2626	.1096 %	.1325	WEIGHT TAKEN	
THYROID/PARATHYR (TT)	.0179	.0075 %	.0090	WEIGHT TAKEN	

OBSERVATIONS

NECROPSY

HISTOPATHOLOGY

LIVER (LI) :
-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MINIMAL

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

001287

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 205

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96029 SEX: FEMALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 239.5 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 7:41 PROSECTOR: NANCY DIEDRICH REORDER: STEVE VAN ADESTINE
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:
BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:
STOMACH, GL (ST), STOMACH, NONGL (SU)

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 206

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96030		SEX: FEMALE	DOSE GROUP: 3	SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE	TERMINAL BODY WEIGHT: 243.6 GRAMS
DATE OF DEATH: 12/02/98		STUDY DAY OF DEATH: 93	STUDY WEEK OF DEATH: 14	RECORDED BY: TERRI L. WILLIAMS	
DATE AND TIME OF NECROPSY: 12/02/98 11:22		PROSECTOR: JODI THIEREN		WEIGHED BY: TONY HANFELD	
POST-FIX WEIGHTER: NOT AVAILABLE		PATHOLOGIST: JOHN EIGHMY			

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS	WEIGHT TAKEN
ADRENAL (AD)	.0640	.0263 %	.0316		WEIGHT TAKEN
BRAIN (BR)	2.0259	.8317 %	1.0000		WEIGHT TAKEN
KIDNEY (KD)	2.1625	.8877 %	1.0674		WEIGHT TAKEN
LIVER (LI)	10.3610	4.2533 %	5.1143		WEIGHT TAKEN
OVARY (OV)	.1272	.0522 %	.0628		WEIGHT TAKEN
SPLEEN (SP)	.5984	.2456 %	.2954		WEIGHT TAKEN
THYMUS (TH)	.2458	.1009 %	.1213		WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0121	.0050 %	.0060		WEIGHT TAKEN

NECROPSY	OBSERVATIONS	HISTOPATHOLOGY
THYMUS (TH) :		
-RED FOCUS(I)/AREA(S) :		
LN, MANDIBULAR (MN) :		
-DIFFUSELY RED; BILATERAL		
GENERAL COMMENT (GC) :		
-BONE MARROW SMEAR TAKEN		
LIVER (LI) :		
-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MINIMAL		
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL		
-NECROSIS, COAGULATIVE, -MINIMAL		
THYMUS (TH) :		
-HEMORRHAGE, -MINIMAL		
LN, MANDIBULAR (MN) :		
-HEMORRHAGE, -MINIMAL		

001289

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 207

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96030 SEX: FEMALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 243.6 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:22 PROSECTOR: JODI THIEREN REORDER: TERRI L. WILLIAMS
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC),
ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG),
HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF), URINARY BLADDER (UB), UTERUS (UT),
CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

STOMACH, GL (ST), STOMACH, NONGL (SU)

631290

APPENDIX 6

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96031	SEX: FEMALE	DOSE GROUP: 3	SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
DATE OF DEATH: 10/01/98	STUDY DAY OF DEATH: 31	STUDY WEEK OF DEATH: 5	TERMINAL BODY WEIGHT: 205.5 GRAMS
DATE AND TIME OF NECROPSY: 10/01/98 11:34	PROSECTOR: WENDY CHERRY		RECORDER: KEVIN BILLINGS
POST-FIX WEIGHTER: NOT AVAILABLE	PATHOLOGIST: JOHN EIGHMY		WEIGHER: MICHELE ZIMA
ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO
ADRENAL (AD)	---	---	---
BRAIN (BR)	---	---	---
KIDNEY (KD)	---	---	---
LIVER (LI)	7.4421	3.6215 %	---
OVARY (OV)	---	---	---
SPLEEN (SP)	---	---	---
THYMUS (TH)	---	---	---
THYROID/PARATHYR (TT)	---	---	---
			ORGAN STATUS
			NOT TAKEN
			NOT TAKEN
			NOT TAKEN
			WEIGHT TAKEN
			NOT TAKEN
			NOT TAKEN
			NOT TAKEN
THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:			
BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),			
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),			
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),			
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),			
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),			
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),			
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI), GENERAL COMMENT (GC)			

001291

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 209

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL-
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96032	SEX: FEMALE	DOSE GROUP: 3	SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
DATE OF DEATH: 10/01/98	STUDY DAY OF DEATH: 31	STUDY WEEK OF DEATH: 5	TERMINAL BODY WEIGHT: 203.4 GRAMS
DATE AND TIME OF NECROPSY: 10/01/98 11:09	PROSECTOR: WENDY CHERRY		RECORDER: KEVIN BILLINGS
POST-FIX WEIGHT: NOT AVAILABLE	PATHOLOGIST: JOHN EIGHMY		WEIGHER: MICHELE ZIMA
ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO

ADRENAL (AD)	-----	-----	NOT TAKEN
BRAIN (BR)	-----	-----	NOT TAKEN
KIDNEY (KD)	-----	-----	NOT TAKEN
LIVER (LI)	6.7736	3.3302 %	WEIGHT TAKEN
OVARY (OV)	-----	-----	NOT TAKEN
SPLEEN (SP)	-----	-----	NOT TAKEN
THYMUS (TH)	-----	-----	NOT TAKEN
THYROID/PARATHYR (TT)	-----	-----	NOT TAKEN
THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:			
BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),			
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),			
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),			
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),			
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),			
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),			
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI), GENERAL COMMENT (GC)			

001292

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 210

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96033	SEX: FEMALE	DOSE GROUP: 3	SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE	
DATE OF DEATH: 10/01/98	STUDY DAY OF DEATH: 31	STUDY WEEK OF DEATH: 5	TERMINAL BODY WEIGHT: 210.3 GRAMS	
DATE AND TIME OF NECROPSY: 10/01/98 11:15	PROSECTOR: WENDY CHERRY		RECORDER: KEVIN BILLINGS	
POST-FIX WEIGHTER: NOT AVAILABLE	PATHOLOGIST: JOHN EIGHMY		WEIGHER: MICHELE ZIMA	
ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	---	---	---	NOT TAKEN
BRAIN (BR)	---	---	---	NOT TAKEN
KIDNEY (KD)	---	---	---	NOT TAKEN
LIVER (LI)	7.7835	3.7011 %	---	WEIGHT TAKEN
OVARY (OV)	---	---	---	NOT TAKEN
SPLEEN (SP)	---	---	---	NOT TAKEN
THYMUS (TH)	---	---	---	NOT TAKEN
THYROID/PARATHYR (TT)	---	---	---	NOT TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TV), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI), GENERAL COMMENT (GC)

001293

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 211

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96034 SEX: FEMALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
DATE OF DEATH: 10/01/98 STUDY DAY OF DEATH: 31 STUDY WEEK OF DEATH: 5 TERMINAL BODY WEIGHT: 234.0 GRAMS
DATE AND TIME OF NECROPSY: 10/01/98 12:01 PROSECTOR: MATTHEW AXTMAN RECORDER: KEVIN BILLINGS
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: MICHELE ZIMA

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	---	---	---	NOT TAKEN
BRAIN (BR)	---	---	---	NOT TAKEN
KIDNEY (KD)	---	---	---	NOT TAKEN
LIVER (LI)	8.8195	3.7690 %	---	WEIGHT TAKEN
OVARY (OV)	---	---	---	NOT TAKEN
SPLEEN (SP)	---	---	---	NOT TAKEN
THYMUS (TH)	---	---	---	NOT TAKEN
THYROID/PARATHYR (TT)	---	---	---	NOT TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LONG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI), GENERAL COMMENT (GC)

601294

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 212

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96035	SEX: FEMALE	DOSE GROUP: 3	SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE	
DATE OF DEATH: 10/01/98	STUDY DAY OF DEATH: 31	STUDY WEEK OF DEATH: 5	TERMINAL BODY WEIGHT: 218.8 GRAMS	
DATE AND TIME OF NECROPSY: 10/01/98 11:21	PROSECTOR: WENDY CHERRY		REORDER: KEVIN BILLINGS	
POST-FIX WEIGHTER: NOT AVAILABLE	PATHOLOGIST: JOHN EIGHMY		WEIGHER: MICHELE ZIMA	
ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	---	---	---	NOT TAKEN
BRAIN (BR)	---	---	---	NOT TAKEN
KIDNEY (KD)	---	---	---	NOT TAKEN
LIVER (LI)	7.2747	3.3248 %	---	WEIGHT TAKEN
OVARY (OV)	---	---	---	NOT TAKEN
SPLEEN (SP)	---	---	---	NOT TAKEN
THYMUS (TH)	---	---	---	NOT TAKEN
THYROID/PARATHYR (TT)	---	---	---	NOT TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI), GENERAL COMMENT (GC)

APPENDIX 6

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 213

ANIMAL NUMBER: C96036		SEX: FEMALE	DOSE GROUP: 3	SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE	
DATE OF DEATH: 12/02/98		STUDY DAY OF DEATH: 93	STUDY WEEK OF DEATH: 14	TERMINAL BODY WEIGHT: 253.4 GRAMS	
DATE AND TIME OF NECROPSY: 12/02/98 7:45		PROSECTOR: GREGORY CODDINGTON		RECORDER: STEVE VAN ADESTINE	
POST-FIX WEIGHER: NOT AVAILABLE		PATHOLOGIST: JOHN EIGHMY		WEIGHER: TONY HANFELD	
		ABSOLUTE ORGAN WEIGHT (GRAMS)		ORGAN TO BRAIN WEIGHT RATIO	
				O R G A N S T A T U S	
ADRENAL (AD)	.0611	.0241 %		WEIGHT TAKEN	
BRAIN (BR)	2.1326	.8416 %		WEIGHT TAKEN	
KIDNEY (KD)	2.3286	.9189 %		WEIGHT TAKEN	
LIVER (LI)	9.7463	3.8462 %		WEIGHT TAKEN	
OVARY (OV)	.1035	.0408 %		WEIGHT TAKEN	
SPLEEN (SP)	.4891	.1930 %		WEIGHT TAKEN	
THYMUS (TH)	.3725	.1470 %		WEIGHT TAKEN	
THYROID/PARATHYR (TT)	.0205	.0081 %		WEIGHT TAKEN	

NECROPSY OBSERVATIONS HISTOPATHOLOGY

LIVER (LI) :
-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MINIMAL
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 214

ANIMAL NUMBER: C96036 SEX: FEMALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 253.4 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 7:45 PROSECTOR: GREGORY CODDINGTON REORDER: STEVE VAN ADESTINE
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

STOMACH, GL (ST), STOMACH, NONGL (SU)

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 215

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96037
SEX: FEMALE
DOSE GROUP: 3
SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98
STUDY DAY OF DEATH: 93
STUDY WEEK OF DEATH: 14
TERMINAL BODY WEIGHT: 251.1 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 7:37
PROSECTOR: MICHELE ZIMA
REORDER: WENDY CHERRY
POST-FIX WEIGHTER: NOT AVAILABLE
PATHOLOGIST: JOHN EIGHMY
WEIGHER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0804	.0320 %	.0432	WEIGHT TAKEN
BRAIN (BR)	1.8602	.7408 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.3767	.9465 %	1.2777	WEIGHT TAKEN
LIVER (LI)	8.8940	3.5420 %	4.7812	WEIGHT TAKEN
OVARY (OV)	.0918	.0366 %	.0493	WEIGHT TAKEN
SPLEEN (SP)	.5405	.2153 %	.2906	WEIGHT TAKEN
THYMUS (TH)	.2792	.1112 %	.1501	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0196	.0078 %	.0105	WEIGHT TAKEN

NECROPSY

OBSERVATIONS

HISTOPATHOLOGY

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

C01298

APPENDIX 6
Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 216

ANIMAL NUMBER: C96037 SEX: FEMALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 251.1 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 7:37 PROSECTOR: MICHELE ZIMA REORDER: WENDY CHERRY
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (UB), HARNERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

LIVER (LI), STOMACH, GL (ST), STOMACH, NONGL (SU)

601239

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 217

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96038 SEX: FEMALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 275.3 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:14 PROSECTOR: JERALD W. JOHNSON RECORDED: WENDY CHERRY
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0554	.0201 %	.0259	WEIGHT TAKEN
BRAIN (BR)	2.1428	.7784 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.1681	.7875 %	1.0118	WEIGHT TAKEN
LIVER (LI)	9.8295	3.5705 %	4.5872	WEIGHT TAKEN
OVARY (OV)	.1411	.0513 %	.0658	WEIGHT TAKEN
SPLEEN (SP)	.4874	.1770 %	.2275	WEIGHT TAKEN
THYMUS (TH)	.2831	.1028 %	.1321	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0153	.0056 %	.0071	WEIGHT TAKEN

OBSERVATIONS

HISTOPATHOLOGY

NECROPSY

LIVER (LI):
-INFILTRATE, LYMPHOHISTIOCYTIC, MINIMAL
LN, MANDIBULAR (MN):
-HEMORRHAGE, SLIGHT
UTERUS (UT):
>UNREMARKABLE

LN, MANDIBULAR (MN):
-DIFFUSELY RED; BILATERAL
UTERUS (UT):
-LUMEN FILLED WITH FLUID; LEFT: CLEAR FLUID
GENERAL COMMENT (GC):
-BONE MARROW SHEAR TAKEN

001300

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 218

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96038 SEX: FEMALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 275.3 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:14 PROSECTOR: JERALD W. JOHNSON REORDER: WENDY CHERRY
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG),
HARCIERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF), URINARY BLADDER (UB), CERVIX (CV),
OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

STOMACH, GL (ST), STOMACH, NONGL (SU)

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 219

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96039		SEX: FEMALE	DOSE GROUP: 3	SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE	
DATE OF DEATH: 12/02/98		STUDY DAY OF DEATH: 93	STUDY WEEK OF DEATH: 14	TERMINAL BODY WEIGHT: 257.5 GRAMS	
DATE AND TIME OF NECROPSY: 12/02/98 8:13		PROSECTOR: NANCY DIEDRICH		RECORDER: STEVE VAN ADESTINE	
POST-FIX WEIGHER: NOT AVAILABLE		PATHOLOGIST: JOHN EIGHMY		WEIGHER: TONY HANFELD	
		O R G A N			
		S T A T U S			

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	WEIGHT TAKEN	
ADRENAL (AD)	.0835	.0324 %	.0415	WEIGHT TAKEN	
BRAIN (BR)	2.0098	.7805 %	1.0000	WEIGHT TAKEN	
KIDNEY (KD)	2.2091	.8579 %	1.0992	WEIGHT TAKEN	
LIVER (LI)	9.1522	3.5543 %	4.5538	WEIGHT TAKEN	
OVARY (OV)	.1115	.0433 %	.0555	WEIGHT TAKEN	
SPLEEN (SP)	.4315	.1676 %	.2147	WEIGHT TAKEN	
THYMUS (TH)	.3296	.1280 %	.1640	WEIGHT TAKEN	
THYROID/PARATHYR (TT)	.0199	.0077 %	.0099	WEIGHT TAKEN	

		O B S E R V A T I O N S			

NECROPSY	HISTOPATHOLOGY				

		LIVER (LI) :			
		-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL			

GENERAL COMMENT (GC) :					
-BONE MARROW SMEAR TAKEN					
-NO MACROSCOPIC LESIONS					

001302

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 220

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96039 SEX: FEMALE DOSE GROUP: 3 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 257.5 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:13 PROSECTOR: NANCY DIEDRICH REORDER: STEVE VAN ADESTINE
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

STOMACH, GL (ST), STOMACH, NONGL (SU)

091303

APPENDIX 6
Individual Anatomic Pathology Data

PAGE: 221

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
TERMINAL BODY WEIGHT: 222.7 GRAMS
REORDER: TERRI L. WILLIAMS
WEIGHER: TONY HANFELD

DOSE GROUP: 3
STUDY WEEK OF DEATH: 14
PROSECTOR: KIMBERLY DURLAND
PATHOLOGIST: JOHN EIGHMY

SEX: FEMALE
STUDY DAY OF DEATH: 93
ANIMAL NUMBER: C96040
DATE OF DEATH: 12/02/98
DATE AND TIME OF NECROPSY: 12/02/98 12:04
POST-FIX WEIGHER: NOT AVAILABLE

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	WEIGHT TAKEN
ADRENAL (AD)	.0650	.0292 %	.0336	WEIGHT TAKEN
BRAIN (BR)	1.9359	.8693 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.3025	1.0339 %	1.1894	WEIGHT TAKEN
LIVER (LI)	9.2333	4.1461 %	4.7695	WEIGHT TAKEN
OVARY (OV)	.1810	.0813 %	.0935	WEIGHT TAKEN
SPLEEN (SP)	.5292	.2376 %	.2734	WEIGHT TAKEN
THYMUS (TH)	.2778	.1247 %	.1435	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0253	.0114 %	.0131	WEIGHT TAKEN

HISTOPATHOLOGY

NECROPSY

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

001304

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 222

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
TERMINAL BODY WEIGHT: 222.7 GRAMS
RECORDED: TERRI L. WILLIAMS
WEIGHER: TONY HANFELD

DOSE GROUP: 3
STUDY WEEK OF DEATH: 14
PROSECTOR: KIMBERLY DURLAND
PATHOLOGIST: JOHN EIGHMY

ANIMAL NUMBER: C96040
SEX: FEMALE
STUDY DAY OF DEATH: 93
DATE OF DEATH: 12/02/98
DATE AND TIME OF NECROPSY: 12/02/98 12:04
POST-FIX WEIGHTER: NOT AVAILABLE

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:
BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TV), PARATHYROID (PT), STOMACH, NONGL (SU),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), RECTUM (RE), LN, MANDIBULAR (MN),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

LIVER (LI), STOMACH, GL (ST), STOMACH, NONGL (SU)

001305

APPENDIX 6

Covance 6329-225
3M T-6314.1

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 223

ANIMAL NUMBER: C96041
 DATE OF DEATH: 12/02/98
 DATE AND TIME OF NECROPSY: 12/02/98 8:59
 POST-FIX WEIGHER: NOT AVAILABLE

SEX: FEMALE
 STUDY DAY OF DEATH: 93
 PROSECTOR: TODD ACKER
 PATHOLOGIST: JOHN EIGHMY

DOSE GROUP: 4
 STUDY WEEK OF DEATH: 14
 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
 TERMINAL BODY WEIGHT: 212.0 GRAMS
 RECORDER: WENDY CHERRY
 WEIGHER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0654	.0308 %	.0363	WEIGHT TAKEN
BRAIN (BR)	1.8020	.8500 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	1.8605	.8776 %	1.0325	WEIGHT TAKEN
LIVER (LI)	10.2446	4.8324 %	5.6851	WEIGHT TAKEN
OVARY (OV)	.1083	.0511 %	.0601	WEIGHT TAKEN
SPLEEN (SP)	.4309	.2033 %	.2391	WEIGHT TAKEN
THYMUS (TH)	.2212	.1043 %	.1228	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0158	.0075 %	.0088	WEIGHT TAKEN

O B S E R V A T I O N S

NECROPSY

HISTOPATHOLOGY

LUNG (LU) :
 -INFILTRATE, MACROPHAGE, ALVEOLAR, -MINIMAL
 LIVER (LI) :
 -HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MODERATE
 -INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
 -NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL
 AUDITORY SEB GL (AS) :
 -ONE EXAMINED, -PRESENT

GENERAL COMMENT (GC) :
 -BONE MARROW SMEAR TAKEN
 -NO MACROSCOPIC LESIONS

001306

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 224

ANIMAL NUMBER: C96041 SEX: FEMALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 212.0 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:59 PROSECTOR: TODD ACKER RECORDED: WENDY CHERRY
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN),
SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC),
ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), SKIN (SK), MAMMARY, FEMALE (MF), URINARY BLADDER (UB), UTERUS (UT),
CERVIX (CV), OVARY (OV), VAGINA (VA)

APPENDIX 6

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 225

ANIMAL NUMBER: C96042		SEX: FEMALE	DOSE GROUP: 4		SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE	
DATE OF DEATH: 12/02/98		STUDY DAY OF DEATH: 93	STUDY WEEK OF DEATH: 14		TERMINAL BODY WEIGHT: 228.8 GRAMS	
DATE AND TIME OF NECROPSY: 12/02/98 12:00		PROSECTOR: TODD ACKER		REORDER: WENDY CHERRY		
POST-FIX WEIGHER: NOT AVAILABLE		PATHOLOGIST: JOHN EIGHMY		WEIGHER: LYNN HORNER		

ORGAN NAME		ABSOLUTE ORGAN WEIGHT (GRAMS)		ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)		ORGAN TO BRAIN WEIGHT RATIO

ADRENAL (AD)		.0619		.0271 %		.0302
BRAIN (BR)		2.0530		.8973 %		1.0000
KIDNEY (KD)		2.2573		.9866 %		1.0995
LIVER (LI)		13.7996		6.0313 %		6.7217
OVARY (OV)		.1247		.0545 %		.0607
SPLEEN (SP)		.4886		.2135 %		.2380
THYMUS (TH)		.2500		.1093 %		.1218
THYROID/PARATHYR (TT)		.0203		.0089 %		.0099

OBSERVATIONS

NECROPSY

HISTOPATHOLOGY

LIVER (LI) :
-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MODERATE
-PIGMENT, HEPATOCELLULAR, -MINIMAL
KIDNEY (KD) :
-INFLAMMATION, CHRONIC, -MINIMAL
THYMUS (TH) :
-HEMORRHAGE, -MINIMAL
STOMACH, GL (ST) :
-EROSION, -MINIMAL

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

001308

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEPOSE, T-6314) IN RATS

PAGE: 226

ANIMAL NUMBER: C96042 SEX: FEMALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 228.8 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 12:00 PROSECTOR: TODD ACKER RECORDER: WENDY CHERRY
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TV), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN),
SPLEEN (SP), LN, MESENTERIC (MS), THYROID (TV), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA),
AORTA (AO), PITUITARY (PI), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL),
CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG),
AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF), URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV),
VAGINA (VA)

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 227

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96043 SEX: FEMALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 218.3 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:54 PROSECTOR: NANCY DIEDRICH
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY RECORDER: STEVE VAN ADESTINE
WEIGHER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0754	.0345 %	.0374	WEIGHT TAKEN
BRAIN (BR)	2.0160	.9235 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.1708	.9944 %	1.0768	WEIGHT TAKEN
LIVER (LI)	9.1440	4.1887 %	4.5357	WEIGHT TAKEN
OVARY (OV)	.1120	.0513 %	.0556	WEIGHT TAKEN
SPLEEN (SP)	.4212	.1929 %	.2089	WEIGHT TAKEN
THYMUS (TH)	.2302	.1055 %	.1142	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0191	.0087 %	.0095	WEIGHT TAKEN

OBSERVATIONS

NECROPSY

HISTOPATHOLOGY

LIVER (LI) :
- HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -SLIGHT
- INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
THYROID (TY) :
- CYST, ULTIMOBRANCHIAL, -PRESENT
STOMACH, GL (ST) :
- EROSION, -MINIMAL
STOMACH, GL (ST) :
- DARK FOCUS(I)/AREA(S); MUCOSA: MULTIPLE DARK BROWN AREAS,
UP TO 4 X 2 MM
GENERAL COMMENT (GC) :
- BONE MARROW SMEAR TAKEN

001310

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 228

ANIMAL NUMBER: C96043 SEX: FEMALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 218.3 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:54 PROSECTOR: NANCY DIEDRICH
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY RECORDER: STEVE VAN ADESTINE
WEIGHER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), PARATHYROID (PT), ADRENAL, CORTEX (AC),
ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE),
ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG),
AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF), URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV),
VAGINA (VA)

001311

APPENDIX 6
Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS PAGE: 229

ANIMAL NUMBER: C96044 SEX: FEMALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 261.0 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 12:19 PROSECTOR: TODD ACKER
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY RECORDER: WENDY CHERRY
WEIGHER: LYNN HORNER

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0833	.0319 %	.0381	WEIGHT TAKEN
BRAIN (BR)	2.1839	.8367 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.4878	.9532 %	1.1392	WEIGHT TAKEN
LIVER (LI)	14.4781	5.5472 %	6.6295	WEIGHT TAKEN
OVARY (OV)	.1201	.0460 %	.0550	WEIGHT TAKEN
SPLEEN (SP)	.5592	.2143 %	.2561	WEIGHT TAKEN
THYMUS (TH)	.2968	.1137 %	.1359	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0194	.0074 %	.0089	WEIGHT TAKEN

NECROPSY OBSERVATIONS HISTOPATHOLOGY

LIVER (LI) :
-HYPERTROPHY, HEPATOCELLULAR, CENTRIOBLULAR, -SLIGHT
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
HEART (HT) :
-DEGENERATION/NECROSIS, -MINIMAL
THYMUS (TH) :
-HEMORRHAGE, -MINIMAL

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 230

ANIMAL NUMBER: C96044 SEX: FEMALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 261.0 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 12:19 PROSECTOR: TODD ACKER RECORDED: WENDY CHERRY
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN),
SPLEEN (SP), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA),
AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE),
ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG),
AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF), URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV),
VAGINA (VA)

001313

APPENDIX 6
Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 231

ANIMAL NUMBER: C96045		SEX: FEMALE		DOSE GROUP: 4		SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE	
DATE OF DEATH: 12/02/98		STUDY DAY OF DEATH: 93		STUDY WEEK OF DEATH: 14		TERMINAL BODY WEIGHT: 251.4 GRAMS	
DATE AND TIME OF NECROPSY: 12/02/98 9:47		PROSECTOR: KIMBERLY DURLAND		REORDER: TERRI L. WILLIAMS			
POST-FIX WEIGHT: NOT AVAILABLE		PATHOLOGIST: JOHN EIGHMY		WEIGHER: TONY HANFELD			
		ABSOLUTE ORGAN WEIGHT (GRAMS)		ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)		O R G A N S T A T U S	
ADRENAL (AD)		.0862		.0343 %		WEIGHT TAKEN	
BRAIN (BR)		2.0735		.8248 %		WEIGHT TAKEN	
KIDNEY (KD)		2.6621		1.0589 %		WEIGHT TAKEN	
LIVER (LI)		11.5067		4.5770 %		WEIGHT TAKEN	
OVARY (OV)		.1738		.0691 %		WEIGHT TAKEN	
SPLEEN (SP)		.5999		.2386 %		WEIGHT TAKEN	
THYMUS (TH)		.2968		.1181 %		WEIGHT TAKEN	
THYROID/PARATHYR (TT)		.0316		.0126 %		WEIGHT TAKEN	

NECROPSY	OBSERVATIONS	HISTOPATHOLOGY
	LIVER (LI) :	
	-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -SLIGHT	
	-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL	
	-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL	
	PANCREAS (PA) :	
	-ATROPHY, ACINAR, -MINIMAL	
	-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL	
	AUDITORY SEB GL (AS) :	
	-ONE EXAMINED, -PRESENT	
	UTERUS (UT) :	
	-DILATATION, -SLIGHT	
	OVARY (OV) :	
	>UNREMARKABLE	
OVARY (OV) :		
-DIFFUSELY RED; BILATERAL		
GENERAL COMMENT (GC) :		
-BONE MARROW SMEAR TAKEN		

601314

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

PAGE: 232

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL-
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96045 SEX: FEMALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 251.4 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 9:47 PROSECTOR: KIMBERLY DURLAND REORDER: TERRI L. WILLIAMS
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), SKIN (SK), MAMMARY, FEMALE (MF), URINARY BLADDER (UB), CERVIX (CV),
VAGINA (VA)

001315

APPENDIX 6

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 233

ANIMAL NUMBER: C96046 SEX: FEMALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 266.9 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:42 PROSECTOR: NANCY DIEDRICH REORDER: TERRI L. WILLIAMS
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0612	.0229 %	.0292	WEIGHT TAKEN
BRAIN (BR)	2.0991	.7865 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.2050	.8262 %	1.0505	WEIGHT TAKEN
LIVER (LI)	11.8714	4.4479 %	5.6555	WEIGHT TAKEN
OVARY (OV)	.1311	.0491 %	.0625	WEIGHT TAKEN
SPLEEN (SP)	.5352	.2005 %	.2550	WEIGHT TAKEN
THYMUS (TH)	.3266	.1224 %	.1556	WEIGHT TAKEN
THYROID/PARATHYR (TPT)	.0186	.0070 %	.0089	WEIGHT TAKEN

NECROPSY OBSERVATIONS HISTOPATHOLOGY

LIVER (LI):
-HYPERTROPHY, HEPATOCELLULAR, CENTRIOBLULAR, -SLIGHT
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL

GENERAL COMMENT (GC):
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

001316

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 234

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96046 SEX: FEMALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 266.9 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:42 PROSECTOR: NANCY DIEDRICH RECORDER: TERRI L. WILLIAMS
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR), (SM),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA)

001317

APPENDIX 6
Individual Anatomic Pathology Data

PAGE: 235

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96047
SEX: FEMALE
DATE OF DEATH: 12/02/98
STUDY DAY OF DEATH: 93
SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
TERMINAL BODY WEIGHT: 213.3 GRAMS
REORDER: TERRI L. WILLIAMS
WEIGHER: TONY HANFELD
DATE AND TIME OF NECROPSY: 12/02/98 11:53
PROSECTOR: NANCY DIEDRICH
PATHOLOGIST: JOHN EIGHY
POST-FIX WEIGHER: NOT AVAILABLE

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0586	.0275 %	.0281	WEIGHT TAKEN
BRAIN (BR)	2.0822	.9762 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.3414	1.0977 %	1.1245	WEIGHT TAKEN
LIVER (LI)	12.2355	5.7363 %	5.8762	WEIGHT TAKEN
OVARY (OV)	.1156	.0542 %	.0555	WEIGHT TAKEN
SPLEEN (SP)	.4873	.2285 %	.2340	WEIGHT TAKEN
THYMUS (TH)	.3388	.1588 %	.1627	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0191	.0090 %	.0092	WEIGHT TAKEN

NECROPSY

LIVER (LI) :
-DIFFUSELY DARK; ALL LOBES: DARK BROWN

UTERUS (UT) :
-LUMEN FILLED WITH FLUID; BILATERAL: CLEAR FLUID
GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN

HISTOPATHOLOGY

LUNG (LU) :
-INFILTRATE, MACROPHAGE, ALVEOLAR, -MINIMAL

LIVER (LI) :
-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MODERATE
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-PIGMENT, HEPATOCELLULAR, -MINIMAL

THYROID (TY) :
-CYST, ULTIMOBRANCHIAL, -PRESENT

UTERUS (UT) :
-DILATATION, -SLIGHT

601318

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.1

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 236

ANIMAL NUMBER: C96047 SEX: FEMALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 213.3 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:53 PROSECTOR: NANCY DIEDRICH REORDER: TERRI L. WILLIAMS
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN),
SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA),
AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE),
ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG),
AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF), URINARY BLADDER (UB), CERVIX (CV), OVARY (OV), VAGINA (VA)

001319

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 237

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96048 SEX: FEMALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 223.8 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:40 PROSECTOR: KIMBERLY DURLAND REORDER: TERRI L. WILLIAMS
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS	WEIGHT TAKEN
ADRENAL (AD)	.0851	.0380 %	.0404		WEIGHT TAKEN
BRAIN (BR)	2.1074	.9416 %	1.0000		WEIGHT TAKEN
KIDNEY (KD)	2.3036	1.0293 %	1.0931		WEIGHT TAKEN
LIVER (LI)	13.9635	6.2393 %	6.6259		WEIGHT TAKEN
OVARY (OV)	.1738	.0777 %	.0825		WEIGHT TAKEN
SPLEEN (SP)	.5093	.2276 %	.2417		WEIGHT TAKEN
THYMUS (TH)	.2971	.1328 %	.1410		WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0252	.0113 %	.0120		WEIGHT TAKEN

OBSERVATIONS

NECROPSY

HISTOPATHOLOGY

BRAIN (BR) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
LUNG (LU) :
-INFLAMMATION, SUBACUTE, -MINIMAL
LIVER (LI) :
-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MODERATE
-PIGMENT, HEPATOCELLULAR, -MINIMAL
-VACUOLATION, HEPATOCELLULAR, CENTRILLOBULAR TO MIDZONAL, -MINIMAL
KIDNEY (KD) :
-MINERALIZATION, TUBULAR, -MINIMAL
THYROID (TY) :
-CYST, ULTIMOBRANCHIAL, -PRESENT
HARDERIAN GLAND (HG) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
OVARY (OV) :
>UNREMARKABLE

OVARY (OV) :
-DIFFUSELY RED; BILATERAL: DARK RED

APPENDIX 6
Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 238

ANIMAL NUMBER: C96048
SEX: FEMALE
DATE OF DEATH: 12/02/98
STUDY DAY OF DEATH: 93
SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
TERMINAL BODY WEIGHT: 223.8 GRAMS
REORDER: TERRI L. WILLIAMS
POST-FIX WEIGHER: NOT AVAILABLE
PATHOLOGIST: JOHN EIGHMY
WEIGHER: TONY HANFELD

O B S E R V A T I O N S (CONTINUED)

HISTOPATHOLOGY

NECROPSY

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), SPINAL CORD (SC),
TRACHEA (TR), ESOPHAGUS (ES), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH),
LN, MESENTERIC (MS), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI),
STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE),
COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), AUDITORY SEB GL (AS), SKIN (SK),
MAMMARY, FEMALE (MF), URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), VAGINA (VA)

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 239

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96049 SEX: FEMALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 191.4 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:55 PROSECTOR: GREGORY CODDINGTON REORDER: TERRI L. WILLIAMS
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0539	.0282 %	.0297	WEIGHT TAKEN
BRAIN (BR)	1.8149	.9482 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	1.7670	.9232 %	.9736	WEIGHT TAKEN
LIVER (LI)	10.6923	5.5864 %	5.8914	WEIGHT TAKEN
OVARY (OV)	.0973	.0508 %	.0536	WEIGHT TAKEN
SPLEEN (SP)	.4078	.2131 %	.2247	WEIGHT TAKEN
THYMUS (TH)	.2422	.1265 %	.1335	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0173	.0090 %	.0095	WEIGHT TAKEN

NECROPSY OBSERVATIONS HISTOPATHOLOGY

STOMACH, GL (ST) :
-DARK FOCUS(1)/AREA(S); MUCOSA: MULTIPLE DARK BROWN AREAS,
UP TO 5 X 1 MM
GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-TAIL - BENT/MISSING; TIP: MISSING; REMAINING PORTION
COLLECTED

LIVER (LI) :
-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MODERATE
-PIGMENT, HEPATOCELLULAR, -MINIMAL
STOMACH, GL (ST) :
-EROSION, -SLIGHT

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 240

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96049 SEX: FEMALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 191.4 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 10:55 PROSECTOR: GREGORY CODDINGTON RECORDER: TERRI L. WILLIAMS
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA)

001323

APPENDIX 6
Individual Anatomic Pathology Data

PAGE: 241

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96050
SEX: FEMALE
DATE OF DEATH: 12/02/98
STUDY DAY OF DEATH: 93
DOSE GROUP: 4
SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
TERMINAL BODY WEIGHT: 216.8 GRAMS
REORDER: TERRI L. WILLIAMS
WEIGHER: TONY HANFELD
DATE AND TIME OF NECROPSY: 12/02/98 11:40
PROSECTOR: GREGORY CODDINGTON
PATHOLOGIST: JOHN EIGHY
POST-FIX WEIGHER: NOT AVAILABLE

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATISTICS
ADRENAL (AD)	.0634	.0292 %	.0313	WEIGHT TAKEN
BRAIN (BR)	2.0287	.9357 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	1.9598	.9040 %	.9660	WEIGHT TAKEN
LIVER (LI)	12.8289	5.9174 %	6.3237	WEIGHT TAKEN
OVARY (OV)	.0985	.0454 %	.0486	WEIGHT TAKEN
SPLEEN (SP)	.3976	.1834 %	.1960	WEIGHT TAKEN
THYMUS (TH)	.1845	.0851 %	.0909	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0195	.0090 %	.0096	WEIGHT TAKEN

NECROPSY
OBSERVATIONS
HISTOPATHOLOGY
LIVER (LI):
-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MODERATE
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL
-PIGMENT, HEPATOCELLULAR, -MINIMAL

GENERAL COMMENT (GC):
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

001324

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 242

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96050 SEX: FEMALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 216.8 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 11:40 PROSECTOR: GREGORY CODDINGTON RECORDED: TERRI L. WILLIAMS
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:
BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:
BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA)

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 243

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96051 SEX: FEMALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
DATE OF DEATH: 10/01/98 STUDY DAY OF DEATH: 31 STUDY WEEK OF DEATH: 5 TERMINAL BODY WEIGHT: 181.0 GRAMS
DATE AND TIME OF NECROPSY: 10/01/98 11:19 PROSECTOR: MATTHEW AXTMAN REORDER: KEVIN BILLINGS
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: MICHELE ZIMA

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	---	---	---	NOT TAKEN
BRAIN (BR)	---	---	---	NOT TAKEN
KIDNEY (KD)	---	---	---	NOT TAKEN
LIVER (LI)	8.8661	4.8984 %	---	WEIGHT TAKEN
OVARY (OV)	---	---	---	NOT TAKEN
SPLEEN (SP)	---	---	---	NOT TAKEN
THYMUS (TH)	---	---	---	NOT TAKEN
THYROID/PARATHYR (TT)	---	---	---	NOT TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TV), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI), GENERAL COMMENT (GC)

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 244

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96052	SEX: FEMALE	DOSE GROUP: 4	SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
DATE OF DEATH: 10/01/98	STUDY DAY OF DEATH: 31	STUDY WEEK OF DEATH: 5	TERMINAL BODY WEIGHT: 202.7 GRAMS
DATE AND TIME OF NECROPSY: 10/01/98 11:52	PROSECTOR: WENDY CHERY		RECORDER: KEVIN BILLINGS
POST-FIX WEIGHTER: NOT AVAILABLE	PATHOLOGIST: JOHN EIGHMY		WEIGHER: MICHELE ZIMA

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO
ADRENAL (AD)	-----	-----	-----
BRAIN (BR)	-----	-----	NOT TAKEN
KIDNEY (KD)	-----	-----	NOT TAKEN
LIVER (LI)	8.9691	4.4248 %	WEIGHT TAKEN
OVARY (OV)	-----	-----	NOT TAKEN
SPLEEN (SP)	-----	-----	NOT TAKEN
THYMUS (TH)	-----	-----	NOT TAKEN
THYROID/PARATHYR (TT)	-----	-----	NOT TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:			
BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),			
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),			
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),			
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),			
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),			
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),			
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI), GENERAL COMMENT (GC)			

C01327

APPENDIX 6

Covance 6329-225
3M T-6314.1

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 245

ANIMAL NUMBER: C96053 SEX: FEMALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
DATE OF DEATH: 10/01/98 STUDY DAY OF DEATH: 31 STUDY WEEK OF DEATH: 5 TERMINAL BODY WEIGHT: 191.0 GRAMS
DATE AND TIME OF NECROPSY: 10/01/98 11:55 PROSECTOR: WENDY CHERRY REORDER: KEVIN BILLINGS
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: MICHELE ZIMA

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	---	---	---	NOT TAKEN
BRAIN (BR)	---	---	---	NOT TAKEN
KIDNEY (KD)	---	---	---	NOT TAKEN
LIVER (LI)	8.3894	4.3924 %	---	WEIGHT TAKEN
OVARY (OV)	---	---	---	NOT TAKEN
SPLEEN (SP)	---	---	---	NOT TAKEN
THYROID (TH)	---	---	---	NOT TAKEN
THYROID/PARATHYR (TT)	---	---	---	NOT TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI), GENERAL COMMENT (GC)

001328

APPENDIX 6

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 246ANIMAL NUMBER: C96054
SEX: FEMALE
DOSE GROUP: 4
SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE
DATE OF DEATH: 10/01/98
STUDY DAY OF DEATH: 31
STUDY WEEK OF DEATH: 5
TERMINAL BODY WEIGHT: 186.7 GRAMS
DATE AND TIME OF NECROPSY: 10/01/98 11:16
PROSECTOR: TONY HANFELD
REORDER: KEVIN BILLINGS
POST-FIX WEIGHT: NOT AVAILABLE
PATHOLOGIST: JOHN EIGHMY
WEIGHER: MICHELE ZIMA

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	---	---	---	NOT TAKEN
BRAIN (BR)	---	---	---	NOT TAKEN
KIDNEY (KD)	---	---	---	NOT TAKEN
LIVER (LI)	8.0181	4.2946 %	---	WEIGHT TAKEN
OVARY (OV)	---	---	---	NOT TAKEN
SPLEEN (SP)	---	---	---	NOT TAKEN
THYMUS (TH)	---	---	---	NOT TAKEN
THYROID/PARATHYR (TT)	---	---	---	NOT TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI), GENERAL COMMENT (GC)

APPENDIX 6

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 247

ANIMAL NUMBER: C96055	SEX: FEMALE	DOSE GROUP: 4	SACRIFICE STATUS: SCHEDULED, FIRST INTERIM SACRIFICE	
DATE OF DEATH: 10/01/98	STUDY DAY OF DEATH: 31	STUDY WEEK OF DEATH: 5	TERMINAL BODY WEIGHT: 178.9 GRAMS	
DATE AND TIME OF NECROPSY: 10/01/98 11:28	PROSECTOR: TONY HANFELD		RECORDER: KEVIN BILLINGS	
POST-FIX WEIGHTER: NOT AVAILABLE	PATHOLOGIST: JOHN EIGHMY		WEIGHER: MICHELE ZIMA	
ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	O R G A N S T A T U S
ADRENAL (AD)	----	----	----	NOT TAKEN
BRAIN (BR)	----	----	----	NOT TAKEN
KIDNEY (KD)	----	----	----	NOT TAKEN
LIVER (LI)	7.8851	4.4075 %	----	WEIGHT TAKEN
OVARY (OV)	----	----	----	NOT TAKEN
SPLEEN (SP)	----	----	----	NOT TAKEN
THYMUS (TH)	----	----	----	NOT TAKEN
THYROID/PARATHYR (TT)	----	----	----	NOT TAKEN

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TV), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI), GENERAL COMMENT (GC)

001330

APPENDIX 6
Individual Anatomic Pathology Data

PAGE: 248

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96056		SEX: FEMALE		DOSE GROUP: 4		SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE	
DATE OF DEATH: 12/02/98		STUDY DAY OF DEATH: 93		STUDY WEEK OF DEATH: 14		TERMINAL BODY WEIGHT: 209.3 GRAMS	
DATE AND TIME OF NECROPSY: 12/02/98 9:49		PROSECTOR: TODD ACKER				REORDER: WENDY CHERRY	
POST-FIX WEIGHER: NOT AVAILABLE		PATHOLOGIST: JOHN EIGHMY				WEIGHER: LYNN HORNER	
		ABSOLUTE ORGAN WEIGHT (GRAMS)		ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)		O R G A N S T A T U S	
ORGAN NAME							
ADRENAL (AD)		.0720		.0344 %		WEIGHT TAKEN	
BRAIN (BR)		1.9559		.9345 %		WEIGHT TAKEN	
KIDNEY (KD)		1.9486		.9310 %		WEIGHT TAKEN	
LIVER (LI)		12.3563		5.9036 %		WEIGHT TAKEN	
OVARY (OV)		.1065		.0509 %		WEIGHT TAKEN	
SPLEEN (SP)		.4034		.1927 %		WEIGHT TAKEN	
THYMUS (TH)		.2087		.0997 %		WEIGHT TAKEN	
THYROID/PARATHYR (TT)		.0187		.0089 %		WEIGHT TAKEN	

NECROPSY	OBSERVATIONS	HISTOPATHOLOGY
	LIVER (LI) :	-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -SLIGHT -INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL -NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL -VACUOLATION, HEPATOCELLULAR, CENTRILOBULAR TO MIDZONAL, -MINIMAL
	THYMUS (TH) :	-HEMORRHAGE, -MINIMAL
	UTERUS (UT) :	-DILATATION, -SLIGHT

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

001331

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 249

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96056 SEX: FEMALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 209.3 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 9:49 PROSECTOR: TODD ACKER RECORDER: WENDY CHERRY
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC),
ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), CERVIX (CV), OVARY (OV), VAGINA (VA)

001332

APPENDIX 6

Individual Anatomic Pathology Data

PAGE: 250

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

ANIMAL NUMBER: C96057 SEX: FEMALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 250.5 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:39 PROSECTOR: GREGORY CODDINGTON REORDER: STEVE VAN ADESTEINE
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN S T A T U S
ADRENAL (AD)	.0614	.0245 %	.0280	WEIGHT TAKEN
BRAIN (BR)	2.1920	.8750 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.2779	.9093 %	1.0392	WEIGHT TAKEN
LIVER (LI)	13.2419	5.2862 %	6.0410	WEIGHT TAKEN
OVARY (OV)	.1134	.0453 %	.0517	WEIGHT TAKEN
SPLEEN (SP)	.3561	.1422 %	.1625	WEIGHT TAKEN
THYMUS (TH)	.2704	.1079 %	.1234	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0195	.0078 %	.0089	WEIGHT TAKEN

O B S E R V A T I O N S

NECROPSY

HISTOPATHOLOGY

LIVER (LI) :
-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -SLIGHT
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
-NECROSIS, INDIVIDUAL HEPATOCYTE, -MINIMAL
-PIGMENT, HEPATOCELLULAR, -MINIMAL
KIDNEY (KD) :
-MINERALIZATION, TUBULAR, -MINIMAL
THYMUS (TH) :
-INVOLUTION, -MINIMAL
STOMACH, GL (ST) :
>UNREMARKABLE
UTERUS (UT) :
-DILATATION, -MINIMAL

STOMACH, GL (ST) :
-DARK FOCUS(I)/AREA(S); MUCOSA: MULTIPLE BROWN AREAS, UP TO
3 X 1 MM

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN

001333

APPENDIX 6
Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 251

ANIMAL NUMBER: C96057 SEX: FEMALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 250.5 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:39 PROSECTOR: GREGORY CODDINGTON REORDER: STEVE VAN ADESTINE
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), HEART (HT), MUSCLE, SKELETAL (SM), NERVE, SCIATIC (SN),
SPLEEN (SP), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA),
AORTA (AO), PITUITARY (PI), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL),
CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG),
AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF), URINARY BLADDER (UB), CERVIX (CV), OVARY (OV), VAGINA (VA)

001334

APPENDIX 6

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 252

ANIMAL NUMBER: C96058 SEX: FEMALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 246.7 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 7:55 PROSECTOR: LAURIE J. SCHULLER
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY REORDER: STEVE VAN ADESTINE
WEIGHER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0527	.0214 %	.0253	WEIGHT TAKEN
BRAIN (BR)	2.0848	.8451 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.0803	.8433 %	.9978	WEIGHT TAKEN
LIVER (LI)	11.7380	4.7580 %	5.6303	WEIGHT TAKEN
OVARY (OV)	.0992	.0402 %	.0476	WEIGHT TAKEN
SPLEEN (SP)	.5123	.2077 %	.2457	WEIGHT TAKEN
THYMUS (TH)	.2728	.1106 %	.1309	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0157	.0064 %	.0075	WEIGHT TAKEN

OBSERVATIONS

NECROPSY

HISTOPATHOLOGY

LIVER (LI) :
-HYPERTROPHY, HEPATOCELLULAR, DIFFUSE, -SLIGHT
THYMUS (TH) :
-HEMORRHAGE, -MINIMAL
-INVOLUTION, -MINIMAL
THYROID (TY) :
-CYST, ULTIMOBRANCHIAL, -PRESENT
PANCREAS (PA) :
-INFILTRATE, LYMPHOHISTIOCYTIC, -MINIMAL
LN, MANDIBULAR (MN) :
-HEMORRHAGE, -MINIMAL

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
GENERAL INFORMATION (XX) :
>NOTE:>ANIMAL NECROPSIED OUT OF ORDER DUE TO MORIBUND
CONDITION FOLLOWING BLOOD COLLECTION

601335

APPENDIX 6

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 253

ANIMAL NUMBER: C96058 SEX: FEMALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 246.7 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 7:55 PROSECTOR: LAURIE J. SCHULLER RECORDER: STEVE VAN ADESTINE
POST-FIX WEIGHT: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHTER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TV), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), LN, MESENTERIC (MS), PARATHYROID (PT), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA),
AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL),
CECUM (CE), COLON (CO), RECTUM (RE), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK),
MAMMARY, FEMALE (MF), URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA)

001336

APPENDIX 6
Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 254

ANIMAL NUMBER: C96059 SEX: FEMALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 207.9 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:25 PROSECTOR: JODI THIEREN REORDER: STEVE VAN ADESTINE
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0551	.0265 %	.0275	WEIGHT TAKEN
BRAIN (BR)	2.0062	.9650 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.0576	.9897 %	1.0256	WEIGHT TAKEN
LIVER (LI)	11.4068	5.4867 %	5.6858	WEIGHT TAKEN
OVARY (OV)	.1254	.0603 %	.0625	WEIGHT TAKEN
SPLEEN (SP)	.4387	.2110 %	.2187	WEIGHT TAKEN
THYMUS (TH)	.1931	.0929 %	.0963	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0137	.0066 %	.0068	WEIGHT TAKEN

OBSERVATIONS

NECROPSY

HISTOPATHOLOGY

LIVER (LI) :
-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -MODERATE
-PIGMENT, HEPATOCELLULAR, -MINIMAL
THYROID (TY) :
-CYST, ULTIMOBRANCHIAL, -PRESENT
UTERUS (UT) :
-DILATATION, -SLIGHT

GENERAL COMMENT (GC) :
-BONE MARROW SMEAR TAKEN
-NO MACROSCOPIC LESIONS

APPENDIX 6

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEPOSE, T-6314) IN RATS
PAGE: 255

ANIMAL NUMBER: C96059 SEX: FEMALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 207.9 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 8:25 PROSECTOR: JODI THIEREN REORDER: STEVE VAN ADESTINE
POST-FIX WEIGHER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: TONY HANFELD

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), PARATHYROID (PT), ADRENAL, CORTEX (AC),
ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA),
DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), CERVIX (CV), OVARY (OV), VAGINA (VA)

001338

APPENDIX 6

Individual Anatomic Pathology Data

Covance 6329-225
3M T-6314.113-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS
PAGE: 256

ANIMAL NUMBER: C96060	SEX: FEMALE	DOSE GROUP: 4	SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE	
DATE OF DEATH: 12/02/98	STUDY DAY OF DEATH: 93	STUDY WEEK OF DEATH: 14	TERMINAL BODY WEIGHT: 249.6 GRAMS	
DATE AND TIME OF NECROPSY: 12/02/98 12:19	PROSECTOR: MICHELE ZIMA	REORDER: WENDY CHERRY		
POST-FIX WEIGHER: NOT AVAILABLE	PATHOLOGIST: JOHN EIGHMY	WEIGHER: LYNN HORNER		
ORGAN NAME	ABSOLUTE ORGAN WEIGHT (GRAMS)	ORGAN WEIGHT RELATIVE TO BODY WEIGHT (%)	ORGAN TO BRAIN WEIGHT RATIO	ORGAN STATUS
ADRENAL (AD)	.0718	.0288 %	.0349	WEIGHT TAKEN
BRAIN (BR)	2.0579	.8245 %	1.0000	WEIGHT TAKEN
KIDNEY (KD)	2.5943	1.0394 %	1.2607	WEIGHT TAKEN
LIVER (LI)	13.5240	5.4183 %	6.5717	WEIGHT TAKEN
OVARY (OV)	.1100	.0441 %	.0535	WEIGHT TAKEN
SPLEEN (SP)	.4646	.1861 %	.2258	WEIGHT TAKEN
THYMUS (TH)	.3518	.1409 %	.1710	WEIGHT TAKEN
THYROID/PARATHYR (TT)	.0313	.0125 %	.0152	WEIGHT TAKEN
OBSERVATIONS				
NECROPSY		HISTOPATHOLOGY		
THYROID (TY) :		LIVER (LI) :		
>NOTE:>TRIMMING COMMENT: ONE THYROID/PARATHYROID NOT FOUND IN WET TISSUE AT TRIMMING		-HYPERTROPHY, HEPATOCELLULAR, CENTRILOBULAR, -SLIGHT		
PARATHYROID (PT) :		THYROID (TY) :		
>NOTE:>TRIMMING COMMENT: ONE THYROID/PARATHYROID NOT FOUND IN WET TISSUE AT TRIMMING		-CYST, ULTIMOBRANCHIAL, -PRESENT		
GENERAL COMMENT (GC) :		-ONE EXAMINED, -PRESENT		
-BONE MARROW SMEAR TAKEN		PARATHYROID (PT) :		
-NO MACROSCOPIC LESIONS		-ONE EXAMINED, -PRESENT		

C01339

APPENDIX 6

Covance 6329-225
3M T-6314.1

Individual Anatomic Pathology Data

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL-
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MEFOSE, T-6314) IN RATS

PAGE: 257

ANIMAL NUMBER: C96060 SEX: FEMALE DOSE GROUP: 4 SACRIFICE STATUS: SCHEDULED, TERMINAL SACRIFICE
DATE OF DEATH: 12/02/98 STUDY DAY OF DEATH: 93 STUDY WEEK OF DEATH: 14 TERMINAL BODY WEIGHT: 249.6 GRAMS
DATE AND TIME OF NECROPSY: 12/02/98 12:19 PROSECTOR: MICHELE ZIMA RECORDER: WENDY CHERRY
POST-FIX WEIGHTER: NOT AVAILABLE PATHOLOGIST: JOHN EIGHMY WEIGHER: LYNN HORNER

THE FOLLOWING ORGANS WERE UNREMARKABLE AT NECROPSY:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), LIVER (LI), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), THYROID (TY), PARATHYROID (PT),
ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA), AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU),
PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE), ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN),
SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG), AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF),
URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV), VAGINA (VA), TAIL (TI)

THE FOLLOWING TISSUES WERE UNREMARKABLE AT MICROSCOPIC EXAMINATION:

BONE, FEMUR (FE), MARROW, FEMUR (FM), MARROW, STERNUM (SE), BONE, STERNUM (SB), EYE (EY), BRAIN (BR),
SPINAL CORD (SC), LUNG (LU), TRACHEA (TR), ESOPHAGUS (ES), KIDNEY (KD), HEART (HT), MUSCLE, SKELETAL (SM),
NERVE, SCIATIC (SN), SPLEEN (SP), THYMUS (TH), LN, MESENTERIC (MS), ADRENAL, CORTEX (AC), ADRENAL, MEDULLA (MA),
AORTA (AO), PITUITARY (PI), STOMACH, GL (ST), STOMACH, NONGL (SU), PANCREAS (PA), DUODENUM (DU), JEJUNUM (JE),
ILEUM (IL), CECUM (CE), COLON (CO), RECTUM (RE), LN, MANDIBULAR (MN), SALIV GL, MANDIB (SG), HARDERIAN GLAND (HG),
AUDITORY SEB GL (AS), SKIN (SK), MAMMARY, FEMALE (MF), URINARY BLADDER (UB), UTERUS (UT), CERVIX (CV), OVARY (OV),
VAGINA (VA)

001340

APPENDIX 7

Cell Proliferation Report

Note: This appendix of the report contains information supplied by Pathology Associates International and has been reviewed by the Quality Assurance Unit of Pathology Associates International.

001341

Covance 6329-225
3M T-6314.1



Pathology Associates International
A Company of Science Applications International Corporation



**FINAL
CELL PROLIFERATION REPORT**

**13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL
PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MeFOSE, T-6314) IN RATS
COVANCE STUDY NUMBER 6329-225**

PREPARED FOR:

**3M
TOXICOLOGY SERVICES
BUILDING 220-2E-02, 3M CENTER
ST. PAUL, MN 55144-1000**

PREPARED BY:

**PATHOLOGY ASSOCIATES INTERNATIONAL
15 WORMAN'S MILL COURT, SUITE I
FREDERICK, MD 21701**

MAY 12, 2000

5 Worman's Mill Court, Suite I * Frederick, Maryland 21701 * (301) 663-1644 * (301) 663-8994 FAX

001342

TABLE OF CONTENTS

- I. CELL PROLIFERATION REPORT**
- II. TABLES**
- III. SIGNATURE PAGE**
- IV. QUALITY ASSURANCE STATEMENT**
- V. APPENDIX I**

001343

CELL PROLIFERATION REPORT

13-WEEK DIETARY TOXICITY STUDY WITH N-METHYL PERFLUOROOCTANESULFONAMIDO ETHANOL (N-MeFOSE, T-6314) IN RATS COVANCE STUDY NUMBER 6329-225

PURPOSE

The purpose of the study was to assess the toxicity of the test material when administered in the diet to rats for at least 13 weeks.

This report, submitted by Pathology Associates International (PAI) to the study Sponsor, 3M, represents the cell proliferation findings and interpretation for Covance Study Number 6329-225 entitled "13-Week Dietary Toxicity Study with N-Methyl Perfluorooctanesulfonamido Ethanol (N-MeFOSE, T-6314) In Rats". All aspects of the tasks associated with PAI's portion of this study were conducted in compliance with the Food and Drug Administration Good Laboratory Practice (GLP) Regulations as set forth in Title 21 of the US Code of Federal Regulations, Part 58, issued December 22, 1978 (effective June 20, 1979), and with any applicable amendments.

MATERIALS AND METHODS

Tissue Collection for Cell Proliferation

Five animals per sex per group in groups 1 through 4 were sacrificed during week 13 for hepatocellular proliferation. A section of the left lateral lobe of the liver from each of 5 rats per group (groups 1-4) was fixed and processed to paraffin block by Covance per protocol specifications. Tissue blocks were shipped to PAI for sectioning and staining. From each block, a slide was prepared for H&E evaluation and immunohistochemical detection of proliferating cell nuclear antigen (PCNA), a marker of cell proliferation.

Immunohistochemistry for Cell Proliferation

Sections of paraffin-embedded tissues were cut at 5 μ m and placed on positively charged slides (Superfrost Plus, Fisher Scientific, Pittsburgh, PA) to ensure adhesion during processing for PCNA. Standard immunohistochemical methods were used to stain tissues for PCNA (PAI's SOP for immunohistochemistry). Briefly, tissue sections were incubated with a monoclonal antibody to PCNA (DAKO, lot #016, PAI No. A1723) and reagents required for the avidin-biotin peroxidase (ABC Kit, lot #PK-6100, PAI No. K 324) method for the detection of the antigen-antibody complex. PCNA expression in cells in all phases of the cell cycle (G_1 , S, G_2 and M) was localized by the chromagen 3,3'-diaminobenzidine (DAB; Sigma Chemical Co., lot #18H8201). Tissue sections were counterstained with hematoxylin. A negative control slide was included in the staining run and consisted of study tissue that was not incubated with the primary antibody.

Cell Proliferation Measurements

For cell proliferation evaluations, slides were first perused at low magnification (100X) to judge quality of staining, processing and sectioning, pattern of cell labeling (e.g., centrilobular or panlobular, extrahepatocellular proliferation, such as Kupffer cells, bile duct epithelium, endothelium), and histomorphologic changes. Cell proliferation was then quantified at higher magnification (200X). The percentage of hepatocytes in S phase of the cell cycle (labeling index, LI) was determined by scoring at least 3000 hepatocytes in 10 randomly selected fields per animal. Histomorphology was further assessed by evaluating the serial H&E slide for each animal evaluated for cell proliferation.

Statistical Analysis

The Student's *t*-test (two-sided, unequal variance) was used to test for statistical significance in LI between control and treatment groups using Microsoft Excel version 5.0. A *P* value of less than 0.05 was judged to be statistically significant.

RESULTS

Cell Proliferation

Individual animal and group mean cell proliferation data are presented in Section II (Table 1). The percentage of proliferating hepatocytes, as determined by the labeling index (LI), was not significantly increased in treated groups compared to that of controls in either male or female rats. The LI was significantly less than controls in female rats from groups 3 and 4.

Histopathology

Sections from the same tissue blocks used for preparation of PCNA-stained slides were stained with hematoxylin and eosin (H&E) for histopathologic evaluation to facilitate the interpretation of the immunostained slides. Individual animal findings and group summaries are presented in Appendix I.

The results showed no changes in the liver tissue of male and female rats that would alter the interpretation of the PCNA staining in this study.

001345

SUMMARY

In the present study, cell proliferation was measured within the liver of male and female rats from control (0 ppm), low dose (3 ppm), mid dose (30 ppm) and high dose (100 ppm) after 13 weeks on study. No statistically significant increases in cell proliferation, as determined by the PCNA labeling index (LI) were observed in male or female rats after 13 weeks of N-methyl perfluorooctanesulfonamido ethanol. Although cell proliferation was statistically less than controls in mid dose (30 ppm) and high dose (100 ppm) females, the decrease was not dose-related, and the individual animal values were within the range observed in control animals. Therefore, this decrease was not considered to be biologically relevant.

001346

II. TABLES

Legend

- **SEM = Standard Error of the Mean**

Table II-1. Cell Proliferation in Rat Liver

Dose Group	Sex	Animal Number	PCNA Labeling Index (%)
1 (Control - 0 ppm)	M	C95902	0.387
1 (Control - 0 ppm)	M	C95905	0.246
1 (Control - 0 ppm)	M	C95906	0.227
1 (Control - 0 ppm)	M	C95909	0.153
1 (Control - 0 ppm)	M	C95918	0.583
		Mean	0.319
		SEM	0.076
2 (Low - 3 ppm)	M	C95924	0.190
2 (Low - 3 ppm)	M	C95925	0.208
2 (Low - 3 ppm)	M	C95927	0.235
2 (Low - 3 ppm)	M	C95936	0.063
2 (Low - 3 ppm)	M	C95937	0.224
		Mean	0.184
		SEM	0.031
3 (Mid - 30 ppm)	M	C95942	0.162
3 (Mid - 30 ppm)	M	C95947	0.160
3 (Mid - 30 ppm)	M	C95948	0.186
3 (Mid - 30 ppm)	M	C95950	0.065
3 (Mid - 30 ppm)	M	C95958	0.160
		Mean	0.147
		SEM	0.021
4 (High - 100 ppm)	M	C95962	0.130
4 (High - 100 ppm)	M	C95965	0.163
4 (High - 100 ppm)	M	C95967	0.881
4 (High - 100 ppm)	M	C95968	0.096
4 (High - 100 ppm)	M	C95977	0.162
		Mean	0.286
		SEM	0.149

001348

Table II-1. Cell Proliferation in Rat Liver


Dose Group	Sex	Animal Number	PCNA Labeling Index (%)
1 (Control - 0 ppm)	F	C95985	0.179
1 (Control - 0 ppm)	F	C95987	0.153
1 (Control - 0 ppm)	F	C95988	0.298
1 (Control - 0 ppm)	F	C95997	0.215
1 (Control - 0 ppm)	F	C96000	0.181
		Mean	0.205
		SEM	0.025
2 (Low - 3 ppm)	F	C96002	0.244
2 (Low - 3 ppm)	F	C96003	0.749
2 (Low - 3 ppm)	F	C96006	0.285
2 (Low - 3 ppm)	F	C96008	0.253
2 (Low - 3 ppm)	F	C96010	0.122
		Mean	0.330
		SEM	0.108
3 (Mid - 30 ppm)	F	C96021	0.088
3 (Mid - 30 ppm)	F	C96023	0.063
3 (Mid - 30 ppm)	F	C96026	0.099
3 (Mid - 30 ppm)	F	C96027	0.181
3 (Mid - 30 ppm)	F	C96038	0.062
		Mean	0.099
		SEM	0.022
4 (High - 100 ppm)	F	C96045	0.032
4 (High - 100 ppm)	F	C96046	0.128
4 (High - 100 ppm)	F	C96048	0.121
4 (High - 100 ppm)	F	C96049	0.152
4 (High - 100 ppm)	F	C96056	0.128
		Mean	0.112
		SEM	0.021

001349

III. SIGNATURE PAGE

Submitted by:

Project Manager:


Sandra R. Eldridge, Ph.D.

5-15-00
Date

Project Pathologist:


Carolyn Moyer, D.V.M., Diplomate, A.C.V.P.

5/15/00
Date

001350

IV. QUALITY ASSURANCE STATEMENT

001351



Pathology Associates International

A Company of Science Applications International Corporation

Covance 6329-225

3M T-6314.1

SAIC
An Employee-Owned Company

Cell Proliferation Report

13-Week Dietary Toxicity Study with N-Methyl Perfluorooctanesulfonamido Ethanol (N-MeFOSE, T-6314) in Rats


Covance Study Number: 6329-225

QUALITY ASSURANCE STATEMENT

This cell proliferation project has been inspected and audited by the PAI Quality Assurance Unit (QAU) as required by the Good Laboratory Practice (GLP) regulations promulgated by the U.S. Food and Drug Administration (FDA). The cell proliferation report is an accurate reflection of the recorded data. The following table is a record of the inspections/audits performed and reported by the QAU.

<u>Date of Inspection</u>	<u>Phase Inspected</u>	<u>Date Findings Reported to PAI Management/Project Manager</u>
12/23/98	Microtomy	12/23/98
04/03/00	Study Data and	
	Supporting Documentation	04/03/00
04/03/00	Draft Cell Proliferation Report	04/03/00
05/12/00	Final Cell Proliferation Report	05/12/00


Karen E. Butler
Quality Assurance Officer


Date

C01352

APPENDIX I

C01353

Individual Animal Findings

13-Week Dietary Toxicity Study with N-Methyl
Perfluorooctanesulfonamido Ethanol (N-MeFOSE, T-6314) in Rats

Tissue: Liver

ANIMAL NUMBER	SEX	DOSE GROUP	HISTOLOGIC FINDINGS
C96027	F	3	No Significant Findings
C96026	F	3	No Significant Findings
C96021	F	3	No Significant Findings
C96023	F	3	No Significant Findings
C96038	F	3	No Significant Findings
C95948	M	3	No Significant Findings
C95950	M	3	No Significant Findings
C95942	M	3	No Significant Findings
C95947	M	3	No Significant Findings
C95958	M	3	No Significant Findings
C95965	M	4	No Significant Findings
C95967	M	4	No Significant Findings
C95968	M	4	No Significant Findings
C95962	M	4	No Significant Findings
C95977	M	4	No Significant Findings
C96048	F	4	No Significant Findings
C96045	F	4	No Significant Findings
C96056	F	4	No Significant Findings
C96049	F	4	No Significant Findings
C96046	F	4	No Significant Findings
C95985	F	1	No Significant Findings
C95997	F	1	No Significant Findings
C95988	F	1	No Significant Findings
C95987	F	1	No Significant Findings
C96000	F	1	No Significant Findings
C95909	M	1	No Significant Findings
C95902	M	1	No Significant Findings
C95918	M	1	No Significant Findings
C95905	M	1	No Significant Findings
C95906	M	1	No Significant Findings

Individual Animal Findings

**13-Week Dietary Toxicity Study with N-Methyl
Perfluorooctanesulfonamido Ethanol (N-MeFOSE, T-6314) in Rats**

Tissue: Liver

ANIMAL NUMBER	SEX	DOSE GROUP	HISTOLOGIC FINDINGS
C95925	M	2	No Significant Findings
C95937	M	2	No Significant Findings
C95936	M	2	No Significant Findings
C95924	M	2	No Significant Findings
C95927	M	2	No Significant Findings
C96010	F	2	No Significant Findings
C96003	F	2	No Significant Findings
C96002	F	2	No Significant Findings
C96006	F	2	No Significant Findings
C96008	F	2	No Significant Findings
C96003	F	2	No Significant Findings
C96002	F	2	No Significant Findings
C96006	F	2	No Significant Findings
C96008	F	2	No Significant Findings

180 ON 4/24/00

C01355