

Wilmington, Delaware
February 2, 1972

RAY DILL
F. & P. DEPARTMENT
FAIRFIELD

SOLVENTS CLASSIFIED ACCORDING TO RULE 66

Following is a list of photochemically reactive solvents classified according to the definition in Rule 66 and the write-up given to Connecticut:

1. Class A - maximum 5% by volume of total solvent

isophorone
mesityl oxide
 α -pinene
 β -pinene
cyclohexen-4-one
hexadiene

2. Class B - maximum 8% by volume of total solvent

o-xylene
m-xylene
p-xylene
propylbenzene
phenyl cellosolve
butyl benzene
mesitylene
phenyl acetate

3. Class C - maximum 20% by volume of total solvent

ethylbenzene
trichloroethylene
toluene
diacetone alcohol
diisobutyl ketone
methyl isobutyl ketone
methyl isononyl ketone
2-methyl cyclohexanone
3-methyl cyclohexanone
4-methyl cyclohexanone

4. Exempt Solvents - not controlled under Rule 66

acetone
amyl acetate
amyl alcohol
benzene
butyl alcohol
cellosolve
cellosolve acetate
cyclohexanol
cyclohexanone
diethyl ketone
dioxane
butyl cellosolve
ethyl acetate
ethyl butyl carbinol
2-ethylhexyl acetate
butyl ether
heptane
hexane
methyl acetate
methyl ethyl ketone
methyl propyl ketone
nitropropane
perchloroethylene
propyl acetate
propyl alcohol
tetrahydrofuran
trichlorofluoromethane

If you need more information for the State of Connecticut, please let me know.

WILLIAM W. MILLER
POLLUTION CONTROL COORDINATOR

WWM:ddn