

ACUTE TOXICITY TO FISH (FATHEAD MINNOW)

TEST SUBSTANCE

Identity: Mixture containing perfluorooctanesulfonate which may also be referred to as PFOS, FC-95, or as a component of FC-203FC. (1-Octanesulfonic acid) (CAS # 2795-39-3).

Remarks: The 3M production lot number was not noted. The test sample is FC-203FC, identified by the laboratory as R1335. Current information indicates it is a mixture of 0.85% PFOS, 17% diethylene glycol butyl ether, 74.46% water, 2.34% acrylic foamer, 1.7% alkyl ether sulfate, 2.55% sodium octyl sulfate, 0.12% sodium nitrite, 0.09% sodium molybdate, 0.85% triethanolamine, and 0.04% tolyltriazole.

The following summary applies to a mixture with incompletely characterized concentrations of impurities. Data may not accurately reflect toxicity of the fluorochemical component of the test sample.

METHOD:

Method: OECD Guideline 203.

Type: Acute static

GLP: No

Year completed: 1997

Species: *Pimephales promelas*

Supplier: In-house cultures, maintained at ASci-DETD, Duluth, MN.

Analytical monitoring: Temperature, pH, conductivity, and dissolved oxygen.

Exposure period: 96-hours

Statistical methods: The LC₅₀ values were calculated by Trimmed Spearman-Kärber method. The NOEC values were calculated by Fisher's Exact test using Toxstat statistical software, Version 3.5.

Test fish age: Juveniles

Length and weight: Average length = 16.6 mm
Average weight = 0.036 g (wet)

Loading: 0.12 g/L

Pretreatment: None

Test conditions:

Dilution water: Dechlorinated City of Duluth, MN tap water, post-carbon filter. The water was aerated for 24 hours prior to the test.

Dilution water chemistry:

Hardness: 48 mg/L as CaCO₃

Alkalinity: 44 mg/L as CaCO₃

Conductivity: 119 µmhos/cm

pH: 8.08

Lighting: Cool-white fluorescent lights. A daily photoperiod of 16 hours light and 8 hours dark was maintained throughout the testing period.

Stock and test solution preparation: The test material was added directly to the dilution water on a weight/volume basis.

Concentrations dosing rate: Once.

Stability of the test chemical solutions: Not noted.

Exposure vessels: 4 L glass jars containing 3 L test solutions.

Number of replicates: 2

Number of fish per replicate: 10.

Number of concentrations: five plus a blank control

Water chemistry during the study:

pH range (0-96 hours):

7.68 – 8.08 (control exposure)

7.45 – 7.81 (1000 mg/L exposure)

Temperature range (0-96 hours): 20.5 – 20.7 °C

Conductivity range (0-96 hours):

119 – 141 µmhos/cm (control exposure)

119 – 143 µmhos/cm (1000 mg/L exposure)

Dissolved oxygen range (0-96 hours):

6.9 – 8.7 mg/L (control exposure)

5.1 – 8.7 mg/L (1000 mg/L exposure)

RESULTS

Nominal concentrations: Blank control, 130, 216, 360, 600, and 1000 mg/L

Element values: 96-hour LC_{50} = >1000 mg/L
96-hour NOEC = 600 mg/L

Element values based on nominal concentrations

Remarks: Testing was conducted on the mixture as described in the Test Substance Remarks field. The values reported apply to that mixture and not the fluorochemical proportion alone.

CONCLUSIONS

The FC-203FC 96-hour LC_{50} for fathead minnow was determined to be >1000 mg/L. The 96-hour No Observed Effect Concentration (NOEC) was 600 mg/L.

Submitter: 3M Company, Environmental Laboratory, P.O. Box 33331, St. Paul, Minnesota, 55133

DATA QUALITY

Reliability: Klimisch ranking = 2. The study meets all the criteria for quality testing. However, sample purity was not properly characterized and the study

lacks analytical confirmation of the amount of fluorochemical proportion in the solution.

REFERENCES

This test was conducted by AScl Corporation, Duluth Environmental Testing Division, Duluth, MN at the request of the 3M Company, Lab Request number R1335, 1997.

OTHER

Last changed: 6/26/00

STUDY TITLE

ACUTE TOXICITY OF R1335 TO
FATHEAD MINNOW (*Pimephales promelas*)

DATA STANDARD

OECD GUIDELINE 203

STUDY COMPLETED

June 26, 1997

TESTING FACILITY

ASCI Corporation/ASCI-Duluth
Environmental Testing Division
4444 Airpark Boulevard
Duluth, MN 55811


Tel. No. (218) 722-4040
Fax No. (218) 722-2592

STUDY IDENTIFICATION NUMBERS

ASCI Study ID# 5010-068

3M Company Lab Request# R1335

1.0 GENERAL INFORMATION

Study Title	Acute Toxicity of R1335 to Fathead Minnow (<i>Pimephales promelas</i>).	
Data Standard	OECD Guideline 203.	
Sponsor	3M Environmental Technology and Safety Services, Building 2-3E-09, 935 Bush Avenue, St. Paul, MN 55133; Tel No. (612) 778-7452.	
Sponsor's Representative	Susan Beach, 3M Environmental Technology and Safety Services, Building 2-3E-09, 935 Bush Avenue, St. Paul, MN 55133; Tel No. (612) 778-7452.	
Sponsor's Lab Request#	R1335.	
Testing Facility	ASCI Corporation/ASCI-Duluth Environmental Testing Division, 4444 Airpark Boulevard, Duluth, MN 55811; Tel. No. (218) 722-4040.	
Principal Investigator	Joe Dierkes	
Project Director	Joe Amato	
Testing Facility Director	Donald Mount	
Definitive Test Dates	June 17-21, 1997.	
Data Validation and Report Review	Alan Mozol	Report Signature: 
Retention of Raw Data and Final Report	Two years from the date study is completed.	
Location of Raw Data and Final Report	ASCI Corporation/ASCI-Duluth Environmental Testing Division, 4444 Airpark Boulevard, Duluth, MN 55811; Tel. No. (218) 722-4040.	

2.0 OBJECTIVE(S)

To determine the 96-hour 10% and 50% effect concentrations (LC_{10} and LC_{50}), and the 96-hour no observable effect concentration (NOEC) of the test substance, R1335, for fathead minnow (*Pimephales promelas*) under static test conditions.

This study was conducted according to the ASCI study plan retained by the Sponsor and OECD Guideline 203 (OECD 1993).

3.0 TEST METHODS

Test Substance	<i>Properties:</i> R1335, orange-yellow liquid, specific gravity = 1.1 (water = 1), stable and miscible with water. <i>Test concentrations:</i> 0 (dilution water control), 130, 216, 360, 600, and 1000 mg/L. Test solutions prepared by mass addition of test substance to 6-L aliquots of dilution water followed by vigorous stirring.
Test Organisms	<i>Source:</i> Healthy, juvenile fathead minnows (<i>Pimephales promelas</i>) obtained from ASCI-DETD in-house culture. <i>Holding/acclimation:</i> The fish used in the tests were obtained from ASCI's in-house culture. The fish used in the study were hatched between May 17 and 23, 1997. For the 48 hours preceeding test initiation mortality was 0%, and the fish were not fed for 24 hours before test initiation. The fish mean length was 16.6 ± 2.5 mm, and mean wet weight was 36 ± 19 mg. Loading rate was calculated to be 120 mg of fish per L test solution. <i>Use in test:</i> Used 10 organisms/replicate and 2 replicates/test concentration.
Dilution Water	<i>Source:</i> Dechlorinated City of Duluth, MN, tap water. The most recent chemical analysis is in Raw Data Package. <i>Characteristics:</i> At test time, the water had a hardness of 48 mg/L (as $CaCO_3$) and a pH of 8.08. Aerated for 24 h prior to use in the test.

Exposure Chambers	<i>Type:</i> 4-L glass jars. <i>Use in test:</i> 3.0 L of dilution water or test solution/chamber. During the test, kept covered, except when experimental observations were made. Solutions were not aerated during the exposures.
Incubation	<i>Duration:</i> 96 hours. <i>Daily photoperiod:</i> 16 h light and 8 h dark, maintained using cool white fluorescent lamps. <i>Temperature:</i> 21.0° ± 1°C.
Observations	<i>Biological:</i> Daily- Test organisms with abnormal behavior and signs of stress, and dead test organisms. <i>Water chemistry:</i> (1) At test initiation- total hardness and alkalinity, (2) daily- temperature, DO, conductivity, and pH. All determinations were made according to APHA methods (1995).
Endpoint Calculations	<i>LC₁₀:</i> Inhibition Concentration (ICp) Method. <i>LC₅₀'s:</i> Trimmed Spearman-Kärber Method. <i>NOEC:</i> Fisher's Exact Test (TOXSTAT statistical software, Version 3.5, West Inc., and University of Wyoming).

3.0 RESULTS

Test Substance Toxicity (Pertinent data are in Table 1)	24-h LC_{50}: > 1000 mg/L (Not Calculable). 48-h LC_{50}: > 1000 mg/L (Not Calculable). 72-h LC_{50}: > 1000 mg/L (Not Calculable). 96-h LC_{10}: 116 mg/L (Not Reliable). 96-h LC_{50}: > 1000 mg/L (Not Calculable). 96-h NOEC: 600 mg/L.
Ranges of Water Chemistry Parameters	Total hardness (as $CaCO_3$ mg/L): 46-48. Alkalinity (as $CaCO_3$ mg/L): 44-48. Temperature ($^{\circ}C$): 20.5-20.7. DO (mg/L): 5.1-8.8. pH: 7.45-8.08. Specific conductivity ($\mu mhos/cm$): 119-143.

4.0 TEST QA/QC

QA/QC Criteria	Data
Mortality in control exposure must not be more than 10%.	0% mortality.
The lowest DO concentration at test termination must be at least 60% of the air saturation value at test temperature.	The lowest DO concentration measured was 57% of the air saturation value at 21.0°C. This value was measured at test termination for the 1000 mg/L concentration. This sole value was below 60% saturation. DO depression did not appear to significantly affect the toxicity measurement for R1335.
Test duration must be 96 h.	The test duration was 96 h.
Reference Substance Toxicity Test (this test is optional).	Control chart for sodium chloride reference toxicity testing is in Raw Data Package.

5.0 REFERENCES

American Public Health Association (APHA). 1995. Standard Methods for the Examination of Water and Wastewater. APHA, Washington, D.C.

Organization for Economic Cooperation and Development (OECD). 1981/1993. OECD Guidelines for Testing of Chemicals. OECD Publication Information Center, Washington, D.C.

Hamilton, M.A., R.C. Russo, and R.V. Thurston. 1977. Trimmed Spearman-Kärber Method for Estimating Median Lethal Concentrations in Toxicity Bioassays. Environ. Sci. Technol. 11(7): 714-719; Correction 12 (4): 417 (1978).

USEPA. 1993. A Linear Interpolation Method for Sublethal Toxicity: The Inhibition Concentration (ICp) Approach. Version 2.0.

West Inc., and University of Wyoming. Toxstat Software Package. Version 3.5. Cheyenne, Wyoming.

TABLE 1. ACUTE TOXICITY OF R1335 TO FATHEAD MINNOWS (*P. promelas*)

Test Substance Concentration (mg/L)	Cumulative Number of Dead Organisms and Percent Mortality ^a			
	24 h	48 h	72 h	96 h
0 (control)	0 (0)	0 (0)	0 (0)	0 (0)
130	0 (0)	0 (0)	0 (0)	3 (15)
216	0 (0)	0 (0)	0 (0)	2 (10)
360	0 (0)	0 (0)	0 (0)	2 (10)
600	0 (0)	1 (5)	1 (5)	2 (10)
1000	3 (15)	5 (25)	5 (25)	5 (25)

^a Percentages are in parenthesis.

**LETHAL AND NO-EFFECT CONCENTRATIONS WITH
CORRESPONDING 95% CONFIDENCE INTERVALS**

LC's and NOEC	Data (mg/L) ^a	95 % Confidence Interval
24-h LC ₅₀	> 1000	Not calculable
48-h LC ₅₀	> 1000	Not calculable
72-h LC ₅₀	> 1000	Not calculable
96-h LC ₁₀	116	Not reliable
96-h LC ₅₀	> 1000	Not calculable
96-h NOEC	600	Not applicable

^a Values based on nominal test concentrations.