



BUSHY RUN RESEARCH CENTER

R. D. 4, Mellon Road, Export, Pennsylvania 15632

Telephone (412) 733-5200

CONFIDENTIAL: Not to be released
outside UCC without the written
consent of the UCC-sponsoring
Division HS&EA Manager.

Project Report 47-182.
10 Pages
February 4, 1985

BAKELITE® Cycloaliphatic Epoxy Resin ERL-4221

Salmonella/Microsome (Ames)

Bacterial Mutagenicity Assay

Sponsor: R. E. Plevan, HS&EA Manager
Union Carbide Corporation
Specialty Polymers & Composites Division

Summary

Epoxy Resin ERL-4221 was tested for potential mutagenic activity using the salmonella/microsome bacterial mutagenicity assay (Ames test). Epoxy Resin ERL-4221 was tested with and without metabolic activation in triplicate at five concentrations, ranging from 0.1 to 10 milligrams per plate. The highest concentration was cytotoxic in the mutagenicity tests and in a preliminary test to choose appropriate doses. Positive dose-related increases of the numbers of revertant colonies above the solvent control level were observed only with metabolic activation with the two bacterial strains, TA100 and A1535, that detect base-pair substitution mutagens. Epoxy Resin ERL-4221 was an active mutagen in this in vitro screening test.

Sample

Chemical Name: BAKELITE® Cycloaliphatic Epoxy Resin ERL-4221

I.D. #: TF3-98171 4124 RCP

BRRC Sample #: 47-204

BRRC Account #: 84-18-18049-46

CAS #: 2386-87-0

Chemical Synonyms: 7-Oxabicyclo[4.1.0]heptane-3-carboxylic acid,
7-oxabicyclo[4.1.0]hept-3-ylmethyl ester;
3,4-Epoxy cyclohexyl methyl-3,4-epoxy-cyclohexyl
carboxylate

Molecular Formula: C₁₄H₂₀O₄

Submitted by: R. J. Cotter, Bound Brook, NJ

Sponsor: R. E. Plevan, Danbury, CT
Division: Specialty Polymers & Composites
Bound Brook, NJ

Bushy Run Research Center
A Joint Mellon Institute—Union Carbide Corporation Operation

UCC
062999

BAKELITE® Cycloaliphatic Epoxy Resin ERL-4221Salmonella/Microsome (Ames)Bacterial Mutagenicity Assay

I. Introduction and Theory

The Salmonella typhimurium/microsome mutation assay is a microbial screening test which detects the ability of chemicals to cause genetic alterations in the histidine gene of selected indicator strains of Salmonella typhimurium. The indicator strains used for this test are all histidine-requiring (his-) bacteria which carry a mutation in the histidine locus. The strains were obtained from Dr. Bruce Ames, University of California, Berkeley, CA. Histidine-independent (his+) strains can arise either spontaneously or by the mutagenic action of a chemical or physical agent. Base pair mutagens cause a base change in the DNA molecule at the site of the original mutation or at a second site in the DNA which suppresses the original mutation. Frameshift mutagens cause the addition or deletion of single or multiple base pairs in the DNA molecule. The relative frequency of induced or spontaneous reversion to his+ can be quantitated by plating the indicator strains on minimal agar and counting the number of revertant colonies, representing bacteria that were able to grow and form colonies in the absence of histidine.

To test the mutagenic potential of a test chemical, the bacterial strains are exposed to several concentrations of the chemical. Tests are done both with and without the addition of a mammalian liver homogenate (S9) because some chemicals require metabolic conversion for detection of their biological effects. Concurrent solvent and positive controls are also run with each assay to determine the responsiveness of the test system. After a suitable period of incubation (48-72 hrs), direct revertant colony counts are made and evaluated. Test chemicals which produce at least a 2-fold and dose-related increase in mutant colonies over the concurrent control value are considered to be bacterial mutagens and suspect mammalian mutagens.

II. Salmonella typhimurium Mutagenicity Assay

A. Objective

The objective was to assess the mutagenic potential of Epoxy Resin ERL-4221 as measured in the Salmonella typhimurium/microsome mutagenicity assay. A more complete discussion of the theoretical basis of this test is presented in Appendix 1.

B. Test Chemical (Available analytical and physical data is attached as Appendix 2)

1. Sample name: BAKELITE® Cycloaliphatic Epoxy Resin ERL-4221
2. BRRRC Sample Number: 47-204
3. CAS Number: 2386-87-0
4. Purity: > 95%; IR spectra consistent with structure (see Appendix 2)
5. Density: 1.175
6. Date Received: July 25, 1984
7. Storage Conditions: Room Temperature
8. Storage Location: Chemical Hood/Genetic Toxicology Dept.

C. Control Substances

1. Solvent: dimethylsulfoxide, Burdick & Jackson, Muskegon, MI; Lot #AD912; CAS #67-68-5
2. Positive Controls:
 - a) 4-nitro-o-phenylenediamine; BRRRC #44-71; CAS #99-56-9
 - b) 9-aminoacridine; BRRRC #44-233; CAS #90-45-9
 - c) sodium azide; BRRRC #44-72; CAS #26628-22-8
 - d) 2-aminoanthracene; BRRRC #44-67; CAS #613-13-8

D. Metabolic Activation

S9 liver homogenate, prepared from Aroclor 1254-induced, Sprague-Dawley male rats, was purchased from Microbiological Associates, Bethesda, MD. For tests with metabolic activation, 0.5 ml of S9 mix containing 50 µl of S9 was added per plate.

E. Test Protocol

The assay was performed according to BRRRC Standard Operating Procedures 7.4.1A through 7.4.7A, 7.4.12A, and 7.4.13. A general description of the test methods is attached to this report as Appendix 1.

F. Test Dates

1. Test Initiated: October 29, 1984
2. Test Completed: November 2, 1984

G. Storage Location of Materials

1. Raw Data: BRRC Archives
2. Final Report: BRRC Archives

H. Experimental Design (detailed procedures in Appendix 1)**1. Sample Preparation:**

The test substance was dissolved in dimethylsulfoxide (DMSO) to a concentration of 100 mg/ml for doses of 10 mg/plate and below. All subsequent dilutions were made in the same solvent. Dilutions of the test substance were made fresh each day of testing. All dilutions for the mutagenicity tests were gravimetrically analyzed.

2. Dose Selection:

A preliminary toxicity test was performed using strain TA100 to determine the level of toxicity of the test substance. Ten doses were tested for toxicity with a plate assay performed in the manner used for mutagenicity determinations. Toxicity was assessed at 24 to 48 hours after treatment by either growth inhibition of the background lawn or a reduction in the number of spontaneous mutants.

3. Testing:

The test chemical was tested in triplicate at five doses chosen to span a range which included moderately toxic to relatively nontoxic concentrations. If the substance was nontoxic in the preliminary toxicity test, 100 µl of liquid or 50 mg of solid was used as the maximum dose, unless limited by solubility. Testing was performed both with and without metabolic activation. Concurrent solvent and positive controls were run in each test.

I. Quality Control

Data from each test is checked for accuracy and integrity by a second investigator not involved with the study. Compliance of testing procedures with Good Laboratory Practices is audited by the Independent Quality Assurance department at the Bushy Run Research Center.

III. Results

Epoxy Resin ERL-4221 was tested in a preliminary toxicity screen at 117.5, 30, 10, 3, 1, 0.3, 0.1, 0.03, 0.01, and 0.003 milligrams per plate with strain TA100 only. The top three doses of 117.5, 30, and 10 milligrams per plate completely inhibited growth of the background lawn. Based on these results, mutagenicity testing was done with 5 doses of 10, 3, 1, 0.3, and 0.1 milligrams per plate in triplicate. The highest dose was expected to produce some degree of cytotoxicity based on preliminary toxicity test results, observed as either a reduction in the number of revertant colonies or an inhibition of growth of the background lawn. The range of doses that was tested followed the exposure concentration recommendations of the U.S. E.P.A. Health Effects Test guidelines (HG Gene Muta-S. typhimurium, August 1982).

Gravimetric analysis of the initial dilutions of the test substance for both tests indicated no more than a 6.0% error from the stated concentration. All subsequent dilutions had no more than a 7.2% error from the stated concentrations.

All plate counts and the respective means and standard deviations are shown in Table 1 (without activation) and in Table 2 (with activation). No indication of mutagenicity was observed at any of the tested doses without metabolic activation, either by evidence of a dose-response relationship or a doubling of the number of colonies over the solvent control value. With metabolic activation, dose-related mutagenic activity was observed with strains TA100 and TA1535 which detect base-pair substitution mutagens. The ratios of the average numbers of revertant colonies to the average number observed in solvent control cultures increased with increasing dose as follows: 1.0, 1.1, 1.4, and 2.5-fold increases with TA100; and 1.6, 2.0, 5.7, and 15.3 with TA1535. No evidence of mutagenic activity was observed with strains TA98, TA1537, or TA1538.

All of the observed mutagenic effects of Epoxy Resin ERL-4221 appear to be base-pair substitutions following metabolic activation of the test agent. The mutagenic activity observed only with metabolic activation suggests the production of mutagenically active metabolites from the parent compound by the S9 activation system.


Dose selection appeared to be in a suitable range in the mutagenicity tests because some toxicity was evident in both tests. In both tests, toxicity was observed at 10 mg/plate with all strains.

All strains exhibited a positive mutagenic response with the positive controls tested both with and without S9 metabolic activation. Negative (solvent) controls were also tested with each strain, and the spontaneous reversion rates were within the historical ranges at this laboratory (see Appendix 3). All positive and negative controls were run concurrently with the test chemical. Concurrently run sterility checks showed that the S9 mix, PBS, the test chemical and all solvents and controls were sterile.

IV. Conclusion

Epoxy Resin ERL-4221 produced a dose-dependent mutagenic response with the Salmonella typhimurium strains TA100 and TA1535 when tested with metabolic activation. No positive effects were observed in tests without a metabolic activation system. Under the conditions of this assay, Epoxy Resin ERL-4221 was an active mutagenic in the Salmonella/microsome mutagenicity assay.

Reviewed and Approved by:

 1/3/85
W. Christopher Hengler, M.S., M.B.A. Date
Study Director

 1/2/85
Ronald S. Slesinski, Ph.D. Date
Manager, Genetic, Biochemical & Acute Toxicology

 1/4/85
Fred R. Frank, Ph.D. Date
Director

Contributors:

Technical Assistance: Eileen R. Morabit, A.S.

WPC/rkk/0619B-2
11-16-84

UCC
063004

TABLE 1
RESULTS OF THE SALMONELLA MUTAGENICITY ASSAYCHEMICAL: BRRC #47-204, Epoxy Resin ERL-4221
EXPERIMENT DATE: October 29 through November 1, 1984WITHOUT ACTIVATION
SOLVENT: DMSO

TEST AGENT	DOSE PER PLATE	PLATE COUNTS			AVERAGE	STANDARD DEVIATION
		1	2	3		
STRAIN TA98						
SOLVENT	110 MG	22	29	27	26	3.6
POS: 01	0.01 MG	819	953	884	885	67.0
BRRC#	0.1 MG	28	35	22	28	6.5
47-204	0.3 MG	27	18	22	22	4.5
	1 MG	27	22	23	24	2.6
	3 MG	18	18	24	20	3.5
	10 MG	T	T	T	T	-
STRAIN TA100						
SOLVENT	110 MG	134	138	129	134	4.5
POS: 02	0.01 MG	2112	2340	2179	2210	117.2
BRRC#	0.1 MG	142	131	126	133	8.2
47-204	0.3 MG	139	128	127	131	6.7
	1 MG	126	143	135	135	8.5
	3 MG	136	121	138	132	9.3
	10 MG	T	T	T	T	-
STRAIN TA1535						
SOLVENT	110 MG	46	31	28	35	9.6
POS: 02	0.01 MG	2095	2383	2290	2256	147.0
BRRC#	0.1 MG	57	43	36	45	10.7
47-204	0.3 MG	40	38	43	40	2.5
	1 MG	25	50	43	39	12.9
	3 MG	46	35	36	39	6.1
	10 MG	T	T	T	T	-
STRAIN TA1537						
SOLVENT	110 MG	4	7	4	5	1.7
POS: 03	0.06 MG	78	102	93	91	12.1
BRRC#	0.1 MG	7	5	4	5	1.5
47-204	0.3 MG	5	6	3	5	1.5
	1 MG	6	10	4	7	3.1
	3 MG	9	1	7	6	4.2
	10 MG	T	T	T	T	-
STRAIN TA1538						
SOLVENT	110 MG	10	7	9	9	1.5
POS: 01	0.01 MG	1016	1023	1058	1032	22.5
BRRC#	0.1 MG	9	17	10	12	4.4
47-204	0.3 MG	8	6	8	7	1.2
	1 MG	5	4	6	5	1.0
	3 MG	9	5	6	7	2.1
	10 MG	T	T	T	T	-

T: TOXIC

POS: POSITIVE CONTROLS

01: 4-NITRO-O-PHENYLENEDIAMINE

03: 9-AMINOACRIDINE

02: SODIUM AZIDE

04: 2-AMINOANTHRACENE

3330

Report 47-182
Pag 8TABLE 2
RESULTS OF THE SALMONELLA MUTAGENICITY ASSAYCHEMICAL: BRRC #47-204, Epoxy Resin ERL-4221
EXPERIMENT DATE: October 30 through November 2, 1984WITH ACTIVATION
SOLVENT: DMSO

TEST AGENT	DOSE PER PLATE	PLATE COUNTS			AVERAGE	STANDARD DEVIATION
		1	2	3		
STRAIN TA98						
SOLVENT	110 MG	20	22	22	21	1.2
POS: 04	0.01 MG	1314	1342	1814	1490	280.9
BRRC#	0.1 MG	20	21	18	20	1.5
47-204	0.3 MG	20	16	17	18	2.1
	1 MG	28	29	31	29	1.5
	3 MG	21	39	22	27	10.1
	10 MG	T	T	T	T	-
STRAIN TA100						
SOLVENT	110 MG	113	100	87	100	13.0
POS: 04	0.01 MG	1268	1217	1183	1223	42.8
BRRC#	0.1 MG	97	109	93	100	8.3
47-204	0.3 MG	108	108	126	114	10.4
	1 MG	138	134	148	140	7.2
	3 MG	248	248	263	253	8.7
	10 MG	T	T	T	T	-
STRAIN TA1535						
SOLVENT	110 MG	11	10	7	9	2.1
POS: 04	0.01 MG	40	95	89	75	10.2
BRRC#	0.1 MG	10	19	12	14	4.7
47-204	0.3 MG	12	15	28	18	8.5
	1 MG	55	53	46	51	4.7
	3 MG	145	121	149	138	15.1
	10 MG	T	T	T	T	-
STRAIN TA1537						
SOLVENT	110 MG	5	8	6	6	1.5
POS: 04	0.01 MG	79	32	79	63	27.1
BRRC#	0.1 MG	7	5	2	5	2.5
47-204	0.3 MG	3	3	2	3	0.6
	1 MG	1	5	3	3	2.0
	3 MG	9	5	10	8	2.6
	10 MG	T	T	T	T	-
STRAIN TA1538						
SOLVENT	110 MG	17	7	20	15	6.8
POS: 04	0.01 MG	184	209	226	206	21.1
BRRC#	0.1 MG	18	15	9	14	4.6
47-204	0.3 MG	8	4	8	7	2.3
	1 MG	13	15	19	16	3.1
	3 MG	13	15	14	14	1.0
	10 MG	T	T	T	T	-

T: TOXIC

POS: POSITIVE CONTROLS

01: 4-NITRO-O-PHENYLENEDIAMINE

02: SODIUM AZIDE

03: 9-AMINOACRIDINE

04: 2-AMINOANTHRACENE

3331



BUSHY RUN RESEARCH CENTER

R. D. 4, Mellon Road, Export, Pennsylvania 15632

Telephone (412) 733-5200


Good Laboratory Practices Compliance

This study was conducted in accordance with the U.S. Environmental Protection Agency's Good Laboratory Practices (GLP) Standards (40 CFR Part 792, effective December 29, 1983) with the following deviations:

1. The sponsor indicated that Epoxy Resin ERL-4221 was soluble and stable in dimethylsulfoxide and no additional analyses of stability in the test system were performed.
2. The content of the test substance and control mixtures were documented and analyzed by weight or by volume, and no additional analyses were performed.

In the opinion of the study director, these deviations do not affect the validity or reliability of this study.

Prepared by:


W. Christopher Hengler, M.S., M.B.A. Date
Study Director

WPC/rkk/0619B-1
11-08-84

UCC
063007

**BUSHY RUN RESEARCH CENTER**

R. D. 4, Mellon Road, Export, Pennsylvania 15632

Telephone (412) 733-5200

Quality Assurance Unit Study Inspection SummaryTest Substance: BAKELITE[®] Cycloaliphatic Epoxy Resin ERL-4221

Study: Ames Bacterial Mutagenicity Assay

Study Director: W. C. Hengler, M.S., M.B.A.

The Quality Assurance Unit of BRRC conducted the inspections listed below and reported the results to the study director and to management on the dates indicated. It is the practice of this Quality Assurance Unit to report the results of each inspection to both the study director and management.

<u>Date</u>	<u>Inspection Type</u>	<u>Date QAU Report Issued</u>	
		<u>To Study Director</u>	<u>To Management</u>
1-31-84	Standard Protocol	2-2-84	2-3-84
10-3-84	Standard Protocol Form	10-3-84	10-3-84
10-30-84	Event-Preparing Agar	11-2-84	11-28-84
1-15 to 1-18-85	Final Data and Final Report	1-18-85	2-1-85

UCC
063008

Linda J. Calisti 2/4/85
Linda J. Calisti, Group Leader Date
Good Laboratory Practices/Quality Assurance

APPENDIX 1

Standard Procedures for the Salmonella/microsome Mutagenicity Assay

A. Introduction

The strains used in this test are all Salmonella typhimurium histidine auxotrophs developed by Dr. Bruce Ames of the University of California at Berkeley. The test is designed to assay for reverse mutations in the histidine locus by plating the bacteria on minimal agar with only a trace of histidine. Bacteria that have undergone a reverse mutation in this locus can form colonies in minimal agar and are counted as mutants. The trace of histidine in the minimal agar allows the bacteria to undergo a small number of cell divisions which then form the background lawn, visible as a cloudiness in the agar. Observation of the lawn permits an estimate of toxicity. Testing with growing cultures also allows potential mutagens to act on replicating DNA which is often a more sensitive test for several classes of mutagens.

B. Indicator Organisms

Salmonella typhimurium strains TA98, TA100, TA1535, TA1537, and TA1538 are used in this assay. They all lack the excision repair mechanism, which increases their sensitivity to potential mutagens. They also have a mutation affecting their lipopolysaccharide layer which allows large molecules to enter the bacteria more easily. Strains TA98 and TA100 were derived from strains TA1538 and TA1535 by the addition of an R factor which imparts ampicillin resistance and also increases sensitivity to several classes of mutagens.

Indicator organisms are stored at -80°C. Working cultures are prepared monthly by inoculating nutrient broth from the frozen cultures and incubating with agitation overnight. Bacteria are then plated onto Vogel-Bonner Medium E agar plates (master plates) with an excess of histidine and biotin (required because of the lipopolysaccharide deficiency). After incubation for 24 hours, the strains are checked for their genetic markers to verify their identity and purity (SOP #7.4.2A).

For testing, the broth cultures are prepared by inoculating from the master plates into nutrient broth and incubated overnight with agitation. The broth cultures are then kept on ice during the day of testing. Fresh cultures are made each day of testing.

C. Toxicity Testing

Sterile tubes are prepared containing 2 ml soft agar (6 g/l agar and 5 g/l NaCl) with a final concentration of 0.05 mM L-histidine and 0.05 mM D-biotin (the entire mixture is called top agar). Dilutions of the test chemical are made in an appropriate solvent so that the correct amount of chemical can be added to each tube in 100 µl amounts. If less than 100 µl amounts of the test chemical is used, phosphate-buffered saline is used to adjust the total volume to 100 µl.

UCC
063009

At least five doses are tested with maximum doses of 50 mg (for solids) or 100 μ l (for liquids) per plate for nontoxic chemicals, unless limited by solubility. Generally, a chemical cannot be considered to be nonmutagenic unless at least 5 mg/plate has been tested, unless limited by toxicity or solubility (deSerres and Shelby, 1979; HG-Gene Muta-S. typhimurium, August 1982). A 100 μ l aliquot of an overnight broth culture of strain TA100 is added followed by the appropriate amount of test chemical. The mixture is vortexed and poured onto the surface of a Vogel-Bonner Medium E agar plate (VB-E plate). The top agar is allowed to harden and the plates are incubated at 37°C for 24 to 48 hours. The plates are examined for the condition of their background lawns and growth is recorded as either confluent, sparse, or absent. Confluency is considered an indication of nontoxicity, sparse growth indicates moderate toxicity, and lack of growth is recorded as toxic.

D. Dose Selection

For the mutagenicity test, a minimum of five half-log doses are tested within a range of moderately toxic to relatively nontoxic doses based on the results of the toxicity test. If all doses in the toxicity test were nontoxic, then a maximum dose of 50 mg or 100 μ l is used in the mutagenicity test, unless limited by solubility.

E. Controls

Concurrent solvent and positive controls are run with each test. The solvent control is the maximum amount of solvent added with the diluted test chemical. For chemicals tested without dilution (i.e., by direct addition), water is used as the solvent control. Both activation-dependent and activation-independent positive controls are used. The activation-independent controls are 4-nitro-o-phenylenediamine for TA98 and TA1538, sodium azide for TA100 and TA1535, and 9-aminoacridine for TA1537. The activation-dependent control is 2-aminoanthracene (2-anthracene) for all strains. The concentrations are determined from prior dose-response experiments on these chemicals and are listed in the tables of results in this report.

F. Metabolic Activation

Each lot of Aroclor-1254 induced, rat-liver homogenate (S9) is prescreened for activity with 7,12-dimethylbenzanthracene to determine the appropriate amount to use for testing. The S9 is added to the "S9 mix" which contains, per ml, S9 (30-250 μ l), $MgCl_2$ (8 μ moles), KCl (33 μ moles), glucose-6-phosphate (5 μ moles), NADP (4 μ moles), and sodium phosphate, pH7.4 (100 μ moles). S9 mix is prepared fresh each day of testing and kept at 0-4°C.

G. Test Chemicals and Solvents

The solvents of choice are water, DMSO or ethanol. A minimum of five concentrations of the test chemical are diluted in the appropriate solvent and tested. The chemical is diluted so that 100 μ l will deliver the required

Report 47-182

Appendix 1

Page 3 of 4

dose. If smaller volumes are used, phosphate-buffered saline is added to adjust the total volume to 100 μ l. Test chemical solutions are prepared on the day of testing and are gravimetrically analyzed. If the analysis indicates > 10% error from the intended concentration, then either the dilution is repeated or the actual concentration is stated in the report.

H. Dosing

To a sterile tube containing 2 ml of top agar, 100 μ l aliquot of the appropriate bacterial culture is added followed by the addition of 100 μ l of the appropriate solvent, control, or test chemical solution. Either 0.5 ml of S9 mix or 0.5 ml of phosphate-buffered saline (PBS) is added for tests with and without metabolic activation, respectively. The top agar mixture is then poured onto a VB-E plate. Each dose is tested in triplicate and with all five bacterial strains. Sterility checks are done on the PBS and/or S9 mix, all solvents, and the highest concentration of each test chemical. The plates are transferred to a darkened 37°C incubator after hardening and incubated for 48-72 hours.

I. Data Collection

An Artek Model #880 Colony Counter is used to count bacterial colonies or handcounts of the plates are done. The counter is calibrated for each test to check the counting accuracy (SOP #7.4.5A). The numbers of colonies per plate are counted and recorded. An examination is also made of the background lawn on each plate. If toxicity is observed as an inhibition of growth of the background lawn, the plate is not counted, but is recorded as toxic. If the background lawn is sparse and the colony count is still recorded, that number is not used in the calculation of the mean number of colonies and its standard deviation.

J. Criteria for Test Validity and for Interpretation of Results

The spontaneous reversion for the solvent controls should be within this laboratory's historical range. The positive controls should demonstrate that the test systems are responsive with known mutagens. A test chemical is considered to be a bacterial mutagen if the number of revertant colonies is at least twice the solvent control for at least one dose level and there is evidence of a dose-related increase in the number of revertant colonies. If a test chemical produces a marginal or weak response that cannot be reproduced in a second test, the test result will be considered negative. If there is no evidence of a dose-related increase in the number of revertant colonies and the number of revertant colonies is not twice the solvent control, then the test chemical is not considered to be a bacterial mutagen.

K. Data Storage and Retrieval

Copies of the final report, statistical analyses, analytical data and raw data are stored in the BRRC Archives.

UCC
063011

References

1. Ames, B.N., J. McCann and F. Yamasaki, Methods for detecting carcinogens and mutagens with the Salmonella/mammalian-microsome mutagenicity test, Mutation Research, 31: 347-364 (1975).
2. Vogel, H.J., and D.M. Bonner, Acetylornithase of Escherichia coli: partial purification and some properties, Journal of Biological Chemistry, 218: 97-106 (1956).
3. deSerres, F.D., and M.D. Shelby, Recommendations on data production and analysis using the Salmonella/microsome mutagenicity assay, Mutation Research, 64: 159-165 (1979).
4. Environmental Protection Agency, Health Effects Test Guidelines, HG-Gene Muta-S. typimurium, EPA Report No. 560/6-82-001, August, 1982.

WPC/rkk/0619B-1
11-08-84UCC
063012

REQUEST FOR STANDARD TOXICOLOGY STUDIES
AT BUSHY RUN RESEARCH CENTERH. H.
L. H.BUSINESS
CONFIDENTIAL

To: Mr. R.C. Myers
Bushy Run Research Center
RD 4 Mellon Road
Export, PA 15632

From: Name R. J. Cotton

Sponsoring Division SECURITY POLICE & COMBAT SITES

Dept. R + D

Location (city) GALE PA

Charge to Account No. 25-506-9440

Technology Manager G. T. Kuhlthorn, Jr.

Study

Authorization: R. J. Cotton 12 July 1984

Signature

Date

COMPLETE COPIES OF THIS FORM
MUST BE SENT (BY THE SUBMITTER) TO:

- A. Dr. Bryan Ballantyne
Technical Center (3005)
South Charleston, WV 25303
- B. Dr. W.C. Kuryla
Technical Center (3005)
South Charleston, WV 25303
- C. Dr. E.R. Homan
Bushy Run Research Center
RD 4 Mellon Road
Export, PA 15632
- D. Sponsoring Division
HS & EA Manager
Name: RE BLANK
Loc.: PA 15632

TESTS REQUESTED (Standard procedures; other tests require specific protocols. Please contact the Corporate Applied Toxicology Dept. for advice and guidance.)

RECEIVED

1 RANGE FINDING:

- ☐ Acute Oral ☐ Other _____
- ☐ Acute Percutaneous
- ☐ Acute Inhalation
- ☐ Skin Irritancy
- ☐ Eye Irritancy

3. IN VITRO:

- ☒ Ames (Salmonella) Test
- ☐ Gene Mutation (CHO)
- ☐ Chromosome Exchange (SCE)
- ☐ Chromosome Damage (Micronucleus)
- ☐ DNA Damage (UDS)
- ☐ Macrophage (RAM) test
- ☐ Other _____

AUG 24 1984

W. C. HENGLER

2 KNOWN PROCEDURAL REQUIREMENT (circle):

OECD, FHSA, DOT, FIFRA, TSCA, RCRA, Other _____



PLEASE PROVIDE THE FOLLOWING INFORMATION. IT IS NEEDED IN ORDER TO EXPEDITE WORK ON THE REQUESTED STUDIES.

PRODUCT/COMPOSITION NAME: ECL-4221ACS NOMENCLATURE: 7-OXABICYCLO(4.1.0)HEPTANE-3-CARBOXYLIC ACID, 7-OXABICYCLO(4.1.0)HEPT-3-YL METHYL ESTERCAS NO.: 2386-87-0

CHEMICAL SYNONYMS/ALTERNATE/TRADE NAMES:

MAJESTIC CY-179 (3,4-EPOXYCYCLOHEXYLMETHYL 3,4-EPOXYCYCLOHEXYLMETHANECARBOXYLATE)

STRUCTURAL FORMULA OR COMPOSITION OF MIXTURES:



DISTRIBUTION OF REPORT: A standard divisional distribution list will be used unless otherwise requested (see page 4).

FOR BRRC USE:

ACKNOWLEDGEMENT OF SAMPLE RECEIPT

Date Received: July 25, 1984By: T. H. HenglerBRRC Sample No. 117-204UCC
063013

CHEMICAL INFORMATION

1. SAMPLE PURITY/COMPOSITION: *Commercial product. Sample is probably at least 95% of the indicated structure. Impurities would include dimer and acetate derivatives from reaction in acetic acid.*
- Analytical test method employed* _____
- *(Send copy of available data, source and location of analyses.)

Percent Composition

Amounts and identities of known impurities: _____ %

Amount of unidentified impurities: _____ %

Test material: _____ %

2. REQUIRED SAMPLE QUANTITY:Acute Tests:

- 32 ounces for full series of tests
- 16 ounces for chemicals with known high toxicity
- 2 ounces for irritancy tests only

In vitro Tests: 4 ounces

NOTE: To reduce problems involving disposal of unused amounts of the sample, supply only the amount specified above3. PHYSICAL CHARACTERISTICS: (Give all which are known):Gas _____ Liquid ☒ Solid _____ Powder _____ Granular _____Molecular Wt. 252

Vapor Pressure (@ 3 temp. between 15° and 150°C)

Density (liquids) 1.175 (25/25°)1. <0.1 mm @ 20°C °C

pH (aqueous sol.) _____

2. _____ mm @ _____ °C

Freezing point -20 °C °C

3. _____ mm @ _____ °C

Boiling point 354°C / 760 mm °CFlash point 118 °C

(Tag, closed cup)

4. SOLUBILITY:

Solvent _____ Percent Solution _____

Water *7.14 g/100 ml at 25°C* 0.05

Acetone _____

Ethanol _____

Dimethylsulfoxide _____

Vegetable Oil _____

Other (specify) _____

5. STABILITY: (Check appropriately):Shelf Life _____ < 6 mo; ☒ > 6 mo.Light sensitive _____ yes; ☒ noTemp. sensitive _____ yes; ☒ noOxidizes _____ yes; ☒ noPolymerizes* ☒ yes; _____ no

Known solvent _____

Incompatibility (specify) _____

HANDLING/STORAGE/DISPOSAL INFORMATION:

1. RECOMMENDED PRECAUTIONS FOR HANDLING*: KEEP AWAY FROM SKIN AND EYES

*Enclose UCC (or other company) Material Safety Data Sheet - if available.

Any unusual fire or explosion hazards: NO

Any chemical incompatibilities (materials to avoid): STRONG ACIDS AND BASES

2. SPECIAL PERSONNEL PROTECTIVE EQUIPMENT NEEDED: WASH EXPOSED AREAS
WITH SOAP AND WATER

3. RECOMMENDED INACTIVATION/DECONTAMINATION PROCEDURE(S):

4. RECOMMENDED STORAGE CONDITIONS:

Temperature (Check appropriately):

Room Temp. ☒

Refrigerate ☐ < 20°C

Freeze ☐ < 0°C

Special Conditions Required (Check appropriately):

Under N₂ gas ☐

Dessicator ☐

Other ☐

5. SAMPLE DISPOSAL: (Check appropriately)

☐ Return unused sample to:
(attach DOT shipping instructions)

☒ Disposal by BRRC is authorized in compliance with applicable regulations.

UCC
063015

OTHER PERTINENT INFORMATION

The following information will be useful in determining or confirming the necessity of specific toxicity tests:

1. Proposed use of material (circle): Industrial Intermediate Home Use Experimental

Industrial Product Food Drug Cosmetic Packaging (specify Food, Drug, Cosmetic)

Other use (specify): _____

2. Potential hazard route(s) (circle): Skin Inhalation Mouth Eye

Other (specify): _____

3. Form shipped in commerce (circle): Solid Liquid Gas Suspension or Slurry

Solvent or Vehicle composition: _____

ATTACHMENTS (TO BE PROVIDED BY SUBMITTER):

- * - Analytical data for sample.
- * - Material Safety Data Sheet of sample material (if available).
- * - DOT shipping information (if applicable).

DISTRIBUTION LIST: Unless indicated below, the standard distribution list for the sponsoring division will be used.

☐ Please use the following divisional distribution list for the final study report:



MATERIAL SAFETY DATA SHEET

EFFECTIVE DATE: January 1, 1982

Report 47-182 3341
Appendix 2
Page 5 of 9

I. IDENTIFICATION

PRODUCT NAME: BAKELITE Cycloaliphatic Epoxy Resin ERL-4221, ERL-4221G
CHEMICAL NAME: 3,4-Epoxy cyclohexyl methyl-
3,4-Epoxy cyclohexyl carboxylate
FORMULA: $C_{24}H_{20}O_4$ CHEMICAL FAMILY: Diepoxide
MOLECULAR WEIGHT: 252.30
SYNONYMS: --
DEPARTMENT OF HAZARD CLASSIFICATION None
TRANSPORTATION SHIPPING NAME None
CAS # 2386-87-0 CAS NAME 7-Oxabicyclo[4.1.0]heptane-3-carboxylic acid, 7-oxabicyclo[4.1.0]hept-3-ylmethyl ester

II. PHYSICAL DATA

BOILING POINT, 760 mm Hg 101.325 kPa	354 °C (669 °F) Extrapolated	FREEZING POINT	Sets to glass below -20 °C (-4 °F)
SPECIFIC GRAVITY (H ₂ O = 1)	1.175 at 25/25 °C	VAPOR PRESSURE at 20 °C	< 0.001 kPa < 0.01 mm Hg
VAPOR DENSITY (air = 1)	8.7	SOLUBILITY IN WATER, % by wt. at 20°	0.05
PER CENT VOLATILES BY VOLUME	Nil	EVAPORATION RATE (Butyl Acetate = 1)	< 0.01
APPEARANCE AND ODOR	Low viscosity liquid; characteristic odor.		

III. INGREDIENTS

MATERIAL	%	TLV (Units)	HAZARD
BAKELITE Epoxy Resin ERL-4221, ERL-4221G	100	Not established	Slight skin and eye irritant

IV. FIRE AND EXPLOSION HAZARD DATA

FLASH POINT (test method(s))	245° F, Pensky-Martens Closed Cup 400° F, Cleveland Open Cup	
FLAMMABLE LIMITS IN AIR, % by volume	LOWER Not determined (Nonvolatile Fluid)	UPPER Not determined (Nonvolatile Fluid)
EXTINGUISHING MEDIA	Use water spray, carbon dioxide, dry chemical, alcohol-type or universal-type foams applied by manufacturer's recommended technique.	
SPECIAL FIRE FIGHTING PROCEDURES	Don't spray pool fires directly, a solid stream of water directed into hot burning liquid can cause frothing. Use supplied breathing air and protective clothing.	
UNUSUAL FIRE AND EXPLOSION HAZARDS	Hazardous polymerization may occur, due to heat from exposure to a fire.	

EMERGENCY PHONE NUMBER

304/744-3487

This number is available days, nights, weekends, and holidays

UCC
063017

While Union Carbide Corporation believes that the data contained herein are factual and the opinions expressed are those of qualified experts regarding the results of the tests conducted, the data are not to be taken as a warranty or representation for which Union Carbide Corporation assumes legal responsibility. They are offered solely for your consideration, investigation, and verification. Any use of these data and information must be determined by the user to be in accordance with applicable Federal, State, and local laws and regulations.

UNION CARBIDE CORPORATION • SPECIALTY CHEMICALS AND PLASTICS DIVISION
OLD RIDGEBURY ROAD, DANBURY, CT 06817

3342

Report 47-182
Appendix 2
Page 6 of 9

V. HEALTH HAZARD DATA

TLV AND SOURCE: None established by ACGIH or OSHA.

ACUTE EFFECTS OF OVEREXPOSURE

SWALLOWING	Moderately toxic. May cause nausea, vomiting, and diarrhea.
SKIN ABSORPTION	None currently known.
INHALATION	None currently known.
SKIN CONTACT	May cause irritation.
EYE CONTACT	May cause irritation.
CHRONIC EFFECTS OF OVEREXPOSURE	None currently known.
OTHER HEALTH HAZARDS	None currently known.

EMERGENCY AND FIRST AID PROCEDURES:

SWALLOWING	Give two glasses of water and induce vomiting by putting finger down throat.
SKIN	Wash with soap and water.
INHALATION	No emergency care anticipated.
EYES	Flush with water.

NOTES TO PHYSICIAN

There is no specific antidote. Treatment of overexposure should be directed at the control of symptoms and the clinical condition.

UCC
063018

VI. REACTIVITY DATA

STABILITY		CONDITIONS TO AVOID	None
UNSTABLE	STABLE		
--	✓		
INCOMPATIBILITY (materials to avoid)		Acids, strong bases, amines.	
HAZARDOUS COMBUSTION OR DECOMPOSITION PRODUCTS		Burning can produce CO and/or CO ₂ .	
HAZARDOUS POLYMERIZATION		CONDITIONS TO AVOID	Acids, strong bases, amines.
May Occur	Will not Occur		
✓	--		

VII. SPILL OR LEAK PROCEDURES

STEPS TO BE TAKEN IF MATERIAL IS RELEASED OR SPILLED	Collect for disposal. Avoid discharge to waterways; see Section IX.
WASTE DISPOSAL METHOD	Dilute with an inert solvent and incinerate in a furnace where permitted under appropriate Federal, State, and local regulations.

VIII. SPECIAL PROTECTION INFORMATION

RESPIRATORY PROTECTION (specify type)	None required in normal use.		
VENTILATION	General (mechanical) room ventilation is expected to be satisfactory.		
PROTECTIVE GLOVES	Plastic	EYE PROTECTION	Safety glasses
OTHER PROTECTIVE EQUIPMENT	Eye bath and/or safety shower.		

IX. SPECIAL PRECAUTIONS

PRECAUTIONS TO BE TAKEN IN HANDLING AND STORING

Avoid contact with eyes, skin, and clothing.
Wash thoroughly after handling.
Do not take internally.

FOR INDUSTRY USE ONLY

OTHER PRECAUTIONS

This product is essentially insoluble in water.
It is resistant to rapid biodegradation and may be toxic to aquatic life. Avoid discharge to natural waters.

UCC
063019

3344



UNION CARBIDE CORPORATION
SPECIALTY POLYMERS & COMPOSITES DIVISION

Report 47-182
Appendix 2
Page 8 of 9

P.O. BOX 570, BOUND BROOK, N.J. 08805
TELEPHONE (201) 326-8000

RECEIVED

SEP 12 1984

W.C. HENGLER

September 6, 1984

Mr. W. C. Hengler
Bushy Run Technical Center
RD4, Mellon Road
Export, Pennsylvania 15632

Dear Mr. Hengler:

Infrared spectra have been run on the samples of cycloaliphatic epoxides that you returned to me recently. The spectra are enclosed for your files. The spectra that were obtained on these samples were compared with standard spectra for these products. In two cases, 4205 and 4299, comparisons were made with other samples of the these products that are being utilized in current laboratory programs.

Comparison of these various spectra by the infrared experts in our Analytical Group (Dr. Clark and Mr. Lewis) led to the conclusion that the samples that you have are authentic samples of the UCC cycloaliphatic epoxides. Thus, the mutagenicity testing will be performed on valid samples of 4221, 4299, 4234, 4206 and 4205.

Very truly yours,


encls.

A handwritten signature in dark ink, appearing to read "R. J. Cotter".
R. J. Cotter

RJC:mjk

cc:Dr. E. M. Clark, Mr. J. Lewis, Dr. R. E. Plevan - DB

UCC
063020



ABSCISSA		ORDINATE		SCAN TIME		REP. SCAN		SINGLE BEAM	
EXPANSION		EXPANSION		MULTIPLIER		TIME DRIVE			
		% T		SLIT PROGRAM		OPERATOR		DATE: 3-8-61	
SAMPLE: ERL-4321		REMARKS		SOLVENT		CELL PATH			
ORIGIN: HUGO from Body				CONCENTRATION		REFERENCE			
HOW: 16.72-16.75 cm									

UCC
063021

APPENDIX 3

Salmonella typhimuriumA. Historical Data on Spontaneous Reversion Rates at BRRC
As of June 30, 1984

<u>Strain</u>	<u>+/-S9</u>	<u>N</u>	<u>Mean</u>	<u>Standard Deviation</u>	<u>Median</u>	<u>Range</u>
TA98	-S9	98	18.2	5.02	18	7, 33
	+S9	98	25.6	7.40	26	14, 44
TA100	-S9	98	123.1	26.89	120	61, 187
	+S9	98	101.5	17.86	100	65, 137
TA1535	-S9	98	32.5	13.61	32	7, 60
	+S9	98	13.4	7.34	14	2, 35
TA1537	-S9	98	4.3	1.90	4	1, 9
	+S9	98	5.0	2.61	5	1, 13
TA1538	-S9	98	7.2	3.72	7	2, 19
	+S9	98	15.0	5.41	16	4, 29

B. Historical Data on Positive Control Values at BRRC
As of January 26, 1984

<u>Strain</u>	<u>Positive Control</u>	<u>+/-S9</u>	<u>N</u>	<u>Mean</u>	<u>Standard Deviation</u>	<u>Median</u>	<u>Range</u>
TA98	NPD, 10 µg	-S9	84	877.1	166.73	870	532, 1301
	2AA, 10 µg	+S9	84	1434.2	327.54	1457	604, 2037
TA100	NZ, 10 µg	-S9	84	1621.8	277.37	1696	869, 2069
	2AA, 10 µg	+S9	84	1181.7	323.54	1201	475, 2225
TA1535	NZ, 10 µg	-S9	84	1622.0	253.42	1666	837, 2050
	2AA, 10 µg	+S9	84	99.3	30.84	98	37, 182
TA1537	9AA, 60 µg	-S9	84	140.6	130.70	82	28, 489
	2AA, 10 µg	+S9	84	122.5	45.52	120	36, 230
TA1538	NPD, 10 µg	-S9	84	1124.2	196.66	1168	573, 1582
	2AA, 10 µg	+S9	84	532.0	230.61	486	146, 1249

S9- rat liver homogenate; NPD- 4-nitrophenylenediamine; 2AA- 2-aminoanthracene;
NZ- sodium azide; 9AA- 9-aminoacridine.