

effect was not
on MC, which
-a- acridine

SCE induction
ever, the differ-
taken into con-
yte test system

lying the acri-
of the results.
technical assis-

er Programme

of sub-populations

G1-prophase of the
Hum. Genet., 35

elds in the human

and kinetics in

n culture measured

unds, XII. Biologic
s, Arch. Immunol.

. A search for anti-
aminoacridines and

chromatids, Nature

nosome aberration
st., 9 (1977) 289-

f phytohaemagglu-
7-376,
anges of restricted

biting compounds,
51 (1977) 1725-

Mutation Research, 67 (1979) 97-100
© Elsevier/North-Holland Biomedical Press

Short Communication

COMPARATIVE EVALUATION OF THE FREQUENCY OF CHROMOSOMAL ABERRATIONS AND THE SCE NUMBERS IN PERIPHERAL LYMPHOCYTES OF WORKERS OCCUPATIONALLY EXPOSED TO VINYL CHLORIDE MONOMER

M. KUČEROVÁ, Z. POLÍVKOVÁ and J. BÁTORA *

*Pediatric Department of Postgraduate Medical Institute, Institute of Hygiene and
Epidemiology, Prague, Šrobárova 48 (Czechoslovakia)*

(Received 31 March 1978)

(Revision received 15 November 1978)

(Accepted 23 November 1978)

There are now available plenty of authoritative data on the carcinogenic [1-3, 13] and mutagenic [1,2,10] activity of vinyl chloride monomer. Increased frequency of chromosomal aberrations detectable in peripheral lymphocytes of chronically exposed workers has been reported by numerous authors [4-6, 11,12].

We have repeatedly analysed, using routine cytogenetic techniques [8], 3 blood samples from each of 9 workers occupationally exposed for 10-27 years to relatively high mean annual doses of vinyl chloride monomer (about 20-150 ppm of air). In the third blood sample series we also used the SCE technique [14] and compared the frequency of detected chromosomal aberrations with the number of SCEs in each sample. The blood samples from 8 healthy persons were used as controls and were collected with the 3rd blood sampling from the vinyl chloride (VC) workers. The control persons were matched for age and sex and they were not exposed to known mutagens during 3 months before the time of blood collection.

The results are summarized in three tables. Tables 1 and 2 show the instability and non-homogeneity in the frequency of chromosomal aberrations over a period of two years. Mostly chromatid and chromosome breaks were detected; chromatid and chromosome exchanges occurred only sporadically.

Table 2 compares the frequency of chromosomal aberrations and the number of SCE per cell in 7 blood samples. There is an apparent increase in both parameters in comparison with control levels.

The mathematical evaluation of our results was calculated only for the 3rd collection. Conventional aberrations in VC workers were non-homogeneous.

* District Hygienic Station, Prievidza.

TABLE 1

CHROMOSOMAL ANALYSES OF 9 WORKERS OCCUPATIONALLY EXPOSED TO VINYL CHLORIDE (TYPES OF ABERRATION)

Exposure (years)	Aberrant cells (%)	Number of chromatid breaks /100 cells	Number of chromatid exchanges /100 cells	Number of chromosome breaks /100 cells	Number of chromosome exchanges /100 cells	Number of despiralizations /100 cells
E.H.	1 11	5	1	4	0	1
25-27	2 0	0	0	0	0	0
	3 11	5	0	4	3	0
J.P.	1 4	4	0	0	0	0
16-18	2 2	0	0	1	1	0
	3 2	1	0	0	1	0
K.K.	1 2	2	0	0	0	0
15-17	2 7	6	1	0	0	0
	3 3	3	0	0	0	0
A.H.	1 1	1	0	0	0	0
14-16	2 1	1	0	0	0	0
	3 6	2	0	4	0	0
M.M.	1 0	0	0	0	0	0
13-15	2 1	0	0	0	1	0
	3 7	3	0	4	0	0
E.J.	1 3	1	1	0	1	0
12-14	2 6	4	0	2	0	0
	3 11	2	1	8	0	0
J.K.	1 1	0	0	1	0	0
12-14	2 4	3	0	1	0	0
	3 3	3	0	0	0	0
F.M.	1 1	1	0	0	0	0
11-13	2 2	1	0	1	0	0
	3 1	1	0	0	0	0
A.T.	1 4	2	0	1	0	1
10-12	2 1	0	0	1	0	0
	3 3	0	0	3	0	0
8 controls (mean)	1.8	1	0	0.8	0.2	0

The frequency of SCEs was significantly higher in cells of VC workers in comparison with controls. The level of significance was $P < 0.001$, according to the *t* test.

The detailed control data are in Table 4. The medical histories, and the smoking and drinking habits of workers tested are presented in Table 3. Smoking and other habits of workers appeared to have no effect on the frequency of any chromosomal changes.

Our positive results are contradictory to findings reported by Hansteen [7] who did not observe any increase either in the frequency of aberrations or in the SCE numbers. However, the subjects examined by this author had been exposed to very low doses of vinyl chloride (0.1 ppm) and only for the relatively short period of two years. This can be taken as a further proof that VCM-

TABLE 2

CHROMOSOMAL ANALYSES OF 9 WORKERS OCCUPATIONALLY EXPOSED TO VINYL CHLORIDE (TYPES OF ABERRATION)

Exposure (years)	Aberrant cells (%)	Number of chromatid breaks /100 cells	Number of chromatid exchanges /100 cells	Number of chromosome breaks /100 cells	Number of chromosome exchanges /100 cells	Number of despiralizations /100 cells
E.H.	25-					
J.P.	16-					
K.K.	15-					
A.H.	14-					
M.M.	13-					
E.J.	12-					
J.K.	12-					
F.M.	11-					
A.T.	10-					

R&S 004368

Our results are in agreement with routine and mutagenicity tests. The sensitivity of the tests about the

TABLE 3

MEDICAL HISTORIES

E.H.

J.P.

K.K.

A.H.

M.M.

E.J.

J.K.

F.M.

A.T.

NYL CHLO-

TABLE 2

CHROMOSOMAL ANALYSES OF 9 WORKERS OCCUPATIONALLY EXPOSED TO VINYL CHLORIDE

		Exposure		1st collection aberrant cells (%) ^a	2nd collection aberrant cells (%) ^a	3rd collection	
		years	ppm			aberrant cells (%) ^a	SCE/cell ^b
1	E.H.	25-27	20-150	11	0	11 ^c	17.50 ^c
0	J.P.	16-18	20-80	4	2	2	9.56
0	K.K.	15-17	15-70	2	7	3	15.02 ^c
0	A.H.	14-16	20-150	1	1	6	—
0	M.M.	13-15	20-90	0	1	7 ^c	15.26 ^c
0	E.J.	12-14	20-150	3	6	11 ^c	16.41 ^c
0	J.K.	12-14	20-90	1	4	3	10.60
0	F.M.	11-13	20-90	1	2	1	12.22 ^c
0	A.T.	10-12	20-90	4	1	3	—
0	Mean					5.2 ± 1.3	13.80 ± 1.13
0	8 controls (mean)					1.8 ± 0.4	9.41 ± 0.40

^a 100 cells were scored per sample.^b 50 cells were scored per sample.^c Significantly increased in comparison with control values.

induced chromosomal changes depend both on dose and length of exposure to the vinyl chloride monomer.

Our results seem to lead to a conclusion that the two cytogenetic methods — routine and SCE — are equally suitable for testing of high-dose vinyl chloride mutagenicity in vivo. The increase of SCEs in our results is more homogeneous. The sensitivities of the two methods to detect mutagenic activity seem to be about the same.

TABLE 3

MEDICAL HISTORY, AND SMOKING AND DRINKING HABITS OF WORKERS TESTED

	Diagnostic radiation	Smoking	Drinking	Drugs
E.H.	—	+	±	Meprobat Papaverin
J.P.	—	±	++	Chloramphenicol Syntophyllin
K.K.	+	—	±	—
A.H.	—	—	±	Solutan Syntophyllin
M.M.	—	—	±	—
E.J.	—	+	±	Bellaspone
J.K.	—	—	±	—
F.M.	—	++	++	Superpyrin Thiospasmin
A.T.	—	—	±	Ketazon

R&S 004369

TABLE 4
CHROMOSOMAL ANALYSIS OF 8 CONTROLS

	Aberrant cells ^a (%)	SCE/cell ^b	Smoking	Drinking
M.K.	3	10.88	—	—
J.K.	0	8.22	—	—
V.B.	3	10.30	++	+
J.S.	3	10.16	++	+
K.B.	1	10.46	—	—
M.Ku.	1	8.26	—	—
Z.P.	2	7.94	±	—
R.S.	1	9.02	—	+
Mean	1.8 ± 0.4	9.41 ± 0.40		

^a 100 cells were scored per sample.

^b 50 cells were scored per sample.

The controls were not exposed to X-rays and drugs at the time of blood collection.

References

- 1 Bartsch, H., C. Malaveille and R. Montesano, Human, rat and mouse liver-mediated mutagenicity of vinyl chloride in *S. typhimurium* strains, *Int. J. Cancer*, 15 (1975) 429-437.
- 2 Bartsch, H., and R. Montesano, Mutagenic and carcinogenic effects of vinyl chloride, *Mutation Res.*, 32 (1975) 93-114.
- 3 Creech, J.L., and M.N. Johnson, Angiosarcoma of the liver in the manufacture of polyvinyl chloride, *J. Occup. Med.*, 16 (1974) 150-151.
- 4 Ducatman, V., K. Hirschhorn and I.J. Selikoff, Vinyl chloride exposure and human chromosome aberrations, *Mutation Res.*, 31 (1975) 163-169.
- 5 Funes-Cravioto, F., B. Lambert, J. Lindsten, L. Ehrenberg, A.T. Natarajan and S. Osterman-Golkar, Chromosome aberrations in workers exposed to vinyl chloride, *Lancet*, 1 (1975) 459.
- 6 Hansteen, I.L., L. Hillestad and E. Thils-Evensen, Chromosome studies in workers exposed to vinyl chloride, *Mutation Res.*, 38 (1976) 112.
- 7 Hansteen, I.L., Chromosome breakage frequency and sister chromatide exchanges in PVC workers two years after exposure, A follow-up study, Abstract of 6th International Chromosome Conference, Helsinki, August 1977.
- 8 Hungerford, D.A., Leukocytes cultured from small inocula of whole blood and preparation of metaphase chromosomes by treatment with hypotonic KCl, *Stain Technol.*, 40 (1965) 333-338.
- 9 Kučerová, M., Cytogenetic analysis of human chromosomes and its value for the estimation of genetic risk, *Mutation Res.*, 41 (1976) 123-130.
- 10 Loprieno, N., R. Barale, S. Baroncelli, C. Bauer, G. Bronzetti, A. Cammellini, G. Cercignani, C. Corsi, G. Gervasi, C. Leporini, R. Nieri, A.M. Rossi, G. Stretti and G. Turchi, Evaluation of the genetic effects induced by vinyl chloride monomer (VCM) under mammalian metabolic activation: in vitro and in vivo studies, *Mutation Res.*, 40 (1976) 85-96.
- 11 Purchase, I.F.H., C. Richardson and D. Anderson, Chromosomal and dominant lethal effects of vinyl chloride, *Lancet*, II (1975) 410-411.
- 12 Purchase, I.F.H., C. Richardson and D. Anderson, Chromosomal effects in peripheral lymphocytes, *Proc. Roy. Soc. Med.*, 69 (1976) 290-291.
- 13 Viola, P.L., A. Bigotti and A. Caputo, Oncogenic response of rat skin, lungs and bones to vinyl chloride, *Cancer Res.*, 31 (1971) 516-522.
- 14 Wolff, S., and P. Perry, Differential Giemsa staining of sister chromatids and the study of sister chromatid exchanges without autoradiography, *Chromosoma*, 48 (1974) 341-353.

R&S 004370

Paper
The
esis.
lanc

Copy
All
mit-
with

Sub
pub
mer
Jun-
out

Subs
Am

Price
posi-
the

Adve