ACUTE TOXICITY OF PYDRAUL® MC (BN-77-412B) TO FATHEAD MINNOW (*Pimephales promelas*).

TOXICITY TEST REPORT

SUBMITTED TO

MONSANTO COMPANY

ST. LOUIS, MISSOURI

REPORT #BW-78-3-093

E G & G, Bionomics
Aquatic Toxicology Laboratory
790 Main Street
Wareham, Massachusetts
March, 1978

DSW 586516

STLCOPCB4091850

INTRODUCTION -

A 96-hour, static, acute toxicity determination was performed at the Aquatic Toxicology Laboratory of E G & G, Bionomics, Wareham, Massachusetts to determine the acute toxicity of Pydraul® MC (BN-77-412B) to fathead minnow (Pimephales promelas).

MATERIALS AND METHODS

Procedures used in the acute toxicity determination followed those described in "Methods for acute toxicity tests with fish, macroinvertebrates, and amphibians" (U.S. EPA, 1975).

Pydraul® MC (BN-77-412B), a green viscous liquid, was tested as received from the Monsanto Company, St. Louis, Missouri. Nominal test concentrations are reported as milligrams of Pydraul® MC per liter of test solution (mg/ℓ).

Fathead minnow used were obtained from a commercial fish supplier in Ohio. Upon receipt at Bionomics, the fish were held in a 1700-ℓ circular metal pool that had previously been coated with an epoxy resin paint to prevent leaching of extraneous materials into the water. Well water characterized as having a total hardness and alkalinity as calcium carbonate ($CaCO_3$) of 32 mg/ℓ and 20 mg/ℓ, respectively (APHA, et al.,

STLCOPCB4091851

1975), a pH of 7.1 and a temperature of 22 ± 1°C flowed through the pool at a minimum of 4 l/minute. This flow rate was adequate to maintain a dissolved oxygen (DO) concentration greater than 70% of saturation. Experimental animals were maintained under these conditions for a minimum of 14 days. During this time period, the observed cumulative mortality of the fish was less than 2% and they were judged to be in excellent condition. All fish were fed a dry, pelleted food, ad libitum, daily and ground liver weekly except during the 48-hours prior to testing.

Two independent tests involving two different series of concentrations were performed for this toxicity determination. The mortality data derived from a preliminary range-finding test were used to define a narrower series of concentrations to be used in the subsequent definitive test.

The tests were conducted in 19.6-ℓ glass jars which contained 15 ℓ of test solution. The diluent water used was soft water reconstituted according to recommended procedures (U.S. EPA, 1975). This water was characterized as having a total hardness and alkalinity as $CaCO_3$ of 40 mg/ℓ and 31 mg/ℓ, respectively (APHA, et al., 1975), a pH of 7.6 and a conductivity of 159 micromhos per centimeter. A concentrated stock solution of Pydraul® MC was prepared by mixing 3.4 grams of the test material

STLCOPCB4091852

in 100 milliliters (mℓ) of acetone, resulting in a 34 mg/mℓ suspension. This suspension was shaken to insure homogeneity and the appropriate amount was pipetted directly into each test jar and mixed by stirring.

Control and solvent control jars were also established. Both jars contained the same dilution water as used in the exposure jars but contained no Pydraul® MC, while the solvent control also contained 4.4 mℓ of acetone. All test solutions were maintained at 22 ± 1°C and were not aerated.

Ten fathead minnow with a mean and standard deviation (N=30) wet weight of 1.1 ± 0.6 grams and a mean standard length of 38 ± 8 millimeters were randomly distributed to each test jar within 30 minutes after the test compound was added.

During the study, the pH and dissolved oxygen concentrations were measured at 0-, 24-, 48- and 96-hours in the control, high, middle and low test concentrations. The pH was measured with an Instrumentation Laboratory pH meter and combination electrode, and the temperature and DO were measured with a YSI combination oxygen-temperature meter and probe.

A 50 mℓ sample was taken from the 2.2 mg/mℓ concentration of Pydraul® MC at 96 hours and preserved with 10 mℓ of hexane and

STLCOPCB4091853

stored for future shipment to Monsanto Company at the request of Ms. Rita Comotto. Control and solvent control samples were also taken and preserved similarly.

Test concentrations and corresponding percentage mortality data derived from the definitive test were used to calculate 24-, 48-, 72- and 96-hour median lethal concentrations (LC50's) and 95% confidence intervals by means of the moving average angle method (Harris, 1959). The LC50 is defined as the calculated nominal concentration of the test compound in diluent water which caused 50% mortality in the test animal population at the stated exposure interval.

Prior to analysis by this method, nominal compound concentrations were transformed to logarithms and corresponding percentage mortalities to angles. Each group of three successive angles was then averaged and the LC50 was estimated by linear interpolation between the successive concentrations whose average angles bracketed 45°.

RESULTS

The 96-hour LC50 for fathead minnow exposed to Pydraul® MC was 1.6 mg/ℓ. The pH and DO concentrations measured during the 96-hours of exposure are presented in Table 1. The cal-

STLCOPCB4091854

culated LC50 values and 95% confidence intervals for 24-, 48-, 72- and 96-hours are presented in Table 2. Table 3 presents a summary of percentage mortalities observed in each individual test concentration at the 24-, 48-, 72- and 96-hour exposure intervals. Also presented in Table 3 are the observed behavioral responses of the fish. Test concentrations turned cloudy immediately after treatment with the stock solution.

STLCOPCB4091855

LITERATURE CITED

APHA, AWWA, WPCF. 1975. Standard Methods for the examination of water and wastewater. 14th Edition, Washington, D.C. 1193 pp.

Harris, E.K. 1959. Confidence limits for the LD50 using the moving average angle method. Biometrics, Vol. 4, #3. pp. 157-164.

U.S. EPA. 1975. Methods for acute toxicity tests with fish, macroinvertebrates, and amphibians. Ecological Research Series (EPA-660/3-75-009), 61 pp.

DSW 586522

STLCOPCB4091856

SUBMITTED BY:

E G & G, Bionomics
Aquatic Toxicology Laboratory
790 Main Street
Wareham, Massachusetts
March, 1978

PREPARED BY:

Joseph V. Sousa

Joseph V Sousa

Aquatic Biologist

APPROVED BY:

Stephen J. Ells

Stephen J. Ells

Aquatic Toxicologist

George A. Cary

George A Cary

Director, Aquatic Biology

DSW 586523

STLCOPCB4091857

Table 1 -- pH and DO concentrations measured during the 96-hour toxicity test with fathead minnow (Pimephales promelas) exposed to Pydraul® MC.

	Nominal concentration (mg/ℓ)	0-hour	24-hour	48-hour	96-hour
pH	10	7.6	6.8	6.7	6.7
	3.6	7.6	6.9	6.8	6.7
	0.46	7.3	6.6	6.6	6.4
	control (acetone)	7.6	6.9	6.9	6.7
	control	7.5	6.9	6.9	6.9
DO (mg/ℓ)	10	8.1	4.2	0.4	0.4
	3.6	8.2	4.0	0.5	1.5
	0.46	8.9	1.4	0.7	1.4
	control (acetone)	8.2	4.7	1.2	1.4
	control	8.1	4.0	2.1	4.0

DSW 586524

Table 2 -- Acute toxicity of Pydraul® MC to fathead minnow (Pimephales promelas) during the 96 hours of exposure.

LC50 (milligrams/liter)				No discernible effect level at 96 hours (mg/ℓ)
24-hour	48-hour	72-hour	96-hour	
2.9 (2.2-3.7)[a]	1.8 (1.4-2.4)	1.6 (1.2-2.1)	1.6 (1.2-2.1)	0.46

[a] 95% confidence interval.

DSW 586525

STLCOPCB4091859

Table 3 -- Concentrations tested and correponding percentage mortalities of fathead minnow (Pimephales promelas) observed after 24-, 48-, 72- and 96-hours exposed to Pyraul® MC.

Nominal concentration (mg/ℓ)	% mortality			
	24-hour	48-hour	72-hour	96-hour
10	100	100	100	100
6	100	100	100	100
3.6	70[a,b]	90[a,c]	100	100
2.2	20[a,c,d]	60[e,f]	60[f,g]	60[f,g]
1.3	10[c,f]	30[e,f]	40[f]	40[f]
0.78	0[c,f]	0[f]	10[f]	10[f]
0.46	0	0	0	0
control (acetone)	0	0	0	0
control	0	0	0	0

[a] Fish showed signs of hemorrhaging.
[b] Fish showed a complete loss of equilibrium.
[c] Some fish showed a complete loss of equilibrium.
[d] Some fish showed a partial loss of equilibrium.
[e] Fish were at the surface.
[f] Some fish showed signs of hemorrhaging.
[g] Some fish were twitching.

DSW 586526

2/14 TEMP RE
0 23° 24-22°
48 23°

OBSERVATIONS

General Behavior

CLE - Complete loss of equilibrium

PLE - Partial loss of equilibrium

HYP - Hyperactive

LETH - Lethargic

PFAE - Pectoral fins anteriorly extended

EXO - Exopthalamus

EA - Extended abdomen

SU - Surfacing

SO - Sounding

TW - Twitching

AS - At the surface

Swimming

ERR - Erratic

GY - Gyrating

SK - Skattering

IN - Inverted

Pigmentation

DRK - Dark

Integument

MS - Mucous shedding

EMP - Excessive mucous production

Respiration

RA - Rapid

SL - Slow

GA - Gulping air

Solution

CLDY - Cloudy

PRE - Precipitate

FOS - Film on surface

UC - Undissolved chemical

Diane M. Sullivan - DMS

Charlotte Stiefel - CS

DSW 586527

STLCOPCB4091861

E G & G, Bionomics

Recon fenvs X

MPC /%	COMPANY	LOT #	DATE TEST STARTED: 2/13/80	DATA BY
Pydraul MC / 100 %	Monsanto	BN 77-412	TIME TEST STARTED: 4:30	WDeliver WDO

SPECIES	SOURCE	MEAN (N=30)	STOCK CONC.	SOLVENT	CONTROL JAR TEMP. C
Pimephales promelas	Tank P1 Lot # 77443 State Ohio	length 38 ± 7.7 mm weight 1.1 ± .63 g	34 mg/l (mg/ml)	ACETONE	0 23° 24 23° 48 2 72 23° 96 23° / hour

DILUTION WATER

SOURCE Recon TYPE RC56 pH 7.6 HARDNESS (mg/l $CaCO_3$) 40 ALKALINITY (mg/l $CaCO_3$) 31 CONDUCTIVITY (µmhos/cm) 159 SALINITY (o/oo) ____

Test Container Volume = 19.6 l
Diluent Volume = 15 l

10 animals / 1 replicate

COMMENTS: sampled 2.2 mg/l 96 hour water samples
samples both controls

H2O cloudy after addition of stock, remained cloudy for length of test
*HEM - fish showed signs of hemorraging

WDO ok MJC 2/20/78

PERCENTAGE MORTALITY

Apparent 96-Hour No Effect Level 0.46 mg/l

CONC. µg/l g/l mg/l	Amount required (ml)	0-hour DO	0-hour pH	CS # dead	24-hour Obser.	24-hour DO	24-hour pH DMS	CS # dead	48-hour Obser.	48-hour DO	48-hour pH DMS	WDO # dead	72-hour Obser.	72-hour DO	72-hour pH	CS # dead	96-hour Obser.	96-hour DO
Control		8.1	7.5	0		4.0	6.9	0		2.1	6.9	0				0		4.0
Sol. (control acetone	4.4	8.2	7.6	0		4.7	6.9	0		1.2	6.9	0				0		1.4
P1 10	4.4	8.1	7.6	10		4.2	6.8	0		0.4	6.7	0				0		0.4
6	2.7			10				0				0				0		
3.6	1.6	8.2	7.6	7	HEM* CLE	4.0	6.9	2	1 CLE HEM	0.5	6.8	1				0		1.5
2.2	0.97			2	HEM. 2CLE 6 PLE			4	As 2HEM			0	TW 2 2hem			0	2TW 2 HEM	
not used 1.3 hour	0.57			0				0	1HEM.			0	1 hem			0	1 hem.	
not used 0.78 2/20	0.34	8.1	7.6 DMS	1		1.6	6.9 DMS	1		0.5	6.8 DMS	1 CS				0		1.6
2/14/78 1.3 Time: 3:38	0.57	8.9	7.3	1	2HEM 3CLE	2.7	6.6	2	aa 2hem	0.6	6.6	1	2HEM			0	2hem	1.9
0.78	0.34			0	1 CLE 2HEM			0	2hem			1	2HEM			0	2hem	

DSW 586528

STLCOPCB4091862