

*Carcinogens
New Jersey
Chambers*

RECEIVED
MAY 5 1978
A. E. MONTAGNA

RECEIVED
MAY 10 1978
R. A. DeCOURT

RECEIVED
MAY 1 1978
DR. E. Q. HULL

RECEIVED
JUN 06 1978
R. N. WHEELER, JR.

A RATIONAL VIEW OF CANCER IN NEW JERSEY

by

RECEIVED
SEP 18 1978
R. N. WHEELER, JR.

Harry B. Demopoulos, M.D.
Associate Professor of Pathology
New York University Medical Center
New York, New York

*In the interest of an informed approach to the problem of cancer
in New Jersey this paper has been reproduced, with permission of
the author, by:*

New Jersey State Chamber of Commerce
5 Commerce Street
Newark, New Jersey 07102

Additional copies are available upon request.

UCC
057550

The following document contains simplifications for ease of understanding. As in most aspects of human disease, there are exceptions, and alternate opinions. The following attempts to condense the contemporary, responsible thoughts on the different aspects of cancer.

This Analysis was prepared while Dr. Demopoulos was Director of the Cancer Institute of New Jersey

UCC
057551

S U M M A R Y

A RATIONAL VIEW OF CANCER IN NEW JERSEY

This summary outlines the essential points that are explained in the attached documents.

I. Definitions

- o A major distinction is made between mortality and incidence rates. Mortality rates simply represent statistics reported from death certificates and provide insufficient data for any solutions; incidence rates are the numbers of new cases/year and relate to living patients. The status of New Jersey as "the number one cancer state" is based on mortality rates.
- o The difference between industrial and environmental cancer is stressed. Industrial exposures in the "work place" cause a minor number of cancers, and there is an even smaller number attributable to industrial carcinogens that "have escaped into the community" (less than 0.00001% of cancer deaths). Environmental cancer refers to cancers induced by an individual's personal environment which includes cigarette smoking, excess alcohol consumption, ingestion of high fat/low fibre diets, use of nitrate/nitrite containing meats, consumption of foods with artificial colors, and other aspects of life-style; the occupational aspects of the environment are important, but relatively less important.

II. How Cancers Start

- o Normal cells have repair mechanisms to undo the damage caused by chemical and physical agents. However, these repair systems can be overloaded.
- o Cancer-causing agents generally take 20-25 years to produce a cancer. The example of the Hiroshima and Nagasaki survivors is given. This 20-25 year period is known as the "lag phase" in carcinogenesis.
- o Many substances can act together, either in an additive way or synergistically (the sum total of the effect is greater than just additive).

III. The Causes of Human Cancers

- o Specific cancers have been linked with definite agents, but the major lethal cancers, lung and large bowel, are linked to personal habits. Only a small percent of cancers are industry-related.
- o Nitrites/nitrates and artificial colors are cancer causing chemicals (nitrites/nitrates are converted into dangerous nitrosamines when preserved meats are heated), but specific human cancers have not yet been linked to these substances.
- o Life-styles that include obesity, and multiple pregnancies, are associated with a greater risk for cancers of the breast and uterus, respectively.
- o The most dangerous human carcinogens are cigarettes, alcohol, dietary fats, nitrites/nitrates, and artificial food colors. These are the most widely distributed in the communities, and are proven to be responsible for the largest percentage of cancers, estimated as high as 80% (by Dr. Frank Rauscher, the recent Director of the National Cancer Institute, and Dr. Theodore Cooper, the recent Assistant Secretary for Health in HEW).

IV. Specific Problems in New Jersey

- o There is a complex array of medical, social, geological and geographic factors that must enter into any analyses and solutions for New Jersey's cancer problems, e.g., the dense urban population in N.J., inadequate educational and medical leadership in the past, etc. The sensitive area of "the quality of medical care in New Jersey" may explain a portion of the high cancer mortality rates during the period covered by the mortality study.
- o New Jersey's unfortunate prominence as the "number one cancer state" is based solely on mortality data from 1950-1969 and does not consider the mobility and shifts in populations from New York City and Philadelphia. These two cities, as well as other non-industrialized urban areas, have death rates as high as those of New Jersey. The density of the urban population of New Jersey confounds any analyses of the data available.

V. New Jersey Industry and Cancer

- o New Jersey had a 17% greater death rate, for white males, and a 14% greater death rate, for white females, compared to the rest of the country. This is the basis of New Jersey's infamy.
- o Other urban centers, even with light industry, share New Jersey's death rates, e.g., New York City, its Northern Westchester suburb, its Eastern Nassau suburb, and San Francisco. The state of New Jersey, when compared to other states, ranks number one. If, however, only populations are compared and state boundaries are ignored, then New Jersey's rates are equal to the rates of urban areas in other parts of the country. The problem lies however in the fact that even the rural areas of New Jersey have "urban rates"; this is the heart of the problem and requires further study. The answer may be as simple as the fact that many New Jersey rural dwellers, who died in 1950-1969, may have been city inhabitants from New York and Philadelphia at some time in the past and carried their damaged cells with them when they moved.
- o The types of cancers that are typically "industry-related" do not account for New Jersey's excess 2,000 cancer deaths each year. All types of cancer deaths are increased, and in some cases there is a negative correlation (benzene causes lymphomas and leukemias, and benzene is a prominent industrial pollutant - yet, the mortality rates from lymphomas and leukemias are not above the national average).
- o Separate studies by Drs. Louria and Demopoulos have suggested that only 600 of the 14,000 deaths in New Jersey might be "industry-related".

VI. The List of Carcinogens in S-3035, Section 6

- o The list includes a mix of substances; some are no longer in use, some are not carcinogens, some are strong and others are weak carcinogens. Apparently, at the present time, all are being handled with adequate precautions.
- o Asbestos and vinyl chloride are weak carcinogens.
- o Research is required to determine permissible exposure levels as has been done with the most powerful carcinogenic agent, x-rays.

A RATIONAL VIEW OF CANCER IN NEW JERSEY

New Jersey has unfortunately achieved infamy because statistics from a National Cancer Institute study reveals that New Jersey had the highest mortality rate, per 100,000 general population, in America in the period 1950-1969. This means one of two things: a) if you developed cancer and lived in New Jersey at that time, you were more likely to die of it because your cancer has been detected at a more advanced stage than in other states, and/or the complicated treatment that was needed was not as available as in other parts of the country; b) the risk of developing a lethal form of cancer was greater in New Jersey than in other states. It is not possible to determine which of these two reasons, or what combination is the truth because reliable incidence data does not exist throughout New Jersey. The data that is available is mortality data, which only shows how many people die each year from cancer. The mortality data, although very inadequate, has triggered massive controversies regarding:

- o industry-related cancer
- o environmental cancer
- o the personal environment
- o life-styles and cancer
- o inadequate health resources
- o toxic substances
- o dietary factors
- o banning of so-called carcinogens
- o politics
- o early detection
- o financial aspects of cancer
- o possible solutions

Cancer is the most complicated disease process, compared to the other major killers such as heart disease, strokes and accidents. There are over a hundred different forms of cancer, and they start in different organs of the body, in different types of individuals, and under poorly understood circumstances. No other disease process is so intricately interwoven with the very fabric of society; as though to emphasize their perplexing, intricate nature, cancer cells remain very similar to the patient's normal cells, thereby frustrating most attempts to "weed" them out.

In order to put cancer into perspective, so that possible solutions can be evaluated, the major complex aspects must be understood.

1. DEFINITIONS

Cancer - a malignant growth, composed of solid masses of disorganized cells that are ever-growing, and are capable of spreading to organs far from the original site, e.g., breast cancer starts as a lump, and some of its component cells will microscopically invade the blood stream which will carry them to the lungs, bones, liver, and brain.

Metastasis - the term used to refer to the spread of cancer cells from its site of origin, to other organs.

Early Detection - the process of detecting the cancerous mass while it is still small and therefore less likely to have metastasized.

Radical Surgery - the principal weapon in use today for treating cancer; the surgeon cuts widely around the cancerous mass hoping that none of the cancer cells have microscopically metastasized; the surgeon and other cancer experts generally have no way of knowing whether an individual case has already spread microscopically; however, the smaller the original cancer mass, the less likely it is to have spread.

Mortality Rate - the number of people dying as a result of cancer each year; it is generally given as the number of cancer deaths per 100,000 general population. In America, the average figure is about 170/100,000.

Incidence Rate - the number of new cases that are diagnosed each year, again given per 100,000; in America the average figure is about 340/100,000; about one-half of this number will eventually die of their disease, but over a period of 2-8 years; the other half of the newly diagnosed cancer cases will live out a normal life expectancy and die of some other cause.

Industry-Related Cancer - a cancer whose cause can be, at least in part, traced back to a distinct exposure to a chemical or sometimes a physical agent that was present in the "work-place"; this generally encompasses employees and, more rarely, members of their families who are exposed to the employees "contaminated" work clothes.

Environmental Cancer - refers to most cancers, possibly 80%; however, the term "environment" is all-encompassing and relates mostly to the personal environment that results from life-styles, habits, and dietary factors; occupational exposures, and industry-related events comprise a minor component of "environmental cancer".

Carcinogen - an agent, chemical or physical, that is capable of initiating irreparable damage to a cell, such that the cell may be ultimately transformed into a cancer cell.

Co-Carcinogen - a chemical which by itself is incapable of causing cancer, but in combination with small doses of carcinogens (doses too small to cause cancer) will cause the development of cancers.

Epidemiology - the science of studying what disease occurs in which types of people, and under what circumstances - it constitutes medical detective work.

II. HOW CANCERS START

Cancer cells are no longer "self-controlled", the way normal cells are. Of the many trillions of cells in our bodies, many of them divide and multiply in order to replace "worn out" cells. Most remarkable is that the normal replacements are exactly like the worn-out ones. Cancers start as a result of some cells losing their normal control mechanisms. This happens as a result of damaging the genetic material and the delicate membranes of a cell beyond repair. It is important to realize that from conception, i.e., when a sperm and egg cell have united to form a unique individual, that single cell and all of its subsequent dividing cells are constantly barraged by damaging agents such as viruses, chemicals, and ionizing radiation. In the overwhelming majority of damaging insults, our cells repair themselves. When repair is inadequate, we see the development of birth defects, cancer, or death. The genetic material of the nucleus, the DNA, has incredible repair machinery, and the membranes of a cell, in spite of their extraordinary delicacy and complexity, are undergoing incessant replacement, 24 hours a day, 365 days a year. In short, evolution has given us resilient cells so that our bodies can take a certain amount of damage. However, the repair systems can be overloaded and some damaging agents can specifically attack the repair systems.

It is clear that there are certain tolerance limits, and if exceeded, the results may be birth defects, cancer or cell death.

Most cancers are the result of a complex sequence of damaging events that have not been repaired. However, the damage need not be expressed; there is a need to have other factors which will "bring out" the cancerous damage. In some instances, it may take 20-25 years to "bring out" the cancerous damage. This is referred to as the "lag phase" and is best exemplified by the survivors of the atomic blasts at Hiroshima and Nagasaki. These large amounts of radiation caused irreparable damage to cells, but the cancerous nature of the damage was not expressed as a lump of uncontrolled cells until 20-25 years later. In these and a few other instances, there are exposures to a single damaging agent and clear cut blame can be affixed to that agent.

In the overwhelming majority of cancers, there are multiple agents which interact and cause cumulative damage. While there are many hundreds of damaging agents, there are only a few different types of parts in a cell; hence, the same type of cell part may be damaged by several different substances. We can therefore see additive effects, and sometimes synergism, wherein the result is more than just additive.

III. THE CAUSES OF HUMAN CANCERS

The causes of many cancers are known and are listed below. The numbers in parentheses represent the percent of total cancer deaths caused by that particular type of cancer.

<u>Type of Cancer</u>	<u>(% of Deaths)</u>	<u>Causes</u>
o lung cancer	(20%)	cigarette smoking*
o mouth cancer	(2%)	cigarette smoking* <u>plus</u> excess alcoholism** <u>plus</u> inadequate mouth care
o larynx cancer ("voice box")	(1%)	cigarette smoking* <u>plus</u> excess alcoholism**
o esophagus cancer ("food tube")	(5%)	cigarette smoking* <u>plus</u> excess alcoholism**
o colon and rectal	(16%)	high dietary fat <u>plus</u> low dietary fibre
o liver cancer (ordinary type)	(0.5%)	excess alcoholism**
o urinary bladder cancer	(5%)	unrestricted use of certain chemicals used in dye production
o mesothelioma	(0.001%)	unrestricted use of asbestos
o hemangiosarcoma (a special type of liver cancer)	(0.001%)	unrestricted use of vinyl chloride

From the preceding list, it is clear that we know what causes over 50% of cancer deaths. There are several major cancer types whose causes are not known and account for a total of 30% of the deaths; these are cancers of the breast, ovaries, pancreas, and prostate. In some of these cancers, we know a spectrum of associated findings that add up to a greater risk, as in cancer of the breast, but this does not mean that a causal factor(s) that can be controlled has been discovered.

In addition to the above known causes of cancer, the following substances have been identified as potent carcinogens that are consumed in significant quantities (milligrams/day) by the majority of Americans; these substances have not yet been linked as causative factors to a specific type of cancer, as in the case of cigarette smoking and lung cancer, but they are nonetheless

* generally more than one pack/day

** generally more than three ounces of distilled liquor/day, or more than sixteen ounces of wine/day

suspected of causing several different types of cancers. They may act as co-carcinogens, or act synergistically.

- o nitrites/nitrates - are converted to dangerous nitrosamines when preserved meats, such as frankfurters, ham, etc., are heated
- o artificial food colors - some colors that are widely used in a multitude of foods and beverages are powerful carcinogens

Some aspects of particular lifestyles are associated with a greater risk for developing cancer and include:

- o obesity associated with breast, colon and rectal cancer
- o multiple pregnancies coupled with inadequate gynecologic care associated with cancer of the uterine cervix
- o repetitive sun exposure causes cancer of the skin

Added to all of the above are many other agents that are carcinogenic, but they are either weak, sparsely distributed, or not well delineated. This includes:

- o hormones used to relieve symptoms of the menopause associated with cancer of the endometrium
- o pharmacologic drugs used to treat high blood pressure associated with cancers of the breast and colon
- o high benzene levels capable of causing leukemias
- o varied organic compounds in the work place thought capable of causing cancers of the skin, stomach, liver, lungs and urinary bladder

The conclusion that should be reached at this point is that everyone is exposed to some combination of carcinogenic substances every day and the most significant ones are uncontrolled. Many agents are carcinogenic and there is a desperate need to study and define which of these agents should be brought under more restrictive control. Logic would direct immediate attention, at this time, to those carcinogens that cause the greatest number of cancer deaths, and are also widely used. This would result in a list, in order of importance, as follows:

- o cigarettes
- o alcohol
- o dietary fat and fibre
- o nitrites/nitrates
- o artificial food colors

IV. SPECIFIC PROBLEMS IN NEW JERSEY

The State of New Jersey is characterized statistically in a number of areas that have a relevance to the high cancer mortality rates, and which may also be involved in the solutions:

- o Most densely urbanized population
- o Lowest average land elevation with respect to sea-level
- o Highest unemployment rates
- o Reliance on local water sources of household consumption, in contradistinction to other states where water sources are not derived from the water table or rivers found in the concentrated urban areas.
- o Ranks 46th among the states in money spent for higher education per capita.
- o Until recent years lacked any credible medical schools; the two existing ones are developmental and reportedly rank 85th and 87th out of the approximate 100 American Schools according to the average scores of the students on the National Medical Board Examinations.
- o 75% of the interns and residents in the hospitals are foreign medical school graduates; this is the highest in the country; in New York, the second highest, the figure is 50%, while in California, this number is 5%; the reliance on foreigners is directly traceable to the absence of a large enough pool of graduating medical students who have a desire to undertake or continue their training in New Jersey hospitals.

The intermingling of medical, social, geographic, and geological factors listed above, provide clues for analyzing New Jersey's complex cancer problems, and the thought that is required for the solutions. There is no simple approach. and to highlight this, a critical examination of the New Jersey cancer data is in order.

The National Cancer Institute, under a program directed by Dr. Fraumeni, conducted a nationwide county-by-county study of cancer death rates per 100,000 general population. This was done by transcribing what was written on the death certificates. The medical records of the patients were not examined. The death rates from 1950-1969 were catalogued from death certificates, and were classified by anatomic site, and sex. Hence, the number of deaths from cancer of the mouth, or of the stomach, in males or females, in Atlantic County, or Essex County, and other such numbers became available. These numbers are interesting, and, in general, there is no county in New Jersey that could be considered "safe" according to these data.

The 1950-1969 mortality study was not designed by Dr. Fraumeni to provide information for definitive solutions. To employ this type of data for directing conclusive solutions is a frank misuse. The study was conducted to provide an overview of the scope of the cancer problem in the nation and nothing more. The reasons that the Fraumeni data of 1950-1969 cannot be used to direct any solutions stem from the lack of medical records data such as:

- o Size of cancer and extent of disease when the patient was first admitted and diagnosed in the hospital. These two

UCC

057559

factors, size and extent of disease at time of diagnosis, are the principal determinants of the patient's prognosis as to life or death; if the cancer is large, chances are very high that the cancer cells have already metastasized.

- o No treatment data is available to judge adequacy of therapy, and of long term, dedicated follow-up.
- o No patient histories with respect to dangerous habits, nature of the diet, previous places of residence, occupation, income and educational level.

If the Fraumeni data were to be used for a direct attack on the cancer problem, without any further extensive studies, several false leads would be pursued based simply on mobility and shifting of the population, especially in the years from 1950-1969. Miami, Florida has cancer death rates equal to those of New Jersey. With a moment's careful thought, the reason becomes obvious - many Miami residents are retired Northeasterners who have come from crowded urban settings. It is critical to remember that a 20-25 year lag period is required to develop a cancer from the numerous environmental exposures (including cigarettes and alcohol).

Further examination of the N.C.I. mortality studies reveals the following death rates of other urbanized areas and their suburbs.

o New York City	215/100,000
o Westchester County (a Northern suburb of New York City) —	200/100,000
o Nassau County (an Eastern suburb of New York City)	212/100,000
o Philadelphia	221/100,000
o Chicago	206/100,000
o St. Louis City, Missouri	220/100,000
o San Francisco	206/100,000

Since New Jersey received former dwellers of New York City and Philadelphia during the post-war years of suburban expansion, circumstances in these two cities may ultimately be responsible. Westchester and Nassau Counties may be high in mortality rates for the same reason that New Jersey is, i.e., the city dwellers led life-styles that predisposed to cancer and when the population shift out of the cities occurred, these individuals carried their irreparably damaged DNA and altered cell parts with them. If many of these individuals were in the lag phase (20-25 years), this could explain, at least in part, the findings. This is a similar explanation for the high rates in Miami, Florida.

There are many other confounding factors that preclude conclusions, and these are as follows:

- o While the population was shifting into the suburbs, including into New Jersey, industry also continued to grow in the state.
- o In the 1930's-1940's (the period of time when people were being exposed in order to develop their cancers and live 20-25 years later in 1950-1969), industry and government were largely ignorant of chemical carcinogens, hence fewer safeguards may have been used in this period.

V. NEW JERSEY INDUSTRY AND CANCER

The county-by-county death statistics of 1950-1969 are divided by anatomic site and by sex. The overall death rates for white males was the highest in the country, but this was not the case for females or non-whites. This can lead to a great deal of speculation, but the data is insufficient for reaching a solution.

The Fraumeni numbers indicate that the national average death rate from cancer was 174/100,000 (for white males), while in New Jersey, it was 205/100,000, a 17% increase. In females, the national average was 130 and, in New Jersey, 148/100,000 about a 14% increase. These are the increases that have precipitated the current controversy. The increases are significant and were consistent through the 19 years (1950-1969) of the study. More detailed numbers are presented in appendix A.

Adding to this data are the estimates of the American Cancer Society, regarding New Jersey incidences and mortalities. These are given in appendix B.

The conclusions pertaining to New Jersey are as follows:

1. General Findings on New Jersey:
Proportionally, New Jersey's figure on cancer mortality is greater than that of any other state in America. Nearly all of New Jersey falls into the highest decile in the U.S. (top 10%) for white males and for white females (Appendix A).
2. Cancer categories in which the mortality, proportional to population, is higher in New Jersey than in any other state:
 - o Rectal cancer (males and females combined)
The rating of New Jersey as having the worst concentration of mortality from rectal cancer is based largely on the extremely high death rates among both men and women in northern New Jersey, the Trenton-Princeton area included. This would not be classified as an industry-related cancer by cancer specialists.
 - o Bladder cancer (males only - women not significantly different from the rest of the United States) A proportion of these are industry-related.
 - o Cancer of the ovary - The mortality figures among the women of New Jersey's northwest corner -- Sussex and Warren Counties -- ranks with the worst in the country. Mortality figures are nearly as high among women in Bergen, Morris, and Passaic Counties. This causes the mortality figures for women in northern New Jersey in this cancer classification to be worse than for any other single concentrated area in the nation. This would not be classified as an industry-related cancer.
3. Cancer categories in which the mortality rates for the state of New Jersey rank approximately with the worst statewide records in the nation:
 - o Cancer of the large intestine (males and females combined)
New Jersey ranks with Rhode Island as the worst in the nation. This is not regarded as industry-related.

UCC
057561

- o Cancer of the Trachea, Bronchus, and Lung - Mortality figures among New Jersey males rank as the highest in the U.S., along with the mortality among males in Louisiana and Florida. A small proportion of these are industry-related.
 - o Breast Cancer among females - The record in this category is worst in New York, with New Jersey, Rhode Island and Massachusetts just behind. This is not classified as industry-related.
4. Cancer categories in which mortality for portions of New Jersey ranks with the worst areas in the nation.
- o Cancer of the Esophagus (males only) - Northern New Jersey, Connecticut, New York City, Long Island, and the Greater Philadelphia Area of Pennsylvania, combined, comprise the worst single area in the U.S. for mortality figures in this category. The rate for females in northern New Jersey is above the national average. This is generally caused by a combination of cigarette smoking and excessive alcoholism. In addition, women with rare benign esophageal problems are predisposed to cancer of this organ. It is not generally regarded as being industry-related.
 - o Cancer of the Larynx (males only) - The highest concentrations of mortality from this type of cancer are in northern New Jersey, New York City, Long Island, the Greater Philadelphia Area and the Pittsburgh area. A small proportion of these are industry-related, but generally are caused by a combination of cigarette smoking and excessive alcoholism.

If an attempt is made to correlate the types of cancers that are known to be "industry-related" (i.e., industrial substances contribute together with other factors to the development of cancer), with the types of cancers occurring in New Jersey, it would be expected that most of the excess cancer deaths in New Jersey would fall into the "industry-related" types - this is not the case.

There are about 14,000 deaths each year in New Jersey, and about 26,000 new cases each year (from appendix B, estimates for 1974). If New Jersey had average U.S. rates, these numbers would be 12,000 and 22,000, respectively. The excess 2,000 deaths each year and the excess 4,000 new cases each year should fall into the classical "industry-related" categories which include a small portion of cancers of the:

- o Urinary bladder
- o Respiratory system
- o Liver
- o Skin
- o Lymph organs and bone marrow (lymphomas and leukemias)

Instead, the "excess" 2,000 deaths are spread across all of the anatomic sites in the N.C.I. mortality study (lip, salivary glands, nasopharynx, mouth, esophagus, stomach, large intestine, rectum, liver, pancreas, upper and lower respiratory tracts, breast, uterine cervix, body of uterus, ovaries, prostate, testis, kidneys, urinary bladder, skin, eye, brain, endocrines, bones, and connective tissues). Further, there are some negative correlations, e.g. benzene is reportedly an industrial pollutant in New Jersey and supposedly is

the highest in the nation, yet the cancer that should be caused by benzene, such as lymphomas and leukemias, occur at the lower national rates.

o Lymphomas N.J. (White males)	4.93
U.S. (White males)	4.89
o Leukemias N.J. (White males)	8.74
U.S. (White males)	8.81

Analyses performed by Dr. Donald Louria, Chairman of the Department of Preventive Medicine and Community Health, New Jersey Medical School, Newark, and presented under the title of "Cancer in New Jersey: An Overview" at the "Seminar for Physicians: Cancer Risk Identification within New Jersey, and Methods of Cancer Control", May 12, 1976, in Cherry Hill, found that only 600 of the 14,000 cancer deaths in New Jersey might be industry-related. A different analytic study conducted by Dr. Harry B. Demopoulos, former Director of the Cancer Institute of New Jersey, also revealed the same types of numbers, i.e., no more than 600 of the 14,000 cancer deaths could be industry-related, among the deaths reported in 1950-1969; this report was given by Dr. Demopoulos to the "Skevin Committee" in testimony on November 5, 1976.

These two independent analyses therefore indicate that 4.3% of the total cancer deaths in New Jersey could be "industry-related". These percentages are important and yet it is essential to realize that these analyses are based on insufficient data and represent the highest possible number of "industry-related" exposures. This does not mean that industrial pollutants were solely responsible. If an analysis is attempted of how many cancer deaths were caused solely by industrial pollutants, the data is found to be totally inadequate and very soft estimates yield fractions of 1%.

While the cited studies and analyses refer to past events, there is meager data since 1969. The American Cancer Society estimates do not provide sufficient information to answer the obvious question - are cancer mortality rates and incidences in New Jersey the same, better, or worse than for the period 1950-1969? Current, but inadequate, "samples" from hospitals that have excellent Tumor Registries indicate that their cancer case workload has increased by 50% in the past 5 years, and that the average age of the cancer patient is younger by 5-6 years. This type of data is fragmentary and may reflect changes in referral patterns to some hospitals, or it may indicate a worsening of the New Jersey cancer problems. Clear cut answers require far more data.

VI. THE LIST OF CARCINOGENESIS IN BILL NO. S-3035, SECTION 6

The list includes a mix of substances

- o Some are no longer in use, e.g., 4 Aminodiphenyl
- o Some are not carcinogenic, e.g., alpha naphthylamine - its carcinogenicity was proven to be due to contamination of alpha naphthylamine
- o Some are very potent carcinogens, e.g., benzidine
- o Some are rather weak carcinogens, e.g., asbestos and vinyl chloride
- o All are currently handled with precautions that lead to low exposures of workers such that cancers will not develop.

Asbestos and vinyl chloride are termed weak carcinogens on the basis of careful analyses of the cancers that they cause. Excessive concerns over asbestos as a carcinogen has been prompted by cases such as that of a 14 year old boy who developed mesothelioma; he apparently was exposed to this when he was helping his father to smooth down the joints of the newly replaced plaster board walls in their home. Asbestos was in the joint material, not in the plaster board. There was no other known asbestos exposure, and the father did not otherwise work with it. Cases such as this are exceedingly rare and form an inadequate data base. Mesothelioma does occur in nature, without asbestos exposure, and it cannot be ascertained whether this 14 year old boy would have been a "Natural" victim, or whether the asbestos was indeed causal. Far more numbers are needed for statistically valid studies in such unusual cases.

The relative weaknesses of asbestos and vinyl chloride are borne out by the fact that workers who were exposed to very large, uncontrolled levels developed relatively few cancers as a result. This is in contrast to a powerful carcinogen such as 3,4-dimethyl 4-aminodiphenyl, wherein 15-20% of exposed workers developed urinary bladder cancer in a short lag phase (7-8 years). When the amounts of asbestos, vinyl chloride, and 4-aminodiphenyl are compared, versus the numbers of cancers developed, then asbestos and vinyl chloride are weak carcinogens.

The idea of a comprehensive ban on all carcinogens would lead to the restriction of many activities and substances. Radiation, by x-rays, is the most powerful carcinogenic agent. There are methods for converting physical carcinogens into chemical equivalents, and when this is done, x-rays are quite potent. When the Manhattan Project (building the A-Bomb during World War II) was in full swing, the Department of Pathology at Rochester University was given the job of determining the ill-effects of radiation. There were some individuals who were so impressed by the pathologic changes that they decreed a "Zero-exposure". When subsequent, rational studies were done, it was found that small doses of radiation could be tolerated, even though the effects were cumulative over a life time. This type of exacting research has made it possible to use x-rays for medical use, nuclear plants for energy, etc. The same type of information must be obtained for chemicals.

UCC
057564

APPENDIX A

NATIONAL CANCER
INSTITUTE

MORTALITY STUDY

1950 - 1969

- EXHIBIT 1 - New Jersey Cancer Mortality 1950-1969
- EXHIBIT 2 - New Jersey Cancer Mortality 1950-1969, by County
- EXHIBIT 3 - States With Highest Cancer Mortality, 1950-1969
- EXHIBIT 4 - All Malignant Neoplasms 1950-1969, by State

UCC
057565

CANCER MORTALITY
1950-1969

All Malignant Neoplasms

(All cancer categories) causing mortality
(Male, Female Combined)

	<u>Number</u> ¹	<u>Annual Rate</u> (per 100,000 pop. whites only)
Total deaths, United States	2,572,035 - M	174.04 - M
	2,253,282 - F	130.10 - F
Total deaths, New Jersey	106,900 - M	205.01 - M
	93,379 - F	147.92 - F

Highest Rates by New Jersey Counties:

Hudson	14,049 - M	231.8 - M
	11,004 - F	153.5 - F
Middlesex	6,556 - M	220.8 - M
	5,251 - F	149.2 - F
Essex	16,975 - M	215.1 - M
	15,258 - F	154.5 - F

Highest Number by New Jersey counties:

Essex	16,975 - M	215.1 - M
	15,258 - F	154.5 - F
Hudson	14,049 - M	231.8 - M
	11,004 - F	153.5 - F
Bergen	12,863 - M	202.1 - M
	11,894 - F	148.1 - F

¹ This number is the total deaths from 1950-1969

APPENDIX A - EXHIBIT 2

ALL MALIGNANT NEOPLASMS (all cancer categories, causing mortality)
New Jersey, by counties

	<u>White</u>				<u>Non-white</u>			
	<u>Male</u>		<u>Female</u>		<u>Male</u>		<u>Female</u>	
	<u>Total</u>	<u>Rate</u>	<u>Total</u>	<u>Rate</u>	<u>Total</u>	<u>Rate</u>	<u>Total</u>	<u>Rate</u>
Atlantic	3,213	195.3	2,897	145.2	598	220.1	484	150.8
Bergen	12,863	202.1	11,894	148.1	373	281.4	344	192.2
Burlington	2,692	188.5	2,395	139.9	196	216.3	138	141.0
Camden	6,329	204.7	5,647	148.4	586	228.0	520	177.1
Cape May	1,264	194.7	1,037	142.9	72	184.2	78	176.8
Cumberland	1,683	181.4	1,551	140.0	150	155.8	141	144.2
Essex	16,975	215.1	15,258	154.5	2,385	219.2	2,155	154.6
Gloucester	1,935	191.1	1,674	141.8	182	183.1	142	146.0
Hudson	14,049	231.8	11,004	153.5	645	289.7	571	197.3
Hunterdon	1,025	175.7	926	143.3	17	297.1	19	303.2
Mercer	4,639	205.4	3,973	145.2	422	203.5	351	150.1
Middlesex	6,556	220.8	5,251	149.2	282	279.0	223	208.2
Monmouth	5,754	199.0	5,340	147.6	515	225.4	360	135.3
Morris	3,851	179.2	3,606	135.5	109	248.6	98	170.1
Ocean	2,577	185.5	2,028	137.2	69	265.3	64	231.6
Passaic	7,981	209.5	6,631	147.8	355	271.5	302	182.7
Salem	848	185.9	731	146.6	147	232.0	97	161.9
Somerset	2,151	182.8	1,820	135.7	74	226.9	66	228.5
Sussex	923	180.8	797	140.5	18	913.3	12	465.4
Union	8,311	203.4	7,757	151.6	620	252.2	534	166.3
Warren	1,281	189.3	1,162	147.7	15	274.2	10	187.7
New Jersey	106,900	205.01	93,379	147.92	7,830	230.33	6,709	163.41

Rates indicated are annual per 100,000 population
 Total deaths, for the period, 1950-1969, are given.

UCC
 057567

STATE	WHITE MALE		NONWHITE MALE			WHITE FEMALE		NONWHITE FEMALE	
	NUMBER	RATE	NUMBER	RATE		NUMBER	RATE	NUMBER	RATE
ALABAMA	29066	152.44	10156	140.24		25643	111.80	10915	127.17
ARIZONA	14782	156.19	1062	128.83		11301	110.48	811	111.76
ARKANSAS	22197	144.14	4915	132.13		17726	108.83	4660	121.99
CALIFORNIA	216761	171.39	15086	170.77		195171	128.09	10785	124.10
COLORADO	21816	140.19	620	169.99		20219	117.29	462	116.37
CONNECTICUT	44501	195.68	1439	231.75		38333	138.64	1047	139.39
DELAWARE	5691	179.75	1100	235.97		5171	134.42	774	162.57
DISTRICT OF COLUMBIA	7915	203.75	6790	264.55		8123	141.73	5445	166.05
FLORIDA	76859	163.58	10683	179.78		56475	110.54	8959	133.14
GEORGIA	33499	153.77	11546	152.23		30383	111.27	12652	130.90
IDaho	8546	139.02	103	121.67		6647	110.15	73	109.27
ILLINOIS	162672	182.81	15012	216.77		142394	137.78	13040	160.98
INDIANA	67635	164.24	4139	210.85		62111	130.60	3408	157.97
IOWA	46897	156.60	516	213.55		42106	124.14	380	151.28
KANSAS	30949	143.89	1647	189.14		28094	115.08	1286	141.1
KENTUCKY	38813	146.39	4220	199.50		36016	121.61	3580	154.2
LOUISIANA	32662	190.39	14574	193.68		24611	118.98	12466	143.77
MAINE	17793	178.53	47	154.11		16135	140.46	48	161.70
MARYLAND	39157	192.43	8099	224.84		35366	138.66	6080	154.73
MASSACHUSETTS	95772	192.23	1971	215.92		90506	139.47	1480	141.86
MICHIGAN	113318	182.42	9800	210.58		92946	135.63	7830	151.26
MINNESOTA	54422	158.24	539	176.18		47219	127.05	380	128.98
MISSISSIPPI	18050	156.40	9846	136.74		15030	113.03	10146	129.44
MISSOURI	70822	164.55	7394	213.58		63213	125.58	5882	159.07
MONTANA	10201	153.07	227	140.69		7453	119.28	207	157.24
NEBRASKA	23955	157.48	576	219.93		20736	123.75	482	175.99
NEVADA	4050	167.39	179	136.61		2654	118.79	180	148.97
NEW HAMPSHIRE	11944	189.19	17	130.20		10655	140.20	16	110.02
NEW JERSEY	106900	205.01	7437	230.33		93179	147.92	6709	163.41
NEW MEXICO	7689	136.30	340	95.50		6915	115.10	352	96.48
NEW YORK	307997	199.24	21572	227.69		273316	148.01	18920	152.80
NORTH CAROLINA	36533	140.11	10543	147.17		33864	106.97	10701	124.39
NORTH DAKOTA	9039	144.34	97	143.57		7084	119.45	110	165.14
OHIO	146265	178.41	12523	226.35		130411	136.25	9762	159.70
OKLAHOMA	34295	155.95	2436	144.80		28878	116.03	2904	131.1
OREGON	28314	155.12	471	158.47		23148	119.27	307	122.88
PENNSYLVANIA	189018	183.08	15046	236.89		170851	140.26	11528	157.10
RHODE ISLAND	16434	203.17	342	236.56		14770	143.37	212	133.64
SOUTH CAROLINA	16754	154.96	7135	145.59		15295	111.59	7647	123.97
SOUTH DAKOTA	10513	149.86	252	137.79		8384	119.98	302	184.07
TENNESSEE	38356	146.28	7374	163.79		35763	115.95	7796	142.51
TEXAS	107557	158.51	15831	167.53		90072	113.34	13452	128.45
UTAH	8369	133.14	198	152.86		7233	102.06	101	108.96
VERMONT	6981	173.02	18	207.22		6551	136.41	11	119.06
VIRGINIA	38218	153.53	11501	189.62		35279	119.12	9254	138.94
WASHINGTON	45015	164.20	1247	169.72		36251	123.43	773	129.39
WEST VIRGINIA	26025	154.62	1771	191.28		22241	123.75	1373	149.77
WISCONSIN	65376	166.23	876	191.54		56898	132.35	707	140.97
WYOMING	3953	138.93	64	123.16		2962	109.09	52	130.08
UNITED STATES	2572035	174.04	264185	184.28		2253292	130.10	226561	139.18

UCC
057568

APPENDIX B

AMERICAN CANCER
SOCIETY

MORTALITY AND INCIDENCE ESTIMATES

1974

UCC
057569

Estimated Cancer Deaths for All Sites,
Plus Major Sites, by State - 1974

State	All Sites		Major Sites								
	Number of Deaths	Death Rate per 100,000 Population	Breast	Colon-Rectum	Lung	Oral	Uterus	Prostate	Stomach	Pancreas	Leukemia
Alabama	5,400	155	425	550	1,200	125	250	325	175	325	200
Alaska	200	62	15	20	50	5	10	10	10	10	10
Arizona	2,600	134	225	300	650	70	50	150	100	150	125
Arkansas	3,500	175	225	400	850	75	100	250	125	200	200
California	33,200	153	3,200	4,200	7,300	750	1,000	1,400	1,400	1,800	1,400
Colorado	2,900	122	275	400	500	60	70	175	100	175	150
Connecticut	5,400	167	500	720	1,000	175	125	250	250	275	250
Delaware	900	152	70	125	225	25	20	30	30	50	30
Dist. of Columbia	1,600	199	175	200	325	60	60	80	60	80	40
Florida	14,600	197	1,200	1,900	3,500	350	400	800	600	700	500
Georgia	6,300	130	550	50	1,500	150	275	375	250	350	275
Hawaii	900	110	60	90	150	30	20	30	90	70	50
Idaho	1,100	150	90	125	175	20	25	80	40	70	70
Illinois	19,900	172	1,900	1,800	4,100	450	700	1,000	850	1,100	900
Indiana	8,600	158	800	1,300	1,800	175	325	450	250	450	350
Iowa	5,200	182	500	850	950	100	125	350	175	250	275
Kansas	4,000	175	375	550	750	90	125	275	100	225	200
Kentucky	5,400	165	425	700	1,200	150	200	300	150	325	275
Louisiana	5,900	156	475	600	1,500	150	200	325	250	300	250
Maine	2,200	212	175	300	425	40	60	125	90	100	80
Maryland	6,400	150	600	850	1,600	175	200	300	200	325	200
Massachusetts	11,200	189	1,200	1,700	2,200	300	300	475	525	600	400
Michigan	14,500	155	1,400	1,900	3,100	300	425	750	525	700	550
Minnesota	6,500	163	600	950	1,100	125	125	450	300	375	275
Mississippi	3,600	162	275	400	750	70	125	250	150	225	200
Missouri	8,900	184	800	1,200	2,100	175	275	550	275	475	400
Montana	1,300	185	100	150	200	25	30	70	50	80	60
Nebraska	2,800	185	250	425	475	60	70	175	100	175	150
Nevada	750	133	60	80	200	15	20	25	10	50	30
New Hampshire	1,500	189	150	250	325	30	50	80	40	80	70
New Jersey	14,000	184	1,400	2,200	3,000	300	400	550	650	700	500
New Mexico	1,200	115	100	125	200	20	30	50	50	70	50
New York	37,700	200	4,000	5,800	7,400	800	1,000	1,500	1,700	2,000	1,400
North Carolina	6,900	132	600	700	1,400	175	300	375	225	375	350
North Dakota	1,100	181	90	150	175	15	25	70	60	70	50
Ohio	19,000	172	1,800	2,700	4,100	425	650	900	700	900	750
Oklahoma	4,500	170	350	550	950	80	125	300	150	275	200
Oregon	3,700	166	325	500	850	80	100	200	125	200	200
Pennsylvania	23,500	197	2,300	3,600	4,600	500	750	1,100	950	1,200	950
Rhode Island	2,000	204	200	350	400	60	50	80	100	90	60
South Carolina	3,500	132	300	375	750	80	150	200	100	200	150
South Dakota	1,200	182	80	175	200	20	40	100	50	90	80
Tennessee	6,300	155	550	750	1,400	150	225	375	200	350	275
Texas	17,100	144	1,400	1,800	4,000	375	600	800	650	950	900
Utah	1,100	97	100	150	175	20	30	80	50	60	50
Vermont	850	182	70	150	175	20	30	50	30	50	40
Virginia	6,800	139	650	800	1,500	175	250	375	225	375	275
Washington	5,600	154	500	700	1,300	125	150	300	225	325	275
West Virginia	3,400	201	250	400	850	70	125	200	125	200	125
Wisconsin	7,800	169	800	1,200	1,300	175	200	450	350	400	325
Wyoming	500	149	40	60	75	10	10	40	15	30	20
United States	355,000	167	33,000	48,000	75,000	8,000	11,000	18,000	14,000	19,000	15,000

**Estimated New Cancer Cases for All Sites,
Plus Major Sites, by State - 1974**

UCC

057571

State	All Sites*	Major Sites								
	Number of Cases	Breast	Colon-Rectum	Lung	Oral	Uterus (Invasive)	Prostate	Stomach	Pancreas	Leukemia
Alabama	10,000	1,100	1,100	1,300	350	1,000	950	300	350	300
Alaska	300	50	50	60	15	20	30	15	10	20
Arizona	4,800	600	600	700	150	350	450	150	150	150
Arkansas	6,400	600	800	1,000	200	500	750	200	200	300
California	61,000	8,700	8,700	8,100	2,300	3,900	4,200	2,300	1,900	2,000
Colorado	5,400	750	800	550	150	350	550	150	200	200
Connecticut	10,000	1,400	1,700	1,100	500	550	750	400	300	350
Delaware	1,700	200	300	250	70	100	90	50	50	40
Dist. of Columbia	3,100	500	400	350	200	250	250	90	80	60
Florida	27,000	3,300	3,900	3,900	1,100	1,600	2,400	1,000	750	700
Georgia	12,000	1,500	1,400	1,700	450	1,100	1,100	400	350	400
Hawaii	1,700	150	200	200	50	80	60	150	70	70
Idaho	2,100	250	250	200	50	100	250	70	70	100
Illinois	37,000	5,200	5,800	4,500	1,400	2,700	3,000	1,400	1,200	1,200
Indiana	16,000	2,200	2,700	2,000	500	1,300	1,400	400	450	500
Iowa	9,500	1,400	1,800	1,000	300	650	1,100	300	250	400
Kansas	7,300	1,000	1,100	800	300	650	800	150	250	300
Kentucky	10,000	1,200	1,400	1,300	450	900	900	250	350	400
Louisiana	11,000	1,300	1,200	1,700	450	850	950	400	300	350
Maine	3,600	500	600	450	125	250	400	150	100	100
Maryland	12,000	1,600	1,800	1,800	500	900	900	350	350	300
Massachusetts	21,000	3,300	3,500	2,400	900	1,200	1,400	850	600	550
Michigan	27,000	3,800	3,900	3,400	900	1,800	2,300	850	750	750
Minnesota	12,000	1,600	2,000	1,200	400	600	1,300	500	400	400
Mississippi	6,600	750	800	800	200	700	750	250	250	300
Missouri	16,000	2,200	2,500	2,300	500	1,200	1,700	450	500	550
Montana	2,000	250	300	250	70	150	200	80	80	80
Nebraska	5,300	700	900	550	150	300	500	150	200	200
Nevada	1,300	150	150	250	50	60	80	15	50	40
New Hampshire	2,800	400	500	350	90	200	250	70	80	100
New Jersey	26,000	3,800	4,500	3,300	900	1,600	1,700	1,100	750	700
New Mexico	2,200	250	250	250	60	150	150	80	70	70
New York	70,000	10,800	12,000	8,200	2,500	4,300	4,500	2,800	2,100	2,000
North Carolina	13,000	1,600	1,400	1,600	500	1,200	1,100	350	400	500
North Dakota	2,000	250	300	200	50	80	200	100	70	70
Ohio	35,000	4,900	5,600	4,500	1,300	2,500	2,700	1,200	950	1,000
Oklahoma	8,200	950	1,100	1,100	250	600	900	250	300	300
Oregon	6,800	900	1,000	950	250	500	600	200	200	300
Pennsylvania	43,000	6,300	7,400	5,100	1,500	2,800	3,300	1,500	1,300	1,300
Rhode Island	3,700	550	700	450	200	200	300	150	90	80
South Carolina	6,300	800	750	850	250	700	600	150	200	200
South Dakota	2,200	250	350	250	50	150	300	80	90	100
Tennessee	12,000	1,500	1,600	1,500	450	1,000	1,100	300	350	400
Texas	31,000	3,800	3,700	4,400	1,100	2,300	2,400	1,100	1,000	1,200
Utah	2,100	300	300	200	50	200	250	80	60	80
Vermont	1,600	200	300	200	60	100	150	50	50	50
Virginia	13,000	1,800	1,700	1,700	500	1,100	1,100	350	400	400
Washington	10,000	1,400	1,500	1,400	400	700	900	350	350	400
West Virginia	6,200	700	800	900	200	600	600	200	200	150
Wisconsin	14,000	2,200	2,500	1,400	500	850	1,300	600	400	450
Wyoming	800	100	100	90	20	60	90	20	30	30
United States	655,000	90,000	99,000	83,000	24,000	46,000	54,000	23,000	20,000	21,000

*Does not include carcinoma-in-situ of the uterine cervix or superficial skin cancers. These estimates are offered as a rough guide and should not be regarded as definitive. They are calculated according to the distribution of estimated 1974 cancer deaths by state. Especially note that year to year