

Health Effects Assessment for 1,1-Dichloroethylene (*Vinylidene Chloride*)

U.S. Environmental Protection Agency

September 1984

U.S. DEPARTMENT OF COMMERCE
National Technical Information Service

NTIS

SL 065456

PB86-134624
EPA/540/1-86/051
September 1984

HEALTH EFFECTS ASSESSMENT
FOR 1,1-DICHLOROETHYLENE

U.S. Environmental Protection Agency
Office of Research and Development
Office of Health and Environmental Assessment
Environmental Criteria and Assessment Office
Cincinnati, OH 45268

U.S. Environmental Protection Agency
Office of Emergency and Remedial Response
Office of Solid Waste and Emergency Response
Washington, DC 20460

REPRODUCED BY
NATIONAL TECHNICAL
INFORMATION SERVICE
U.S. DEPARTMENT OF COMMERCE
SPRINGFIELD, VA 22161

SL 065457

DISCLAIMER

This report has been funded wholly or in part by the United States Environmental Protection Agency under Contract No. 68-03-3112 to Syracuse Research Corporation. It has been subject to the Agency's peer and administrative review, and it has been approved for publication as an EPA document. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

PREFACE

This report summarizes and evaluates information relevant to a preliminary interim assessment of adverse health effects associated with 1,1-dichloroethylene. All estimates of acceptable intakes and carcinogenic potency presented in this document should be considered as preliminary and reflect limited resources allocated to this project. Pertinent toxicologic and environmental data were located through on-line literature searches of the Chemical Abstracts, TOXLINE, CANCERLINE and the CHEMFATE/DATALOG data bases. The basic literature searched supporting this document is current up to September, 1984. Secondary sources of information have also been relied upon in the preparation of this report and represent large-scale health assessment efforts that entail extensive peer and Agency review. The following Office of Health and Environmental Assessment (OHEA) sources have been extensively utilized:

U.S. EPA. 1980b. Ambient Water Quality Criteria for Dichloroethylenes. Environmental Criteria and Assessment Office, Cincinnati, OH. EPA 440/5-80-041. NTIS PB 81-117525.

U.S. EPA. 1983b. Health Assessment Document for Vinylidene Chloride. Environmental Criteria and Assessment Office, Research Triangle Park, NC. EPA 600/8-83-031A. NTIS PB 84-126762.

The intent in these assessments is to suggest acceptable exposure levels whenever sufficient data were available. Values were not derived or larger uncertainty factors were employed when the variable data were limited in scope tending to generate conservative (i.e., protective) estimates. Nevertheless, the interim values presented reflect the relative degree of hazard associated with exposure or risk to the chemical(s) addressed.

Whenever possible, two categories of values have been estimated for systemic toxicants (toxicants for which cancer is not the endpoint of concern). The first, the AIS or acceptable intake subchronic, is an estimate of an exposure level that would not be expected to cause adverse effects when exposure occurs during a limited time interval (i.e., for an interval that does not constitute a significant portion of the lifespan). This type of exposure estimate has not been extensively used or rigorously defined, as previous risk assessment efforts have been primarily directed towards exposures from toxicants in ambient air or water where lifetime exposure is assumed. Animal data used for AIS estimates generally include exposures with durations of 30-90 days. Subchronic human data are rarely available. Reported exposures are usually from chronic occupational exposure situations or from reports of acute accidental exposure.

The AIC, acceptable intake chronic, is similar in concept to the ADI (acceptable daily intake). It is an estimate of an exposure level that would not be expected to cause adverse effects when exposure occurs for a significant portion of the lifespan [see U.S. EPA (1980a) for a discussion of this concept]. The AIC is route specific and estimates acceptable exposure for a given route with the implicit assumption that exposure by other routes is insignificant.

Composite scores (CSs) for noncarcinogens have also been calculated where data permitted. These values are used for ranking reportable quantities; the methodology for their development is explained in U.S. EPA (1983a).

For compounds for which there is sufficient evidence of carcinogenicity, AIS and AIC values are not derived. For a discussion of risk assessment methodology for carcinogens refer to U.S. EPA (1980a). Since cancer is a process that is not characterized by a threshold, any exposure contributes an increment of risk. Consequently, derivation of AIS and AIC values would be inappropriate. For carcinogens, q1*s have been computed based on oral and inhalation data if available.

ABSTRACT

In order to place the risk assessment in proper context, the reader is referred to the preface of this document. The preface outlines limitations applicable to all documents of this series as well as the appropriate use and interpretation of the quantitative estimates.

Evidence for the carcinogenicity of 1,1-dichloroethylene in animals is limited. There are essentially no useful human data. A number of oral exposure experiments have yielded negative results, as have inhalation studies. One inhalation study showed a significant increase in the incidence of kidney adenocarcinomas in mice exposed to 1,1-dichloroethylene vapors. Mutagenicity evaluations in numerous in vitro systems have yielded positive results.

The U.S. EPA (1983b) has used the kidney adenocarcinoma data in male mice for the computation of a human q_1^* for inhalation exposure to 1,1-dichloroethylene and derived a q_1^* of $1.47 \times 10^{-1} \text{ (mg/kg bw/day)}^{-1}$.

ACKNOWLEDGEMENTS

The initial draft of this report was prepared by Syracuse Research Corporation under Contract No. 68-03-3112 for EPA's Environmental Criteria and Assessment Office, Cincinnati, OH. Dr. Christopher DeRosa and Karen Blackburn were the Technical Project Monitors and Helen Ball was the Project Officer. The final documents in this series were prepared for the Office of Emergency and Remedial Response, Washington, DC.

Scientists from the following U.S. EPA offices provided review comments for this document series:

- Environmental Criteria and Assessment Office, Cincinnati, OH
- Carcinogen Assessment Group
- Office of Air Quality Planning and Standards
- Office of Solid Waste
- Office of Toxic Substances
- Office of Drinking Water

Editorial review for the document series was provided by:

- Judith Olsen and Erma Durden
- Environmental Criteria and Assessment Office
- Cincinnati, OH

Technical support services for the document series was provided by:

- Bette Zwyer, Pat Daunt, Karen Mann and Jacky Bohanon
- Environmental Criteria and Assessment Office
- Cincinnati, OH

TABLE OF CONTENTS

	<u>Page</u>
1. ENVIRONMENTAL CHEMISTRY AND FATE.	1
2. ABSORPTION FACTORS IN HUMANS AND EXPERIMENTAL ANIMALS	2
2.1. ORAL	2
2.2. INHALATION	2
3. TOXICITY IN HUMANS AND EXPERIMENTAL ANIMALS	3
3.1. SUBCHRONIC	3
3.1.1. Oral.	3
3.1.2. Inhalation.	3
3.2. CHRONIC.	6
3.2.1. Oral.	6
3.2.2. Inhalation.	8
3.3. TERATOGENICITY AND OTHER REPRODUCTIVE EFFECTS.	8
3.3.1. Oral.	8
3.3.2. Inhalation.	8
3.4. TOXICANT INTERACTIONS.	9
4. CARCINOGENICITY	10
4.1. HUMAN DATA	10
4.1.1. Oral.	10
4.1.2. Inhalation.	10
4.2. BIOASSAYS.	10
4.2.1. Oral.	10
4.2.2. Inhalation.	10
4.3. OTHER RELEVANT DATA.	13
4.4. WEIGHT OF EVIDENCE	17
5. REGULATORY STANDARDS AND CRITERIA	18

TABLE OF CONTENTS (cont.)

	<u>Page</u>
6. RISK ASSESSMENT	19
6.1. ACCEPTABLE INTAKE SUBCHRONIC (AIS)	19
6.2. ACCEPTABLE INTAKE CHRONIC (AIC).	19
6.3. CARCINOGENIC POTENCY (q_1^*)	19
6.3.1. Oral.	19
6.3.2. Inhalation.	19
7. REFERENCES.	20
APPENDIX A: Summary Table for 1,1-Dichloroethylene	31
APPENDIX B1: Cancer Data Sheet for Derivation of q_1^*	32
APPENDIX B2: Calculation of q_1^*	33

LIST OF TABLES

<u>No.</u>	<u>Title</u>	<u>Page</u>
3-1	Effect on Experimental Animals of Long Term Inhalation of 1,1-Dichloroethylene.	4
3-2	Pathologic Effects of Long-Term Ingestion of 1,1-Dichloroethylene Incorporated in the Drinking Water of Sprague-Dawley Rats	7
4-1	Results of Oral Carcinogenicity Bioassays of 1,1-Dichloroethylene.	11
4-2	Results of Inhalation Carcinogenicity Bioassay of 1,1-Dichloroethylene.	12
4-3	Distribution of the Different Types of Mammary Tumors After Exposure by Inhalation to 1,1-Dichloroethylene in Air After 137 Weeks.	14
4-4	Distribution of the Different Types of Tumors After Exposure by Inhalation to 1,1-Dichloroethylene in Air After 121 Weeks	15

LIST OF ABBREVIATIONS

ADI	Acceptable daily intake
AIC	Acceptable intake chronic
AIS	Acceptable intake subchronic
BCF	Bioconcentration factor
bw	Body weight
CAS	Chemical Abstract Service
CS	Composite score
LOEL	Lowest-observed-effect level
ppm	Parts per million
STEL	Short-term exposure limit
TLV	Threshold limit value
TWA	Time-weighted average

1. ENVIRONMENTAL CHEMISTRY AND FATE

The relevant physical and chemical properties and environmental fate of 1,1-dichloroethylene (CAS No. 75-35-4), commonly known as vinylidene chloride, are as follows:

Chemical class:	halogenated aliphatic hydrocarbon
Molecular weight:	96.94 (Mabey et al., 1981)
Vapor pressure:	600 mm Hg at 25°C (U.S. EPA, 1983b)
Water solubility:	2250 mg/l at 25°C (U.S. EPA, 1983b)
Octanol/water partition coefficient:	69 (estimated) (Mabey et al., 1981)
BCF:	7 (estimated from the equation of Veith et al., 1979)
Half-lives in	
Air:	2 days (Cupitt, 1980)
Water:	1-6 days (estimated)

The half-life of 1,1-dichloroethylene in aquatic media has been estimated from the reaeration rate ratio of 0.601 and the oxygen reaeration rate constant of 0.19-0.96 day⁻¹ as given by Mabey et al. (1981).

The half-life of 1,1-dichloroethylene in soil could not be located in the literature available; however, evaporation is expected to be the predominant loss mechanism from the soil surface. Based on the octanol/water partition coefficient and aqueous solubility it can be speculated that leaching may play a significant role in determining the fate of this chemical in soils. In fact, the detection of this compound in several groundwaters (U.S. EPA, 1983b) is indicative of the leachability of this compound from soils.

2. ABSORPTION FACTORS IN HUMANS AND EXPERIMENTAL MAMMALS

2.1. ORAL

A number of investigators have reported the rapid appearance of labeled 1,1,-dichloroethylene in the urine and expired air of rats given an intragastric dose of [^{14}C] 1,1-dichloroethylene (position of label unspecified) (McKenna et al., 1978; Reichert et al., 1979; Jones and Hathway, 1978). These investigators concluded that the systemic absorption of 1,1-dichloroethylene following intragastric administration is rapid and fairly complete.

2.2. INHALATION

Andersen et al. (1979) exposed fasted male rats to atmospheres containing various concentrations of 1,1-dichloroethylene in a closed chamber. They observed an initial rapid phase followed by a slow phase of uptake. The rapid phase was believed to represent whole body equilibrium and was first-order with a rate constant of 2.2 hour^{-1} ($t_{1/2}=0.315 \text{ hour}$). The slow phase was believed to represent metabolism.

3. TOXICITY IN HUMANS AND EXPERIMENTAL ANIMALS

3.1. SUBCHRONIC

3.1.1. Oral. Rampy et al. (1977) administered 50, 100 or 200 mg 1,1-dichloroethylene/l drinking water to groups of 10 male and 10 female Sprague-Dawley rats for 90 days. There was an increased incidence of cytoplasmic vacuolization of hepatocytes in the high dose groups.

Quast et al. (1983) administered capsules containing 1,1-dichloroethylene in peanut oil (0, 6.25, 12.5 or 25 mg/kg bw/day) to groups of four male and four female beagle dogs for 97 days. No effects were observed on general appearance or demeanor, body weight, food consumption, hematology, clinical chemistry, urinalysis, organ weights, gross pathology or histopathology.

3.1.2. Inhalation. Subchronic exposure to 1,1-dichloroethylene results predominantly in damage to the liver and kidneys (Irish, 1962; Prendergast et al., 1967; Gage, 1970; Rampy et al., 1977). The most thorough of these studies was the study by Prendergast et al. (1967) in which rats, guinea pigs, rabbits, dogs and monkeys were exposed to atmospheric concentrations ranging from 20-395 mg/m³ 1,1-dichloroethylene for up to 90 days (Table 3-1). Intermittent exposure (8 hours/day, 5 days/week, for 6 weeks) to 395 mg/m³ produced no deaths, visible signs of toxicity or histopathological changes in any species. At each dose level, the inhalation chamber typically contained 50 rats (Long-Evans or Sprague-Dawley), 15 Hartley guinea pigs, 3 squirrel monkeys, 3 New Zealand rabbits and 2 beagle dogs. The controls consisted of 304 rats, 314 guinea pigs, 57 monkeys, 48 rabbits and 34 dogs. Continuous exposure to concentrations up to 189 mg/m³ produced dose-related mortality in guinea pigs and monkeys. For guinea pigs, mortality was significantly increased over controls in the three highest dose

TABLE 3-1

Effect on Experimental Animals of Long Term Inhalation of 1,1-Dichloroethylene*

Concentration	Schedule	Species	Mortality	Significant Findings
100 ppm (395±32 mg/m ³)	30 exposures, 8 hours/day, 5 days/week	rat	0/15	None
		guinea pig	0/15	None
		rabbit	0/3	Weight loss in treated animals
		dog	0/2	None
		monkey	0/3	Weight loss in treated animals
48 ppm (189±6.2 mg/m ³)	90 days, 24 hours/day	rat	0/15	Animals gained less weight than controls. Hepatic lesions. Renal lesions.
		guinea pig	7/15	Mortality occurred between day 4 and day 9 of exposure. Slight elevation of liver alkaline phosphatase activity and serum glutamic-pyruvic transaminase activity.
		dog	0/2	Animals lost weight. Hepatic lesions. One dog developed an adrenal cortical adenoma.
		monkey	3/9	Mortality occurred on days 26, 60 and 64. Animals lost weight. Hepatic lesions.
26 ppm (101±4.4 mg/m ³)	90 days, 24 hours/day	rat	0/15	None
		guinea pig	3/15	Mortality occurred between day 3 and day 5 of exposure.
		rabbit	0/3	Animals lost weight.
		dog	0/2	Animals lost weight.
		monkey	2/3	Mortality occurred between day 3 and day 6 of exposure.

SL 065470

TABLE 3-1 (cont.)

Concentration	Schedule	Species	Mortality	Significant Findings
16 ppm (61±5.7 mg/m ³)	90 days, 24 hours/day	rat	0/15	Animals gained less weight than controls.
		guinea pig	3/15	Mortality occurred on day 3 and day 4.
		dog	0/2	None
		monkey	0/9	Animals lost weight.
5 ppm (20±2.1 mg/m ³)	90 days, 24 hours/day	rat	2/45	Animals gained less weight than controls.
		guinea pig	2/45	None
		dog	0/6	Animals lost weight.
		monkey	1/21	None
Control	NR	rat	7/304	None
		guinea pig	2/314	
		rabbit	2/48	
		dog	0/34	
		monkey	1/57	

*Source: Prendergast et al., 1967

NR = Not reported

groups (61, 101 and 189 mg/m³) and was of borderline significance at the 20 mg/m³ exposure level ($p=0.078$, 1-tailed Fisher exact test). Growth depression was noted in all species at the high dose level. Renal lesions were observed in rats and hepatic lesions and/or enzyme changes were observed in all species at the high dose level; however, no histological lesions were found at concentrations of <101 mg/m³. Depressed weight gain and increased mortality were observed in some species at all exposure levels.

3.2. CHRONIC

3.2.1. Oral. 1,1-Dichloroethylene was administered in the drinking water (50, 100 or 200 mg/L) to groups of 48 male and 48 female Sprague-Dawley rats for 2 years (Quast et al., 1983; Humiston et al., 1978; Rampy et al., 1977). These water concentrations provided doses of 5-12, 8-20 and 16-40 mg/kg bw/day, respectively. The control groups consisted of 80 males and 80 females. 1,1-Dichloroethylene had no significant effect on general appearance, body weight, food consumption, water consumption, hematology, urinalysis, clinical chemistry or organ weights. Gross and histopathological examination revealed a number of statistically significant lesions (Table 3-2), the most important of which were hepatocellular fatty changes and periportal hepatocellular hypertrophy.

The chronic toxicity of 1,1-dichloroethylene has also been determined in F344/N rats and B6C3F1/N mice. Fifty animals/sex/dose were given 0, 1 or 5 mg/kg bw/day (rats) or 0, 2 or 10 mg/kg bw/day (mice) by gavage, 5 days/week for 2 years (NTP, 1982). Mortality and growth rates were not affected at either dose level in either species. Increased incidences of chronic renal inflammation were observed in the high dose rats (males, 43/48 vs. 26/50 controls; females, 9/44 vs. 3/49 controls) and liver necrosis in high dose male mice (7/49 vs. 1/46 controls).

TABLE 3-2

Pathologic Effects of Long-Term Ingestion of 1,1-Dichloroethylene
Incorporated in the Drinking Water of Sprague-Dawley Rats*

Effect	Dose Level					
	50 ppm		100 ppm		200 ppm	
	M	F	M	F	M	F
Increased incidence of intra-abdominal fluid or blood in abdominal cavity			✓			
Increased incidence in the total number of rats with hepatocellular fatty change or fatty degeneration				✓	✓	✓
Increased incidence of hepatocellular fatty change with location in lobule not specified		✓		✓		✓
Increased incidence in periportal hepatocellular fatty change						✓
Increased incidence of periportal hepatocellular hypertrophy		✓		✓	✓	✓
Increased incidence of hepatic centrilobular atrophy						✓
Increased incidence of mammary gland fibroadenomas/adenofibromas		✓				

*Source: U.S. EPA, 1983b; Humiston et al., 1978

M = male; F = female

3.2.2. Inhalation. Rampy et al. (1977) and McKenna et al. (1982) reported the results of a Dow Chemical Company 2-year inhalation study in which groups of 104 male and 104 female Sprague-Dawley rats were exposed to 0, 10 or 40 ppm (0, 39.7 or 158.6 mg/m³) 1,1-dichloroethylene 6 hours/day, 5 days/week, for 5 weeks, after which the exposures were increased to 0, 25 or 75 ppm (0, 99.1 or 297.4 mg/m³) for the remainder of the 18-month exposure period (~73 weeks). No dose-related changes were observed in mortality, body weight, hematology or clinical chemistry. Hepatocellular fatty changes were observed in both sexes at both dose levels. This effect was reversible after treatment was discontinued. Similar hepatic changes have been reported in mice and rats exposed to 55 ppm (218.1 mg/m³) 6 hours/day, 5 days/week for 6-12 months (TWA=38.9 mg/m³) (Lee et al., 1977; Hong et al., 1981).

3.3. TERATOGENICITY AND OTHER REPRODUCTIVE EFFECTS

3.3.1. Oral. Nitschke et al. (1980, 1983) administered 1,1-dichloroethylene in the drinking water (50, 100 or 200 mg/l) to Sprague-Dawley rats for 3 generations. Each parental generation consisted of 10 males and 20 females. There were a number of statistically significant effects on reproduction; however, these effects were not dose-related and occurred sporadically throughout the 3 generations. These effects were, therefore, probably not related to exposure to 1,1-dichloroethylene. At doses of 100 and 200 mg/l, statistically significant increases in hepatocellular fatty changes were observed in the F₁ males and females and the F₂ females.

3.3.2. Inhalation. The teratogenicity of inhaled 1,1-dichloroethylene has been tested in rats, rabbits and mice (Short et al., 1977a; Murray et al., 1979). Signs of fetal toxicity, minor skeletal alterations and soft-tissue alterations were observed at doses that produced maternal toxicity

and were considered to be fetotoxic and embryotoxic manifestations of maternal toxicity. Maternal toxicity in rats was observed by Short et al. (1977a) at exposure levels as low as 15 ppm.

3.4. TOXICANT INTERACTIONS

The metabolism of dichloroethylenes involves the production of reactive epoxide intermediates that bind covalently to cellular macromolecules (Bonse et al., 1975; Hathway, 1977; McKenna et al., 1978). Compounds such as disulfiram decrease the covalent binding of 1,1-dichloroethylene and protect against lethality and hepatotoxicity (Short et al., 1977b). Pretreatment with inducers of microsomal enzyme systems also decreased the hepatotoxicity of 1,1-dichloroethylene (Reynolds et al., 1975; Jenkins et al., 1972), but increased mortality (Carlson and Fuller, 1972). Compounds that deplete cellular glutathione increase the hepatotoxicity of 1,1-dichloroethylene (Jaeger et al., 1973a,b, 1974, 1977).

4. CARCINOGENICITY

4.1. HUMAN DATA

4.1.1. Oral. Pertinent data regarding the oral carcinogenicity of 1,1-dichloroethylene in humans could not be located in the available literature.

4.1.2. Inhalation. Ott et al. (1976) investigated the effects of occupational exposure to 1,1-dichloroethylene (<0.2% vinyl chloride) among 138 Dow Chemical Company workers. TWA concentrations were estimated based on job descriptions and industrial hygiene surveys; the subjects were divided into groups exposed to <10 ppm (assumed 5 ppm average), 10-24 ppm (assumed 17 ppm average) and ≥ 25 ppm (assumed 43 ppm average). There were no statistically significant differences between the exposed groups and controls matched for age and smoking habits; however, the population examined in the study may not be adequate for the detection of cancer.

4.2. BIOASSAYS

4.2.1. Oral. The available data regarding the oral carcinogenicity of 1,1-dichloroethylene in experimental animals are summarized in Table 4-1. These studies have failed to demonstrate a carcinogenic potential for 1,1-dichloroethylene in either rats or mice following oral exposure.

4.2.2. Inhalation. The available data on the inhalation carcinogenicity of 1,1-dichloroethylene in experimental animals are summarized in Table 4-2. The only studies in which 1,1-dichloroethylene has produced positive results are those of Maltoni et al. (1977, 1980) in which Sprague-Dawley rats and Swiss mice were used. There were at least 30 animals/sex/dose at the beginning of the exposure period, with 90-100 animals/sex in the controls. Rats were exposed to atmospheres containing 0, 10, 25, 50, 100 or 150 ppm (0, 39.7, 99.1, 198.3, 396.5 or 594.8 mg/m³) 1,1-dichloroethylene, 4 hours/day, 4-5 days/week, for 12 months.

TABLE 4-1
Results of Oral Carcinogenicity Bioassays of 1,1-Dichloroethylene*

Species	Dose	Route of Administration	Total Duration of Observation (weeks)	Findings	Reference
Sprague-Dawley rats	20, 10, 5, 0.5 mg/kg for 12 months	gavage, daily	147	No statistically significant increase	Maltoni et al., 1977
Sprague-Dawley rats	50, 100, 200 ppm in drinking water	Ingestion	104	No statistically significant increase	Quast et al., 1983
Fischer 344 rats	5 mL/kg of a 1000 or 200 ppm solution	gavage, 5 days/week	103	No statistically significant increase	NTP, 1982
B6C3F1 mice	10 mL/kg of a 1000 or 200 ppm solution	gavage, 5 days/week	103	No statistically significant increase	NTP, 1982
Sprague-Dawley rats	0.5, 5, 10, 20 mg/kg/day	gavage, 5 days/week	52-59	No brain tumors	Maltoni et al., 1982

*Source: U.S. EPA, 1983b

SL 065477

TABLE 4-2

Results of Inhalation Carcinogenicity Bioassays of 1,1-Dichloroethylene*

Species	Dose	Route of Administration	Total Duration of Observation	Findings	Reference
Sprague-Dawley rats	10, 25, 50, 100, 150 ppm, 4-5 days/week for 12 months	Inhalation, 4 hours/day	137 weeks	Statistically significant increase in total mammary tumors, but not carcinomas alone, only at 10 and 100 ppm; no dose response	Malloni et al., 1977, 1980
Swiss mice	10, 25 ppm, 4-5 days/week for 12 months	Inhalation, 4 hours/day	121 weeks	Kidney carcinomas at 25 ppm in males (none in controls) Statistically significant increase in mammary carcinomas in females; no dose response	Malloni et al., 1977, 1980
Chinese hamsters	25 ppm, 4-5 days/week for 12 months	Inhalation, 4 hours/day	157 weeks	No statistically significant increase	Malloni et al., 1977, 1980
Wistar rats	200 ppm for 6 months, followed by 100 ppm for 6 months, 5 days/week	Inhalation, 4 hours/day	Lifetime	No statistically significant increase	Viola and Caputo, 1977
Sprague-Dawley rats	100, 75 ppm, 5 days/week for 12 months	Inhalation, 4 hours/day	Lifetime	No statistically significant increase	Viola and Caputo, 1977
CD-1 mice	55 ppm, 5 days/week	Inhalation, 6 hours/day	12 months	No statistically significant increase	Lee et al., 1978
CD rats	55 ppm, 5 days/week for 12 months	Inhalation, 6 hours/day	12 months	No statistically significant increase	Lee et al., 1978
Sprague-Dawley rats	25, 75 ppm for 24 months	Inhalation	104 weeks	No statistically significant increase	McKenna et al., 1982
CD mice	55 ppm, 5 days/week 1, 3 or 6 months	Inhalation, 6 hours/day	13, 15 or 18 months	No statistically significant increase	Hong et al., 1981
CD rats	55 ppm, 5 days/week 1, 3, 6 or 10 months	Inhalation, 6 hours/day	13, 15, 18 or 22 months	No statistically significant increase	Hong et al., 1981
Sprague-Dawley rats	10, 25, 50, 100, 150 ppm	Inhalation, 4-5 days/week	52 weeks	No brain tumors	Malloni et al., 1982

*Source: U.S. EPA, 1983b

SL 065478

The results of these studies are summarized in Tables 4-3 and 4-4. There were indications that 1,1-dichloroethylene induced mammary tumors in both rats and mice; however, there was no clear dose-response relationship and these tumors could not positively be attributed to exposure to 1,1-dichloroethylene. The only tumors that the authors considered related to the treatment were kidney adenocarcinomas in male mice.

4.3. OTHER RELEVANT DATA

1,1-Dichloroethylene has been tested for yeast and bacterial mutagenicity in the Ames assay, the liquid suspension assay, the host-mediated assay and exposure of bacteria to atmospheres containing 1,1-dichloroethylene, both with and without mammalian metabolic activating systems (Bartsch et al., 1975, 1979; Malaveille et al., 1977; Simon et al., 1977, 1979; Simon, 1978; Baden et al., 1976, 1977, 1978; Waskell, 1978; Greim et al., 1975; Cerna and Kypenova, 1977; Laumbach et al., 1977; Bartsch, 1976; Barbin et al., 1978; Bonse et al., 1975). 1,1-Dichloroethylene is mutagenic to Escherichia coli, Salmonella typhimurium, Bacillus subtilis and Saccharomyces cerevisiae, in the presence, but not the absence, of a mammalian metabolic activating system.

1,1-Dichloroethylene vapors have been demonstrated to be mutagenic to the plant, Tradescantia, following a 24-hour exposure to concentrations as low as 22 ppm (87.2 mg/m³); however, a 6-hour exposure to 1288 ppm (5107 mg/m³) did not produce a mutagenic effect (Van't Hof and Schairer, 1982). In contrast, negative results have been obtained in assays using cultured mammalian cells (Drevon and Kuroki, 1979) and in dominant lethal assays in mice (Andersen and Jenkins, 1977) and rats (Short et al., 1977c).

TABLE 4-3

Distribution of the Different Types of Mammary Tumors After Exposure by Inhalation to 1,1-Dichloroethylene in Air After 137 Weeks^{a,b}

Group No.	Concen- trations	Animals (Sprague-Dawley rats, 16 Weeks Old at Start)					Average Latency Time ^f (weeks)	No. of Tumors/ Tumor- Bearing Animals	X ^g	No.	Histologically Examined Histotype ^c				
		Sex	No. at Start	Corrected ^d Number	X ^e	Fibromas and Fibroadenomas					Carcinomas				
						Average Latency Time ^f (weeks)					X ^h	No.	X ^h	Average Latency Time ^f (weeks)	
I	150 ppm	M	60	60	13.3	97±14	1.0	100.0	6	75.0	109±8	1	12.5	26	
		F	60	60	73.3	82±3	1.5	97.7	38 ⁱ	88.4	83±3	9	20.9	78±8	
		M/F	120	120	43.3	82±3	1.4	98.1	44	86.3	86±3	10	19.6	73±8	
II	100 ppm	M	30	30	16.7	104±9	1.0	100.0	5	100.0	104±9	0	0	0	
		F	30	30	83.3	82±4	1.7	92.0	21 ⁱ	91.3	83±5	3	13.0	102±10	
		M/F	60	60	50.0	85±4	1.6	93.3	26	92.8	87±4	3	10.7	102±10	
III	50 ppm	M	30	30	23.3	106±5	1.0	100.0	7	100.0	106±5	0	0	0	
		F	30	30	76.7	79±4	1.9	95.6	21 ⁱ	95.4	82±4	1	4.5	68	
		M/F	60	60	50.0	86±4	1.7	96.7	28	96.5	88±4	1	3.4	68	
IV	25 ppm	M	30	28	14.3	103±10	1.0	100.0	4	100.0	103±10	0	0	0	
		F	30	30	70.0	86±4	1.6	95.2	20 ⁱ	100.0	87±4	4	20.0	82±10	
		M/F	60	58	43.1	88±4	1.5	96.0	24	100.0	90±4	4	16.7	82±10	
V	10 ppm	M	30	29	10.3	81±23	1.0	100.0	3	100.0	81±23	0	0	0	
		F	30	30	93.3	83±4	1.6	85.7	24 ⁱ	100.0	85±4	5	20.8	90±14	
		M/F	60	59	52.5	81±4	1.5	87.1	27	100.0	85±4	5	18.5	90±14	
VI	no treat- ment (controls)	M	100	87	12.6	115±6	1.0	100.0	11	100.0	115±6	0	0	0	
		F	100	99	61.6	87±2	1.5	91.8	44	78.6	88±3	16	28.6	95±5	
		M/F	200	186	38.7	91±3	1.4	93.0	55	82.1	93±3	16	23.9	95±5	

^aSource: Maltoni et al., 1980^bExposure was for 4 hours/day, 4-5 days/week for 52 weeks^cTwo or more tumors of the same and/or different types (fibroadenomas, carcinomas, sarcomas, carcinosarcomas) may be present in the same animals.^dA carcinoma was found in one male in the 150 ppm group, and no animals were observed to have sarcomas.^eFive animals after 10 weeks, when the first tumor (a leukemia) was observed.^fThe percentages refer to the corrected numbers.^gAverage age at the onset of the first mammary tumor per animal, detected at the periodic control or at autopsy.^hThe percentages refer to total numbers of animals bearing mammary tumors.ⁱThe percentages refer to total numbers of animals bearing mammary tumors, histologically examined.^jStatistically significant increase compared to control by chi-square test ($p < 0.05$). Comparisons are made between numbers with tumors/corrected numbers.

SL 065480

TABLE 4.4

Distribution of the Different Types of Tumors After Exposure by Inhalation to 1,1-Dichloroethylene in Air After 121 Weeks^{a,b}

Groups No.	Treatment		Animals (Swiss mice 16 weeks old [(Groups I,II,III,IV, V,VI) and 9 weeks old (Groups IV bis, VII at start)])		Animals with Tumors											
					Kidney Adenocarcinomas				Mammary Tumors ^c				Pulmonary Adenomas ^d			
	Concentrations	Length	Sex	No. at Start	Corrected Number ^e	No.	%	Average Latency Time ^f (weeks)	Corrected Number ^e	No.	%	Average Latency Time ^e (weeks)	Corrected Number ^e	No.	%	Average Latency Time ^f (weeks)
I	200 ppm	2 days	M	60	1	0	0	0	6	0	0	0	5	0	0	0
			F	60	28	0	0	0	53	1	1.9	87	46	1	2.2	57
			M/F	120	29	0	0	0	59	1	1.7	87	53	1	1.9	57
II	100 ppm	2 days	M	30	12	0	0	0	21	0	0	0	18	2	11.1	62±7
			F	30	13	0	0	0	28	3	10.7	46±3	26	2	7.7	53±2
			M/F	60	25	0	0	0	49	3	6.1	46±5	44	4	9.1	58±4
III	50 ppm	1 week	M	30	17	1	5.9	64	27	0	0	0	26	1	3.8	62
			F	30	14	0	0	0	28	2	7.1	39±13	27	3	11.1	80±8
			M/F	60	31	1	3.2	64	55	2	3.6	39±13	53	4	7.5	75±7
IV	25 ppm	52 weeks	M	30	21	39	14.3	71±5	29	0	0	0	28	7 ^h	25.0	73±6
			F	30	26	0	0	0	30	4 ^h	13.3	68±11	29	7 ^h	24.1	85±6
			M/F	60	47	3	6.4	71±5	59	4	6.8	68±11	57	14	24.6	30±4
IV bis	25 ppm	52 weeks	M	120	98	259	25.5	75±2	117	1	0.8	46	113	16 ^h	14.2	77±3
			F	120	112	1	0.9	77	118	12 ^h	10.2	69±4	118	11 ^h	9.3	78±6
			M/F	240	210	26	12.4	75±2	235	13	5.5	67±4	231	27	11.7	77±3
V	10 ppm	52 weeks	M	30	25	0	0	0	30	0	0	0	28	11 ^h	39.3	71±5
			F	30	26	0	0	-	30	6 ^h	20.0	63±5	30	3 ^h	16.0	68±4
			M/F	60	51	0	0	0	60	6	10.0	63±5	58	14	24.1	70±4
VI	no treatment (controls)	NA	M	100	56	0	0	0	92	1	1.1	25	80	3	3.7	66±7
			F	100	73	0	0	0	97	2	2.1	49±7	92	4	4.3	56±7
			M/F	200	129	0	0	0	189	3	1.6	41±9	172	7	4.1	60±4

SL 065481

TABLE 4-4 (cont.)

Groups No.	Treatment		Animals (Swiss mice 16 weeks old (Groups I,II,III,IV, V,VI) and 9 weeks old (Groups IV bis, VII at start))		Animals with Tumors											
					Kidney Adenocarcinomas				Mammary Tumors ^c				Pulmonary Adenomas ^d			
					Corrected Number ^e	No.	%	Average Latency Time ^f (weeks)	Corrected Number ^e	No.	%	Average Latency Time ^e (weeks)	Corrected Number ^e	No.	%	Average Latency Time ^f (weeks)
VII	no treatment (controls)	NA	M	90	70	0	0	0	80	0	0	0	73	3	4.1	56.11
			F	90	85	0	0	0	88	0	0	0	86	2	2.3	75.12
			M/F	180	155	0	0	0	168	1	0.6	83	159	5	3.1	64.8

^aSource: Maltoni et al., 1980

^bExposure was for 4 hours/day, 4-5 days/week for 52 weeks

^cAll mammary tumors in females were histologically diagnosed as carcinomas.

^dSome pulmonary adenomas were cellular atypias.

^eAlive animals when the first tumor was observed: kidney adenocarcinoma, 55 weeks; mammary tumor, 27 weeks; pulmonary adenoma, 36 weeks. The percentages refer to the corrected numbers.

^fAverage time from the start of the experiment to the detection (at the periodic control or at autopsy).

^g $p < 0.01$, combined 25 ppm (28/119) males vs. control males (0/196) by chi-square test. Based on corrected numbers.

^h $p < 0.01$ combined control males (6/153) vs. 10 ppm males (11/28) and vs. combined 25 ppm males (29/294). Also, combined control females (6/178) vs. 10 ppm females (3/30) and vs. combined 25 ppm females (18/147)

ⁱ $p < 0.01$ combined control females (3/185) vs. 10 ppm females (6/30) and vs. combined 25 ppm females (16/148). Based on corrected numbers.

NA = Not applicable

SL 065482

4.4. WEIGHT OF EVIDENCE

IARC (1982) has evaluated the evidence for carcinogenicity of 1,1-dichloroethylene and concluded that the evidence for carcinogenicity in humans is "inadequate," the evidence for carcinogenicity in animals is "limited," and the evidence for activity in short-term tests is "sufficient." Applying the criteria for weight of evidence proposed by the Carcinogen Assessment Group of the U.S. EPA (Federal Register, 1984) 1,1-dichloroethylene is most appropriately classified in Group C - Possible Human Carcinogen.

5. REGULATORY STANDARDS AND CRITERIA

The ACGIH (1980) has established a TLV of 5 ppm ($\sim 20 \text{ mg/m}^3$) and a STEL of 20 ppm ($\sim 80 \text{ mg/m}^3$), which are believed low enough to prevent overt toxicity in exposed workers.

The U.S. EPA (1980b) has estimated that an ambient water concentration of $0.33 \text{ } \mu\text{g/l}$ would result in excess carcinogenic potency over a lifetime exposure. Both NIOSH and OSHA consider 1,1-dichloroethylene to be a potential carcinogen and have established an exposure limit of 1.0 ppm ($\sim 4 \text{ mg/m}^3$) as a TWA or 5 ppm ($\sim 20 \text{ mg/m}^3$) as a 15-minute ceiling (ACGIH, 1980).

6. RISK ASSESSMENT

6.1. ACCEPTABLE INTAKE SUBCHRONIC (AIS)

1,1-Dichloroethylene is a chemical associated with cancer in animals and for which data are sufficient for computing a q_1^* . It is, therefore, inappropriate to calculate an oral or inhalation AIS for 1,1-dichloroethylene.

6.2. ACCEPTABLE INTAKE CHRONIC (AIC)

1,1-Dichloroethylene is a chemical associated with cancer in animals and for which data are sufficient for computing a q_1^* . It is, therefore, inappropriate to calculate an oral or inhalation AIC for 1,1-dichloroethylene.

6.3. CARCINOGENIC POTENCY (q_1^*)

6.3.1. Oral. No data were located in the available literature that indicated a carcinogenic potential for orally administered 1,1-dichloroethylene. Therefore, no q_1^* could be derived.

6.3.2. Inhalation. Only one study was located in the available literature that indicated a carcinogenic response to inhaled 1,1-dichloroethylene (Maltoni, et al., 1977, 1980). In this study, groups of at least 30 Swiss mice/sex/dose were exposed to 0, 39.7 or 99.1 mg 1,1-dichloroethylene/m³, 4 hours/day, 4-5 days/week, for 12 months. Kidney adenocarcinomas were observed in 28/119 male mice in the high dose groups as compared with 0/126 control male mice. The U.S. EPA (1983b) has analyzed these data and derived a q_1^* of 1.47×10^{-2} (mg/kg bw/day)⁻¹. The data base from which this q_1^* is calculated is presented in Appendix B. This assessment uses the same study as U.S. EPA (1980b); however, when the water quality document was developed, only interim results of this study were available, hence the difference in estimates.

7. REFERENCES

ACGIH (American Conference of Governmental Industrial Hygienists). 1980. Documentation of the Threshold Limit Values, 4th ed. (Includes Supplemental Documentation, 1981, 1982, 1983). Cincinnati, OH.

Andersen, M.E. and L.J. Jenkins, Jr. 1977. Oral toxicity of 1,1-dichloroethylene in the rat: Effects of sex, age and fasting. Environ. Health Perspect. 21: 157-163. (Cited in U.S. EPA, 1983b)

Andersen, M.E., M.L. Gargas, R.A. Jones and L.J. Jenkins, Jr. 1979. The use of inhalation techniques to assess the kinetic constants of 1,1-dichloroethylene metabolism. Toxicol. Appl. Pharmacol. 47(2): 395-409. (Cited in U.S. EPA, 1983b)

Baden, J.M., M. Brinkenhoff, R.S. Wharton, B.A. Hitt, V.F. Simmon and R.I. Mazze. 1976. Mutagenicity of volatile anesthetics: Halothane. Anesthesiology. 45(3): 311-318. (Cited in U.S. EPA, 1983b)

Baden, J.M., M. Kelley, R.S. Wharton, B.A. Hitt, V.F. Simmon and R.I. Mazze. 1977. Mutagenicity of halogenated ether anesthetics. Anesthesiology. 46: 346-350. (Cited in U.S. EPA, 1983b)

Baden, J.M., M. Kelley, V.F. Simmon, S.A. Rice and R.I. Mazze. 1978. Fluroxene mutagenicity. Mutat. Res. 58: 183-191. (Cited in U.S. EPA, 1983b)

Barbin, A., G. Planche, A. Croisy, C. Malaveille and H. Bartsch. 1978. Detection of electrophilic metabolites of halogenated olefins with 4-(4-nitrobenzyl) pyridine or with Salmonella typhimurium. Mutat. Res. 53: 150. (Cited in U.S. EPA, 1983b)

Bartsch, H. 1976. Mutagenicity tests in chemical carcinogenesis. In: Environmental Pollution and Carcinogenic Risk, IARC Scientific Publications No. 13. INSERM Symposia Series. IARC, Lyon, France. Vol. 52, p. 229-240. (Cited in U.S. EPA, 1983b)

Bartsch, H., C. Malaveille, R. Montesano and L. Tomatis. 1975. Tissue mediated mutagenicity of vinylidene chloride and 2-chlorobutadiene in Salmonella typhimurium. Nature (London). 225(5510): 641-643. (Cited in U.S. EPA, 1983b)

Bartsch, H., C. Malaveille, A. Barbin and G. Planche. 1979. Mutagenic and alkylating metabolites of haloethylenes, chlorobutadienes and dichlorobutenes produced by rodent or human liver tissues. Evidence for oxirane formation by P-450 linked microsomal oxygenase. Arch. Toxicol. 41: 249-277. (Cited in U.S. EPA, 1983b)

Bonse, G., T. Urban, D. Reichert and D. Henschler. 1975. Chemical reactivity, metabolic oxirane formation and biological reactivity of chlorinated ethylenes in the isolated perfused rat liver preparation. Biochem. Pharmacol. 24: 1829-1834. (Cited in U.S. EPA, 1983b)

Carlson, G.P. and G.C. Fuller. 1972. Interaction of modifiers of hepatic microsomal drug metabolism and the inhalation toxicity of 1,1-dichloroethylene. Res. Commun. Chem. Pathol. Pharmacol. 4(3): 553-560. (Cited in U.S. EPA, 1983b)

Cerna, M. and H. Kypenova. 1977. Mutagenic activity of chloroethylenes analyzed by screening system tests. Mutat. Res. 46(3): 214-215. (Cited in U.S. EPA, 1983b)

Cupitt, L.T. 1980. Fate of Toxic and Hazardous Materials in the Air Environment. U.S. EPA, Environmental Sciences Research Laboratory, ORD, Research Triangle Park, NC. EPA 600/3-80-084. NTIS PB 80-221948.

Drevon, C. and T. Kuroki. 1979. Mutagenicity of vinyl chloride, vinylidene chloride and chloroprene in V79 Chinese-hamster cells. Mutat. Res. 67(20): 173-182. (Cited in U.S. EPA, 1983b)

Federal Register. 1984. Environmental Protection Agency. Proposed guidelines for carcinogenic risk assessment. 49 FR 46294-46299.

Gage, J.C. 1970. The subacute inhalation toxicity of 109 industrial chemicals. Br. J. Ind. Med. 27(1): 1-18. (Cited in U.S. EPA, 1983b)

Greim, H., G. Bonse, Z. Radwan, D. Reichert and D. Henschler. 1975. Mutagenicity in vitro and potential carcinogenicity of chlorinated ethylenes as a function of metabolic oxirane formation. Biochem. Pharmacol. 24(21): 2013-2017. (Cited in U.S. EPA, 1983b)

Hathway, D.E. 1977. Comparative mammalian metabolism of vinyl chloride and vinylidene chloride in relation to oncogenic potential. Environ. Health Perspect. 21: 55-59. (Cited in U.S. EPA, 1983b)

Hong, C.B., J.M. Winston, L.P. Thornburg, C.C. Lee and J.S. Woods. 1981. Follow-up study on the carcinogenicity of vinyl chloride and vinylidene chloride in rats and mice. Tumor incidence and mortality subsequent to exposure. J. Toxicol. Environ. Health. 7: 909-924. (Cited in U.S. EPA, 1983b)

Humiston, C.G., J.F. Quast, C.E. Wade, J. Ballard, J.E. Beyer and R.W. Lisowe. 1978. Results of a two-year toxicity and oncogenicity study with vinylidene chloride incorporated in the drinking water of rats. MCA Report No. VCD 1.3-Tox-0r1-Dow. Toxicology Research Laboratory Health and Environmental Research, Dow Chemical USA, Midland, MI. (Cited in U.S. EPA, 1983b)

IARC (International Agency for Research on Cancer). 1982. Results and conclusions. In: Chemicals, Industrial Processes and Industries Associated with Cancer in Humans. IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans. WHO, IARC, Lyon, France. Vol. 1-29 (Suppl 4).

Irish, D.D. 1962. Aliphatic halogenated hydrocarbons. In: Patty's Industrial Hygiene and Toxicology, 2nd ed., Vol. 2, F.A. Patty, Ed. John Wiley and Sons, Inc., NY. p. 1305-1308. (Cited in U.S. EPA, 1983b)

Jaeger, R.J., R.B. Conolly and S.D. Murphy. 1973a. Diurnal variation of hepatic glutathione concentration and its correlation with 1,1-dichloroethylene inhalation toxicity in rats. Res. Commun. Chem. Pathol. Pharmacol. 6(2): 465-471. (Cited in U.S. EPA, 1983b)

Jaeger, R.J., M.J. Trabulus and S.D. Murphy. 1973b. Biochemical effects of 1,1-dichloroethylene in rats. Dissociation of its hepatotoxicity from a lipoperoxidative mechanism. Toxicol. Appl. Pharmacol. 24(3): 457-467. (Cited in U.S. EPA, 1983b)

Jaeger, R.J., R.B. Conolly and S.D. Murphy. 1974. Effect of 18-hour fast and glutathione depletion on 1,1-dichloroethylene-induced hepatotoxicity and lethality in rats. Exp. Mol. Pathol. 20(2): 187-198. (Cited in U.S. EPA, 1983b)

Jaeger, R.J., S. Szabo and L.J. Coffman. 1977. 1,1-Dichloroethylene hepatotoxicity: Effect of altered thyroid function and evidence for the subcellular site of injury. J. Toxicol. Environ. Health. 3(3): 545-555. (Cited in U.S. EPA, 1983b)

Jenkins, L.J., Jr., M.J. Trabulus and S.D. Murphy. 1972. Biochemical effects of 1,1-dichloroethylene in rats. Comparison with carbon tetrachloride and 1,2-dichloroethylene. Toxicol. Appl. Pharmacol. 23(3): 501-510. (Cited in U.S. EPA, 1983b)

Jones, B.K. and D.E. Hathway. 1978. The biological fate of vinylidene chloride in rats. Chem. Biol. Interact. 20(1): 27-41. (Cited in U.S. EPA, 1983b.)

Laumbach, A.D., S. Lee, J. Wong and U.N. Streips. 1977. Studies on the mutagenicity of vinyl chloride metabolites and related chemicals. Prev. Detect. Cancer Proc. Int. Symp. 1: 155-170. (Cited in U.S. EPA, 1983b)

Lee, C.C., J.C. Bhandari, J.M. Winston, et al. 1977. Inhalation toxicity of vinyl chloride and vinylidene chloride. Health Perspect. 21: 25-32. (Cited in U.S. EPA, 1983b)

Lee, C.C., J.C. Bhandari, J.M. Winston, W.B. House, R.L. Dixon and J.S. Woods. 1978. Carcinogenicity of vinyl chloride and vinylidene chloride. J. Toxicol. Environ. Health. 4(1): 15-30. (Cited in U.S. EPA, 1983b)

Mabey, W.R., J.H. Smith, R.T. Podoll, et al. 1981. Aquatic Fate Process Data for Organic Priority Pollutants. U.S. EPA, Monitoring and Data Support Division, Office of Water Regulations and Standards, Washington, DC. EPA 440/4-81-014.

Malaveille, C., G. Planche and H. Bartsch. 1977. Factors for efficiency of the Salmonella microsome mutagenicity assay. Chem. Biol. Interact. 17(2): 129-136. (Cited in U.S. EPA, 1983)

Maltoni, C., G. Cotti, L. Morisi and P. Chieco. 1977. Carcinogenicity bioassays of vinylidene chloride. Research plan and early results. Med. Lav. 68(4): 241-262. (Cited in U.S. EPA, 1983b)

Maltoni, C., G. Cotti, L. Morisi and P. Chieco. 1980. Toxicity and Carcinogenicity Bioassay of Vinylidene Chloride II. Chronic Toxicity and Carcinogenicity Report of Institute of Oncology and Tumor Center. Bologna, Italy. (Cited in U.S. EPA, 1983b)

Maltoni, C., A. Ciliberti and D. Carretti. 1982. Experimental contributions in identifying brain potential carcinogens in the petrochemical industry. Ann. NY Acad. Sci. 381: 216-249. (Cited in U.S. EPA, 1983b)

McKenna, M.J., J.A. Zemple, E.O. Madrid, W.H. Braun and P.J. Gehring. 1978. Metabolism and pharmacokinetic profile of vinylidene chloride in rats following oral administration. Toxicol. Appl. Pharmacol. 45(3): 821-835. (Cited in U.S. EPA, 1983b)

McKenna, M.J., J.F. Quast, H.O. Yakel, M.F. Balmer and L.W. Rampy. 1982. Vinylidene Chloride: A Chronic Inhalation Toxicity and Oncogenicity Study in Rats. Toxicology Research Laboratory, Health and Environmental Sciences, Dow Chemical USA, Midland, MI. (Cited in U.S. EPA, 1983b)

Murray, F., K. Nitschke, L. Rampy and B. Schwetz. 1979. Embryotoxicity and fetotoxicity of inhaled or ingested vinylidene chloride in rats and rabbits. Toxicol. Appl. Pharmacol. 48: 189-202. (Cited in U.S. EPA, 1983b)

Nitschke, K.D., B.A. Schwetz, C.G. Humiston, et al. 1980. A Multiple Generation Reproduction Study in Rats Maintained on Drinking Water Containing Vinylidene Chloride. Toxicology Research Laboratory, Health and Environmental Research, Dow Chemical USA, Midland, MI. (Cited in U.S. EPA, 1983b)

Nitschke, K.D., F.A. Smith, J.F. Quast, J.M. Norris and B.A. Schwetz. 1983. A three-generation rat reproductive toxicity study of vinylidene chloride in the drinking water. Fundam. Appl. Toxicol. 3: 75-79. (Cited in U.S. EPA, 1983b)

NTP (National Cancer Institute/National Toxicology Program). 1982. NTP Technical Report on the Carcinogenesis Bioassay of Vinylidene Chloride in F344/N Rats and B6C3F1/Mice (Gavage Study). NPT No. 80-82. NIH Publ. No. 82-1784. NTP, Research Triangle Park, NC and Bethesda, MD. U.S. DHHS, PHS, National Institute of Health. (Cited in U.S. EPA, 1983b)

Ott, M.G., W.A. Fishbeck, J.C. Townsend and E.J. Schneider. 1976. A health study of employees exposed to vinylidene chloride. J. Occup. Med. 18(11): 735-738. (Cited in U.S. EPA, 1983b)

Prendergast, J.A., R.A. Jones, L.J. Jenkins, Jr. and J. Siegel. 1967. Effects on experimental animals of long-term inhalation of trichloroethylene, carbon tetrachloride, 1,1,1-trichloroethane, dichlorodifluoromethane and 1,1-dichloroethylene. Toxicol. Appl. Pharmacol. 10(2): 270-289.

Quast, J.F., C.G. Humiston, C.E. Wade, et al. 1983. A chronic toxicity and oncogenicity study in rats and subchronic toxicity study in dogs on ingested vinylidene chloride. Fund. Appl. Toxicol. 3(1): 55-62. (Cited in U.S. EPA, 1983b)

Rampy, L.W., J.F. Quast, C.G. Humiston, M.F. Balmer and B.A. Schwetz. 1977. Interim results of two-year toxicological studies in rats of vinylidene chloride incorporated in the drinking water or administered by repeated inhalation. Environ. Health Perspect. 21: 33-43. (Cited in U.S. EPA, 1983b)

Reichert, D., H.W. Werner, M. Metzler and D. Henschler. 1979. Molecular mechanism of 1,1-dichloroethylene toxicity: Excreted metabolites reveal different pathways of reactive intermediates. Arch. Toxicol. 42(3): 159-169. (Cited in U.S. EPA, 1983b)

Reynolds, E.S., M.T. Moslen, S. Szabo, R.J. Jaeger and S.D. Murphy. 1975. Hepatotoxicity of vinyl chloride and 1,1-dichloroethylene: Role of mixed function oxidase system. Am. J. Pathol. 81: 219. (Cited in U.S. EPA, 1983b)

Short, R.D., Jr., J.L. Minor, J.M. Winston, B. Ferguson and T. Unger. 1977a. Toxicity Studies of Selected Chemicals. Task II. The Developmental Toxicity of Vinylidene Chloride Inhaled by Rats and Mice During Gestation. Prepared by Midwest Research Institute, Kansas City, MO, under Contract 68-01-3242. U.S. EPA, Office of Toxic Substances, Washington, DC. EPA 560/6-77-022. (Cited in U.S. EPA, 1983b)

Short, R.D., J.M. Winston, J.L. Minor, C.D. Hong, J. Seifter and C.C. Lee. 1977b. Toxicity of vinylidene chloride in mice and rats and its alterations by various treatments. J. Toxicol. Environ. Health. 3(5-6): 913-921. (Cited in U.S. EPA, 1983b)

Short, R.D., J.L. Minor, J.M. Winston and C.C. Lee. 1977c. A dominant lethal study in male rats after repeated exposures to vinyl chloride or vinylidene chloride. J. Toxicol. Environ. Health. 3: 5-6. (Cited in U.S. EPA, 1983b)

Simmon, V.F. 1978. Structural correlations of carcinogenic and mutagenic alkyl halides. In: Structural Correlations between Carcinogenicity and Mutagenicity, DHEW (ISS-FDA) Publication No. 78-1046, p. 163-171. (Cited in U.S. EPA, 1983b)

Simmon, V.F., K. Kauhanen and R.C. Tardiff. 1977. Mutagenic activity of chemicals identified in drinking water. In: Progress in Genetic Toxicology. (Cited in U.S. EPA, 1983b)

Simmon, V.F., H.S. Rosenkranz, E. Zeiger and L.A. Poirier. 1979. Mutagenic activity of chemical carcinogens and related compounds in the intraperitoneal host-mediated assay. J. Natl. Cancer Inst. 62: 911-918. (Cited in U.S. EPA, 1983b)

U.S. EPA. 1980a. Guidelines and Methodology Used in the Preparation of Health Effects Assessment Chapters of the Consent Decree Water Quality Criteria. Federal Register. 45:79347-79357.

U.S. EPA. 1980b. Ambient Water Quality Criteria for Dichloroethylenes. Environmental Criteria and Assessment Office, Cincinnati, OH. EPA 440/5-80-041. NTIS PB 81-117525.

U.S. EPA. 1983a. Methodology and Guidelines for Reportable Quantity Determinations Based on Chronic Toxicity Data. Prepared by the Environmental Criteria and Assessment Office, Cincinnati, OH, OHEA for the Office of Solid Waste and Emergency Response, Washington, DC.

U.S. EPA. 1983b. Health Assessment Document for Vinylidene Chloride. Environmental Criteria and Assessment Office, Research Triangle Park, NC. EPA 600/8-83-031A. NTIS PB 84-126762.

Van't Hof, J. and L.A. Schairer. 1982. Tradescantia assay systems for gaseous mutagens. A report of the U.S. Environmental Protection Agency Gene Tox Program. Mutat. Res. 99(3): 303-315. (Cited in U.S. EPA, 1983b)

Veith, G.D., D.L. DeFoe and B.V. Bergstedt. 1979. Measuring and estimating the bioconcentration factors of chemicals in fish. J. Fish Res. Board Can. 36: 1040-1048.

Viola, P.L. and A. Caputo. 1977. Carcinogenicity studies on vinylidene chloride. Environ. Health Perspect. 21: 45-47. (Cited in U.S. EPA, 1983b)

Waskell, L. 1978. Study of the mutagenicity of anesthetics and their metabolites. Mutat. Res. 57: 141-153. (Cited in U.S. EPA, 1983b)

APPENDIX A

Summary Table for 1,1-Dichloroethylene

	Species	Experimental Dose/Exposure	Effect	q1*	Reference
Inhalation					
				ND	
				ND	
	mouse	39.7 or 99.1 mg/m ³	kidney adenocarcinomas	1.47×10^{-1} (mg/kg bw/day) ⁻¹	Maltoni et al., 1977, 1980; U.S. EPA, 1983b
Oral					
				ND	
				ND	
				ND	

ND = Not derived

SL 065497

APPENDIX B1

Cancer Data Sheet for Derivation of q_1^*

Compound: 1,1-dichloroethylene

Reference: Maltoni et al., 1980

Species, Strain, Sex: mice, Swiss, male

Body weight: 0.03 kg (assumed)

Length of exposure (t_e) = 52 weeks

Length of experiment (L_e) = 104 weeks

Lifespan of animal (L) = 104 weeks

Tumor site and type: kidney, adenocarcinoma

Route, vehicle: inhalation

Experimental Doses or Exposures (ppm)	Transformed Dose† (ppm)	Incidence
		No. Responding/No. Tested or Examined
0	0	0/126
10	0.54	0/25
25	1.35	28/119

†Total dose period = 1/2 total lifetime

Unadjusted q_1^* from study = $1.7 \times 10^{-1} \text{ (ppm)}^{-1}$

Human $q_1^* = 1.47 \times 10^{-1} \text{ (mg/kg/day)}^{-1}$ (see Appendix B2)

APPENDIX B2

Calculation for q_1^*

Lifetime risk of cancer associated with 1 ppm, p:

$$p = 1 - e^{-0.17}$$

For 1,1-dichloroethylene: $1 \mu\text{g}/\text{m}^3 = 0.25 \text{ ppm}$ by the formula

$$C (\text{mg}/\text{m}^3) = C (\text{ppm}) \times \text{MW (molecular weight of chemical)} \div 24.45 (\text{moles/l of air})$$

Lifetime risk of cancer associated with $1 \text{ mg}/\text{m}^3$, p:

$$p = 1 - e^{-(0.17)(0.25)} = 4.2 \times 10^{-2}$$

$$q_1^* (\text{mg}/\text{kg}/\text{day})^{-1} = 70 \text{ kg} \times 4.2 \times 10^{-2} \div 20 \text{ m}^3/\text{day} \cdot 1 \text{ mg}/\text{m}^3 = 0.147$$

where:

70 = assumed body weight of humans in kg

20 = human inhalation rate in m^3/day .