

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



February 19, 1992

David Anselment
16317 28th NE
Seattle, WA 98155

Dear Mr. Anselment:

While you were on board the S/S Arco Fairbanks from December 8 to December 14, 1991, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Fairbanks showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Fairbanks 20 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from 0.001 to 0.01 f/cc.

The complete report is on board the S/S Arco Fairbanks and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



February 19, 1992

Gilbert Avecilla
7378 Santa Valera Street
Bucana Park, CA 90620

Dear Mr. Avecilla:

While you were on board the S/S Arco Fairbanks from December 8 to December 14, 1991, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Fairbanks showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Fairbanks 20 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from 0.001 to 0.01 f/cc.

The complete report is on board the S/S Arco Fairbanks and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



February 19, 1992

Arthur Berkins
Route 3, Box 314
Texarkana, TX 75501

Dear Mr. Berkins:

While you were on board the S/S Arco Fairbanks from December 8 to December 14, 1991, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Fairbanks showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Fairbanks 20 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from 0.001 to 0.01 f/cc.

The complete report is on board the S/S Arco Fairbanks and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



February 19, 1992

Jesse Brewer
Arco Marine, Inc.
c/o Arco Fairbanks

Dear Mr. Brewer:

While you were on board the S/S Arco Fairbanks from December 8 to December 14, 1991, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Fairbanks showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Fairbanks 20 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from 0.001 to 0.01 f/cc.

The complete report is on board the S/S Arco Fairbanks and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



February 19, 1992

Bowen Carpenter
P. O. Box 1382
Lyman, WA 98263

Dear Mr. Carpenter:

While you were on board the S/S Arco Fairbanks from December 8 to December 14, 1991, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Fairbanks showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Fairbanks 20 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from 0.001 to 0.01 f/cc.

The complete report is on board the S/S Arco Fairbanks and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



February 19, 1992

Deborah Carpenter
399 Campbell Lk Rd.
Anacortes, WA 98221

Dear Ms. Carpenter:

While you were on board the S/S Arco Fairbanks from December 8 to December 14, 1991, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Fairbanks showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Fairbanks 20 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from 0.001 to 0.01 f/cc.

The complete report is on board the S/S Arco Fairbanks and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



February 19, 1992

Sam Chan
45400 Deluz Road
Temecula, CA 92590

Dear Mr. Chan:


While you were on board the S/S Arco Fairbanks from December 8 to December 14, 1991, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Fairbanks showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Fairbanks 20 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from 0.001 to 0.01 f/cc.

The complete report is on board the S/S Arco Fairbanks and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



February 19, 1992

Donald Davis
Route One Box 200
Wing, AL 36483

Dear Mr. Davis:

While you were on board the S/S Arco Fairbanks from December 8 to December 14, 1991, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Fairbanks showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Fairbanks 20 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from 0.001 to 0.01 f/cc.

The complete report is on board the S/S Arco Fairbanks and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,



Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



February 19, 1992

Matthew Frazier
823 N. Gaffey Place
San Pedro, CA 90731

Dear Mr. Frazier:

While you were on board the S/S Arco Fairbanks from December 8 to December 14, 1991, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Fairbanks showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Fairbanks 20 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from 0.001 to 0.01 f/cc.

The complete report is on board the S/S Arco Fairbanks and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

500 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



February 19, 1992

Robert Goodwin, Jr.
464 Westree Lane
Plantation, FL 33324

Dear Mr. Goodwin, Jr.:

While you were on board the S/S Arco Fairbanks from December 8 to December 14, 1991, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Fairbanks showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Fairbanks 20 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from 0.001 to 0.01 f/cc.

The complete report is on board the S/S Arco Fairbanks and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



February 19, 1992

Richard Hearn
Arco Marine, Inc.
c/o Arco Spirit

Dear Mr. Hearn:

While you were on board the S/S Arco Fairbanks from December 8 to December 14, 1991, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Fairbanks showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Fairbanks 20 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from 0.001 to 0.01 f/cc.

The complete report is on board the S/S Arco Fairbanks and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



February 19, 1992

Craig Imsland
5691 Valletta Way
Sacramento, CA 95820

Dear Mr. Imsland:

While you were on board the S/S Arco Fairbanks from December 8 to December 14, 1991, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Fairbanks showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Fairbanks 20 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from 0.001 to 0.01 f/cc.

The complete report is on board the S/S Arco Fairbanks and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

ARCO Marine, Inc.
300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



February 19, 1992

James Muller
4416 Cambridge Ave.
El Paso, TX 79903

Dear Mr. Muller:

While you were on board the S/S Arco Fairbanks from December 8 to December 14, 1991, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Fairbanks showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Fairbanks 20 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from 0.001 to 0.01 f/cc.

The complete report is on board the S/S Arco Fairbanks and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



February 19, 1992

Kathie Pavese
80 Huntington, St., #7
Huntington Beach, CA 92648

Dear Mr. Pavese:

While you were on board the S/S Arco Fairbanks from December 8 to December 14, 1991, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Fairbanks showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Fairbanks 20 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from 0.001 to 0.01 f/cc.

The complete report is on board the S/S Arco Fairbanks and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



February 19, 1992

Jorge Pineda
405 East 3rd St., #163
Long Beach, CA 90802

Dear Mr. Pineda:

While you were on board the S/S Arco Fairbanks from December 8 to December 14, 1991, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Fairbanks showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Fairbanks 20 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from 0.001 to 0.01 f/cc.

The complete report is on board the S/S Arco Fairbanks and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



February 19, 1992

Geoffrey Robinson
3530 Wise Street
Bremerton, WA 98310

Dear Mr. Robinson:

While you were on board the S/S Arco Fairbanks from December 8 to December 14, 1991, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Fairbanks showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Fairbanks 20 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from 0.001 to 0.01 f/cc.

The complete report is on board the S/S Arco Fairbanks and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



February 19, 1992

Carol Taylor
2634 Superior Street
Bellingham, WA 98226

Dear Ms. Taylor:

While you were on board the S/S Arco Fairbanks from December 8 to December 14, 1991, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Fairbanks showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Fairbanks 20 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from 0.001 to 0.01 f/cc.

The complete report is on board the S/S Arco Fairbanks and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



February 19, 1992

Wade Vandervacht
Arco Marine, Inc.
c/o Arco Fairbanks

Dear Mr. Vandervacht:


While you were on board the S/S Arco Fairbanks from December 8 to December 14, 1991, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Fairbanks showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Fairbanks 20 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from 0.001 to 0.01 f/cc.

The complete report is on board the S/S Arco Fairbanks and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



February 19, 1992

Christopher Brooks
Arco Marine, Inc.
c/o Arco Spirit

Dear Mr. Brooks:

While you were on board the S/S Arco Fairbanks from December 8 to December 14, 1991, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Fairbanks showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Fairbanks 20 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from 0.001 to 0.01 f/cc.

The complete report is on board the S/S Arco Fairbanks and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



February 19, 1992

Don Compton
1712 Merritt Park Drive
Orlando, FL 32803

Dear Mr. Compton:

While you were on board the S/S Arco Fairbanks from December 8 to December 14, 1991, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Fairbanks showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Fairbanks 20 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from 0.001 to 0.01 f/cc.

The complete report is on board the S/S Arco Fairbanks and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Seagate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



February 19, 1992

Robert Doran III
304 West Masters Lane
Melbourne, FL 32901

Dear Mr. Doran:

While you were on board the S/S Arco Fairbanks from December 8 to December 14, 1991, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Fairbanks showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Fairbanks 20 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from 0.001 to 0.01 f/cc.

The complete report is on board the S/S Arco Fairbanks and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



February 19, 1992

Erica Iseninger
5301 Northwest Trail
Corpus Christi, TX 78410

Dear Ms. Iseninger:

While you were on board the S/S Arco Fairbanks from December 8 to December 14, 1991, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Fairbanks showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Fairbanks 20 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from 0.001 to 0.01 f/cc.

The complete report is on board the S/S Arco Fairbanks and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



February 19, 1992

Gary Pierce
Arco Marine, Inc.
c/o Arco Fairbanks

Dear Mr. Pierce:

While you were on board the S/S Arco Fairbanks from December 8 to December 14, 1991, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Fairbanks showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Fairbanks 20 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from 0.001 to 0.01 f/cc.

The complete report is on board the S/S Arco Fairbanks and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



February 19, 1992

Jeffrey Portillo
1932 Aztec Ct.
West Linn, OR 97068

Dear Mr. Portillo:


While you were on board the S/S Arco Fairbanks from December 8 to December 14, 1991, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Fairbanks showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Fairbanks 20 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from 0.001 to 0.01 f/cc.

The complete report is on board the S/S Arco Fairbanks and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cc

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



February 19, 1992

Dean Quirion
Arco Marine, Inc.
c/o Arco California

Dear Mr. Quirion:

While you were on board the S/S Arco Fairbanks from December 8 to December 14, 1991, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Fairbanks showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Fairbanks 20 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from 0.001 to 0.01 f/cc.

The complete report is on board the S/S Arco Fairbanks and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



February 19, 1992

Albert Scara Jr.
1004 Sea Girt Avenue
Sea Girt, NJ 08750

Dear Mr. ~~Scara~~ Jr.: *AL*

While you were on board the S/S Arco Fairbanks from December 8 to December 14, 1991, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Fairbanks showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Fairbanks 20 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from 0.001 to 0.01 f/cc.

The complete report is on board the S/S Arco Fairbanks and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay
Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



February 19, 1992

Clinton Strout
5250 Cabrito Drive
Las Vegas, NV 89103

Dear Mr. Strout:

While you were on board the S/S Arco Fairbanks from December 8 to December 14, 1991, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Fairbanks showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Fairbanks 20 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from 0.001 to 0.01 f/cc.

The complete report is on board the S/S Arco Fairbanks and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



February 19, 1992

Dennis Weeks
P. O. Box 64, State Street
Castine, ME 04421

Dear Mr. Weeks:

While you were on board the S/S Arco Fairbanks from December 8 to December 14, 1991, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Fairbanks showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Fairbanks 20 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from 0.001 to 0.01 f/cc.

The complete report is on board the S/S Arco Fairbanks and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,

Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

300 Oceangate
Post Office Box 22617
Long Beach, California 90801-5617
Telephone 213 590 4400
TWX 910 341 6829



February 19, 1992

Stephen Snyder
736 Waverford Ave.
Maple Shade, NJ 08052

Dear Mr. Snyder:

While you were on board the S/S Arco Fairbanks from December 8 to December 14, 1991, Clayton Environmental Services, Inc. was taking samples to determine airborne levels of asbestos and conducting employee awareness training. During the training classes we indicated that the results of the survey would be provided to all crew members on board at the time. Recently the results of the independent lab analysis were received. The full report has been sent to the vessel and we will ask the Master to make the report available to all crew members. At the completion of the monitoring program, a summary report of the findings will be mailed to all vessels.

Airborne asbestos levels are measured by drawing a volume of air through a filter and counting the number of fibers that are retained in the filter. All air samples collected on the Fairbanks showed concentrations below the applicable permissible exposure limits (PELs) established by the U.S. Coast Guard of 0.1 fibers/cubic centimeters (f/cc) and the U.S. Occupational Safety and Health Administration (OSHA) PEL of 0.2 f/cc. These samples were also below the Environmental Protection Agency (EPA) recommended concentration of 0.01 f/cc for general occupancy.

On board the S/S Arco Fairbanks 20 samples were taken and analyzed to determine the presence of airborne fibers. Samples were taken from both working and living spaces. Samples analyzed by an independent lab measured airborne fiber levels ranging from 0.001 to 0.01 f/cc.

The complete report is on board the S/S Arco Fairbanks and available for your review. If you would like any additional information, please feel free to call me at (310) 590-4521.

Sincerely,


Jay C. Kitchener
Manager, Safety, Health and
Environmental Protection

JCK/cec

STEW
THANKS FOR YOUR COMMENTS
LAST SUMMER RE: FAIRBANKS
SHIPYARD. WE WILL CONTINUE
TO MONITOR AND ENSURE
PROPER PRECAUTIONS ARE TAKEN
BY SHIPYARD WORKERS HANDLING
ASBESTOS ON OUR VESSELS.