

Reprinted from the *A. M. A. Archives of Industrial Health*
June 1958, Vol. 17, pp. 634-653
Copyright 1958, by American Medical Association

An Epidemiological Study of Lung Cancer in Asbestos Miners

DANIEL C. BRAUN, M.D., and T. DAVID TRUEN, M.A., Pittsburgh

FROM
INDUSTRIAL HYGIENE FOUNDATION
4400 Fifth Ave., Pittsburgh 13, Pa

Ever since the pronounced increase in the incidence of lung cancer among males became apparent, there have been attempts to associate it with one or another of the various elements in the environment of man. The approach used by some workers has been to suspect one or several substances and then set about in an intensive search for lung cancer among persons who have had any exposure to those materials. In this connection Smith¹ writes: "The tendency of authors reporting the coincidental occurrence of primary lung cancer with silicosis or with any other theoretical etiologic conditions, has been to emphasize the percentage relationship in extremely small series of cases, with control cases which are not in any way comparable."

It would seem inevitable that asbestos should come under scrutiny in this manner, because prolonged exposure to this material is known to cause a specific type of pneumoconiosis, and because persons who show this form of pneumoconiosis often come to autopsy and provide a ready source of material for study. It was in this way that reports of the simultaneous occurrence of lung cancer and asbestosis began to accumulate after the report of a case by Lynch and Smith² in 1935. Within the next 10 years, about 15 additional cases were reported, and in 1954 Merewether³ reviewed all deaths from asbestosis recorded in England since March, 1924. Lung cancer occurred in 16% of these cases. Ginyne,⁴

whose work is also frequently referred to as establishing a connection between asbestosis and cancer of the lung, reported in 1951 that cancer of the lung was present in 14.1% of asbestosis cases examined by him. In 1941, Nordmann and Sorge⁵ claimed to have produced lung cancer in mice which they exposed to asbestos dust.

Since 1951, additional cases of cancer of the lung coexisting with asbestosis have been reported, and, according to Huenner⁶ about 100 such cases had been reported up to 1955. As a result, an association between the two diseases appears to have been accepted by many authors, and several writers were using the term "asbestosis cancer" of the lung. Werber,⁷ in 1952, stated categorically that in 7% to 17% of cases of asbestosis, after a latent period of about 1½ to 20 years, carcinoma becomes established in the lung.

On the other hand, not all authors accepted this alleged association without reservation. Saupe⁸ in 1939 reported that he had discovered no cases of lung cancer among 620 cases of asbestosis which he had examined; and in 1942, Holleb and Angrist⁹ expressed the opinion that the number of cases of asbestosis with lung cancer was too small for statistical evaluation. In 1947, Wegelius¹⁰ reported 126 radiologically diagnosed cases of asbestosis among 476 workers in Finland, and found no cases of lung cancer in this group. Goldblatt and Goldblatt in their section of Merewether's latest book,¹¹ state: "But at no stage in all these impressive researches was any clue obtained which might have offered any support to the possibility that asbestos could act as a..."

Accepted for publication Jan. 20, 1958.

This study was made possible through a grant from the Quebec Asbestos Mining Association. Medical Director (Dr. Braun) and Statistical Consultant (Mr. Truen), Industrial Hygiene

84 32/
1/20
Braun Exp
11/13/57

LUNG CANCER IN ASBESTOS MINERS

is no reliable criterion by which one can anticipate carcinogenicity and, as is well known, relatively minute changes in the structure of a chemical carcinogen are sufficient to diminish or eliminate carcinogenic action.

If asbestos is indeed to be regarded as a carcinogen, the need is felt to demonstrate some property which can be regarded as something more than inertness."

These authors advance the theory that, until some more experimental evidence of direct carcinogenesis by asbestos or a decomposition product of it can be obtained, asbestos might be considered as a "co-carcinogen" which only induces a further development of a preneoplastic condition brought about by something independent of the asbestos, such as an endogenous factor.

Thus the literature, while tending to support the thesis that asbestosis is in some way related to the development of lung cancer, is by no means unanimous. Altogether, it is perhaps more confusing than enlightening. A careful review shows that the majority of the reports are clinical and not epidemiological. They lack many elements necessary for the application of epidemiological techniques to their content, and most of the authors do not make claim to having done so. What has happened is that succeeding authors have drawn conclusions and generalized beyond the scope of the works which they quote. Nowhere, for example, have we found references to a population of asbestos workers, although several authors who have quoted the observed incidence of lung cancer in autopsies of persons who also had asbestosis imply that this incidence applies to asbestos workers, generally. We have likewise been unable to find any study which actually calculated the incidence of lung cancer among a population of persons who had asbestosis, and not just those who came to autopsy. With the exception of a paper by Doll,²³ none of those reviewed give any data on exposure and dust concentrations, and even Doll's paper merely mentions

"scheduled" areas, by which is meant, "those areas where processes are carried on which were scheduled under the Asbestos Industry Regulations of 1931 as being dusty."

There is, furthermore, a complete lack of definition of terms as used in the published literature. For example, the term "asbestosis," as used, may refer to changes observable only by microscopic examination of the lung tissue, or it may mean a radiologically detectable condition.

Most of the published reports obviously included women among their cases, but none of them do not give the number or proportion of women involved in the study.

There is also a lack of uniformity as to what type of exposure most studies have dealt with. Of 99 cases enumerated by Hueper²⁴ in 1955, only 10 appear to have originated in the United States, and 7 in Canada. Some of the earlier reports apparently included asbestos miners, but it can be assumed, since 82 of the 99 cases had originated in England, and since no asbestos mining operations are carried on in that country, that most of the reported cases have involved workers in the textile or fabricating industries.

Such factors as smoking habits, family history of cancer, length of time in the industry, and age of the individual case are also notably absent in the majority of these reports.

With this understanding of the limitations of the existing literature with respect to epidemiological generalization, it may be of value to consider in somewhat more detail some representative earlier publications, a few of which were referred to briefly above.

One of the most detailed studies and one which deserves the most serious consideration is that reported by Doll²⁵ in 1955. This study reviews causes of death among asbestos workers based on coroners' records. It also attempts to estimate the risk by studying records of men who worked for at least 20 years in exposed situations.

Doll concluded that lung cancer was a specific industrial hazard of certain asbestos workers and that, after 20 years of exposure, the risk is 10 times as great as for the general population.

This article is important for several reasons, in addition to the definite conclusions at which it arrives. For example, it begins by stating that "in view of the infrequency of asbestosis, this large number of cases (161 cases of lung cancer) suggests—but does not prove—that lung cancer is an occupational hazard of asbestos workers." Neither this article nor any previous one which we have examined presents any figures to prove that asbestosis is an infrequent occurrence. Estimates of the number of persons potentially exposed to asbestos dust in the United States alone vary from 10,000 to 35,000, and the incidence of asbestosis of any degree might be higher than Doll imagines.

This study, like so many others, involves autopsy records. The number of persons involved in the statistical analysis is only 113, representing only 1,042.25 man-years of life. It is also true that in selecting men who had been employed for at least 20 years, the study automatically excluded those who died from other causes after shorter employment.

Another reason why this publication is of importance is a statement which it contains to the effect that "the strongest evidence that it (lung cancer) may be a hazard (in asbestos workers) has been produced by Merewether and by Gloyne."

In 1951 Gloyne¹¹ presented a review of 1205 autopsies on persons who had worked in various dusty occupations. This number included 132 asbestos workers, of whom 121 showed "pneumoconiosis"—presumably asbestosis. Primary cancer of the lung occurred 17 times in this group, an incidence rate of 14.1% for lung cancer among asbestosis cases coming to autopsy. There were in his series 796 cases with silicosis and 69% of these also showed primary cancer of the lung. The incidence of

lung cancer in other forms of pneumoconiosis was 6.7%, and in 169 cases which proved not to have any type of pneumoconiosis it was 8.3%. Gloyne considered "the mortality of the asbestos workers" to be "disturbing." First of all, it is obvious that the paper does not deal with the "mortality of asbestos workers," and secondly, it must be borne in mind that all of Dr. Gloyne's cases were submitted to him for study because the findings were unusual for uncomplicated pneumoconiosis. It can reasonably be assumed that cases, including those of asbestosis, in which the findings were not considered unusual were not sent to Dr. Gloyne for examination. As a matter of fact, in the same paragraph in which he expresses concern over the incidence rate in asbestosis, Dr. Gloyne himself points out that the rate for lung cancer based on necropsies at the London Chest Hospital was 21.3% while the figures of the Registrar-General showed only 2.4%. He thus recognized that autopsies on a certain selected group of cases were not representative of the general population. It would seem, then, that notwithstanding the value of Dr. Gloyne's work, its importance as an index of the prevalence of lung cancer in asbestotics has been misinterpreted by some who have quoted him. All that it really shows is the fact that in a group of 121 cases, selected for special study primarily because they seemed abnormal by preliminary examination, 17, or 14.1%, had lung cancer.

Merewether¹² in 1947, in the report of the Chief Inspector of Factories, reviewed all cases reported between 1924 and 1946 in which asbestosis was the cause of death or a coexisting condition. This work was later extended to include all such cases reported up to December, 1954, by which time there were 344 deaths, including 205 males and 139 females. Among them were 55 cases (16%) of cancer of the lung, 41 in males and 14 in females. It is quite possible that a large number of asbestotics who did not die of their asbestosis, or in whose

P21
3720

LUNG CANCER IN ASBESTOS MINERS

death certificate it was not mentioned, may have been missed. The import of this 16% is enhanced by the simultaneous statement that the incidence of lung cancer in autopsies of the general population is only 1%. The danger of attempting to compare a rate found in 344 cases with the rate for the general population without respect to age, occupation, and many other variables, such as smoking habits, is obvious.

Lynch,¹⁰ who with Smith¹¹ had reported the first case in 1935, reported 4 cases of carcinoma of the lung in a series of 49 autopsies on workers in an asbestos manufacturing plant who were shown to have "demonstrable deposits of asbestos in the lungs." This, of course, is not necessarily identical with the disease asbestosis. Lynch himself points out that, although this is an incidence of 8.2%, "both figures are too small for very serious statistical types of calculation." Nevertheless, later writers have used this paper to strengthen the case for an association of carcinoma of the lung with asbestosis. It is also of interest that Klotz¹² found only the same number of cases of lung cancer in a series nearly 10 times as large, i. e., 4 in 478 cases of asbestosis.

Behrens, as cited by Merewether,¹³ estimated that, of 309 cases of asbestosis in the literature, 44 showed associated cancer of the lung—giving an incidence of 14.2%. This is an illustration of generalizing an incidence obtained in a group of cases which were undoubtedly reported only because some of them showed lung cancer, to possibly hundreds of asbestotics whose cases were never reported. The same applies to the conclusion of Teleky,¹⁴ who appears to have reviewed reports of 39 autopsies on persons with asbestosis among which 6 cases of lung cancer occurred. Information from sources such as these does not justify generalizations with regard to mortality rates.

Perhaps no one has written so extensively on the subject as has Hueper.^{15, 16} In 1935 he reviewed the cases¹⁷ reported

prior to that date and enumerated a total of 99. Eleven of these were those discussed by Doll¹⁸ and may have been cases covered by other authors. Eight were discovered by Kennaway and Kennaway¹⁹ in an analysis of death certificates, and, unless Merewether's study was incomplete, these cases should have been included in his report. Of the remaining 80, it is quite possible that the 31 contributed by Merewether and the 17 by Gloyne contain some duplication with each other or with those of other English authors.

Principles of the Epidemiological Method

Dorn²⁰ has pointed out that much of what is now thought to be pertinent concerning the comparative frequency of lung cancer in different population groups has been developed from the analysis of clinical material, particularly surgical and autopsy records, supplemented to some extent by the reported impressions of various clinicians based upon their personal observations. More recently, however, attention has turned to the systematic investigation of this problem by the same methods that have proved so successful previously in the study of communicable diseases, that is to say, by epidemiological methods.

In order to apply this method of investigation to the problem under discussion, we were of the opinion that a study should be planned to as to provide (1) a well-defined population group; (2) available data for all members of this population, including the healthy as well as the ill; (3) a sample which is truly representative of the population; (4) reliable and valid observations relating to the problem of the study.

A serious defect, common to most of the studies which have been reported, is that little or no information concerning the healthy people in the group seems to have been available to the author. In order to draw a generalization regarding all asbestos workers, it is necessary for a study to include living persons as well as the dead

Limiting the investigation to cases coming to autopsy, as has been frequently done in earlier studies, still further restricts its use in generalization. The problem with which we are concerned is whether asbestos miners experience more lung cancer than does the general population. The answer necessitates the collection of reliable information on asbestos miners as a group, as well as on the general population.

It seems advisable to discuss the differences between the epidemiological approach and that used in the studies which have been reported to date. A very important consideration is the fact that lung cancer, in spite of its increasing numbers, is still a disease of low incidence; that is, in a given population not many persons will contract this particular disease. This fact requires that large samples or groups must be studied to provide meaningful results.

Recognizing the difficulty of obtaining such large samples, most earlier writers deviated from the epidemiological method and sought to circumvent the requirement of observing well persons by (1) comparing the relative frequency of cancer in various sites; (2) comparing the relative frequency of cancer in a group of hospitalized patients; (3) comparing the relative frequency of cancer in a group of cases coming to autopsy.

Attempting to compare two population groups, looking only at the relative frequency of cancer in various body sites may result in finding a higher percentage (relative frequency) in one of the groups, when, in fact, the mortality rate of cancer of a particular organ is exactly the same in both groups. This is very clearly demonstrated in the excellent article by Dorn.²⁰ The mortality rate from a particular cause is the true measure of comparison.

It is apparent that selected groups such as hospitalized patients or autopsy cases may not be in any way representative of a larger group, and that in dealing with such samples, the observer may easily find more cases of a given disease than would

be found in another group of the same size, but representative of the general population. It is true that investigation of cases from such a sample can furnish information valuable for research, but the use of this information in drawing generalizations is necessarily restricted. It is the obligation of both the investigator and of those who read his report to make proper comparisons and to draw only those conclusions which are valid and justified. A good statistical study of cases of cancer of the lung occurring in a group of autopsies can lead to a proper inference concerning the frequency of lung cancer among cases coming to autopsy, but only to such cases. For information from such a study to be projected to some larger group, it is necessary that the autopsies represent a good sample of that larger group. To assume that such is the case in any particular series is dangerous and likely to be false.

There is some danger that the figures reported by some authors may be misconstrued as applying to asbestos workers or even asbestos miners, when, in fact, the authors in question do not make this generalization, nor can the generalization be made for the reasons stated. Close study of the reports reveals that the percentages cited relate only to the group of autopsies covered by the particular investigation.

The present study, in contrast to the earlier works, has been planned to utilize the epidemiological method. A well-defined group of asbestos miners has been established in such a way that it constitutes a good sample of the whole population of asbestos miners in Quebec. Data for all members of this group have been collected and analyzed. Those concerning lung cancer have received most careful consideration. Details of the methods employed will be set forth later, but the type of approach is considered to permit of fair comparisons and valid generalizations.

Collection and Analysis of Data

A preliminary survey of potential sources of information in February, 1956, involved

LUNG CANCER IN ASBESTOS MINERS

discussions with the physicians in charge of the asbestos companies' programs and with clinicians, pathologists, representatives of City and Provincial health departments and of the Canadian Cancer Society, and other interested persons. It was found that morbidity data, although somewhat limited, were available from such sources as the hospitals in Montreal and Quebec City, and the 13 cancer detection centers in the Province. However, because of the high mortality in lung cancer, it seemed advisable to depend upon data relating to deaths. These we found to be obtainable at the vital statistics department of the Ministry of Health in Quebec City. From the preliminary survey, it was apparent that extensive and detailed information could be gathered with respect to both the persons employed in the asbestos mining industry and the mortality figures for the general population.

Following this exploratory survey, the initial effort was directed to the collection of data relating to all workers who had been processed through the clinic at Thetford Mines since its inception in 1947 and similar information regarding all workers at Asbestos, Que. Data from the clinical records included the age, family and personal medical histories, smoking habits, number of years of exposure, an estimate of weighed exposure, and the course of the individual's health status or the cause of his death.

From this information it was possible to formulate a "cohort" which could be well defined, should be representative of the whole group, and could be followed for a definite period of time. All of the available experience indicates that the development of asbestosis in less than five years of exposure must be somewhat rare. Accordingly, the cohort was defined as including every miner who had a total exposure of five or more years, and who was on the employment rolls in 1950. Office and other nonexposed personnel, regardless of length of employment, were not included. This

cohort was then followed by means of the annual physical examination records through a six-year interval, 1950 through 1955. All data regarding this group were then tabulated in order to determine the characteristics of the cohort. For those who survived the entire period, reference was made to the physical examination results and x-ray findings at the end of the period. Those who had died were tabulated separately, and the cause of death was corroborated by examination of the death certificates. A further search was made concerning those in the original cohort who remained unaccounted for when the living and the known dead had been tabulated. They represent men who had left employment through retirement or resignation. Eventually all but a small number of these were accounted for as either living or dead, and in the latter event, the cause of death was substantiated in a similar manner, and the results added to the original list of deaths.

Death certificates for the Province of Quebec for the years 1952 to 1955, inclusive, were reviewed in the department of vital statistics of the Provincial Health Ministry, together with statistical summaries of the causes of deaths in the Province by counties. All cases in which death was certified as having been due to primary cancer of the lung were examined for such information as place of residence, occupation, date of death, hospital in which death occurred, and whether or not an autopsy was performed. Cases in which lung cancer was given as a cause of death, but in which it was not specified as to whether the cancer originated in the lung, were also reviewed in an effort to include all instances of primary carcinoma of the lung in the study.

The statistics for the Province of Quebec relate to population, total deaths from all causes, total deaths from cancer of all sites, and deaths from lung cancer. These were collected and tabulated by counties and by sex for the years 1950 to 1955, inclusive.

From them, death rates for the general population of Quebec and of individual counties were calculated for specific years and analyzed by cause.

Practically all employees of one company are covered by a group policy of life insurance which, fortunately, nearly all of them continue to carry when they retire. A very few are not covered by this policy, and those who leave the industry for one reason or another except retirement usually are no longer covered, but this is likewise a small number. As an additional check upon the information obtained from the clinical records on this group, the records of the life insurance company were examined for all death claims paid under the policy, and particular notice was taken of the claims in which the proof of death was based on cancer of the lung.

Deaths from lung cancer among asbestos miners were thus determined from the clinical records in the medical service of the industry and checked by means of the death certificates and insurance company records. The deaths were then verified individually by reviewing them with the physicians in charge of the medical services. In this manner, there was established a list of cases in which primary cancer of the lung is considered to have been proved as the cause of death. A few cases in which lung cancer is strongly suspected but not proved as the cause of death were considered separately. Mortality rates have been calculated using both the "proved" and the total of "proved" and "suspected" cases during the years under observation. Comparisons were then made between the death rates from the same cause among specific segments of unexposed persons. All lung cancer deaths, both suspected and proved, were carefully analyzed to determine possible relationship or correlations between the development of lung cancer and any factor known from the clinical records, such as family history of cancer, personal history of heavy smoking, existence of asbestosis, or exposure to asbestos.

In addition to this analysis of deaths occurring in the cohort and during the years under observation, every known death from cancer of the lung, as well as every case diagnosed but still living, has been tabulated and analyzed. They will be discussed separately from those included in the population and time-interval under study.

A comparison of lung cancer mortality in the asbestos-producing counties has been made with that in counties which are far removed from the asbestos mines and in which, presumably, no asbestos miners live.

Finally, in order to broaden the comparison of death rates in different population groups, the rates have been collected for Canada generally, and for the United States, according to the most recent published and unpublished material.

Results and Interpretation

The cohort which was constructed according to the criteria described in the preceding section has been considered individually and compared with the general population. Description of the cohort will be presented here as a preface to the results of the study:

Original Cohort	6,091
Lost Persons Deducted	133
Final Cohort	5,958
Living in 1955 (working and retired)	5,271
Dead by 1955	187
Cancer of lung	9
Questionable cancer of lung	3
Other causes	169
Unknown causes	6
Smokers	4,672
Non-smokers	1,285
Unknown	20

Tables 1, 2, 3, and 4 present age, number of years of employment, weighted average exposure, and smoking habits of the cohort.

A comparison of the exposure to asbestos dust is presented in Table 5. All members of the cohort were placed in one of three categories, representing increasing degrees of exposure based on a weighted average of the years spent at various levels of dustiness. The degree of dustiness for each job category was determined after consultation with persons familiar with the environment and conditions in the various work

LUNG CANCER IN ASBESTOS MINERS

TABLE 1.—Number and Percentage Distribution by Age

Age	Number	Per Cent
20-24	3 00*	46
25-29	1 121	18
30-34	615	10
35-39	313	5
40-44	2	—
Unknown	—	—
Totals	3,556	100
Average Age	41	—

* Less than 0.5%.

areas. For the purposes of calculation, the assumption has been made that the relationship between these categories is linear, and that Category II is twice as dusty, and Category III three times as dusty as Category I.

TABLE 3.—Number and Percentage Distribution by Exposure Category

Exposure Category	Number	Per Cent
I	2,031	57
II	2,130	60
III	1,772	50
Unknown	2	—
Totals	5,935	100
Average Exposure	2.0	—

* Less than 0.5%.

The fourth variable, smoking habits, was similarly tabulated and is shown in Table 4. This was included because the information was available and because smoking was regarded as one of the variables which, besides the environment, could conceivably influence the development of lung cancer. As used in this presentation, the term smoker refers to a cigarette smoker, habit-

TABLE 2.—Number and Percentage Distribution by Length of Employment

Length of Employment	Number	Per Cent
5-9	1,795	50
10-19	2,276	64
20-29	723	20
30-39	403	11
40-49	183	5
50+	33	1
Totals	5,356	100
Average Years of Employment	18	—

ually smoking more than five cigarettes per day. Persons who smoke pipes or cigars exclusively were not considered to be smokers for the purpose of this study.

Table 5 presents the year-by-year experience of the cohort and indicates the

TABLE 4.—Number and Percentage Distribution by Smoking Habits

Smoking Habits	Number	Per Cent
Smoker	4,473	75
Nonsmoker	1,263	21
Unknown	30	—
Totals	5,956	100

* Less than 0.5%.

number of deaths each year from specified causes. In general, a case was considered to be "proved" as one of primary cancer of the lung when the records showed that the diagnosis had been supported by an autopsy or surgical resection of the lung with microscopic examination of the removed tissue. In one case so considered, however, diagnosis was confirmed by bron-

TABLE 5.—Year-by-Year Experience of Cohort and Death Rates per 100,000 Man-Years of Risk

Year	No. Alive At Beginning of Year	Cause of Death				Man-Years of Risk
		Proved Primary Ca. of Lung	Supported Primary Ca. of Lung	Other Causes	Unknown	
1930	5,956	1	7	13	1	5,956
1931	5,943	3	9	17	1	5,923
1932	5,922	6	9	27	0	5,903.5
1933	5,885	7	1	34	0	5,866.5
1934	5,866	1	0	33	4	5,879
1935	5,810	3	0	36	0	5,793.5
Totals	—	21	17	160	6	53,713.5
"Proved" rate per 100,000 man-years of risk = $\frac{21}{53,713.5} \times 100,000 = 39.3$						
"Supported" rate per 100,000 man-years of risk = $\frac{17}{53,713.5} \times 100,000 = 31.8$						

Brown—Tupper

541
P21
8720

TABLE 6.—"Proved" Cases of Primary Cancer of the Lung

Case No.	Age	Smoker	Exposure	Died	Autopsy	Asbestos
1	48	Yes	25 yr. in Cat. II	10-12-51	Yes	Yes
2	43	Yes	21 yr. in Cat. I	2-2-52	Yes	No
3	43	Yes	27 yr. in Cat. I	7-20-52	Yes	No
4	39	Yes	22 yr. in Cat. I	8-20-52	Yes	No
5	48	Yes	22 yr. in Cat. II	5-31	Yes	Yes
6	61	Yes	23 yr. in Cat. III	6-30-53	Yes	Yes
7	36	Yes	20 yr. in Cat. II	11-22-53	Yes	Yes
8	37	Yes	16 yr. in Cat. I	8-2-54	Yes	No
9	46	Yes	28 yr. in Cat. III	2-2-55	Bronch.	No

choscopy with visualization and biopsy. In another, although there was no autopsy, the diagnosis of primary cancer of the lung seems to have been beyond question.

The term "suspected" primary cancer of the lung was applied to those cases in which the diagnosis remains in doubt but some of the evidence points to cancer of the lung. There were three such cases.

The term "man-years of risk" has been used to mean the number of men at risk for the year under observation. A person who lived throughout the year was counted as a full man-year of risk, but one who died during the year was counted as one-half a man-year.

Deaths occurring in the cohort, and in which lung cancer is considered to have been proved as a cause, are shown in Table 6. In Table 7 are shown three deaths which have been considered as "suspected" lung cancer cases.

An indication of the importance of these "suspected" cases in interpreting the results of the calculations is desirable before further discussion of the mortality rates which are derived in later tables. For example, it happens that the rate found for the proved cases is close to the "expected" rate based on the general population figures for the Province, as will be shown later, and on this basis we should find eight deaths from lung cancer among the cohort. Actually, nine cases were observed. If, however, the 3 additional "suspected" cases were included, increasing this figure to 12, the total

would be very close to the 95% level of significance. However, having found just 12 cases, we are not above this level, and therefore the hypothesis that asbestos miners do not have a higher mortality from lung cancer than does the general population cannot be rejected. Nevertheless, the occurrence of 12 cases in this sample would increase the rate to a point which approaches the significant level. Because of the tremendous importance of the questionable cases in this respect, some detail regarding them will be given here.

In one of these cases, the suspicion of cancer of the lung is based upon the x-ray interpretation, and, although no autopsy was performed, the death certificate indicates that death was due to lung cancer. It is well known that the x-ray appearance of fibrosis, especially if a localized density or a superimposed tuberculous lesion is present, can simulate that of a tumor, and by itself, does not justify the inclusion of this case as one of "proved" cancer of the lung. A second case was certified as having died by reason of hydrothorax, possibly due to lung cancer, but again there was no surgery and no postmortem examination. In the third case, although it was subjected to autopsy, two pathologists disagreed as to whether lung cancer was present. The death was certified as having been due to chronic myocarditis with nephritis and pulmonary congestion, and possibly cancer of the lung. On the basis of these facts, it seems unwarranted to include these three cases among

TABLE 7.—"Suspected" Primary Cancer of the Lung

Case No.	Age	Smoker	Exposure	Died	Autopsy	Asbestos
P	49	Yes	22 yr. in Cat. III	10-25-50	No	No
1	46	Yes	23 yr. in Cat. III	7-1-54	Yes	Yes
2	55	Yes	43 yr. in Cat. II	8-30-55	No	No

LUNG CANCER IN ASBESTOS MINERS

"proved" instances of lung cancer. On the other hand, they cannot, in fairness, be disregarded completely. It is for this reason that mortality rates have been calculated both ways.

Table 8 gives the rates by age group.

The rates by length of employment are shown in Table 9. During the first 40 years of employment, the rate rises, an observation which seems plausible since the men were growing older. However, after 40 years of exposure there are no "proved" cases reported for a total of 240 men during the six years, or about 1440 man-years of exposure. When the "suspected" cases are added, one case does show up in this

TABLE 8—Lung Cancer Deaths by Age Group

Age Group	No. of Persons	No. of Deaths	
		Proved	Total
20-24	1,901	1	1
25-34	1,121	1	2
35-44	415	1	1
45-54	313	4	5
Unknown	2	0	0
Total	3,352	7	12

Annual Lung Cancer Death Rates per 100,000 Man-Years of Exposure

Age Group	Proved	Total
20-24	4	4
25-34	15	44
35-44	42	41
45-54	212	263
Unknown	0	0
Overall	273	348

period. This would produce a rate of 69 per 100,000, again demonstrating the importance of these questionable, but unproved, cases to the final conclusion. Because, if there were no cases in this number of men with long exposure, and if asbestos is a carcinogenic agent, it must be concluded that these 240 men have demonstrated considerable resistance. This is a biological phenomenon which has been observed previously and is consistent with the theory of an intrinsic or endogenous factor in cancer. The only other explanation would be that the susceptible members of this age group had died earlier of lung cancer. Table 18, which appears later in this section, in-

Brune—Truen

TABLE 9—Lung Cancer Deaths by Length of Employment

Length of Employment	No. of Persons	No. of Deaths	
		Proved	Total
1-9	1,194	0	0
10-19	2,248	1	1
20-29	70	1	2
30-39	643	3	7
40-49	163	0	0
50-59	33	0	0
Total	3,338	4	12

Annual Lung Cancer Death Rates per 100,000 Man-Years of Exposure

Length of Employment	Proved	Total
1-9	0	0
10-19	1	2
20-29	14	34
30-39	134	182
40-49	0	0
50-59	0	0
Overall	149	218

icates that the members of the cohort did not die from lung cancer at a younger age than the general population.

The rates by weighted exposure are shown in Table 10, and it will be noted that they present strong evidence against asbestos being a carcinogenic agent, for, if exposure to asbestos is in any way connected to lung cancer, we would expect that the longer and heavier the exposure, the higher the rate that would be found. The only possible error in this interpretation could occur if the weighted exposures were inversely related to years of employment,

TABLE 10—Lung Cancer Deaths by Exposure Category

Exposure Category	No. of Persons	No. of Deaths	
		Proved	Total
I	7,031	4	4
II	2,170	3	4
III	1,772	2	4
Unknown	3	0	0
Total	13,976	9	12

Annual Lung Cancer Death Rates per 100,000 Man-Years of Exposure

Exposure Category	Proved	Total
I	4	23
II	23	31
III	19	37
Unknown	0	0
Overall	46	91

643
P21
11720

TABLE 11—Number of Persons in Various Weighted Exposure Categories by Length of Employment

Length of Employment	Weighted Exposure Categories			Unknown	Total
	I	II	III		
5-9	37	399	190	0	1,793
10-19	643	563	331	1	2,538
20-29	314	363	243	7	927
30-39	247	218	127	1	603
40-49	74	87	41	1	183
50+	24	21	10	0	55
Total	2,621	2,130	1,772	1	6,524
Average years of exposure	17.3	17.3	18.5	29.0	17.3

in which case the heaviest weighted exposure (Category III) would show the shortest length of employment. Table 11, which lists the number of persons in various exposure categories by length of employment indicates that this error has not occurred. In fact, the average number of years of employment for each exposure category is almost identical.

Table 12, which develops the rates for smokers and nonsmokers, is most striking. It shows that not a single case of lung cancer developed among the 1263 nonsmokers and that all cases of lung cancer, both "proved" and "suspected," occurred in smokers.

Table 12 was so striking that it was felt that further verification was necessary. It was possible that some abnormal distribution may have occurred, e.g., the non-

TABLE 13—Number and Percentage Distribution of Smokers and Nonsmokers by Age Groups

Age Group	Number		
	Smoker	Nonsmoker	Unknown
20-29	1,301	489	12
30-39	1,096	724	4
40-49	411	707	2
50+	184	150	1
Unknown	2	0	1
Total	3,094	1,970	20
Average age	29.3	44.2	40.7

Age Group	Percentage Distribution		
	Smoker	Nonsmoker	Unknown
20-29	48.5	34.5	40.0
30-39	35.3	36.7	39.0
40-49	13.3	35.4	10.0
50+	5.9	7.6	5.0
Unknown	0.1	0.0	0.0
Total	100%	100%	100%

* Less than 0.05%.

TABLE 12—Lung Cancer Death for Smokers and Nonsmokers

Number of Persons and Number of Lung Cancer Deaths by Smoking Habit	No. of Deaths	
	Proven	Total
Smokers	1,673	12
Nonsmokers	1,263	0
Unknown	20	0
Total	2,956	12

Annual Lung Cancer Death Rates per 100,000 Man-Years of Exposure by Smoking Habit

	No. of Deaths	
	Proven	Total
Smokers	22	43
Nonsmokers	0	0
Unknown	0	0
Overall	22	43

smokers may have included a larger percentage of young men. Consequently additional Tables 13, 14, and 15 were constructed to show the distribution of smokers and nonsmokers by age, length of employment, and degree of exposure. Although there are slight differences they do not account for the fact that all observed cases of lung cancer were in smokers. In respect to age (Table 13) the combined average age of the smokers was 49 years less than that of the nonsmokers. Table 14 shows that as far as length of employment is con-

TABLE 14—Number and Percentage Distribution of Smokers and Nonsmokers by Length of Employment

Length of Employment	Number		
	Smokers	Nonsmokers	Unknown
5-9	1,408	177	10
10-19	1,917	136	4
20-29	1,096	199	2
30-39	411	146	1
40-49	110	71	0
50+	37	22	0
Total	3,071	1,261	20
Average length of employment	17.3	19.3	16.0

Length of Employment	Percentage Distribution		
	Smoker	Nonsmoker	Unknown
5-9	30.1	29.3	30.0
10-19	42.4	33.7	30.0
20-29	35.6	33.7	10.0
30-39	13.4	13.1	20.0
40-49	3.6	5.7	0.0
50+	1.2	1.8	0.0
Total	100%	100%	100%

LUNG CANCER IN ASBESTOS MINERS

TABLE 15.—Number and Percentage Distribution of Smokers and Nonsmokers by Exposure Category

Exposure Category	Number		
	Smokers	Nonsmokers	Unknown
I.....	1,333	474	4
II.....	1,661	432	7
III.....	1,423	338	6
Unknown.....	4	1	0
Totals.....	4,421	1,245	17
Average exposure category.....	2.0	1.9	2.3

Exposure Category	Percentage Distribution		
	Smokers	Nonsmokers	Unknown
I.....	32.2	35.5	20.0
II.....	36.7	34.7	33.0
III.....	30.3	26.7	43.0
Unknown.....	0.1	0.1	0.0
Totals.....	100%	100%	100%

cerned, the smokers had worked about 2.3 years less on the average than the nonsmokers. With longer exposure and greater age, one would expect the nonsmoking group to show a higher rate of lung cancer were due to asbestos. Table 15 shows that the average exposure category was almost the same for the two groups. Therefore, this variable seems to be of no importance in accounting for this difference.

The result of this additional analysis is that none of these factors appears to lessen the effect of Table 12.

Comparison of the Cohort Experience with that of the Province of Quebec, Dominion of Canada, and the United States.—In order to make a comparison of the experience among asbestos miners with that of the general population of the Province of Quebec, statistics were gathered, as stated earlier, in the office of the Division of

Demography in the Provincial Ministry of Health. The data on total deaths, deaths from all forms of cancer, and deaths from cancer of the lung were obtained by sex and by county for the years 1950 through 1955. In addition, all death certificates which specified primary cancer of the lung, and all those which indicated lung cancer but did not specify the origin, were examined for the years 1952 through 1955.

Table 16 gives a tabulation of the number of deaths from lung cancer in the Province and in the cohort for the years 1950 through 1955, and shows the annual rate per 100,000 in these segments. It will be noted from the table that the mortality rate for the "proved" cases in the cohort is only slightly higher than the rate for the Province. When the "suspected" cases are included in the calculation, the rate for the cohort rises to 33.8 per 100,000, which is about 50% higher than the rate for the Province. This, it will be recalled from the previous discussion of the effect of the "suspected" cases on the results, approaches but does not exceed the significant level.

One further interesting observation from Table 16 is the rather marked increase in the total number of cases for the Province between 1950 and 1955. It is assumed that at least part of this increase is due to improved recognition and reporting of lung cancer during the interval. For this reason, the years 1954 and 1955 were thought to be more nearly representative of actual conditions. Even so, it is quite likely that the general population is not studied for lung cancer with the same diligence with which

TABLE 16.—Comparison of Cohort with Province of Quebec

	Number of Persons	Lung Cancer Deaths						Annual Rate per 100,000
		1950	1951	1952	1953	1954	1955	
Province	1,196,000*	196	220	243	202	203	337	1,624
Cohort	3,723†							
Total		3	2	8	2	1	3	12
Proved		1	1	2	1	1	1	5
Province (estimated on basis of 1954-55)	1,192,000	182	216	243	202	202	334	1,612

In the Province figures it has been assumed all these lung cancer deaths are for males only.
 * Approximate mid-point of the enumerated population for 1951, and the estimated population for 1954 (Rapport, Div. de la Démographie).
 † Number alive at beginning of 1952.

Brown—Truman

645
 P21
 12 7 20

TABLE 17.—Lung Cancer Deaths for the Province of Quebec*

Age Group	Estimated Population	Number of Deaths							
		1954				1955			
		Total Certified		Specified Primary		Total Certified		Specified Primary	
		Total	Proved	Total	Proved	Total	Proved	Total	Proved
20-44	777,500	17	13	11	9	17	9	17	9
45-54	322,000	47	36	43	29	50	33	56	29
55-64	127,000	100	58	89	22	117	57	116	57
65+	121,000	111	33	65	29	148	34	148	34
Totals	1,347,500	275	139	208	89	342	153	337	153

Age Group	Death Rates per 100,000							
	1954				1955			
	Total Certified		Specified Primary		Total Certified		Specified Primary	
	Total	Proved	Total	Proved	Total	Proved	Total	Proved
20-44	2.2	1.7	1.4	1.2	2.2	1.2	2.2	1.2
45-54	14.6	11.2	13.5	9.1	15.5	10.3	17.2	10.8
55-64	77.9	45.7	70.1	17.8	92.1	45.8	94.5	46.3
65+	91.7	27.3	53.8	24.0	123.2	28.2	123.2	28.2
Overall	20.3	13.9	15.4	9.9	27.6	12.3	27.2	12.3

* Data from death certificates.
 † Estimated for population obtained by applying 1951 percentages for age groups for males to the total population for 1954, as given in Report, Div. 21a, December 1956.

this disease is looked for in the miners, and it seems probable that the mortality rates for the Province may be low. This would appear to be substantiated by the fact that the reporting of cases in the cohort showed no such increase over the same period.

Table 17 was compiled to show the annual age-specific lung cancer rate of cases in which the death certificate merely read "cancer of the lung," as distinguished from those in which the diagnosis was confirmed by autopsy, surgery, or biopsy. The term "specified primary" refers to those cases in these two categories in which the tumor was specified as having originated in the lung. It will be noted that, of the total cases reported in 1955, a much higher percentage than in 1954 were specified as primary. The table also shows that a higher percentage of the total cases certified in 1955 were proved, again indicating increasing interest in this disease.

A comparison has been made between the age-specific rates shown in Table 17, and those for the cohort, shown in Table 8. An average of the 1954 and 1955 rates for the Province has been used, since the 1955 figure was higher and may have been exceptional.

This comparison, summarized in Table 18, shows that the observed number of deaths in our sample is not significantly greater than the expected number of deaths, based on the average of the 1954 and 1955 figures for the Province. It is true that, in the case of the age group of 65 and over, the five deaths provide a figure which is almost significant at the 95% level. However, it should be noted that this number includes one of the suspected but unproved cases previously referred to. Furthermore, it is rather likely that the rate for the general population is understated in this age group for the obvious reason that the exact cause of death in the very old is not

TABLE 18.—Comparison of the Actual and Expected Number of Lung Cancer Deaths by Age Among Asbestos Miners

Age Group	Province Total Certified Primary Rate per 100,000	No. of Miners	Expected No. of Deaths*	Observed No. of Deaths	
				Proved	Total
20-44	1.8	3,801	6.8	1	1
45-54	14.6	1,124	16.5	2	2
55-64	77.9	619	48.1	3	3
65+	91.7	215	19.7	4	5
Unknown	..	..	..	0	0

* The expected number is based on the average of the 1954 and 1955 age-specific rates for the Province of Quebec.
 † Actual 5.0

LUNG CANCER IN ASBESTOS MINERS

TABLE 19—Age Distribution of Adult Males for the Province of Quebec, 1931*

Age Group	Province		Percentage in Cohort
	Number	Percentage	
20-24	777,133	43	16
25-29	1,048,922	14	19
30-34	1,238,004	11	10
35-39	1,113,147	10	3
Total	1,156,196	100	100

* Report, 1934.

a matter of the same intensity of interest as it is in younger persons.

Table 18 also answers a question previously raised. It shows that the members of the cohort have not died from lung cancer at an age earlier than the general population, and that such an explanation cannot be offered for the absence of lung cancer in 240 men with more than 40 years of employment referred to on page 643.

Before leaving this comparison of the Province with the miners, it should be shown that their age distributions are reasonably the same. That this is the case can be observed from Table 19.

It should be remembered that the miners retire and consequently, it can be expected that the oldest age group will be larger in the general population. The data presented in Table 20 indicate that the lung cancer rate generally decreases after age 70. Therefore, we could expect the rate for all people over 65 to be smaller than the rate for the group between 65 and 75, which would apply to the oldest group of miners. It is felt that by using the whole adult male population, we have developed rates for the general population which are somewhat lower than if we had been able to exclude the people in the general population over 75.

A comparison between the asbestos miners and the population of the Dominion as a whole was made, using statistical material from several sources. In one source, Phillips¹⁹ gave age- and sex-specific rates for Canada for three periods between 1931 and 1952. The rates for males are given in Table 20.

These figures show strikingly the increase in rates between 1931 and 1952, and this

Brown—Truman

TABLE 20—Annual Death Rates per 100,000 for Cancer of the Lung in Canada*

Age Group	Groups of Years		
	1931-1933	1941-1943	1950-1952
Under 20	0.2	0.3	1.3
20-24	0.9	0.9	3.8
25-29	1.9	1.2	3.0
30-34	3.0	3.4	4.3
35-39	5.2	12.3	16.1
40-44	8.2	16.9	27.2
45-49	12.7	27.0	36.1
50-54	13.3	33.7	57.8
55-59	13.1	44.3	102.9
60-64	10.7	30.6	94.3
65-69	13.4	30.8	83.9
70-74	8.2	27.9	56.1
75-79	11.4	44.3	71.0
All Ages	2.8	7.1	23.8

* Mortality from Lung Cancer in Canada, 1931 to 1952.

increase is particularly marked after age 50, confirming an observation previously made, to the effect that until recently lung cancer has probably been underdiagnosed in the older age groups in the general population.

To use these figures for purposes of comparison, it is necessary to combine the rates for certain age groups in order to conform to the age distributions used in this study. Since the exact populations in each age group for the years indicated is not known, this must be an approximation. However, the rates would be somewhat as follows:

Age Group	Rate
20-44	1
45-54	27
55-64	68
65-	90.95

These rates are, in general, lower than those developed for the total (proved and suspected) cases of lung cancer among the asbestos miners. The only large difference, however, is in the age group of 65 years and over, and it is quite possible that the rate for this group may have increased for Canada between 1952 and 1954 as it did for the Province of Quebec (Table 17).

A further comparison has been made with an over-all rate obtained from the American Cancer Society for respiratory cancer deaths in Canada in 1953. This rate, for males, is 20.8 per 100,000, or 3 more per 100,000 than Phillips' 1950-1952 rate, and compares with 23.2 per 100,000 for proved cases and 33.3 per 100,000 for total cases among the

647
P21
18-20

TABLE 21.—Number of Deaths and Death Rates per 100,000 by Age Groups for the Adult Male Population of the United States*

Age Group	Population	Cases	Rate per 100,000
20-24	21,348,000	663	3.1
25-34	18,000,000	797	4.4
35-44	15,300,000	825	5.4
45-54	14,700,000	843	5.7
Total	69,348,000	3,128	4.5

* Data from "Vital Statistics of the United States," Vol. 1 and 2, 1952.

asbestos miners in this study. It is therefore obvious that there are no important differences between the rates for asbestos miners and those for the general population of Quebec and the Dominion of Canada.

Since it is probable that figures for the United States are more complete and, therefore, possibly more comparable to the data for the miners, age-specific rates were computed from "Vital Statistics of the United States," Volumes I and II, for 1952. These rates have been tabulated in Table 21.

It is apparent that these rates compare favorably with those for the asbestos miners as shown in Table 8. Still other rates for the United States were obtained from the American Cancer Society, and for males, these were 25.3 per 100,000 in 1953, and 28.0 per 100,000 in 1955. They are not identical with the rate calculated from the figures of the office of Vital Statistics, but this is possibly because the American Cancer Society rates are for males of all ages. Nevertheless, they too, compare favorably with the rates of 25 (or 34 for total cases) obtaining among the asbestos miners.

Turning for a moment to a comparison between the asbestos miners and persons who are exposed to asbestos in one form or another (as distinguished from the general population groups just discussed, who have no exposure) an interesting observation can be developed by deduction. Hueper²² has stated that there are about 35,000 persons exposed in the United States, and we have found that the Canadian mines employ about 8000. Elsewhere, it has been estimated that the workers in England who have exposure total between 3000 and 5000. With workers in Africa, Denmark, Nor-

way, and other countries, at least 50,000 persons must be exposed throughout the world, and it can be assumed that this number has been fairly constant in the 20 years since 1935 when the first case of asbestosis with lung cancer was reported. At least 1,000,000 man-years of exposure has thus been accumulated, and this figure can be divided by the approximately 150 cases of lung cancer with asbestosis reported during the 20-year period. This gives a rate of 15 per 100,000, which is at least indicative that any lung cancer rate which can be calculated for workers exposed to asbestos dust is not much greater than that for the unexposed population.

Comparison Between Eight Counties Adjacent to the Asbestos-Producing Areas and Eight Selected Counties—To compare lung cancer mortality rates in the counties surrounding the asbestos-producing areas with another group of counties in which no asbestos miners are likely to reside, the rates were computed on the basis of figures for the years 1950 through 1955. The eight counties selected for comparison were Argenteuil, Chateaugay, Montmagny, Portneuf, Richlieu, Riviere-du-Loup, St. Hyacinthe, and Terrebonne, mainly because they represent a wide geographic distribution throughout the Province. The counties selected because of their proximity to the asbestos mines include Arthabaska, Beauce, Drummond, Frontenac, Megantic, Richmond, Sherbrooke, and Wolfe. Table 22 shows the number of lung cancer deaths for the years 1950 through 1955 for each of these counties, and a mortality rate, based on the adult male population in 1952. To emphasize the comparison, Megantic County has been shown separately, as has the Province of Quebec and also the Province with the eight "asbestos-producing" counties subtracted. Because of its unique lung cancer death rate, Montreal et Isle de Jesus has also been listed in order to provide further comparison.

It is apparent from the table that the lung cancer death rate for the eight counties

LUNG CANCER IN ASBESTOS MINERS

TABLE 22.—Number of Lung Cancer Deaths and Rate per 100,000 Man-Years

Countries	Adult Male Population 1957	Male Lung Cancer Deaths *						Total	Rate per 100,000
		1950	1951	1952	1953	1954	1955		
Megantic County	13,100	3	1	3	3	1	4	15	18.9
Eight "Adjacent" Countries	87,400	6	3	9	16	1	16	51	9.5
Eight Selected Countries	53,000	7	10	3	18	3	9	48	18.5
Province of Quebec	1,146,000	194	230	245	303	303	337	1,612	22.5
Province of Quebec less eight "Adjacent" countries	1,100,000	188	217	238	287	299	341	1,570	22.5
Province of Quebec less eight "Adjacent" countries	104,000	3	7	10	19	16	23	78	22.5

* It is assumed that all male lung cancer deaths occurred after age 30.

immediately surrounding the asbestos-producing areas is practically identical with that of eight countries selected for comparison. While Megantic County has a rate nearly twice that of the combined eight selected countries, it is lower than the rate for the Province, and considerably lower than the rate for Montreal. The figure for Montreal would certainly be higher except for the very low numbers of deaths reported for 1950 and 1951, and it would appear that in those years some error in reporting has undoubtedly been made. On the basis of the other years, 1950 and 1951 deaths would be expected to be about 200 greater. This would result in a rate of 40 per 100,000.

The only possible conclusion from this comparison is that there is no evidence that the persons who live and work in the countries surrounding and adjacent to the asbestos-producing areas have any greater incidence of lung cancer than those who live elsewhere in the Province.

Comment on All Recorded Lung Cancer Cases, Living and Dead, among the Asbestos Miners—Although a simple enumeration of all the known or suspected cases of cancer of the lung in these areas has no particular value from a statistical point of view, it is of interest to summarize such cases for the record. There were nine deaths prior to the beginning of the time period covered by the study, including one in which the diagnosis was mediastinal lymphosarcoma. During the period covered by this investigation, there were nine proved cases and three suspected cases in the cohort. Through 1956 and to date in 1957, there were eight deaths, six of which were merely

suggestive of cancer of the lung and included such diagnoses as mediastinal lymphosarcoma, mesothelioma, cancer of the leg with metastases to lung, abscess of lung, and cancer of the pancreas. One other was diagnosed on the basis of x-ray only. In addition, there are now living four cases in which the diagnostic evidence is strongly suggestive of lung cancer. This is a total of 33 cases of all types, including 10 "suspected" but unproved cases, and 4 that are still living. The remaining 19 constitute the total of proved cases of cancer of the lung among the asbestos miners since 1940.

The proved cases averaged 59 years of age at death, and varied between 37 years and 68 years. Their working span covered periods varying from a minimum of 14 years to a maximum of 37 years. Only three men had less than 25 years of employment in the industry. Seven among those on whom such information is available had a weighed exposure placing them in Category III, and six worked in an exposure represented by Category I.

There were only 17 among these proved lung cancer cases in which we have information regarding the presence of asbestosis. Asbestosis was present in nine, although it was minimal in two. Two pathologists disagreed regarding its presence in another. At least seven of the 19 proved lung cancers, therefore, were not accompanied by asbestosis.

Summary and Conclusions

Interest in the question of whether there may be an association between lung cancer

Drawn—Tread

549
P2/
16/20

and exposure to asbestos has been evident since the report in 1933 by Lynch and Smith of a case in which lung cancer and asbestosis were both present. As additional cases in which the two diseases coexisted were reported, a causal association appears to have been gradually accepted by many authors, although some workers considered the correlation to be inconclusive. The present study was undertaken in an effort to determine whether a causal relationship did, in fact, exist between exposure to asbestos and cancer of the lung.

Since most earlier studies had been limited to enumerating the lung cancers found in certain selected samples, such as cases coming to autopsy or death certificates in which asbestosis was mentioned, it was apparent that they could not fulfill the requirements of an epidemiological and statistical approach to the problem. The present study was, therefore, designed to meet the requirements of this method.

After a preliminary survey to explore the availability of reliable information, data were gathered on workers in the asbestos mines in Quebec, based on their medical records. A cohort was defined as a group of asbestos miners having at least five years of exposure and who were in the industry in 1950. Data relative to their characteristics were collected and their status at the end of a six-year period of observation was determined. In the case of those who had died, an exhaustive search of death certificates and insurance records was carried out in order to determine as nearly as possible the exact cause of death. Mortality rates from lung cancer for the general population of the Province of Quebec and its various counties and for the Dominion of Canada, as well as the United States were calculated from statistics collected in the appropriate places. Comparisons of the rates obtained for asbestos workers and for the other population groups were made according to accepted statistical methods.

Records were obtained on 6091 persons who fulfilled the criteria of the cohort. It

was not possible to trace 133 of these for the whole period, but 5771 of the remaining 5958 were found to be still living in 1956 or later. Of the 187 known dead, cancer of the lung was considered to have been reasonably proved in 9 and to be strongly suggested in 3.

The members of the cohort were studied with respect to age, length of employment, a weighted average of their exposure, and their smoking habits. It was found that 4673 were smokers within the definition of that term as used in this study. Thirty-four per cent of the cohort were more than 45 years of age, and thirty per cent had been employed for longer than 20 years. Thirty per cent had a weighted exposure which placed them in the category of highest exposure.

The mortality rate for lung cancer, as computed on the basis of nine "proved" deaths among the cohort was 25.3 per 100,000. When the three "suspected" cases were added, the "total" rate for the cohort rose to 33.8. The importance of the suspected but unproved cases in determining these rates has been reiterated because it is likely that such cases would not be included in the statistics for the general population and because they influence the results so markedly.

According to the findings in this study, the mortality rate from lung cancer does not appear to increase with length of exposure or with degree of exposure, a fact which presents strong evidence against the carcinogenicity of asbestos.

Comparison of the experience among the asbestos miners with that of various segments of the unexposed, comparable population shows that the observed number of deaths among the miners is not significantly greater than the expected number. The rate for proved cases among the asbestos miners (25.3 per 100,000) compares well with the rate of 22.5 per 100,000 for the rest of the Province and 20.8 per 100,000 for adult males throughout the Dominion of Canada. It also compares satisfactorily with rates

LUNG CANCER IN ASBESTOS MINERS

of 37.2, 25.3, and 28.0 obtained from various sources for adult males in the United States. Finally, in this matter of comparison, it would appear that the world-wide experience of persons exposed to asbestos dust is not worse with respect to lung cancer than that of the unexposed population.

The countries surrounding the asbestos-producing areas, in which it is presumed most of the asbestos miners live, have almost identical mortality rates with those of eight countries widely scattered through the Province, and are lower than those for the remainder of the Province, and much lower than the rate for Montreal.

Since 1940 there have been 19 cases in which the diagnosis of primary cancer of the lung may be considered to have been proved. Approximately half of these cases were associated with asbestosis. All but one died in the recognized "cancer-age" and at least one-third had only the lightest exposure (Category I) to asbestos dust.

On the basis of what are believed to be complete and reliable data, it seems fair to conclude that the asbestos miners in the Province of Quebec do not have a significantly higher death rate from lung cancer than do comparable segments of the general population.

Furthermore, the death rate from lung cancer in the areas contiguous to the asbestos operations is comparable to that in areas widely scattered throughout the Province of Quebec and is lower than in some urbanized areas within the Province.

REFERENCES

1. Alarcio Escobar, R.: Bronchial Carcinoma: Review of 200 Cases. *J. Internat. Coll. Surgeons* 26:375-379 (Sept.) 1956.
2. Allen, M. L.: Bronchiogenic Carcinoma Associated with Pneumonoconiosis: Report of 2 Cases. *J. Indust. Hyg* 16:346-347 (Nov.) 1934.
3. Cartier, P.: A Contribution to the Study of Asbestosis. *Arch. mal. profess.* 10:389-395, 1949.
4. Anderson, C. S., and Dible, J. H.: Silicosis and Carcinoma of the Lung. *J. Hyg.* 38:185-204 (March) 1938.
5. Baader, E. W.: Asbestosis. *Deutsche med. Wchnschr.* 65:407-408 (March 17) 1939.
6. Behrens, W.: Experimental Asbestosis. *Schweiz. Ztschr. allg. Path.* 14:275-277, 1931.
7. Behrens, W., Jr.: The Clinical Picture and Pathology of Asbestosis. *Ztschr. Unfallmed. u. Berufskrankh.* 45:129-140 (June 15) 1952.
8. Berblinger, W.: Increase of Lung Cancer and Diseases Due to Dust Inhalation. *Med. Klin* 27:1337-1342 (Sept. 11) 1931.
9. Berenolium, I.: Irritation and Carcinogenesis. *Arch. Path.* 33:233-244 (Oct.) 1944.
10. Bohme, A.: Asbestosis. *Deutsche med. Wchnschr.* 62:928-930 (June 5) 1936.
11. Bohme, A.: Results of Periodical Examinations of Workers in an Asbestos Factory. *Beitr. Silikose Forsch.* 11:34 1931.
12. Bowles, O.: Asbestos-Milling, Marketing and Fabrication. Information Circular No. 6869. U. S. Department of the Interior, Bureau of Mines, 1935, pp. 1-26.
13. Breslow, L.; Hoaglin, L.; Rasmussen, G., and Abrams, H. K.: Occupations and Cigarette Smoking as Factors in Lung Cancer. *Am. J. Pub. Health* 44:171-181 (Feb.) 1954.
14. Bristol, L. F.: Roentgenologic Aspects of Silicosis and Asbestosis. *A. M. A. Arch. Indust. Health* 11:189-195 (March) 1955.
15. Cancer of the Lung: An Evaluation of the Problem. Proceedings of the Scientific Section, Annual Meeting, American Cancer Society, Inc., Nov. 3-4, 1953. New York, American Cancer Society, Inc., 1956.
16. Cartier, P.: Asbestosis Cancer of the Lung, in discussion on Smith, W. E.: Survey of Some Current British and European Studies of Occupational Tumor Problems. *A. M. A. Arch. Indust. Hyg.* 5:262-263, 1952.
17. Cartier, P.: Some Clinical Observations of Asbestosis in Mine and Mill Workers. *A. M. A. Arch. Indust. Health* 11:204-207 (March) 1955.
18. Clerens, J.: Research into Pulmonary Asbestosis in Belgium. *Arch. belges med. Sociale* 8:557-565 (Nov.) 1951.
19. Climes, Asbestosis and Silicosis. *Brit. M. J.* 1:379, 1931.
20. Coffin, G. L.; Durfee, H. C.; Maier, H. C.; Pardee, H. E. B., and Winter, E. L.: The Effects of Tobacco Smoking. Panel Meeting, Bull. New York Acad. Med. 32:133-156 (Feb.) 1956.
21. Cohart, E. M.: Lung Cancer and Economic Status. *Cancer* 8:1126-1129 (Nov.-Dec.) 1955.
22. Cooke, W. E.: Pulmonary Asbestosis. *Brit. M. J.* 2:1024-1025 (Dec. 3) 1927.
23. Cureton, R. J. R.: Squamous Cell Carcinoma Occurring in Asbestosis of the Lung. *Brit. J. Cancer* 2:219-223 (Sept.) 1948.
24. Currier, S. J.; Schneiderman, M. A., and Greenhouse, S. W.: Some Statistical Considerations.

Brian—Truman

P2/
15720

- trends in the Study of Cancer in Industry, *Am. J. Pub. Health* 44:1159-1166 (Sept.) 1954.
35. Davies, D. F.: Current Status of Lung Cancer Research: Some Pathogenetic Aspects, *CA* 6:169-174 (Sept.) 1956.
36. Desmeules, R.; Rousseau, L.; Giroux, M., and Sirois, A.: Asbestosis and Pulmonary Cancer, *Semaine de hôp. Paris* 23:1820-1823 (Aug. 7) 1947.
37. Doll, R.: Bronchial Carcinoma: Incidence and Aetiology (Milroy Lectures, abridged), *Brit. M. J.* 2:321-327 (Sept. 5); 323-390 (Sept. 12) 1953.
38. Doll, R.: Mortality from Lung Cancer Among Asbestos Workers, *Brit. J. Indust. Med.* 12:81-86, 1955.
39. Downes, J.: Pulmonary Asbestosis, *Am. J. Pub. Health* 23:1275-1281 (Dec.) 1933.
40. Dorn, H. F., and Cudde, S. J.: Morbidity from Cancer in the United States, *Pub. Health Monograph No. 29*, P. H. S. Publication No. 418, U. S. Public Health Service, 1955, 121 pp.
41. Dorn, H. F.: Cancer Morbidity Surveys: A Tool for Testing Theories of Cancer Etiology, *Am. J. Pub. Health* 45:615-621 (May) 1955.
42. Egbert, D. S., and Genger, A. J.: Pulmonary Asbestosis and Carcinoma: Report of a Case with Necropsy Findings, *Am. Rev. Tuberc.* 34:143-150 (July) 1936.
43. Ellman, P.: Pulmonary Asbestosis: Its Clinical, Radiological, and Pathological Features and Associated Risk of Tuberculosis Infection, *J. Indust. Hyg* 15:165-183 (July) 1933.
44. Ellman, P.: Pulmonary Asbestosis, *Proc. Roy. Soc. Med.* 34:537 (July) 1941.
45. Felt, A.: Pneumoconiosis in Asbestos Workers, *Presse med.* 39:1872-1874 (Dec. 19) 1931.
46. Fulton, W. B.; Dooley, A.; Matthews, J. L., and Houza, R. L.: Asbestosis: Part III. The Effects of Exposure to Dust Encountered in Asbestos Fabricating Plants on the Health of a Group of Workers, *Special Bulletin No. 42*, Pennsylvania Department of Labor and Industry, Bureau of Indust. Standards, Sept. 20, 1935.
47. Gardner, L. U., and Cummings, D. E.: Studies on Experimental Pneumoconiosis: Inhalation of Asbestos Dust: Its Effect upon Primary Tuberculous Infection, *J. Indust. Hyg* 13:65-81 (Feb.); 97 (March) 1931.
48. Gilliam, A. G.: Mortality Trends in Lung Cancer, *Cancer* 8:1130-1136 (Nov-Dec.) 1953.
49. Gilliam, A. G.: Mortality Attributed to Lung Cancer in the Large Cities of the United States in 1948 and 1949, *J. Nat. Cancer Inst.* 15:1307-1312, 1955.
50. Gloyne, S. R.: Two Cases of Squamous Carcinoma of the Lung Occurring in Asbestosis, *Tubercle* 17:5-10 (Oct.) 1935.
51. Gloyne, S. R.: Pneumoconiosis: A Histological Survey of Necropsy Material in 1205 Cases, *Lancet* 1:810-814 (April 14) 1951.
52. Gloyne, S. R.: A Case of Oat-Cell Carcinoma of the Lung Occurring in Asbestosis, *Tubercle* 18:100-101 (Dec.) 1936.
53. Gloyne, S. R., and Merewether, E. R. A.: Asbestos, Occupation and Health, Supplement to *Encyclopedia on Occupation and Health*, Geneva, International Labor Office, 1938.
54. Goldblatt, M. W., and Goldblatt, J.: Industrial Carcinogenesis and Toxicology, in *Industrial Medicine and Hygiene*, edited by E. R. A. Merewether, London, London, Butterworth & Co., Ltd. 1956, Vol. 3, pp. 185-188.
55. Haenszel, W. M.: Epidemiological Tests of Theories on Lung Cancer Etiology, *Pub. Health Rep.* 71:163-172 (Feb.) 1956.
56. Hammond, E. C.: Lung Cancer and Common Inhalants, *Cancer* 7:1100-1108 (Nov.) 1954.
57. Hammond, E. C.: Etiology of Bronchiogenic Carcinoma, *CA* 6:156-168 (Sept.) 1956.
58. Hillel, H. B., and Angnst, A.: Bronchiogenic Carcinoma in Association with Pulmonary Asbestosis: Report of 2 Cases, *Am. J. Path.* 18:123-125 (Jan.) 1942.
59. Homburger, F.: The Co-Incidence of Primary Carcinoma of Lungs and Pulmonary Asbestosis: Analysis of Literature and Report of 2 Cases, *Am. J. Path.* 19:797-807 (Sept.) 1943.
60. Horn, D.: Is Lung Cancer on the Increase? Evaluation of Present Day Evidence: Cancer of the Lung, Proceedings of the Scientific Section, Annual Meeting, American Cancer Society, Inc., Nov. 3-4, 1953.
61. Hornig, F.: Clinical Considerations on the Question of Industrial Cancer of Asbestos Workers, *Ztschr. Krebsforsch.* 47:281-287, 1938.
62. Hueper, W. C.: Cancer in Its Relation to Occupation and Environment, *Bull. Am. Soc. Control Cancer* 25:61-69 (June) 1943.
63. Hueper, W. C.: Significance of Industrial Cancer in the Problem of Cancer, *Occup. Med.* 2:190-200 (Sept.) 1946.
64. Hueper, W. C.: Environmental and Occupational Cancer, *Public Health Service, Supp.* 209, U. S. Public Health Service, 1949.
65. Hueper, W. C.: A Methodology for Environmental and Occupational Cancer Surveys, *Public Health Monograph No. 1*, P. H. S. Publication No. 12, U. S. Public Health Service, 1950.
66. Hueper, W. C.: Environmental Lung Cancer, *Indust. Med.* 20:49-62 (Feb.) 1951.
67. Hueper, W. C.: Occupational and Environmental Pulmonary Cancers with Special Reference to Pneumoconiosis, *Proceedings, 7th Saranac Symposium on Pneumoconiosis*, 1952.
68. Hueper, W. C.: A Quest into the Environmental Causes of Cancer of the Lung, *Pub.*

AUTOPSY PROTOCOL

LUTIGS, Mary

LABORATORY ACCESSION NO. 10-59-255

DECEASED: October 27, 1959 around
12 noon

AUTOPSY: Performed on the same day
at 7 P.M. in Manville, N. J.

PROSECUTOR: Michel Jaris, M. D.

P2
17-
Brain
12/14/79 87
120"

LUNG CANCER IN ASBESTOS MINERS

Health Monograph No. 36, P. H. S. Publication No. 432, U. S. Public Health Service, 1955.

59. Hueper, W. C.: Environmental Causes of Cancer of the Lung other than Tobacco Smoke, *Dis. Chest*, 30:141-153 (Aug.) 1956.

60. Isseibacher, K. J.; Klaus, H., and Hardy, H. L.: Asbestosis and Bronchogenic Carcinoma: Report of One Autopsied Case and Review of Available Literature, *Am. J. Med.* 15:721-732 (Nov.) 1953.

61. Kennaway, E. L. and Kennaway, N. M.: A Study of the Incidence of Cancer of the Lung and Larynx, *J. Hyg.* 36:236-267 (June) 1936.

62. Kennaway, E. L. and Kennaway, N. M.: Studies of Incidence of Cancer of the Lung and Larynx, *Brit. J. Cancer* 5:153-158 (June) 1951.

63. Klotz, M. O.: Association of Silicosis and Carcinoma of the Lung, *Am. J. Cancer* 35:38-49 (Jan.) 1939.

64. Lanza, A. J.; McConnell, W. I., and Fehnel, J. W.: The Effects of the Inhalation of Asbestos Dust on the Lungs of Asbestos Workers. Preliminary Study, *Pub. Health Rep.* 30:1-12 (Jan. 4) 1935.

65. Lanza, A. J., editor: *Silicosis and Asbestosis*. New York and London, Oxford University Press, 1938.

66. Lew, E. A.: Use of Life Insurance Company Records for Cancer Studies, *A.M.A. Arch. Indust. Hyg.* 5:198-203 (March) 1952.

67. Lutzbach, A. J., and Wedler, H. W.: Occupational Cancer among Asbestos Workers, *Arch. path. Anat.* 307:387-409, 1941.

68. Lynch, K. M., and Smith, W. A.: Pulmonary Asbestosis: Carcinoma of Lung in Asbestos-Silicosis, *Am. J. Cancer* 24:56-64 (May) 1935.

69. Lynch, K. M., and Smith, W. A.: Pulmonary Asbestosis: A Report of Bronchial Carcinoma and Epithelial Metaplasia, *Am. J. Cancer* 36:567-573 (Aug.) 1939.

70. Lynch, K. M., and Cannon, W. M.: Asbestosis: Analysis of 40 Necropsied Cases, *Dis. Chest* 14:874-889 (Nov.-Dec.) 1948.

71. McPheeters, S. B.: A Survey of a Group of Employees Exposed to Asbestos Dust, *J. Indust. Hyg.* 18:229-239 (April) 1936.

72. Merewether, E. R. A.: The Occurrence of Pulmonary Fibrosis and Other Pulmonary Affections in Asbestos Workers, *J. Indust. Hyg.* 12:198 (May); 239 (June) 1930.

73. Merewether, E. R. A., and Price, C. W.: Report on Effects of Asbestos Dust on the Lungs and Dust Suppression in the Asbestos Industry, London, His Majesty's Stationery Office, 1930.

74. Merewether, E. R. A.: A Memorandum on Asbestosis, *Tubercle* 15:109; (Dec.) 1933; 15:152 (Jan.) 1934.

75. Merewether, E. R. A.: Annual Reports of the Chief Inspector of Factories, London, His Majesty's Stationery Office, 1947.

76. Merewether, E. R. A., editor: *Industrial Medicine and Hygiene*, London, Butterworth & Co., Ltd., 1956, Vol. 1.

77. Nordmann, M.: The Industrial Cancer of Workers in Asbestos, *Ztschr. Krebsforsch.* 47:288-302, 1938.

78. Nordmann, M., and Sorge, A.: Pulmonary Cancer Produced by Asbestos Dust in Experimental Animals, *Ztschr. Krebsforsch.* 51:168-182, 1941.

79. Phillips, A. J.: Mortality from Cancer of the Lung in Canada (1931-1952), *Canad. M. A. J.* 71:242-244 (Sept.) 1954.

80. Saupe, E.: Further Contributions to the Roentgenological Diagnosis of Asbestosis, *Arch. Gewerbepath. u. Gewerbehyg.* 9:391-406, 1939.

81. Smith, K. W.: Pulmonary Disability in Asbestos Workers, *A.M.A. Arch. Indust. Health* 12:198-203 (Aug.) 1955.

82. Smith, K. W.: Pneumoconiosis and Lung Cancer with Special Reference to Silicosis and Asbestosis, *Comptes. Med.* 2:3-10 (Nov.) 1949.

83. Smith, W. E.: Survey of Some Current British and European Studies of Occupational Tumor Problems: Part III Asbestos, *A.M.A. Arch. Indust. Hyg.* 5:242-263 (March) 1952.

84. Stoll, R., Bass, R., and Anger, A.: Asbestosis Associated with Bronchogenic Carcinoma, *A.M.A. Arch. Int. Med.* 38:831-834 (Dec.) 1951.

85. Teleky, L.: Occupational Lung Cancer, *Acta Union internat. contre Cancer* 3:253-273, 1938, also, *Zentralbl. Gewerbehyg.* 27:33, 1940.

86. Vorwald, A. J.; Durkan, T. M., and Pratt, P. C.: Experimental Studies of Asbestosis, *A.M.A. Arch. Indust. Hyg.* 3:1-43 (Jan.) 1951.

87. Wenler, H. W.: Asbestosis and Lung Cancer, *Deutsche med. Wochenschr.* 69:375-376 (Aug. 6) 1943.

88. Wegelius, C.: Changes in the Lungs in 126 Cases of Asbestosis Observed in Finland, *Acta radiol.* 25:139-152, 1947.

89. Werber, M.: Pulmonary Asbestosis Associated with Carcinoma, *Zentralbl. Arbeitsmed. u. Arbeitsschutz* 2:179-180 (Nov.) 1952.

90. Wood, W. B., and Giovne, S. R.: Pulmonary Asbestosis Complicated by Pulmonary Tuberculosis, *Lancet* 2:954-956 (Oct. 31) 1931.

91. Wyers, H.: Asbestosis, *Postgrad. Med.* 25:631-635 (Dec.) 1949.

92. Wyniler, E. L., and Graham, E. A.: Etiologic Factors in Bronchogenic Carcinoma and Special Reference to Industrial Exposures: Report of 537 Proved Cases, *A.M.A. Arch. Indust. Hyg.* 4:221-235 (Sept.) 1951.