

PROJECT NO: B7L-21-69  
REPORT FILE

*Industrial* BIO-TEST *Laboratories, Inc.*  
1810 FRONTAGE ROAD  
NORTHBROOK, ILLINOIS 60062

REPORT TO  
MONSANTO COMPANY  
TWO-YEAR CHRONIC ORAL TOXICITY WITH  
AROCOR 1260  
IN ALBINO RATS

NOVEMBER 12, 1971

IBT NO. B7298



GBRNO02349

REPORT TO  
MONSANTO COMPANY  
TWO-YEAR CHRONIC ORAL TOXICITY WITH  
AROCOR 1260  
IN ALBINO RATS

NOVEMBER 12, 1971

IBT NO. B7298

I. Introduction

At the request of Monsanto Company, a two-year chronic oral toxicity study was conducted using albino rats to determine the potential toxicity of Aroclor 1260, Lot No. AK-3. The following report presents the results of this investigation.

6BRN002350

LEXOLDMON002687

## II. Summary

A two-year chronic toxicity study was conducted using albino rats fed diets containing 1 (T-I), 10 (T-II), or 100 (T-III) ppm Aroclor 1260.

Food consumption and body weight gains were not altered at any level of Aroclor 1260 fed. The number of animals dying and the time at which the deaths occurred did not vary among the test and control groups.

Hematological and clinical blood chemistry studies conducted after 3, 6, 9, 12, 18, or 24 months did not reveal any affects related to Aroclor 1260. Urine analyses failed to reveal any differences between test and control animals.

At sacrifice after 3, 6, or 12 months on test organ weights, organ to body weight and organ to brain weight ratios disclosed several randomly occurring intergroup differences. The lack of any consistent dose related response and the absence of any deleterious histopathologic change confirm that these differences were not related to the ingestion of Aroclor 1260.

At the final sacrifice after 24 months on test, the liver weights and liver to body weight or brain weight ratios were significantly elevated in the rats from the T-III group. Histologic examination of the livers from the T-III group revealed several animals with vacuolar change. This lesion is morphologically indicative of fatty degeneration. Specific fat stains confirmed the presence of fat in these vacuoles. Focal hypertrophy and focal hyperplasia were also found in the livers from animals fed Aroclor 1260.

GBRNO02351

Hyperplasia of the urinary bladder was found in an animal from the control group but not in any of the test animals.

The incidences and types of all tumors were about the same in all groups, including the control group, and are considered normal for rats of this age.

Respectfully submitted,

INDUSTRIAL BIO-TEST LABORATORIES, INC.

Report prepared by:

Philip S. Smith

Philip Smith, B. S.  
Assistant Toxicologist  
Rat Toxicity

Report approved by:

James B. Plank

James B. Plank  
Senior Group Leader  
Rat Toxicity

Paul L. Wright

Paul L. Wright, Ph. D.  
Section Head, Toxicology

M. L. Keplinger

M. L. Keplinger, Ph. D.  
Manager, Toxicology

November 12, 1971

lam:sjn

GBRND02352

### III. Procedure

#### A. Experimental Animals

The animals employed in the test were Charles River strain\* albino rats. Four hundred rats (200 males and 200 females) were selected for the experiment, ear-punched with the animal number assigned and housed individually in standard wire-bottomed steel rat cages. Each cage bore a color-coded card identifying the rat with respect to project number, dose level assignment, individual animal number and sex.

#### B. Organization of Groups

A structural outline of the experiment is given in Table I.

TABLE I  
TEST MATERIAL: Aroclor 1260  
Two-Year Chronic Oral Toxicity Study - Albino Rats  
Outline of Experiment

| Group   | Number of Animals |        | Dietary Level<br>(ppm) |
|---------|-------------------|--------|------------------------|
|         | Male              | Female |                        |
| Control | 50                | 50     | None Administered      |
| T-I     | 50                | 50     | 1                      |
| T-II    | 50                | 50     | 10                     |
| T-III   | 50                | 50     | 100                    |

\* Charles River Breeding Laboratories, North Wilmington, Mass.

### C. Diets, Feeding and Food Consumption

All diets were prepared in the central diet room of this laboratory. The basic ration from which all diets were constructed was a standard, pulverized stock rat ration\*. The diet for any given test group was prepared by blending the calculated amount of Aroclor 1260 with a pre-weighed portion of the stock ration in a Hobart mixer.

Fresh diets were prepared each week and every rat was offered an amount of food sufficient for ad libitum feeding.

Food consumption was recorded for five rats of each sex in every group weekly for the first three months and monthly for the next nine months. Thereafter, periodic spot checks were made.

### D. Body Weights and Weight Gains

Initially, the body weight of each rat in every group was determined and recorded. Thereafter, individual weighings were made weekly for the first 13 weeks and monthly thereafter until the conclusion of the investigation.

### E. Mortality and Reactions

Checks for mortality and abnormal behavioral reactions were made daily throughout the investigation.

### F. Hematologic Studies and Urine Analyses

Blood studies, including determinations of hemoglobin concentrations, hemotocrit value, erythrocyte count and both total and differential

\* Purina Rat Chow, Ralston Purina Co., St. Louis, Mo.

leukocyte counts, were conducted upon five males and five females from both the control and 100 ppm groups.

Urine analyses for the presence of glucose, albumin, microscopic elements, and determinations of pH and specific gravity were conducted upon urine samples at the same time intervals and from the same animals as those employed for the blood studies.

These studies were conducted after 3, 6, 9, 12, 18, and 24 months of feeding.

#### G. Clinical Blood Chemistry Studies

Determinations of blood urea nitrogen concentration (BUN), serum alkaline phosphatase activity (SAP), fasted blood glucose concentration and serum glutamic-pyruvic transaminase activity (SGPT) were conducted after 3, 6, 9, 12, 18, and 24 months of feeding upon the same rats as were the routine blood studies.

#### H. Pathologic Studies

When an animal succumbed during the test period, a gross autopsy was performed. In those cases when postmortem autolysis was not advanced, representative tissues were taken and preserved in ten percent formalin solution for possible future histopathologic study.

After 3, 6, and 12 months of testing, five animals of each sex from each group were sacrificed and subjected to complete gross pathologic examinations.

Absolute organ weights were recorded and organ to body weight and organ to brain weight ratios computed. The following organs were included: liver, kidneys, spleen, gonads, heart and brain. These data were then subjected to statistical analysis. An Analysis of Variance was conducted first and significant effects disclosed by that treatment were further studied by "t"-tests.

Complete microscopic examinations were conducted upon the tissues and organs from all control and 100 ppm group rats sacrificed after 3, 6, or 12 months of testing. The following tissues and organs were included in the examinations: heart, trachea, lungs, liver, pancreas, esophagus, stomach, small intestine (duodenum, jejunum and ileum), caecum, colon, spleen, lymph nodes, kidneys, urinary bladder, gonads, prostate gland, seminal vesicles, uterus, pituitary gland, adrenal glands, salivary glands, thyroid gland, parathyroid glands, skeletal muscle, sternum, bone, peripheral nerve, spinal cord and brain (cerebrum, cerebellum and pons).

The aforementioned tissues were prepared and stained with Hematoxylin-Eosin stain.

After two years of feeding, all surviving animals were sacrificed and subjected to complete gross pathologic examination.

The absolute weights of the liver, kidneys, spleen, gonads, heart and brain were recorded immediately after sacrifice.

GBRN002356

Microscopic examinations were conducted upon tissues and organs from selected animals dying during the experiment and upon all animals from the final sacrifice.

Tissues and organs examined were the same as those previously mentioned.

I. Tumor Incidence and Classification

A tabulation of the incidence of tumor formation in each group was made at the conclusion of the investigation. In addition, tumor data for individual animals including location, weight, size and pathologic classification were recorded.

GBRN002357

IV. Results

A. Body Weights and Weight Gains

The male and female body weight data are assembled in Table II.

The data in this table also include the 3, 12 and 24 month total weight gains.

There were no significant effects due to ingestion of the compound

6BRN002358

LEXOLDMON002695

**TABLE II**  
**TEST MATERIAL: Aroclor 1260**  
**Two-Year Chronic Oral Toxicity Study - Albino Rats**  
**Body Weight and Total Weight Gain Data**  
**Summary of Mean Values**

| Dietary Level of<br>Aroclor 1260<br>(ppm) | Sex | Body Weight<br>(grams)<br>Week: |     |     |     |     |     |     |     |     |     |     |     |     | 90-Day<br>Total Weight<br>Gain |             |
|-------------------------------------------|-----|---------------------------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|--------------------------------|-------------|
|                                           |     | 0                               | 1   | 2   | 3   | 4   | 5   | 6   | 7   | 8   | 9   | 10  | 11  | 12  | 13                             | (grams/rat) |
| Control                                   | M   | 133                             | 203 | 245 | 313 | 362 | 373 | 391 | 409 | 424 | 440 | 440 | 449 | 467 | 474                            | 341         |
|                                           | F   | 136                             | 178 | 190 | 219 | 244 | 254 | 259 | 265 | 270 | 276 | 281 | 278 | 290 | 288                            | 152         |
| 1                                         | M   | 133                             | 192 | 250 | 322 | 361 | 374 | 394 | 416 | 428 | 448 | 450 | 474 | 479 | 490                            | 357         |
|                                           | F   | 136                             | 167 | 189 | 218 | 241 | 252 | 256 | 261 | 268 | 273 | 277 | 281 | 290 | 288                            | 152         |
| 10                                        | M   | 133                             | 204 | 245 | 317 | 368 | 382 | 400 | 417 | 436 | 448 | 461 | 467 | 475 | 481                            | 348         |
|                                           | F   | 136                             | 174 | 190 | 217 | 242 | 254 | 260 | 263 | 268 | 274 | 281 | 279 | 285 | 288                            | 152         |
| 100                                       | M   | 133                             | 208 | 247 | 319 | 363 | 375 | 394 | 425 | 437 | 451 | 460 | 479 | 486 | 500                            | 367         |
|                                           | F   | 136                             | 173 | 188 | 222 | 240 | 253 | 258 | 262 | 265 | 267 | 270 | 275 | 275 | 280                            | 144         |

GBRND02359

LEXOLDMON002696

TABLE II continued

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Body Weight and Total Weight Gain Data

Summary of Mean Values

| Dietary Level of<br>Aroclor 1260<br>(ppm) | Sex | Body Weight<br>(grams)<br>Month: |     |     |     |     |     |     |     |     | 12-Month<br>Total Weight<br>Gain<br>(grams/rat) |
|-------------------------------------------|-----|----------------------------------|-----|-----|-----|-----|-----|-----|-----|-----|-------------------------------------------------|
|                                           |     | 4                                | 5   | 6   | 7   | 8   | 9   | 10  | 11  | 12  |                                                 |
| Control                                   | M   | 498                              | 533 | 565 | 593 | 622 | 642 | 667 | 688 | 701 | 568                                             |
|                                           | F   | 297                              | 310 | 322 | 333 | 345 | 359 | 374 | 383 | 390 | 254                                             |
| 1                                         | M   | 516                              | 547 | 578 | 600 | 628 | 643 | 668 | 688 | 699 | 566                                             |
|                                           | F   | 298                              | 307 | 314 | 321 | 333 | 340 | 353 | 367 | 379 | 243                                             |
| 10                                        | M   | 496                              | 526 | 551 | 581 | 606 | 633 | 660 | 685 | 709 | 576                                             |
|                                           | F   | 291                              | 306 | 314 | 331 | 340 | 358 | 370 | 381 | 388 | 252                                             |
| 100                                       | M   | 522                              | 555 | 580 | 607 | 633 | 650 | 681 | 707 | 717 | 584                                             |
|                                           | F   | 289                              | 302 | 307 | 314 | 326 | 333 | 342 | 347 | 363 | 227                                             |

GBRND02360

LEXOLDMON002697

TABLE II continued

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Body Weight and Total Weight Gain Data

Summary of Mean Values

| Dietary Level of<br>Aroclor 1260<br>(ppm) | Sex | Body Weight<br>(grams)<br>Month: |     |     |     |     |     |     |     |     |     |     |     | 24-Month<br>Total Weight<br>Gain<br>(grams/rat) |
|-------------------------------------------|-----|----------------------------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-------------------------------------------------|
|                                           |     | 13                               | 14  | 15  | 16  | 17  | 18  | 19  | 20  | 21  | 22  | 23  | 24  |                                                 |
| Control                                   | M   | 728                              | 746 | 759 | 768 | 769 | 767 | 760 | 758 | 739 | 726 | 687 | 642 | 509                                             |
|                                           | F   | 426                              | 444 | 466 | 481 | 500 | 515 | 533 | 547 | 551 | 550 | 564 | 551 | 415                                             |
| 1                                         | M   | 721                              | 740 | 760 | 768 | 776 | 774 | 762 | 753 | 733 | 718 | 715 | 654 | 521                                             |
|                                           | F   | 390                              | 427 | 466 | 433 | 445 | 476 | 481 | 493 | 514 | 530 | 484 | 525 | 385                                             |
| 10                                        | M   | 728                              | 745 | 776 | 761 | 768 | 780 | 774 | 761 | 735 | 715 | 683 | 645 | 512                                             |
|                                           | F   | 416                              | 434 | 482 | 449 | 473 | 508 | 519 | 530 | 536 | 542 | 536 | 584 | 448                                             |
| 100                                       | M   | 746                              | 760 | 772 | 775 | 774 | 769 | 760 | 750 | 733 | 714 | 711 | 641 | 508                                             |
|                                           | F   | 380                              | 386 | 417 | 414 | 435 | 453 | 471 | 481 | 500 | 507 | 485 | 537 | 401                                             |

GBRND02361

LEXOLDMON002698

**B. Food Consumption**

The data obtained from the food consumption measurements conducted during the first 12 months of feeding are summarized in Table III.

These data revealed no significant difference between test and control rats. The periodic checks conducted during the remainder of the investigation also revealed no outstanding differences between test and control animals.

GBRN002362

LEXOLDMON002699

**TABLE III**

**TEST MATERIAL: Aroclor 1260**

**Two-Year Chronic Oral Toxicity Study - Albino Rats**

**Food Consumption Data**

**Summary of Mean Values**

| Dietary Level of<br>Aroclor 1260<br>(ppm) | Sex | Food Consumption<br>(grams/rat/seven days)<br>Week: |     |     |     |     |     |     |     |     |     |     |     |     | Average Food<br>Consumption<br>(grams/rat/seven days) |
|-------------------------------------------|-----|-----------------------------------------------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-------------------------------------------------------|
|                                           |     | 1                                                   | 2   | 3   | 4   | 5   | 6   | 7   | 8   | 9   | 10  | 11  | 12  | 13  |                                                       |
| Control                                   | M   | 94                                                  | 160 | 175 | 163 | 137 | 137 | 225 | 197 | 204 | 188 | 191 | 194 | 196 | 174                                                   |
|                                           | F   | 73                                                  | 140 | 144 | 107 | 104 | 133 | 208 | 154 | 165 | 151 | 146 | 150 | 151 | 140                                                   |
| 1                                         | M   | 118                                                 | 154 | 151 | 169 | 201 | 116 | 200 | 198 | 201 | 193 | 189 | 187 | 194 | 175                                                   |
|                                           | F   | 95                                                  | 118 | 97  | 139 | 160 | 107 | 162 | 139 | 165 | 149 | 150 | 143 | 147 | 136                                                   |
| 10                                        | M   | 106                                                 | 146 | 144 | 142 | 190 | 136 | 196 | 214 | 214 | 192 | 191 | 199 | 200 | 175                                                   |
|                                           | F   | 94                                                  | 124 | 128 | 124 | 179 | 90  | 190 | 167 | 155 | 151 | 150 | 149 | 152 | 143                                                   |
| 100                                       | M   | 121                                                 | 150 | 156 | 117 | 187 | 124 | 121 | 190 | 196 | 186 | 184 | 199 | 189 | 163                                                   |
|                                           | F   | 89                                                  | 93  | 123 | 151 | 151 | 101 | 193 | 136 | 140 | 144 | 144 | 147 | 146 | 135                                                   |

GBRND02363

TABLE III continued

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Food Consumption Data

Summary of Mean Values

| Dietary Level of<br>Aroclor 1260<br>(ppm) | Sex | Food Consumption<br>(grams/rat/seven days)<br>Month: |     |     |     |     |     |     |     |     | Average Food<br>Consumption<br>(grams/rat/seven days) |
|-------------------------------------------|-----|------------------------------------------------------|-----|-----|-----|-----|-----|-----|-----|-----|-------------------------------------------------------|
|                                           |     | 4                                                    | 5   | 6   | 7   | 8   | 9   | 10  | 11  | 12  |                                                       |
| Control                                   | M   | 196                                                  | 189 | 168 | 147 | 163 | 184 | 179 | 170 | 187 | 176                                                   |
|                                           | F   | 148                                                  | 154 | 136 | 121 | 138 | 152 | 160 | 141 | 127 | 142                                                   |
| 1                                         | M   | 196                                                  | 200 | 189 | 192 | 179 | 181 | 182 | 172 | 184 | 186                                                   |
|                                           | F   | 148                                                  | 163 | 141 | 163 | 160 | 153 | 157 | 143 | 132 | 151                                                   |
| 10                                        | M   | 186                                                  | 191 | 188 | 193 | 197 | 197 | 168 | 164 | 196 | 186                                                   |
|                                           | F   | 142                                                  | 138 | 145 | 144 | 144 | 151 | 136 | 139 | 141 | 142                                                   |
| 100                                       | M   | 179                                                  | 170 | 180 | 188 | 186 | 166 | 181 | 188 | 180 | 180                                                   |
|                                           | F   | 156                                                  | 127 | 147 | 144 | 134 | 132 | 162 | 149 | 139 | 143                                                   |

GBRNO02364

C. Mortality and Reactions

A frequency distribution of natural deaths occurring during the investigation appears in Table IV.

No untoward behavioral reactions were noted among any of the animals employed in the investigation.

GBRN002365

LEXOLDMON002702

Dr. Richard BLO-TEST Laboratories, Inc.

**TABLE IV**

**TEST MATERIAL: Aroclor 1260**

**Two-Year Chronic Oral Toxicity Study - Albino Rats**

**Mortality Data**

**Frequency Distribution of Deaths**

| Dietary Level<br>(ppm) | Sex | Frequency Distribution of Deaths<br>Months: |     |     |       |       |       |       |       | Total Dead<br>Number Tested* |
|------------------------|-----|---------------------------------------------|-----|-----|-------|-------|-------|-------|-------|------------------------------|
|                        |     | 1-3                                         | 4-6 | 7-9 | 10-12 | 13-15 | 16-18 | 19-21 | 22-24 |                              |
| Control                | M   | 0                                           | 0   | 0   | 2     | 1     | 7     | 6     | 9     | 25/35                        |
|                        | F   | 0                                           | 0   | 0   | 1     | 3     | 5     | 7     | 3     | 19/35                        |
| 1                      | M   | 0                                           | 0   | 1   | 1     | 4     | 7     | 5     | 8     | 26/35                        |
|                        | F   | 0                                           | 0   | 0   | 2     | 3     | 3     | 6     | 5     | 19/35                        |
| 10                     | M   | 1                                           | 1   | 0   | 2     | 4     | 7     | 8     | 6     | 29/35                        |
|                        | F   | 0                                           | 0   | 0   | 3     | 2     | 4     | 5     | 3     | 17/35                        |
| 100                    | M   | 1                                           | 0   | 0   | 2     | 2     | 8     | 6     | 9     | 29/35                        |
|                        | F   | 0                                           | 2   | 2   | 0     | 3     | 5     | 7     | 4     | 23/35                        |

\* Does not include the five animals of each sex from every group sacrificed for interim pathologic studies.

D. Hematologic and Clinical Blood Chemistry Studies

Mean male and female hematologic and clinical blood chemistry data are summarized in Tables V through IX.

Values for all parameters investigated were within the normal range for the albino rat. No significant differences between test and control values were observed.

GBRND02367

LEXOLDMON002704

**TABLE V**  
**TEST MATERIAL: Aroclor 1260**  
**Two-Year Chronic Oral Toxicity Study - Albino Rats**  
**Hematologic Data**  
**Summary of Mean Values**

| Dietary<br>Level<br>(ppm) | Sex | Total Leukocyte Count<br>(thousands/mm <sup>3</sup> ) |      |      |      |     |      | Erythrocyte Count<br>(millions/mm <sup>3</sup> ) |     |     |     |     |     |
|---------------------------|-----|-------------------------------------------------------|------|------|------|-----|------|--------------------------------------------------|-----|-----|-----|-----|-----|
|                           |     | Month:                                                |      |      |      |     |      | Month:                                           |     |     |     |     |     |
|                           |     | 3                                                     | 6    | 9    | 12   | 18  | 24   | 3                                                | 6   | 9   | 12  | 18  | 24  |
| Control                   | M   | 14.0                                                  | 15.4 | 10.1 | 11.1 | 9.9 | 11.2 | 8.1                                              | 8.2 | 8.7 | 8.8 | 7.6 | 6.5 |
|                           | F   | 9.1                                                   | 9.4  | 6.7  | 7.6  | 8.8 | 7.6  | 7.5                                              | 7.7 | 7.3 | 7.8 | 7.7 | 6.8 |
| 100                       | M   | 10.8                                                  | 13.1 | 7.9  | 12.8 | 8.6 | 8.4  | 7.5                                              | 8.1 | 8.0 | 7.6 | 8.0 | 7.6 |
|                           | F   | 13.2                                                  | 8.2  | 7.9  | 6.5  | 7.8 | 7.4  | 7.3                                              | 7.5 | 7.3 | 7.5 | 7.4 | 6.4 |

**TABLE VI**  
**TEST MATERIAL: Aroclor 1260**  
**Two-Year Chronic Oral Toxicity Study - Albino Rats**  
**Hematologic Data**  
**Summary of Mean Values**

| Dietary<br>Level<br>(ppm) | Sex | Hemoglobin Concentration<br>(g/100 ml) |      |      |      |      |      | Hematocrit Value<br>(%) |    |    |    |    |    |
|---------------------------|-----|----------------------------------------|------|------|------|------|------|-------------------------|----|----|----|----|----|
|                           |     | 3                                      | 6    | 9    | 12   | 18   | 24   | 3                       | 6  | 9  | 12 | 18 | 24 |
| Control                   | M   | 17.1                                   | 17.7 | 15.7 | 17.6 | 16.1 | 13.6 | 51                      | 49 | 48 | 50 | 41 | 35 |
|                           | F   | 16.9                                   | 17.4 | 14.1 | 16.7 | 16.4 | 14.6 | 50                      | 47 | 43 | 46 | 42 | 38 |
| 100                       | M   | 16.1                                   | 17.2 | 14.0 | 15.3 | 15.9 | 14.5 | 49                      | 47 | 43 | 43 | 41 | 38 |
|                           | F   | 16.3                                   | 17.0 | 13.7 | 15.6 | 15.3 | 12.9 | 48                      | 47 | 41 | 44 | 39 | 32 |

GBRNO02369

**TABLE VII**  
**TEST MATERIAL: Aroclor 1260**  
**Two-Year Chronic Oral Toxicity Study - Albino Rats**  
**Hematologic Data**  
**Summary of Mean Values**

| Dietary<br>Level<br>(ppm) | Sex | Differential Leukocyte Count (Number of Cells per Hundred) |      |      |      |      |      |             |      |      |      |      |      |
|---------------------------|-----|------------------------------------------------------------|------|------|------|------|------|-------------|------|------|------|------|------|
|                           |     | Lymphocytes                                                |      |      |      |      |      | Neutrophils |      |      |      |      |      |
|                           |     | Month:                                                     |      |      |      |      |      | Month:      |      |      |      |      |      |
|                           |     | 3                                                          | 6    | 9    | 12   | 18   | 24   | 3           | 6    | 9    | 12   | 18   | 24   |
| Control                   | M   | 82.0                                                       | 86.6 | 80.4 | 78.8 | 79.0 | 65.4 | 16.0        | 11.4 | 18.6 | 20.6 | 20.0 | 32.2 |
|                           | F   | 83.0                                                       | 86.6 | 86.8 | 89.0 | 70.2 | 61.4 | 15.4        | 11.6 | 11.8 | 10.0 | 29.2 | 36.0 |
| 100                       | M   | 85.8                                                       | 85.4 | 83.4 | 72.4 | 73.8 | 66.6 | 12.6        | 13.8 | 15.8 | 22.6 | 24.6 | 31.6 |
|                           | F   | 92.0                                                       | 89.6 | 78.8 | 82.2 | 80.6 | 65.6 | 7.6         | 10.0 | 19.4 | 17.0 | 17.4 | 34.2 |

GBRNO02370

TABLE VII continued  
**TEST MATERIAL: Aroclor 1260**  
**Two-Year Chronic Oral Toxicity Study - Albino Rats**  
**Hematologic Data**  
**Summary of Mean Values**

| Dietary<br>Level<br>(ppm) | Sex | Differential Leukocyte Count (Number of Cells per Hundred) |     |     |     |     |     |                       |     |     |     |     |     |
|---------------------------|-----|------------------------------------------------------------|-----|-----|-----|-----|-----|-----------------------|-----|-----|-----|-----|-----|
|                           |     | Monocytes<br>Month:                                        |     |     |     |     |     | Eosinophils<br>Month: |     |     |     |     |     |
|                           |     | 3                                                          | 6   | 9   | 12  | 18  | 24  | 3                     | 6   | 9   | 12  | 18  | 24  |
| Control                   | M   | 0.6                                                        | 0.0 | 0.0 | 0.4 | 0.2 | 1.2 | 1.4                   | 2.0 | 1.0 | 0.2 | 0.8 | 1.6 |
|                           | F   | 0.6                                                        | 0.0 | 0.0 | 0.0 | 0.2 | 1.0 | 1.0                   | 1.8 | 1.4 | 0.2 | 0.6 | 1.6 |
| 100                       | M   | 0.0                                                        | 0.0 | 0.0 | 0.0 | 0.4 | 0.8 | 1.4                   | 0.8 | 0.8 | 1.0 | 1.2 | 1.0 |
|                           | F   | 0.0                                                        | 0.0 | 0.0 | 0.0 | 0.4 | 0.2 | 0.4                   | 0.4 | 0.8 | 0.8 | 1.6 | 0.0 |

TABLE VII continued  
 TEST MATERIAL: Aroclor 1260  
 Two-Year Chronic Oral Toxicity Study - Albino Rats  
 Hematologic Data  
 Summary of Mean Values

| Dietary<br>Level<br>(ppm) | Sex | Differential Leukocyte Count (Number of Cells per Hundred) |     |     |     |     |     |
|---------------------------|-----|------------------------------------------------------------|-----|-----|-----|-----|-----|
|                           |     | Basophils<br>Month:                                        |     |     |     |     |     |
|                           |     | 3                                                          | 6   | 9   | 12  | 18  | 24  |
| Control                   | M   | 0.0                                                        | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
|                           | F   | 0.0                                                        | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 100                       | M   | 0.2                                                        | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
|                           | F   | 0.0                                                        | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |

**TABLE VIII**

**TEST MATERIAL: Aroclor 1260**

**Two-Year Chronic Oral Toxicity Study - Albino Rats**

**Clinical Blood Chemistry Data**

**Summary of Mean Values**

| Dietary<br>Level<br>(ppm) | Sex | Serum Glutamic-Pyruvic<br>Transaminase Activity<br>(Dade Units) |    |    |    |    |    | Blood Urea Nitrogen<br>Concentration<br>(mgs %) |    |    |    |    |    |
|---------------------------|-----|-----------------------------------------------------------------|----|----|----|----|----|-------------------------------------------------|----|----|----|----|----|
|                           |     | Month:                                                          |    |    |    |    |    | Month:                                          |    |    |    |    |    |
|                           |     | 3                                                               | 6  | 9  | 12 | 18 | 24 | 3                                               | 6  | 9  | 12 | 18 | 24 |
| Control                   | M   | 38                                                              | 48 | 45 | 45 | 42 | 36 | 17                                              | 15 | 12 | 10 | 14 | 15 |
|                           | F   | 33                                                              | 39 | 38 | 42 | 32 | 30 | 14                                              | 16 | 15 | 9  | 13 | 11 |
| 100                       | M   | 33                                                              | 39 | 42 | 43 | 31 | 36 | 14                                              | 15 | 17 | 16 | 16 | 15 |
|                           | F   | 29                                                              | 37 | 50 | 40 | 32 | 33 | 15                                              | 15 | 14 | 13 | 12 | 14 |

**TABLE IX**  
**TEST MATERIAL: Aroclor 1260**  
**Two-Year Chronic Oral Toxicity Study - Albino Rats**  
**Clinical Blood Chemistry Data**  
**Summary of Mean Values**

| Dietary<br>Level<br>(ppm) | Sex | Fasted Blood Glucose<br>Concentration<br>(mgs%)<br>Month: |     |     |     |     |     | Serum Alkaline Phosphatase<br>Activity<br>(King-Armstrong Units)<br>Month: |      |      |      |      |      |
|---------------------------|-----|-----------------------------------------------------------|-----|-----|-----|-----|-----|----------------------------------------------------------------------------|------|------|------|------|------|
|                           |     | 3                                                         | 6   | 9   | 12  | 18  | 24  | 3                                                                          | 6    | 9    | 12   | 18   | 24   |
| Control                   | M   | 128                                                       | 123 | 89  | 88  | 138 | 99  | 28.6                                                                       | 24.7 | 21.5 | 20.3 | 16.2 | 22.3 |
|                           | F   | 130                                                       | 118 | 115 | 92  | 134 | 109 | 18.9                                                                       | 17.8 | 10.6 | 11.9 | 16.8 | 20.2 |
| 100                       | M   | 118                                                       | 118 | 116 | 135 | 146 | 98  | 37.8                                                                       | 21.8 | 21.9 | 23.9 | 11.9 | 16.1 |
|                           | F   | 117                                                       | 116 | 115 | 138 | 146 | 94  | 12.0                                                                       | 12.1 | 7.6  | 11.7 | 7.8  | 12.4 |

**E. Urine Analyses**

Mean male and female urine analyses data are summarized in Tables X through XII.

Determinations for glucose concentration were negative for all rats at every interval of examination.

No differences were noted between the urine from control and test animals at the intervals of examination.

GBRN002375

LEXOLDMON002712

**TABLE X**  
**TEST MATERIAL: Aroclor 1260**  
**Two-Year Chronic Oral Toxicity Study - Albino Rats**  
**Urine Analyses Data**  
**Summary of Mean Values**

| Dietary Level<br>(ppm) | Sex | Glucose<br>Month; |   |   |    | Albumin<br>Month; |    |   |   |   |    |    |    |
|------------------------|-----|-------------------|---|---|----|-------------------|----|---|---|---|----|----|----|
|                        |     | 3                 | 6 | 9 | 12 | 18                | 24 | 3 | 6 | 9 | 12 | 18 | 24 |
| Control                | M   | n                 | n | n | n  | n                 | n  | n | n | t | t  | +2 | +3 |
|                        | F   | n                 | n | n | n  | n                 | n  | n | n | n | n  | t  | +2 |
| 100                    | M   | n                 | n | n | n  | n                 | n  | n | t | t | t  | +1 | +3 |
|                        | F   | n                 | n | n | n  | n                 | n  | n | n | n | n  | +1 | +4 |

**Key:**  
n = negative  
t = trace; less than 30 mg/100 ml urine  
+1 = 30 to 100 mg/100 ml urine  
+2 = 100 to 300 mg/100 ml urine  
+3 = 300 to 500 mg/100 ml urine  
+4 = more than 500 mg/100 ml urine

TABLE XI

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Urine Analyses Data

Summary of Mean Values

| Dietary<br>Level<br>(ppm) | Sex | Microscopic Elements |   |   |    |    |    | pH |   |   |    |    |    |
|---------------------------|-----|----------------------|---|---|----|----|----|----|---|---|----|----|----|
|                           |     | 3                    | 6 | 9 | 12 | 18 | 24 | 3  | 6 | 9 | 12 | 18 | 24 |
| Control                   | M   | N                    | N | N | N  | N  | +1 | 7  | 7 | 6 | 7  | 7  | 6  |
|                           | F   | N                    | N | N | N  | t  | t  | 8  | 8 | 7 | 8  | 6  | 6  |
| 100                       | M   | N                    | N | N | N  | N  | t  | 7  | 7 | 6 | 7  | 6  | 6  |
|                           | F   | N                    | N | N | N  | t  | t  | 7  | 8 | 6 | 7  | 6  | 6  |

## Key:

N= Normal

t = trace; less than 5 per low power field

+1 = 5 to 10 per low power field

+2 = 10 to 15 per low power field

+3 = more than 15 per low power field

Industrial Division, Inc.

**TABLE XII**  
**TEST MATERIAL: Aroclor 1260**  
**Two-Year Chronic Oral Toxicity Study - Albino Rats**  
**Urine Analyses Data**  
**Summary of Mean Values**

| Dietary<br>Level<br>(ppm) | Sex | Specific Gravity<br>Month: |       |       |       |       |       |
|---------------------------|-----|----------------------------|-------|-------|-------|-------|-------|
|                           |     | 3                          | 6     | 9     | 12    | 18    | 24    |
| Control                   | M   | 1.029                      | 1.022 | 1.033 | 1.027 | 1.042 | 1.051 |
|                           | F   | 1.020                      | 1.019 | 1.025 | 1.024 | 1.032 | 1.039 |
| 100                       | M   | 1.021                      | 1.025 | 1.031 | 1.030 | 1.043 | 1.039 |
|                           | F   | 1.016                      | 1.023 | 1.035 | 1.029 | 1.028 | 1.045 |

GBRND02378

LEXOLDMON002715

F. Pathologic Studies

1. Three-Month Sacrifice

a. Gross Pathologic Findings

No outstanding differences were noted between test and control rats upon gross pathological examination.

b. Organ Weight and Organ to Body and Organ to Brain Weight Ratio Data

The results of the statistical analyses conducted on absolute organ weights, organ to body weight and organ to brain weight ratios are summarized in Tables XIII through XVIII.

Significant differences between a test group and the control group are designated by asterisks following the test values.

Organ weights, organ to body weight and organ to brain weight ratios disclosed several randomly occurring intergroup differences. The lack of any consistent dose related response and the absence of any deleterious histopathologic change confirm that these differences were not related to the ingestion of Aroclor 1260.

**TABLE XIII**  
**TEST MATERIAL: Aroclor 1260**  
**Two-Year Chronic Oral Toxicity Study - Albino Rats**  
**Organ Weight and Ratio Data**  
**Three-Month Sacrifice**  
**Summary of Mean Values**  
**Organ: Liver**

| Dietary Level<br>of<br>Aroclor 1260<br>(ppm) | Organ Weight<br>(g) |         | Organ/Body Weight Ratio<br>(g/100 g) |         | Organ/Brain Weight Ratio<br>(g/g) |         |
|----------------------------------------------|---------------------|---------|--------------------------------------|---------|-----------------------------------|---------|
|                                              | Males               | Females | Males                                | Females | Males                             | Females |
| Control                                      | 21.4                | 13.0    | 4.13                                 | 4.16    | 10.3                              | 6.82    |
| 1                                            | 20.4                | 13.2    | 3.90                                 | 4.59    | 9.98                              | 6.86    |
| 10                                           | 24.0**              | 12.2    | 4.48                                 | 3.61    | 11.8                              | 6.38    |
| 100                                          | 29.5**              | 14.5    | 5.73**                               | 4.58    | 14.4**                            | 7.72    |

\*\* Statistically significant difference at the 99 percent confidence level.

GBRNO02380

INDUSTRIAL DIVISION LABORATORIES, INC.

**TABLE XIV**  
**TEST MATERIAL: Aroclor 1260**  
**Two-Year Chronic Oral Toxicity Study - Albino Rats**  
**Organ Weight and Ratio Data**  
**Three-Month Sacrifice**  
**Summary of Mean Values**  
**Organ: Kidneys**

| Dietary Level<br>of<br>Aroclor 1260<br>(ppm) | Organ Weight<br>(g) |         | Organ/Body Weight Ratio<br>(g/100 g) |         | Organ/Brain Weight Ratio<br>(g/g) |         |
|----------------------------------------------|---------------------|---------|--------------------------------------|---------|-----------------------------------|---------|
|                                              | Males               | Females | Males                                | Females | Males                             | Females |
| Control                                      | 4.03                | 2.49    | 0.782                                | 0.802   | 1.95                              | 1.31    |
| 1                                            | 4.37                | 2.62    | 0.838                                | 0.911   | 2.14                              | 1.36    |
| 10                                           | 4.00                | 2.61    | 0.747                                | 0.782   | 1.96                              | 1.37    |
| 100                                          | 4.00                | 2.52    | 0.779                                | 0.793   | 1.96                              | 1.35    |

GBRND002381

LEXOLDMON002718

**TABLE XV**  
**TEST MATERIAL: Aroclor 1260**  
**Two-Year Chronic Oral Toxicity Study - Albino Rats**  
**Organ Weight and Ratio Data**  
**Three-Month Sacrifice**  
**Summary of Mean Values**  
**Organ: Spleen**

| Dietary Level<br>of<br>Aroclor 1260<br>(ppm) | Organ Weight<br>(g) |         | Organ/Body Weight Ratio<br>(g/100 g) |         | Organ/Brain Weight Ratio<br>(g/g) |         |
|----------------------------------------------|---------------------|---------|--------------------------------------|---------|-----------------------------------|---------|
|                                              | Males               | Females | Males                                | Females | Males                             | Females |
| Control                                      | 0.766               | 0.578   | 0.150                                | 0.184   | 0.371                             | 0.302   |
| 1                                            | 0.846               | 0.516   | 0.161                                | 0.180   | 0.412                             | 0.269   |
| 10                                           | 0.838               | 0.632   | 0.156                                | 0.189   | 0.411                             | 0.331   |
| 100                                          | 0.718               | 0.556   | 0.140                                | 0.177   | 0.352                             | 0.297   |

TABLE XVI

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Organ Weight and Ratio Data

Three-Month Sacrifice

Summary of Mean Values

Organ: Gonads

| Dietary Level<br>of<br>Aroclor 1260<br>(ppm) | Organ Weight<br>(g) |         | Organ/Body Weight Ratio<br>(g/100 g) |         | Organ/Brain Weight Ratio<br>(g/g) |         |
|----------------------------------------------|---------------------|---------|--------------------------------------|---------|-----------------------------------|---------|
|                                              | Males               | Females | Males                                | Females | Males                             | Females |
| Control                                      | 3.60                | 0.114   | 0.698                                | 0.0370  | 1.74                              | 0.0599  |
| 1                                            | 3.67                | 0.128   | 0.704                                | 0.0450  | 1.80                              | 0.0666  |
| 10                                           | 3.83                | 0.121   | 0.715                                | 0.0368  | 1.89                              | 0.0636  |
| 100                                          | 3.71                | 0.138   | 0.725                                | 0.0429  | 1.81                              | 0.0728  |

TABLE XVII

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Organ Weight and Ratio Data

Three-Month Sacrifice

Summary of Mean Values

Organ: Heart

| Dietary Level<br>of<br>Aroclor 1260<br>(ppm) | Organ Weight<br>(g) |         | Organ/Body Weight Ratio<br>(g/100 g) |         | Organ/Brain Weight Ratio<br>(g/g) |         |
|----------------------------------------------|---------------------|---------|--------------------------------------|---------|-----------------------------------|---------|
|                                              | Males               | Females | Males                                | Females | Males                             | Females |
| Control                                      | 1.59                | 1.05    | 0.307                                | 0.339   | 0.767                             | 0.551   |
| 1                                            | 1.57                | 1.01    | 0.300                                | 0.353   | 0.765                             | 0.526   |
| 10                                           | 1.58                | 1.04    | 0.296                                | 0.311   | 0.781                             | 0.546   |
| 100                                          | 1.52                | 0.946   | 0.298                                | 0.297   | 0.743                             | 0.505   |

TABLE XVIII

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Organ Weight and Ratio Data

Three-Month Sacrifice

Summary of Mean Values

Organ: Brain

| Dietary Level<br>of<br>Aroclor 1260<br>(ppm) | Organ Weight<br>(g) |         | Organ/Body Weight Ratio<br>(g/100 g) |         | Final Body Weights<br>of Sacrificed Rats<br>(g) |         |
|----------------------------------------------|---------------------|---------|--------------------------------------|---------|-------------------------------------------------|---------|
|                                              | Males               | Females | Males                                | Females | Males                                           | Females |
| Control                                      | 2.07                | 1.91    | 0.402                                | 0.616   | 518                                             | 314     |
| 1                                            | 2.05                | 1.92    | 0.392                                | 0.672   | 523                                             | 287     |
| 10                                           | 2.04                | 1.91    | 0.380                                | 0.572   | 536                                             | 337     |
| 100                                          | 2.05                | 1.89    | 0.400                                | 0.593   | 517                                             | 321     |

c. Histopathologic Findings

Histopathologic examination of tissues and organs taken from all animals from both the control and 100 ppm groups was conducted.

Tables XIX and XX list all histopathologic changes noted.

All of the lesions noted in the microscopic examination of tissues were those of spontaneous disease and are not unusual for the albino rat. The most frequent findings were lesions in the trachea and lungs, indicating chronic murine pneumonia. These occurred in the control as well as the rats fed Aroclor 1260.

GBRN002386

LEXOLDMON002723

## TABLE XDX

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Histopathologic Changes

Three-Month Sacrifice

Group: Control

| Number of<br>Animals | Organ Examined  | Findings                       | Incidence | Average<br>Grade |
|----------------------|-----------------|--------------------------------|-----------|------------------|
| 5 Males              | Trachea         | Tracheitis                     | 1         | 1.0              |
|                      | Lung            | Chronic respiratory<br>disease | 1         | 1.0              |
|                      | Urinary bladder | Hyperplasia                    | 3         | 1.0              |
| 5 Females            | Lung            | Chronic respiratory<br>disease | 1         | 1.0              |

All other tissues and organs were normal histologically.

Grading System

0.5 = minimal  
1.0 = slight  
2.0 = mild  
3.0 = moderate  
4.0 = severe  
5.0 = extreme

GBRN002387

LEXOLDMON002724

TABLE XX

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Histopathologic Changes

Three-Month Sacrifice

Group: 100 ppm

| Number of<br>Animals | Organ Examined | Findings                       | Incidence | Average<br>Grade |
|----------------------|----------------|--------------------------------|-----------|------------------|
| 5 Males              | Trachea        | Tracheitis                     | 4         | 1.0              |
|                      | Lung           | Chronic respiratory<br>disease | 2         | 1.0              |
| 5 Females            | Trachea        | Tracheitis                     | 3         | 1.0              |
|                      | Lung           | Chronic respiratory<br>disease | 2         | 1.0              |
|                      | Kidney         | Focal lymphoid<br>infiltration | 1         | 1.0              |

All other tissues and organs were normal histologically.

Grading System

0.5 = minimal  
1.0 = slight  
2.0 = mild  
3.0 = moderate  
4.0 = severe  
5.0 = extreme

GBRN002388

LEXOLDMON002725

2. Six-Month Sacrifice

a. Gross Pathologic Findings

No outstanding differences were noted between test and control animals.

b. Organ Weight and Organ to Body and  
Organ to Brain Weight Ratio Data

The results of the statistical analyses conducted on absolute organ weights, organ to body weight and organ to brain weight ratios are summarized in Tables XXI through XXVI.

Organ weights, organ to body weight and organ to brain weight ratios disclosed several randomly occurring intergroup differences..

The lack of any consistent dose related response and the absence of any deleterious histopathologic change confirm that these differences were not related to the ingestion of Aroclor 1260.

GBRN002389

LEXOLDMON002726

313-1531 Laboratory, Inc.

TABLE XXI  
 TEST MATERIAL: Aroclor 1260  
 Two-Year Chronic Oral Toxicity Study - Albino Rats  
 Organ Weight and Ratio Data  
 Six-Month Sacrifice  
 Summary of Mean Values  
 Organ: Liver

| Dietary Level<br>of<br>Aroclor 1260<br>(ppm) | Organ Weight<br>(g) |         | Organ/Body Weight Ratio<br>(g/100 g) |         | Organ/Brain Weight Ratio<br>(g/g) |         |
|----------------------------------------------|---------------------|---------|--------------------------------------|---------|-----------------------------------|---------|
|                                              | Males               | Females | Males                                | Females | Males                             | Females |
| Control                                      | 22.9                | 13.1    | 3.74                                 | 3.77    | 11.7                              | 6.71    |
| 1                                            | 14.8**              | 15.1    | 2.92**                               | 3.74    | 7.05**                            | 7.86    |
| 10                                           | 19.0*               | 10.5**  | 3.31*                                | 3.08*   | 9.45*                             | 6.07    |
| 100                                          | 24.4                | 11.6    | 4.69*                                | 4.03    | 12.3                              | 6.17    |

\* Statistically significant difference at the 95 percent confidence level.  
 \*\* Statistically significant difference at the 99 percent confidence level.

**TABLE XXII**

**TEST MATERIAL: Aroclor 1260**

**Two-Year Chronic Oral Toxicity Study - Albino Rats**

**Organ Weight and Ratio Data**

**Six-Month Sacrifice**

**Summary of Mean Values**

**Organ: Kidneys**

| Dietary Level<br>of<br>Aroclor 1260<br>(ppm) | Organ Weight<br>(g) |         | Organ/Body Weight Ratio<br>(g/100 g) |         | Organ/Brain Weight Ratio<br>(g/g) |         |
|----------------------------------------------|---------------------|---------|--------------------------------------|---------|-----------------------------------|---------|
|                                              | Males               | Females | Males                                | Females | Males                             | Females |
| Control                                      | 4.04                | 2.79    | 0.664                                | 0.806   | 2.06                              | 1.43    |
| 1                                            | 3.64                | 2.73    | 0.716                                | 0.674*  | 1.73                              | 1.42    |
| 10                                           | 3.85                | 2.35    | 0.670                                | 0.684*  | 1.92                              | 1.36    |
| 100                                          | 3.92                | 2.35    | 0.753                                | 0.814   | 1.97                              | 1.25    |

\* Statistically significant difference at the 95 percent confidence level.

TABLE XXIII

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Organ Weight and Ratio Data

Six-Month Sacrifice

Summary of Mean Values

Organ: Spleen

| Dietary Level<br>of<br>Aroclor 1260<br>(ppm) | Organ Weight<br>(g) |         | Organ/Body Weight Ratio<br>(g/100 g) |         | Organ/Brain Weight Ratio<br>(g/g) |         |
|----------------------------------------------|---------------------|---------|--------------------------------------|---------|-----------------------------------|---------|
|                                              | Males               | Females | Males                                | Females | Males                             | Females |
| Control                                      | 0.752               | 0.684   | 0.122                                | 0.196   | 0.384                             | 0.350   |
| 1                                            | 0.754               | 0.908   | 0.148*                               | 0.220   | 0.359                             | 0.472   |
| 10                                           | 0.836               | 0.566*  | 0.146*                               | 0.167   | 0.416                             | 0.327   |
| 100                                          | 0.636               | 0.443*  | 0.122                                | 0.151   | 0.320                             | 0.236*  |

\* Statistically significant difference at the 95 percent confidence level.

GBRND02392

LEXOLDMON002729

TABLE XXIV

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Organ Weight and Ratio Data

Six-Month Sacrifice

Summary of Mean Values

Organ: Gonads

| Dietary Level<br>of<br>Aroclor 1260<br>(ppm) | Organ Weight<br>(g) |         | Organ/Body Weight Ratio<br>(g/100 g) |         | Organ/Brain Weight Ratio<br>(g/g) |         |
|----------------------------------------------|---------------------|---------|--------------------------------------|---------|-----------------------------------|---------|
|                                              | Males               | Females | Males                                | Females | Males                             | Females |
| Control                                      | 3.55                | 0.0960  | 0.593                                | 0.0279  | 1.81                              | 0.0492  |
| 1                                            | 3.23                | 0.120*  | 0.636                                | 0.0297  | 1.54                              | 0.0625* |
| 10                                           | 3.56                | 0.0968  | 0.620                                | 0.0287  | 1.77                              | 0.0560  |
| 100                                          | 3.53                | 0.0710* | 0.686                                | 0.0250  | 1.77                              | 0.0378* |

\* Statistically significant difference at the 95 percent confidence level.

GBRND02393

TABLE XXV

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Organ Weight and Ratio Data

Six-Month Sacrifice

Summary of Mean Values

Organ: Heart

| Dietary Level<br>of<br>Aroclor 1260<br>(ppm) | Organ Weight<br>(g) |         | Organ/Body Weight Ratio<br>(g/100 g) |         | Organ/Brain Weight Ratio<br>(g/g) |         |
|----------------------------------------------|---------------------|---------|--------------------------------------|---------|-----------------------------------|---------|
|                                              | Males               | Females | Males                                | Females | Males                             | Females |
| Control                                      | 1.70                | 1.05    | 0.282                                | 0.302   | 0.867                             | 0.538   |
| 1                                            | 1.45                | 1.12    | 0.286                                | 0.277   | 0.690                             | 0.583   |
| 10                                           | 1.63                | 1.05    | 0.283                                | 0.307   | 0.810                             | 0.606   |
| 100                                          | 1.49                | 1.06    | 0.286                                | 0.368** | 0.749                             | 0.564   |

\*\* Statistically significant difference at the 99 percent confidence level.

**TABLE XXVI**

**TEST MATERIAL: Aroclor 1260**

**Two-Year Chronic Oral Toxicity Study - Albino Rats**

**Organ Weight and Ratio Data**

**Six-Month Sacrifice**

**Summary of Mean Values**

**Organ: Brain**

| Dietary Level<br>of<br>Aroclor 1260<br>(ppm) | Organ Weight<br>(g) |         | Organ/Body Weight Ratio<br>(g/100 g) |         | Final Body Weights<br>of Sacrificed Rats<br>(g) |         |
|----------------------------------------------|---------------------|---------|--------------------------------------|---------|-------------------------------------------------|---------|
|                                              | Males               | Females | Males                                | Females | Males                                           | Females |
| Control                                      | 1.96                | 1.95    | 0.320                                | 0.560   | 613                                             | 348     |
| 1                                            | 2.10                | 1.92    | 0.412*                               | 0.474*  | 509                                             | 405     |
| 10                                           | 2.01                | 1.73    | 0.349                                | 0.504   | 575                                             | 343     |
| 100                                          | 1.99                | 1.88    | 0.382                                | 0.650*  | 522                                             | 289     |

\* Statistically significant difference at the 95 percent confidence level.

GBRNO02395

LEXOLDMON002732

c. Histopathologic Findings

Histopathologic examination of tissues and organs taken from all animals from both the control and 100 ppm groups was conducted.

Tables XXVII and XXVIII list all histopathologic changes noted.

All of the lesions noted in the microscopic examination of tissues were those of spontaneous disease and are not unusual for the albino rat. The most frequent findings were lesions in the trachea and lungs, indicating chronic murine pneumonia. These occurred in the control as well as the rats fed Aroclor 1260.

GBRN002396

LEXOI DMON002733

TABLE XXVII

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Histopathologic Changes

Six-Month Sacrifice

Group: Control

| Number of<br>Animals | Organ Examined  | Findings              | Incidence | Average<br>Grade |
|----------------------|-----------------|-----------------------|-----------|------------------|
| 5 Males              | Trachea         | Tracheitis            | 2         | 1.5              |
|                      |                 | Focal tracheitis      | 2         | 1.0              |
|                      | Lung            | Focal pneumonitis     | 1         | 0.5              |
|                      |                 | Hyperemia             | 2         | 1.0              |
|                      | Liver           | Focal pericholangitis | 1         | 1.0              |
|                      | Skeletal muscle | Focal inflammation    | 1         | 1.0              |
| 5 Females            | Trachea         | Focal tracheitis      | 1         | 1.0              |
|                      | Lung            | Hyperemia             | 2         | 2.5              |
|                      | Liver           | Hyperemia             | 1         | 1.0              |
|                      |                 | Pericholangitis       | 1         | 0.5              |
|                      | Kidney          | Focal inflammation    | 1         | 1.0              |

All other tissues and organs were normal histologically.

Grading System

0.5 = minimal

1.0 = slight

2.0 = mild

3.0 = moderate

4.0 = severe

5.0 = extreme

GBRN002397

LEXOLDMON002734



3. Twelve-Month Sacrifice

a. Gross Pathologic Findings

Gross pathologic findings among test animals were not significantly different from those noted in control animals.

b. Organ Weight and Ratio Data

The mean organ weight and ratio data collected at the twelve month sacrifice are presented in Tables XXIX through XXXIV.

Statistically significant differences are designated by asterisks following the test group value.

Organ weights, organ to body weight and organ to brain weight ratios disclosed several randomly occurring intergroup differences.

The lack of any consistent dose related response and the absence of any deleterious histopathologic change confirm that these differences were not related to the ingestion of Aroclor 1260.

GBRND02399

LEXOLDMON002736

TABLE XXIX

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Twelve-Month Sacrifice

Organ Weight and Ratio Data

Summary of Mean Values

Organ: Liver

| Dietary Level<br>(ppm) | Organ Weight<br>(g) |         | Organ/Body Weight Ratio<br>(g/100 g) |         | Organ/Brain Weight Ratio<br>(g/g) |         |
|------------------------|---------------------|---------|--------------------------------------|---------|-----------------------------------|---------|
|                        | Males               | Females | Males                                | Females | Males                             | Females |
| Control                | 22.1                | 15.5    | 3.79                                 | 3.52    | 10.8                              | 7.72    |
| 1                      | 17.1*               | 10.7*   | 2.79**                               | 3.01*   | 9.23                              | 5.76    |
| 10                     | 20.0                | 13.6    | 3.30                                 | 3.54    | 10.8                              | 7.28    |
| 100                    | 28.8*               | 14.1    | 4.26                                 | 3.79    | 13.4*                             | 8.52    |

\* Statistically significant difference at the 95 percent confidence level.

\*\* Statistically significant difference at the 99 percent confidence level.

GBRNO02400

LEXOLDMON002737

TABLE XXX

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Twelve-Month Sacrifice

Organ Weight and Ratio Data

Summary of Mean Values

Organ: Kidneys

| Dietary Level<br>(ppm) | Organ Weight<br>(g) |         | Organ/Body Weight Ratio<br>(g/100 g) |         | Organ/Brain Weight Ratio<br>(g/g) |         |
|------------------------|---------------------|---------|--------------------------------------|---------|-----------------------------------|---------|
|                        | Males               | Females | Males                                | Females | Males                             | Females |
| Control                | 4.04                | 3.19    | 0.692                                | 0.738   | 1.97                              | 1.58    |
| 1                      | 3.84                | 2.58**  | 0.625                                | 0.724   | 2.06                              | 1.39    |
| 10                     | 4.00                | 2.78*   | 0.660                                | 0.736   | 2.16                              | 1.48    |
| 100                    | 5.11*               | 2.81    | 0.758                                | 0.755   | 2.38*                             | 1.69    |

\* Statistically significant difference at the 95 percent confidence level.

\*\* Statistically significant difference at the 99 percent confidence level.

RECEIVED NOV 19 1970

GBRND02401

LEXOLDMON002738

**TABLE XXXI**

**TEST MATERIAL: Aroclor 1260**

**Two-Year Chronic Oral Toxicity Study - Albino Rats**

**Twelve-Month Sacrifice**

**Organ Weight and Ratio Data**

**Summary of Mean Values**

**Organ: Spleen**

| Dietary Level<br>(ppm) | Organ Weight<br>(g) |         | Organ/Body Weight Ratio<br>(g/100 g) |         | Organ/Brain Weight Ratio<br>(g/g) |         |
|------------------------|---------------------|---------|--------------------------------------|---------|-----------------------------------|---------|
|                        | Males               | Females | Males                                | Females | Males                             | Females |
| Control                | 0.778               | 0.796   | 0.133                                | 0.183   | 0.380                             | 0.396   |
| 1                      | 0.756               | 0.460*  | 0.122                                | 0.129*  | 0.408                             | 0.247*  |
| 10                     | 0.836               | 0.578   | 0.139                                | 0.150   | 0.457                             | 0.308   |
| 100                    | 0.785               | 0.474*  | 0.116                                | 0.127*  | 0.365                             | 0.284   |

\* Statistically significant difference at the 95 percent confidence level.

GBRND02402

LEXOLDMON002739

TABLE XXXII

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Twelve-Month Sacrifice

Organ Weight and Ratio Data

Summary of Mean Values

Organ: Gonads

| Dietary Level<br>(ppm) | Organ Weight<br>(g) |         | Organ/Body Weight Ratio<br>(g/100 g) |         | Organ/Brain Weight Ratio<br>(g/g) |         |
|------------------------|---------------------|---------|--------------------------------------|---------|-----------------------------------|---------|
|                        | Males               | Females | Males                                | Females | Males                             | Females |
| Control                | 3.46                | 0.116   | 0.596                                | 0.0275  | 1.70                              | 0.0574  |
| 1                      | 3.38                | 0.103   | 0.551                                | 0.0287  | 1.82                              | 0.0552  |
| 10                     | 3.51                | 0.112   | 0.578                                | 0.0294  | 1.89                              | 0.0584  |
| 100                    | 4.03**              | 0.088*  | 0.598                                | 0.0238  | 1.88                              | 0.0527  |

\* Statistically significant difference at the 95 percent confidence level.

\*\* Statistically significant difference at the 99 percent confidence level.

GBRND02403

LEXOLDMON002740

**TABLE XXXIII**

**TEST MATERIAL: Aroclor 1260**

**Two-Year Chronic Oral Toxicity Study - Albino Rats**

**Twelve-Month Sacrifice**

**Organ Weight and Ratio Data**

**Summary of Mean Values**

**Organ: Heart**

| <b>Dietary Level<br/>(ppm)</b> | <b>Organ Weight<br/>(g)</b> |                | <b>Organ/Body Weight Ratio<br/>(g/100 g)</b> |                | <b>Organ/Brain Weight Ratio<br/>(g/g)</b> |                |
|--------------------------------|-----------------------------|----------------|----------------------------------------------|----------------|-------------------------------------------|----------------|
|                                | <b>Males</b>                | <b>Females</b> | <b>Males</b>                                 | <b>Females</b> | <b>Males</b>                              | <b>Females</b> |
| <b>Control</b>                 | <b>1.71</b>                 | <b>1.23</b>    | <b>0.293</b>                                 | <b>0.287</b>   | <b>0.835</b>                              | <b>0.611</b>   |
| <b>1</b>                       | <b>1.68</b>                 | <b>1.07</b>    | <b>0.273</b>                                 | <b>0.301</b>   | <b>0.902</b>                              | <b>0.575</b>   |
| <b>10</b>                      | <b>1.59</b>                 | <b>1.16</b>    | <b>0.261</b>                                 | <b>0.307</b>   | <b>0.845</b>                              | <b>0.620</b>   |
| <b>100</b>                     | <b>1.86</b>                 | <b>1.27</b>    | <b>0.275</b>                                 | <b>0.340</b>   | <b>0.868</b>                              | <b>0.760</b>   |

GBRNO02404

LEXOLDMON002741

TABLE XXXIV

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Organ Weight and Ratio Data

Twelve-Month Sacrifice

Summary of Mean Values

Organ: Brain

| Dietary Level<br>(ppm) | Organ Weight<br>(g) |         | Organ/Body Weight Ratio<br>(g/100 g) |         | Final Body Weight of<br>Sacrificed Rats (grams) |         |
|------------------------|---------------------|---------|--------------------------------------|---------|-------------------------------------------------|---------|
|                        | Males               | Females | Males                                | Females | Males                                           | Females |
| Control                | 2.04                | 2.02    | 0.349                                | 0.460   | 584                                             | 439     |
| 1                      | 1.87                | 1.86    | 0.304                                | 0.522   | 616                                             | 356     |
| 10                     | 1.88                | 1.90    | 0.310                                | 0.493   | 607                                             | 385     |
| 100                    | 2.15                | 1.67**  | 0.318                                | 0.449   | 675                                             | 372     |

\*\* Statistically significant difference at the 99 percent confidence level.

c. Histopathologic Findings

Histopathologic examination of tissues taken from each rat sacrificed from the control and 100 ppm groups after twelve months of testing was conducted.

Tables XXXV and XXXVI list all hisopathologic changes noted.

All of the lesions noted in the microscopic examination of tissues were those of spontaneous disease and are not unusual for the albino rat. The most frequent findings were lesions in the trachea and lungs, indicating chronic murine pneumonia. These occurred in the control as well as the rats fed Aroclor 1260.

GBRNO02406

LEXOLDMON002743

TABLE XXXV

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Histopathologic Changes

Twelve-Month Sacrifice

Group: Control

| Number of<br>Animals | Organ Examined  | Findings                       | Incidence | Average<br>Grade |
|----------------------|-----------------|--------------------------------|-----------|------------------|
| 5 Males              | Trachea         | Tracheitis                     | 1         | 1.0              |
|                      | Lung            | Chronic respiratory<br>disease | 2         | 1.0              |
|                      | Urinary Bladder | Calculus                       | 1         | 1.0              |
| 5 Females            | Trachea         | Tracheitis                     | 1         | 1.0              |
|                      | Lung            | Chronic respiratory<br>disease | 1         | 1.0              |
|                      | Colon           | Parasites                      | 1         | 1.0              |

All other tissues and organs were normal histologically.

Grading System

0.5 = minimal  
1.0 = slight  
2.0 = mild  
3.0 = moderate  
4.0 = severe  
5.0 = extreme

GBRN002407

LEXOLDMON002744

TABLE XXXVI

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Histopathologic Changes

Twelve-Month Sacrifice

Group: 100 ppm

| Number of<br>Animals | Organ Examined | Findings                           | Incidence | Average<br>Grade |
|----------------------|----------------|------------------------------------|-----------|------------------|
| 5 Males              | Trachea        | Tracheitis                         | 1         | 1.0              |
|                      | Lung           | Chronic respiratory<br>disease     | 3         | 1.0              |
|                      | Kidney         | Focal lymphoid<br>infiltration     | 1         | 1.0              |
|                      | Liver          | Vacuolar change of<br>random cells | 1         | 1.0              |
| 5 Females            | Trachea        | Tracheitis                         | 1         | 1.0              |
|                      | Lung           | Chronic respiratory<br>disease     | 1         | 1.0              |

All other tissues and organs were normal histologically.

Grading System

0.5 = minimal  
1.0 = slight  
2.0 = mild  
3.0 = moderate  
4.0 = severe  
5.0 = extreme

GBRN002408

LEXOLDMON002745

4. Final Sacrifice

a. Gross Pathologic Findings

Gross pathologic findings among test animals were not significantly different from those noted in control animals.

b. Organ Weight and Ratio Data

The mean organ weight and ratio data collected at the final sacrifice are presented in Tables XXXVII through XLII. Statistically significant differences are designated by asterisks following the test group value.

Absolute liver weights and liver to body weight or liver to brain weight ratios were significantly elevated in those rats fed 100 ppm Aroclor 1260. Other organ weights, organ to body weight and organ to brain weight ratios disclosed several randomly occurring intergroup differences. The lack of any consistent dose related response and the absence of any deleterious histopathologic change confirm that these differences were not related to the ingestion of Aroclor 1260

**TABLE XXXVII**  
**TEST MATERIAL: Aroclor 1260**  
**Two-Year Chronic Oral Toxicity Study - Albino Rats**  
**Organ Weight and Ratio Data**  
**Summary of Mean Values**

Final Sacrifice

Organ: Liver

| Dietary Level<br>(ppm) | Organ Weight<br>(g) |         | Organ/Body Weight Ratio<br>(g/100 g) |         | Organ/Brain Weight Ratio<br>(g/g) |         |
|------------------------|---------------------|---------|--------------------------------------|---------|-----------------------------------|---------|
|                        | Males               | Females | Males                                | Females | Males                             | Females |
| Control                | 22.0                | 16.3    | 3.47                                 | 3.05    | 10.10                             | 8.42    |
| 1                      | 22.0                | 16.6    | 3.40                                 | 3.35    | 9.93                              | 8.62    |
| 10                     | 24.1                | 18.1    | 2.87*                                | 3.19    | 11.20                             | 9.26    |
| 100                    | 29.9**              | 22.7**  | 4.67**                               | 4.42**  | 13.40*                            | 11.60*  |

\* Statistically significant difference at the 95 percent confidence level.

\*\* Statistically significant difference at the 99 percent confidence level.

**TABLE XXXIX**

**TEST MATERIAL: Aroclor 1260**

**Two-Year Chronic Oral Toxicity Study - Albino Rats**

**Organ Weight and Ratio Data**

**Summary of Mean Values**

**Final Sacrifice**

**Organ: Spleen**

| Dietary Level<br>(ppm) | Organ Weight<br>(g) |         | Organ/Body Weight Ratio<br>(g/100 g) |         | Organ/Brain Weight Ratio<br>(g/g) |         |
|------------------------|---------------------|---------|--------------------------------------|---------|-----------------------------------|---------|
|                        | Males               | Females | Males                                | Females | Males                             | Females |
| Control                | 1.15                | 0.96    | 0.182                                | 0.181   | 0.520                             | 0.492   |
| 1                      | 1.17                | 0.72    | 0.178                                | 0.148   | 0.529                             | 0.370   |
| 10                     | 1.09                | 0.83    | 0.131                                | 0.142   | 0.509                             | 0.428   |
| 100                    | 0.89                | 0.75    | 0.138                                | 0.146   | 0.400                             | 0.379   |

GBRND02411

**TABLE XL**  
**TEST MATERIAL: Aroclor 1260**  
**Two-Year Chronic Oral Toxicity Study - Albino Rats**

**Organ Weight and Ratio Data**

**Summary of Mean Values**

**Final Sacrifice**

**Organ: Gonads**

| Dietary Level<br>(ppm) | Organ Weight<br>(g) |         | Organ/Body Weight Ratio<br>(g/100 g) |         | Organ/Brain Weight Ratio<br>(g/g) |         |
|------------------------|---------------------|---------|--------------------------------------|---------|-----------------------------------|---------|
|                        | Males               | Females | Males                                | Females | Males                             | Females |
| Control                | 2.52                | 0.123   | 0.397                                | 0.0222  | 1.15                              | 0.0628  |
| 1                      | 3.56*               | 0.140   | 0.550*                               | 0.0317  | 1.61*                             | 0.0718  |
| 10                     | 3.36*               | 0.177   | 0.400                                | 0.0338  | 1.56*                             | 0.0900  |
| 100                    | 2.97                | 0.113   | 0.464                                | 0.0229  | 1.33                              | 0.0584  |

\* Statistically significant difference at the 95 percent confidence level.

GBRND02412

LEXOLDMON002749

**TABLE XLI**  
**TEST MATERIAL: Aroclor 1260**  
**Two-Year Chronic Oral Toxicity Study - Albino Rats**

**Organ Weight and Ratio Data**

**Summary of Mean Values**

**Final Sacrifice**

**Organ: Heart**

| Dietary Level<br>(ppm) | Organ Weight<br>(g) |         | Organ/Body Weight Ratio<br>(g/100 g) |         | Organ/Brain Weight Ratio<br>(g/g) |         |
|------------------------|---------------------|---------|--------------------------------------|---------|-----------------------------------|---------|
|                        | Males               | Females | Males                                | Females | Males                             | Females |
| Control                | 1.87                | 1.54    | 0.294                                | 0.288   | 0.860                             | 0.795   |
| 1                      | 1.98                | 1.45    | 0.304                                | 0.294   | 0.893                             | 0.747   |
| 10                     | 1.90                | 1.58    | 0.225**                              | 0.279   | 0.884                             | 0.809   |
| 100                    | 2.00                | 1.49    | 0.313                                | 0.288   | 0.904                             | 0.764   |

\*\* Statistically significant difference at the 99 percent confidence level.

GBRND02413

**TABLE XLII**

**TEST MATERIAL: Aroclor 1260**

**Two-Year Chronic Oral Toxicity Study - Albino Rats**

**Organ Weight and Ratio Data**

**Summary of Mean Values**

**Final Sacrifice**

**Organ: Brain**

| Dietary Level<br>(ppm) | Organ Weight<br>(g) |         | Organ/Body Weight Ratio<br>(g/100 g) |        | Final Body Weight of<br>Sacrificed Animals (grams) |         |
|------------------------|---------------------|---------|--------------------------------------|--------|----------------------------------------------------|---------|
|                        | Males               | Females | Males                                | Female | Males                                              | Females |
| Control                | 2.18                | 1.95    | 0.342                                | 0.362  | 642                                                | 551     |
| 1                      | 2.21                | 1.94    | 0.343                                | 0.407  | 654                                                | 525     |
| 10                     | 2.15                | 1.96    | 0.255**                              | 0.347  | 845**                                              | 584     |
| 100                    | 2.22                | 1.95    | 0.346                                | 0.378  | 641                                                | 644     |

\*\* Statistically significant difference at the 99 percent confidence level.

GBRNO02414

c. Histopathologic Findings

Histopathologic examination of tissues and organs taken from all animals sacrificed after 24 months on test was conducted.

Tables XLIII through XLVI list all histopathologic changes noted. Table XLVII contains the results of the histopathologic examination of the urinary bladders.

Significant liver injury was found in animals fed 100 ppm Aroclor 1260 (T-III). The compound associated lesions consisted of vacuolar change, focal hypertrophy and focal hyperplasia. In addition, there were lesions of inflammation, necrosis, fibrosis and minor degeneration in this group which, although seen in the controls and lower test groups, were more severe and frequent in the T-III group.

The vacuolar change was not seen in the control animals but was observed occasionally in livers from T-I (1 ppm) and T-II (10 ppm) animals. It was frequent in the T-III animals. This lesion is morphologically indicative of fatty degeneration. Formalin-frozen sections of livers from representative animals which displayed vacuolar changes were stained with Oil Red O to reveal the presence of fat. The vacuolar lesion in the cytoplasm of these cells was positively identified as fat.

The hypertrophic change found in the liver was focal and often limited to the central lobular area where groups of cells were swollen to two or three times their normal size with clear pink homogenous cytoplasm. The hyperplasia was associated with the same cells and appeared to be an

extension of the hypertrophic lesion. The hyperplastic cells were also usually hypertrophic. The most severe examples of hyperplasia appeared as nodular growths with limited compression of the surrounding normal hepatic tissue.

The minor lesions of degeneration, hepatitis, ductule cell proliferation, necrosis and focal lymphoid infiltration seen in the control animals and the T-I and T-II groups are lesions of spontaneous disease and are not related to the Aroclor 1260. This level of liver disease is not unusual in old animals.

Hyperplasia of the urinary bladder was found in an animal from the control group. This finding was not present in any of the rats fed Aroclor 1260. All of the other lesions in other tissues found in these animals are related to spontaneous disease and they are not unusual for old rats.



Donovan E. Gordon, D. V. M., Ph. D.  
Diplomate, American College of  
Veterinary Pathologists

GBRN002416

LEXOLDMON002753

TABLE XLIII

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Histopathologic Changes

Final Sacrifice

Group: Control

| Number of<br>Animals | Organ Examined | Findings                    | Incidence | Average<br>Grade |
|----------------------|----------------|-----------------------------|-----------|------------------|
| 11 Males             | Heart          | Chronic focal myocarditis   | 3         | 1.0              |
|                      | Trachea        | Tracheitis                  | 3         | 1.0              |
|                      | Lung           | Chronic respiratory disease | 9         | 1.0              |
|                      | Liver          | Focal degeneration          | 1         | 1.0              |
|                      |                | Ductile hyperplasia         | 2         | 1.0              |
|                      | Stomach        | Chronic gastritis           | 1         | 1.0              |
|                      | Spleen         | Hemosiderosis               | 3         | 1.0              |
|                      | Kidney         | Chronic nephritis           | 8         | 1.0              |
|                      | Testes         | Testicular degeneration     | 2         | 1.0              |
|                      |                | Periarteritis               | 2         | 1.0              |
|                      |                | Inflammation                | 1         | 1.0              |
|                      | Prostate       | Telangiectasis              | 1         | 1.0              |
|                      | Adrenal Gland  | Nodular hyperplasia         | 1         | 1.0              |
|                      |                | Focal fatty change          | 1         | 1.0              |
|                      |                | Focal encephalitis          | 1         | 1.0              |
|                      | Brain          | Inflammation                | 2         | 1.0              |
|                      | Eye            |                             |           |                  |

GBRNO02417



TABLE XLIV

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Histopathologic Changes

Final Sacrifice

Group: 1 ppm

| Number of<br>Animals | Organ Examined  | Findings                    | Incidence | Average<br>Grade |
|----------------------|-----------------|-----------------------------|-----------|------------------|
| 11 Males             | Heart           | Chronic focal myocarditis   | 4         | 1.0              |
|                      | Trachea         | Tracheitis                  | 7         | 1.0              |
|                      | Lung            | Chronic respiratory disease | 10        | 2.0              |
|                      | Liver           | Focal hepatitis             | 1         | 1.0              |
|                      |                 | Focal scarring              | 1         | 1.0              |
|                      |                 | Hemosiderosis               | 1         | 1.0              |
|                      | Pancreas        | Chronic focal pancreatitis  | 2         | 1.0              |
|                      | Stomach         | Acute focal gastritis       | 1         | 3.0              |
|                      | Spleen          | Hemosiderosis               | 1         | 1.0              |
|                      | Lymph Node      | Congestion                  | 1         | 1.0              |
|                      | Kidney          | Chronic nephritis           | 7         | 2.0              |
|                      | Testes          | Testicular degeneration     | 1         | 2.0              |
|                      |                 | Periarteritis               | 2         | 1.0              |
|                      |                 | Inflammation                | 4         | 2.0              |
|                      | Prostate        | Inflammation                | 4         | 2.0              |
|                      | Seminal Vesicle | Inflammation                | 1         | 2.0              |

GBRND02419

TABLE XLIV continued

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Histopathologic Changes

Final Sacrifice

Group: 1 ppm

| Number of<br>Animals | Organ Examined | Findings                    | Incidence | Average<br>Grade |
|----------------------|----------------|-----------------------------|-----------|------------------|
| 11 Males             | Adrenal Gland  | Nodular hyperplasia         | 2         | 1.0              |
|                      |                | Focal fatty change          | 2         | 1.0              |
| 14 Females           | Heart          | Chronic focal myocarditis   | 3         | 1.0              |
|                      |                | Chronic focal myocarditis   | 1         | 3.0              |
|                      | Trachea        | Tracheitis                  | 9         | 1.0              |
|                      | Lung           | Chronic respiratory disease | 13        | 1.0              |
|                      |                | Focal fibrosis              | 1         | 1.0              |
|                      | Liver          | Focal lymphoid infiltration | 1         | 1.0              |
|                      |                | Ductile hyperplasia         | 1         | 0.5              |
|                      |                | Congestion                  | 1         | 1.0              |
|                      |                | Vacuolar change (fatty)     | 1         | 2.0              |
|                      | Stomach        | Chronic gastritis           | 1         | 1.0              |
|                      |                | Acanthosis                  | 2         | 1.0              |

GBRND02420

LEXOLDMON002757

TABLE XLIV continued  
 TEST MATERIAL: Aroclor 1260  
 Two-Year Chronic Oral Toxicity Study - Albino Rats  
 Histopathologic Changes

Final Sacrifice

Group: 1 ppm

| Number of<br>Animals | Organ Examined | Findings                  | Incidence | Average<br>Grade |
|----------------------|----------------|---------------------------|-----------|------------------|
| 14 Females           | Stomach        | Ulceration and acanthosis | 1         | 1.0              |
|                      | Colon          | Parasites                 | 1         | 1.0              |
|                      | Spleen         | Hemosiderosis             | 11        | 1.0              |
|                      |                | Hyperplasia               | 1         | 1.0              |
|                      | Lymph Node     | Congestion                | 1         | 1.0              |
|                      | Kidney         | Chronic nephritis         | 3         | 2.0              |
|                      |                | Vacuolar change (fatty)   | 1         | 1.0              |
|                      | Uterus         | Metritis                  | 4         | 1.0              |
|                      | Adrenal Gland  | Telangiectasis            | 9         | 1.0              |
|                      |                | Nodular hyperplasia       | 3         | 1.0              |
|                      |                | Congestion                | 1         | 1.0              |

GBRND02421

All other tissues and organs were normal histologically.

Grading System

0.5 = minimal      2.0 = mild      4.0 = severe  
 1.0 = slight      3.0 = moderate      5.0 = extreme

TABLE XLV

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Histopathologic Changes

Final Sacrifice

Group: 10 ppm

| Number of<br>Animals | Organ Examined | Findings                    | Incidence | Average<br>Grade |
|----------------------|----------------|-----------------------------|-----------|------------------|
| 7 Males              | Heart          | Chronic focal myocarditis   | 3         | 1.0              |
|                      | Trachea        | Tracheitis                  | 2         | 1.0              |
|                      | Lung           | Chronic respiratory disease | 5         | 1.0              |
|                      |                | Suppurative pneumonia       | 1         | 3.0              |
|                      | Liver          | Vacuolar change (fatty)     | 2         | 1.0              |
|                      | Stomach        | Chronic gastritis           | 1         | 1.0              |
|                      |                | Acanthosis                  | 2         | 1.0              |
|                      | Spleen         | Hemosiderosis               | 3         | 1.0              |
|                      | Kidney         | Chronic nephritis           | 6         | 1.0              |
|                      | Prostate       | Inflammation                | 2         | 1.0              |
|                      | Adrenal Gland  | Nodular hyperplasia         | 1         | 1.0              |
| 14 Females           | Heart          | Chronic focal myocarditis   | 3         | 1.0              |
|                      | Trachea        | Tracheitis                  | 8         | 1.0              |
|                      | Lung           | Chronic respiratory disease | 14        | 1.0              |

GBRND02422

TABLE XLV continued  
TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Histopathologic Changes

Final Sacrifice

Group: 10 ppm

| Number of<br>Animals | Organ Examined | Findings                   | Incidence | Average<br>Grade |
|----------------------|----------------|----------------------------|-----------|------------------|
| 14 Females           | Liver          | Vacuolar change (fatty)    | 1         | 1.0              |
|                      |                | Focal hyperplasia          | 1         | 1.0              |
|                      | Pancreas       | Chronic focal pancreatitis | 1         | 3.0              |
|                      |                | Periarteritis              | 1         | 3.0              |
|                      | Stomach        | Acanthosis                 | 1         | 1.0              |
|                      | Spleen         | Hemosiderosis              | 12        | 1.0              |
|                      | Kidney         | Chronic nephritis          | 10        | 1.0              |
|                      | Uterus         | Metritis                   | 7         | 2.0              |
|                      | Pituitary      | Congestion                 | 1         | 1.0              |
|                      | Adrenal Gland  | Telangiectasis             | 7         | 1.0              |
|                      |                | Nodular hyperplasia        | 1         | 1.0              |
|                      |                | Congestion                 | 1         | 1.0              |

GBRND02423

All other tissues and organs were normal histologically.

.5 = minimal      Grading System      2.0 = mild      4.0 = severe

TABLE XLVI

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Histopathologic Changes

Final Sacrifice

Group: 100 ppm

| Number of<br>Animals | Organ Examined | Findings                        | Incidence                   | Average<br>Grade |     |
|----------------------|----------------|---------------------------------|-----------------------------|------------------|-----|
| 10 Males             | Heart          | Chronic focal myocarditis       | 5                           | 1.0              |     |
|                      |                | Trachea                         | Tracheitis                  | 8                | 1.0 |
|                      |                | Lung                            | Chronic respiratory disease | 9                | 2.0 |
|                      | Liver          |                                 | Congestion                  | 1                | 1.0 |
|                      |                |                                 | Vacuolar change (fatty)     | 2                | 2.0 |
|                      |                |                                 | Focal degeneration          | 1                | 1.0 |
|                      |                |                                 | Congestion                  | 3                | 2.0 |
|                      |                |                                 | Centrolobular hypertrophy   | 2                | 1.0 |
|                      |                |                                 | Telangiectasis              | 1                | 1.0 |
|                      |                | Pancreas                        | Chronic focal pancreatitis  | 2                | 1.0 |
|                      |                | Stomach                         | Inflammation                | 2                | 2.0 |
|                      |                |                                 | Acanthosis                  | 2                | 1.0 |
|                      | Spleen         | Hemosiderosis                   | 7                           | 1.0              |     |
|                      | Kidney         | Focal lymphoid infiltration     | 1                           | 1.0              |     |
|                      |                | Calcified concretions in pelvis | 1                           | 1.0              |     |

GBRND02424

LEXOLDMON002761

TABLE XLVI continued

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Histopathologic Changes

Final Sacrifice

Group: 100 ppm

| Number of<br>Animals | Organ Examined     | Findings                        | Incidence | Average<br>Grade |
|----------------------|--------------------|---------------------------------|-----------|------------------|
| 10 Males             | Kidney (continued) | Calcified concretions in pelvis | 1         | 1.0              |
|                      | Testes             | Aspermia                        | 1         | 1.0              |
|                      | Prostate           | Inflammation                    | 1         | 1.0              |
|                      | Thymus             | Cyst                            | 1         | 1.0              |
|                      | Adrenal Gland      | Telangiectasis                  | 3         | 1.0              |
|                      |                    | Nodular hyperplasia             | 1         | 1.0              |
| 15 Females           | Heart              | Chronic focal myocarditis       | 5         | 1.0              |
|                      | Trachea            | Tracheitis                      | 8         | 1.0              |
|                      | Lung               | Chronic respiratory disease     | 11        | 1.0              |
|                      |                    | Congestion                      | 1         | 1.0              |
|                      |                    | Edema                           | 1         | 1.0              |
|                      | Liver              | Vacuolar change (fatty)         | 5         | 1.0              |

GBRND02425

TABLE XLVI continued

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Histopathologic Changes

Final Sacrifice

Group: 100 ppm

| Number of<br>Animals                                                     | Organ Examined    | Findings                                                                                                     | Incidence | Average<br>Grade |
|--------------------------------------------------------------------------|-------------------|--------------------------------------------------------------------------------------------------------------|-----------|------------------|
| 15 Females<br><br><br><br><br><br><br><br><br><br><br><br><br>GBRND02426 | Liver (continued) | Focal hyperplasia                                                                                            | 4         | 1.0              |
|                                                                          |                   | Chronic inflammation of liver capsule                                                                        | 1         | 1.0              |
|                                                                          |                   | Centrolobular hypertrophy                                                                                    | 5         | 1.0              |
|                                                                          |                   | Necrosis                                                                                                     | 2         | 2.0              |
|                                                                          |                   | Ductile cell hyperplasia, portal fibrosis<br>and widespread eosinophil inclusions<br>in hepatocyte cytoplasm | 1         | 1.0              |
|                                                                          | Pancreas          | Chronic focal pancreatitis and<br>periarteritis                                                              | 1         | 3.0              |
|                                                                          | Spleen            | Hemosiderosis                                                                                                | 9         | 2.0              |
|                                                                          | Kidney            | Chronic nephritis                                                                                            | 8         | 1.0              |
|                                                                          | Uterus            | Metritis                                                                                                     | 7         | 1.0              |
|                                                                          | Pituitary         | Congestion                                                                                                   | 1         | 1.0              |
|                                                                          | Adrenal Gland     | Telangiectasis                                                                                               | 6         | 1.0              |
|                                                                          |                   | Nodular hyperplasia                                                                                          | 1         | 1.0              |
|                                                                          | Parathyroid       | Congestion                                                                                                   | 1         | 1.0              |
|                                                                          |                   | Hyperplasia                                                                                                  | 1         | 1.0              |

All other tissues and organs were normal histologically.

Grading System

TABLE XLVII

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Histopathologic Changes of the Urinary Bladder

Final Sacrifice

| Group              | Sex | Number<br>Examined | Findings*                             | Incidence | Average<br>Grade |
|--------------------|-----|--------------------|---------------------------------------|-----------|------------------|
| C                  | M   | 11                 | Proteinaceous plug (34)               | 1         | -                |
|                    | F   | 14                 | Focal epithelial hyperplasia (62)     | 1         | 1.0              |
| T-I<br>(1 ppm)     | M   | 10                 | Proteinaceous plug (86, 105, and 107) | 3         | -                |
|                    | F   | 11                 | None                                  | -         | -                |
| T-II<br>(10 ppm)   | M   | 7                  | Proteinaceous plug (187)              | 1         | -                |
|                    | F   | 14                 | None                                  | -         | -                |
| T-III<br>(100 ppm) | M   | 10                 | Proteinaceous plug (271)              | 1         | -                |
|                    | F   | 14                 | None                                  | -         | -                |

\* Individual animal numbers are shown in parentheses.

Grading System

1.0 = minimal or trace      4.0 = severe  
 2.0 = mild                      5.0 = extreme  
 3.0 = moderate

GBRND02427

LEXOLDMON002764

G. Detailed Description of Tumor Findings

Tumor data for individual animals including location, weight and pathologic classification are presented in Tables XLVIII through LI.

None of these tumors could be related to the ingestion of Aroclor 1260 and are considered normal for a random population of rats this age.

GBRN002428

LEXOLDMON002765

TABLE XLVIII

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Description of Tumor Findings

Group: Control

| Animal<br>Number<br>and Sex | Location                 | Weight<br>(grams) | Size<br>(cm)           | Classification                               | Months<br>on<br>Test |
|-----------------------------|--------------------------|-------------------|------------------------|----------------------------------------------|----------------------|
| 1-M                         | Abdomen                  | -                 | -                      | Mammary fibroadenoma                         | 24                   |
| 2-M                         | Pancreas                 | 11.76             | -                      | Adenocarcinoma                               | 24                   |
| 46-F                        | Abdomen                  | 64.98<br>35.09    | 8 x 1 x 2<br>5 x 2 x 2 | Mammary fibroadenoma<br>Mammary fibroadenoma | 24                   |
| 47-F                        | Lymph node               | -                 | -                      | Reticulum cell sarcoma                       | 24                   |
| 48-F                        | Abdomen                  | -                 | 2 x 2 x 1              | Mammary fibroadenoma                         | 24                   |
| 50-F                        | Abdomen                  | -                 | 2 x 2 x 3              | Mammary fibroadenoma                         | 7                    |
| 52-F                        | Abdomen                  | -                 | -                      | Mammary fibroadenoma                         | 24                   |
| 59-F                        | Abdomen<br>Adrenal gland | -<br>-            | -<br>-                 | Mammary fibroadenoma<br>Pheochromocytoma     | 24                   |
| 64-F                        | Abdomen                  | 88.92             | 13 x 11 x 4            | Fibroadenoma                                 | 24                   |
| 66-F                        | Abdomen                  | 230.00            | 13 x 11 x 4            | Mammary adenoma                              | 24                   |

GBRND02429

TABLE XLIX

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Description of Tumor Findings

Group: 1 ppm

| Animal<br>Number<br>and Sex | Location             | Weight<br>(grams) | Size<br>(cm)   | Classification                                  | Months<br>on<br>Test |
|-----------------------------|----------------------|-------------------|----------------|-------------------------------------------------|----------------------|
| 99-M                        | Pituitary            | -                 | -              | ✓Chromophobe adenoma                            | 20                   |
| 127-F                       | Pituitary<br>Abdomen | -<br>25.00        | -<br>6 x 3 x 1 | ✓ Chromophobe adenoma<br>✓ Mammary fibroadenoma | 24                   |
| 128-F                       | Pituitary<br>Abdomen | -<br>60.80        | -<br>5 x 6 x 2 | ✓ Chromophobe adenoma<br>✓ Mammary fibroadenoma | 21                   |
| 138-F                       | Abdomen<br>Pituitary | 20.07<br>-        | -<br>-         | ✓ Mammary fibroadenoma<br>✓ Chromophobe adenoma | 24                   |
| 141-F                       | Ovary                | -                 | -              | ✓Adenoma                                        | 24                   |
| 151-F                       | Abdomen              | 14.89             | -              | ✓ Mammary fibroadenoma                          | 24                   |
| 152-F                       | Abdomen              | 540.00            | 13 x 15 x 2    | ✓ Mammary fibroadenoma                          | 22                   |

GBRNO02430

LEXOLDMON002767

**TABLE L**

**TEST MATERIAL: Aroclor 1260**

**Two-Year Chronic Oral Toxicity Study - Albino Rats**

**Description of Tumor Findings**

**Group: 10 ppm**

| <b>Animal<br/>Number<br/>and Sex</b> | <b>Location</b>        | <b>Weight<br/>(grams)</b> | <b>Size<br/>(cm)</b> | <b>Classification</b>  | <b>Months<br/>on<br/>Test</b> |
|--------------------------------------|------------------------|---------------------------|----------------------|------------------------|-------------------------------|
| 205-F                                | Abdomen                | 190.00                    | 9 x 5 x 4            | ✓ Mammary fibroadenoma | 24                            |
| 208-F                                | Pituitary              | -                         | -                    | ✓ Chromophobe adenoma  | 24                            |
| 213-F                                | Abdomen                | -                         | -                    | ✓ Mammary fibroadenoma | 24                            |
| 216-F                                | Right shoulder         | 138.00                    | -                    | ✓ Subcutaneous tumor   | 23                            |
| 220-F                                | Abdomen                | 9.57                      | -                    | ✓ Mammary fibroadenoma | 24                            |
|                                      | Pituitary              | -                         | -                    | ✓ Chromophobe adenoma  |                               |
| 221-F                                | Submaxillary<br>region | 14.44                     | 2 x 3 x 1            | ✓ Fibroadenoma         | 24                            |
|                                      | Abdomen                | 29.0                      | 3 x 5 x 2            | ✓ Mammary fibroadenoma |                               |

GBRND02431

LEXOLDMON002768

**TABLE L continued**

**TEST MATERIAL: Aroclor 1260**

**Two-Year Chronic Oral Toxicity Study - Albino Rats**

**Description of Tumor Findings**

**Group: 10 ppm**

| <b>Animal<br/>Number<br/>and Sex</b> | <b>Location</b>        | <b>Weight<br/>(grams)</b> | <b>Size<br/>(cm)</b> | <b>Classification</b>                          | <b>Months<br/>on<br/>Test</b> |
|--------------------------------------|------------------------|---------------------------|----------------------|------------------------------------------------|-------------------------------|
| 22-F                                 | Left femoral<br>region | 56.00                     | 6 x 8 x 2            | ✓ Subcutaneous tumor                           | 6                             |
| 233-F                                | Pituitary              | -                         | -                    | ✓ Chromophobe adenoma                          | 24                            |
| 234-F                                | Abdomen<br>Pancreas    | 60.82<br>-                | 6 x 4 x 3<br>-       | ✓ Mammary fibroadenoma<br>✓ Islet cell adenoma | 24                            |

6BRN002432

LEXOLDMON002769

TABLE LI

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Description of Tumor Findings

Group: 100 ppm

| Animal<br>Number<br>and Sex | Location            | Weight<br>(grams) | Size<br>(cm) | Classification         | Months<br>on<br>Test |
|-----------------------------|---------------------|-------------------|--------------|------------------------|----------------------|
| 244-M                       | Pancreas            | -                 | -            | Adenoma                | 24                   |
| 264-M                       | Pancreas            | -                 | -            | ✓ Islet cell adenoma   | 21                   |
| 281-F                       | Abdomen             | -                 | -            | ✓ Mammary fibroadenoma | 24                   |
| 293-F                       | Liver               | 35.04             | 6 x 4 x 2    | Lymphosarcoma          | 23                   |
| 294-F                       | Abdominal<br>cavity | 50.75             | 6 x 4 x 3    | ✓ Fibroma              | 24                   |
| 301-F                       | Liver               | -                 | -            | Lymphosarcoma          | 23                   |

GBRND02433

LEXOLDMON002770

TABLE LI continued

TEST MATERIAL: Aroclor 1260

Two-Year Chronic Oral Toxicity Study - Albino Rats

Description of Tumor Findings

Group: 100 ppm

| Animal<br>Number<br>and Sex | Location   | Weight<br>(grams) | Size<br>(cm)           | Classification                                   | Months<br>on<br>Test |
|-----------------------------|------------|-------------------|------------------------|--------------------------------------------------|----------------------|
| 302-F                       | Lymph node | -                 | -                      | ✓ Lymphosarcoma                                  | 24                   |
| 308-F                       | Pituitary  | -                 | -                      | ✓ Chromophobe adenoma                            | 24                   |
| 311-F                       | Abdomen    | 14.00             | 4 x 3 x 2<br>1 x 2 x 1 | ✓ Mammary fibroadenoma<br>✓ Mammary fibroadenoma | 24                   |
| 312-M                       | Pituitary  | -                 | -                      | ✓ Chromophobe adenoma                            | 24                   |
| 314-F                       | Abdomen    | 26.00             | 5 x 4 x 3              | ✓ Mammary fibroadenoma                           | 24                   |

GBRNO02434

LEXOLDMON002771