2ND DRAFT

POLYCHLORINATED BIPHENYLS

CRITERIA:

Aquatic Life - 0.001 ug/l in freshwater;
0.01 ug/l in marine water

INTRODUCTION:

Polychlorinated biphenyls (PCB's) are a class of compounds produced by the chlorination of biphenyls and are known in the United States commercially as Aroclors (R). The degree of chlorination determines their chemical properties and generally their composition can be identified by the numerical nomenclature, e.g., Aroclor 1242, Aroclor 1254, etc. The first two digits represent the molecular type and the last two digits the average percentage by weight of chlorine (NTIS, 1972).

Identification of PCB's in the presence of organochlorine pesticides such as DDT and DDE has been difficult in the past because of their similar chromatographic characteristics (Risebrough et al., 1968). In PCB analysis today, the interference of organochlorine hydrocarbons is overcome by sequential column chromatography on Florsil and silica gel (Armour and Burke, 1970; FDA, 1971). Gas-liquid chromatography with highly sensitive and selective detectors has been employed successfully in their detection at low levels (Nebecker and Puglisi, 1971).

PCB compounds are slightly soluble in water; soluble in fats, oils, and organic solvents, and resistant to both heat and biological degradation (NTIS, 1972). Typically, the specific gravity, boiling point, and melting point of PCB's increase with their chlorine content. PCB's are relatively non-flammable

have useful cooling, insulating, and dielectric properties, and principally are used in the electrical industry in capacitors and transformers.

RATIONALE:

Exposure to PCB is known to cause skin lesions (Schwartz and Peck, 1943) and to increase liver enzyme activity that may have a secondary effect on reproductive processes (Risebrough, 1969; Street et al., 1968; Wasserman et al., 1970). It is not clear whether the effects are due to the PCB's or their contaminants, the chlorinated dibenzofurans, that are highly toxic (Bauer et al., 1961; Schultz, 1968; Varrett, 1970). While chlorinated dibenzofurans are a byproduct of PCB production, it is not known whether they are also produced by the degradation of PCB's (NAS, 1974).

Analyses of 40 fish from the 1970 National Pesticide Monitoring Program indicated only one fish to contain less than 1 ug/g PCB with the ten highest residues ranging from 19 ug/g to 213 ug/g whole body weight (Stalling and Mayer, 1972). Studies of the Milwaukee River revealed ambient water concentrations of 2.0 to 2.8 ug/l and residues in fish as high as 405 ug/g (Veith and Lee, 1971). Open water Lake Michigan concentrations have been reported to be less than 0.01 ug/l with mean residues in coho salmon of about 15 ug/g (Veith, 1973). The Food and Drug Administration guideline for protecting the health of human consumers of fish is 5 ug/g in tissue residues of fish. Based on Lake Michigan data, which indicate that at a concentration of 0.01 ug/l the fish tissue residues exceed a level found to be non-hazardous to man, a criterion of 0.001 ug/l in freshwater is warranted. Bluegill sunfish exposed to Aroclors 1248 and 1254, exhibited a bioaccumulation factor of 7.1×10^4 (Stalling and Mayer, 1972). The bioaccumulation factor for gizzard shad in the Saginaw River (Michigan)

varied between 0.6 x 10^5 and 1.5 x 10^5 for Aroclor 1254 (Michigan Water Res. Comm., 1973). A residue level of 2 ug/g in fish consumed by commercial ranch mink has been shown to prevent survival of offspring (Ringer et al., 1972). Reproduction was almost totally eliminated in ranch mink fed a beef diet containing 0.64 ug/g of Aroclor 1254 (Platonow and Kalstad, 1973). This suggests that a mink-food tissue level of not more than 0.5 ug/g would be required to protect the wildlife consumer.

Median PCB concentrations in whole fish of eight species from Long Island Sound obtained in 1970 were reported to be in the order of 1 ug/g, as were comparable concentrations off the coast of Southern California. (Hays and Risebrough, 1972; Risebrough, 1969). Generally, residues in ocean fish have been below 1 ug/g (Risebrough, 1970).

Surveys of Escambia Bay (Florida) during the period September 1969 to December 1971 have produced data on the pathways and effects of PCB's in the estuarine and marine environments. Although the major PCB source, accidental leakage from a PCB manufacturing plant has been terminated, residues continue to be observed in aquatic organisms of the bay (Duke, 1974). The sediment reservoir of Aroclor 1254 is thought to be a continuing source of PCB to biota. The initial survey of Escambia Bay biota revealed fish, shrimp, and crabs with levels as high as 12 ug/g. Higher levels were detected in higher trophic levels than shrimp, which could implicate a chain transfer from sediment to large animals.

From the Escambia Bay data, which include flow-through bioassays with residue analyses where possible, the following conclusions were reached: (1) all of the Aroclors tested are acutely toxic for certain estuarine organisms;

(2) bioassays lasting longer than 96 hours demonstrated that Aroclor 1254 is toxic to commercial shrimp at less than 1 ug/l; (3) fish, particularly sheepshead minnows, are extremely sensitive to Aroclor 1254 with 0.1 ug/l being lethal to fry; and, (4) acute toxicity of Aroclor 1016 to estuarine organisms is similar to the toxicity of other Aroclors but appears less toxic to fish in long-term exposures than does Aroclor 1254 (Duke, 1974).

Oysters were sensitive to Aroclor 1260 with growth diminished by 44 percent in 10 ug/l and by 52 percent in 100 ug/l. Approximately 10 percent of the pink shrimp died in 100 ug/l, but no apparent effects on pinfish were noted at 100 ug/l. Aroclor 1254 had no apparent effect on juvenile pinfish at 100 ug/l in 48-hour flow-through tests, but killed 100 percent of the pink shrimp. At 100 ug/l of Aroclor 1254 for 96 hours, shell growth of oysters was inhibited and decreased only 41 percent at levels of 10 ug/l. The toxicity of Aroclor 1248 and 1242 to shrimp and pinfish was similar to that of Aroclor 1254. Aroclor 1242 was toxic to oysters at 100 ug/l. Killfish exposed to 25 mg/l of Aroclor 1221 suffered an 85 percent mortality. In 96-hour bioassays, Aroclor 1016 was toxic to an estimated 50 percent of the oysters, brown shrimp, and grass shrimp at 10 ug/l; it was toxic to 18 percent of the pinfish at 100 ug/l (Duke, 1974).

Young oysters exposed to Aroclor 1254 in flowing sea water for 24 weeks experienced reduction in growth rates at 4.0 ug/l, but apparently were not affected by 1.0 ug/l. Oysters accumulated as much as 100,000 times the test-water concentration of 1.0 ug/l. General tissue alterations in the vesicular around the diverticula of the hepatopancreas were noted in the oysters exposed to 5.0 ug/l. No significant mortality was observed in oysters exposed continuously to 0.01 ug/l of Aroclor 1254 for 56 weeks (Duke, 1974).

Blue crabs apparently were not affected by 20 days' exposure to 5.0 ug/l of Aroclor 1254. Pink shrimp exposed under similar conditions experienced a 72 percent mortality. In subsequent flow-through bioassays, 51 percent of the juvenile shrimp were killed by Aroclor 1254 in 16 days and 50 percent of the adult shrimp were killed at 3.0 ug/l in 35 days. From pathological examinations of the exposed pink shrimp, it appears that Aroclor 1254 facilitates or enhances the expression of latent viral infections. Aroclor 1254 was lethal to grass shrimp at 4.0 ug/l in 16 days, to amphipods at 10 ug/l in 30 days, and to juvenile spot at 5.0 ug/l after 20 to 45 days. Sheephead minnows were the most sensitive estuarine organisms to Aroclor 1254 with 0.3 ug/l being lethal to the fry within 2 weeks. Aroclor 1016 in two different 42-day flow-through bioassays caused significant mortalities of pinfish at 32 ug/l and 21 ug/l. Pathological examination of those exposed to 32 ug/l revealed several liver and pancreatic alterations. Sheephead minnows in 28-day Aroclor 1016 flow-through bioassays were not affected by concentrations of 10 ug/l or less, but died at 32 and 100 ug/l (Duke, 1974). The bioaccumulation factors for the different flow-through bioassays are:

Aroclor	Organism	Time	Accumulation Facts (as a multiplier of test water concentrations)
1254	Oyster	30 days	1.01×10^5
1254	Blue Crab	20 days	4×10^3
1254	Grass Shrimp	7 days	3.2×10^3 to 11×10^3
1254	Spot	1 -28 days	37×10^3
1254	Pinfish	35 days	21.8×10^3
1016	Pinfish	42 days	11×10^3 to 24×10^3
1016	Sheephead Minnow		2.5×10^3 to 8.1×10^3

Based upon an accumulation factor of 100,000 in the oyster, it is necessary to limit the marine water concentration of PCB's to a maximum of 0.01 ug/l to protect the human consumer.

Evidence is accumulating that PCB's do not contribute to shell thinning of bird eggs (NAS, 1974). Dietary PCB's produced no shell thinning in eggs of Mallard ducks (Heath et al., 1972). PCB's may increase susceptibility to infectous agents such as viral diseases (Friend and Trainer, 1970), and increase the activity of liver enzymes that degrade steroids, including sex hormones (Risebrough et al., 1968; Street et al., 1968). Laboratory studies have indicated that PCB with its derivatives or metabolites, causes embryonic death of birds (Voss and Koeman, 1970).

REFERENCES CITED:

Armour, J.A., and J.A. Burke, 1970. Method for separating polychlorinated biphenyls from DDT and its analogs. Assoc. Official Anal. Chem. 53:761.

Bauer, H., et al., 1961. Occupational poisonings in the production of chlorophenol compounds. Arch. Gewerbepath 18:538.

Duke, T.W., 1974. Testimony in the matter of proposed toxic pollutant effluent standards for Aldrin-Dieldrin et al. FWPCA (307) Docket #1.

FDA, 1971. U.S. Dept. of H.E.W. Pesticide Analytical Manual, Vol. I and II.

Friend, H., and D.O. Trainer, 1970. Polychlorinated biphenyl interaction with duck hepatitis virus. Science, 170:1314.

Hays, H., and R.W. Risebrough, 1972. Pollutant concentrations in abnormal spring tern from Long Island Sound. AUK 89:19.

Heath, R.G. et al., 1972. Effects of polychlorinated biphenyls on birds. Proceedings IV International Ornithological Congress (In Press).

Michigan Water Res. Comm., 1973. Monitoring for polychlorinated biphenyls in the aquatic environment.

National Academy of Sciences, 1974. Water quality criteria, 1972. (In Press).

Nebeker, A.V., and F.A. Puglisi, 1971. PCB Newsletter, p. 19.

Nebeker, A.V., and F.A. Puglisi. 1974. Effect of polychlorinated biphenyls (PCB's) on survival and reproduction of Daphnia gammarus and Tanytorsus. Trans. Amer. Fish. Soc. (In Press).

National Technical Information Service, 1972. Polychlorinated biphenyls and the environment. NTIS Interdepartmental Task Force, Washington, D.C.

Platonow, N.S. and L.H. Kalstad, 1973. Dietary effects of polychlorinated biphenyls on mink. Can. Jour. Comp. Med. 37:391.

Ringer, R.K. et al., 1972. Effect of dietary polychlorinated biphenyls on growth and reproduction of mink. Amer. Chem. Soc. National Meeting Reprints of Papers, 12:149.

Risebrough, R.W., 1969. Chlorinated hydrocarbons in marine ecosystems. In: Chemical Fallout, M.W. Miller and G.G. Berg, eds., C.C. Thomas, Springfield.

Risebrough, R.W., 1970. More letters in the world. Environ. 12:16.

Risebrough, R.W. et al., 1968. Polychlorinated biphenyls in the global ecosystem. Nature, 220:1098.

Schulz, K.H., 1968. Clinical picture and etiology of chloracne. Arbeitsmed. Sozialmed. Arbeitshgy. 3:25. Reprinted in translation in U.S., Congress. Senate, Committee on Commerce, Effects of 2, 4, 5-T on man and the environment: Hearings, 91st Cong., 2nd sess., pp. 336-341.

Schwartz, L. and S.M. Peck, 1943. Occupational acne. New York State Med. 43:1711.

Stalling, D.L.,and F.L. Mayer, Jr., 1972. Toxicities of PCB's to fish and environmental residues. Env. Health Perspectives, 1:54.

Street, J.C. et al., 1968. Comparative effects of polychlorinated biphenyls and organochlorine pesticides in induction of hepatic microsomal enzymes. Amer. Chem. Soc., 158th National Meeting, Sept. 8-12, 1968.

Varrett, J., 1970. Statement in U.S. Congress, Senate, Committee on Commerce, Effects of 2, 4, 5-T on man and the environment: Hearings, 91st Cong., 2nd sess., pp. 190-360.

Veith, G.D., 1973. Chlorinated hydrocarbons in fish from lake fish. Final report, EPA Grant 1602 PBE.

Veith, G.D. and G.F. Lee, 1971. Chlorobiphenyls (PCB's) in the Milwaukee River. Water Res. 5:1107.

Voss, J.G. and J.H. Koeman, 1970. Comparative toxicoyogic study with polychlorinated biphenyls in chickens with special reference to porphyria, edema, formation, liver nebrosis, and tissue residues. Toxicol. Appl. Pharmacol. 17:656.

Wasserman, M. et al., 1970. The effect of organochlorine insecticides on serum colesterol level in people occupationally exposed. Bull. Environ. Contam. Toxicol. 5:368.