

BIO-MEDICAL RESEARCH
DOCUMENT DESCRIPTION FORM

Duplicate in all cards: →

63 68 69 76
1980 0000082
year as-1961- File number
[Right justify
[Numeric only]

77 78
VC
Sub-Index
Code

Author(s), as Last Name FS (No Punctuation) and
coden for journal as JAMA preceded by one blank space

1	20 21	40 41	60	61 62
DUCHATMAN, A	HIRSCHHORN, K	SOLZGROFF, I J		11
				12
				13

Title of Report; end with space-hyphen-hyphen-space. Follow with Index Terms,
separated from each other with comma-space. Avoid other punctuation;
do not abbreviate.

1 2	61 62
Vinyl Chloride Exposure and Human	21
Chromosome Aberrations - - Metagenesis, etc.	22
	23
	24

Source (Journal, Vol., Number, Pages, Date)

1 2	61 62
Mutation Research 31, 163-168 (1975)	31
	32

Brief Summary

1 2 10	61 62
SUMMARY:	61
	62
	63
	64

R&S 106681

0000082

Schwartz
cc Gehri
163 Koeb
Rosen*Mutation Research*, 31 (1975) 163-168

© Elsevier Scientific Publishing Company, Amsterdam—Printed in The Netherlands

VINYL CHLORIDE EXPOSURE AND HUMAN CHROMOSOME ABERRATIONS*

ALAN DUCATMAN, KURT HIRSCHHORN** AND IRVING J. SELIKOFF

Environmental Sciences Laboratory, Department of Community Medicine and the Division of Medical Genetics, Department of Pediatrics, Mount Sinai School of Medicine of the City University of New York, New York, N.Y. 10029 (U.S.A.)

(Received October 25th, 1974)

(Revision received January 15th, 1975)

SUMMARY

Examination of lymphocyte cultures from 11 vinyl chloride polymerization workers and 10 controls revealed a significantly higher incidence of aberrations in the exposed population. Most of the excess damage was of the "unstable" variety and involved the grossest kinds of changes such as fragments or rearrangements. When these complex changes were regarded as the product of two breaks, the incidence of all breaking events was also significantly increased in the workers. The results indicate the presence of chromosome damage in vinyl chloride exposed workers.

INTRODUCTION

Since the pioneering radiation studies of the early 1960's it has been clear that chromosome morphology examination could uncover evidence of genetic damage in man¹⁻³. The earliest of these studies discovered the now well-known link between carcinogenesis and chromosome changes (for recent review, see ref. 9). These were soon followed by the discovery that a number of chemicals are clastogenic (chromosome breaking), notably alkylating agents and cytostatic drugs, and DNA base analogues⁶. In recent years, occupational exposure to various chemicals has been recognized as another potential source of genetic change in man. Chromosome examination following human exposures had revealed the probable clastogenicity of some pesticides and herbicides^{10,17}. Industrial exposure to benzene has also been implicated by a number of investigators in both leukemogenesis and clastogenesis (refs. 7, 8, 16).

Within the last year, a new industrial carcinogen has been identified⁴. Vinyl chloride, the monomer used for production of the common polymer polyvinyl chloride, has been held responsible for at least 22 angiosarcomas of the liver in exposed wor-

* Supported by USPHS Center Grants ES 00928 and GM 19443, and Research Grant HD 02552.

** Kurt Hirschhorn, M. D. (to whom reprint requests should be addressed) is a Career Scientist of the Health Research Council of the City of New York, 1-513.

R&S 106682

kers¹⁰. There is additional suggestion from both animal and human data of an increased risk for cancers of the central nervous system, the respiratory system, and of the blood forming tissues^{10,12,15}. With an association between clastogenicity and carcinogenesis established for other agents, it appeared that this new carcinogen might also cause chromosome damage. We therefore undertook a blind study of lymphocyte chromosomes from vinyl chloride-exposed workers and nonexposed controls.

EXPOSURE TO VINYL CHLORINE

The 11 subjects studied were male workers who had received repeated exposure to vinyl chloride in an upstate New York polyvinyl chloride polymerization plant. Major exposures came from leaks of unreacted vinyl chloride gas, fumes from polyvinyl chloride slurry, and from polymerization reactor cleaning operations. Reactor cleaning involved skin contact to polyvinyl chloride and inhalation of vinyl chloride gas residues; this operation accounted for the most intense exposures. Duration of recurrent occupational exposure in the 11 men ranged from 4–28 years with an average of 15 years (Table I). There is no record of ambient gas levels in the factory, but it is assumed that these must have exceeded 500 ppm at times, based on reports of odor detection, dizziness, and headaches.

TABLE I

AGE AND DEGREE OF VINYL CHLORIDE EXPOSURE OF SELECTED CASES AND CONTROLS

Case No.	Age	Years exposed	Cells examined	Control No.	Age	Years exposed	Cells examined
1	61	23	50	1	43	0	50
2	50	28	50	2	30	0	50
3	47	19	50	3	29	0	50
4	44	11	50	4	29	0	50
5	40	13	50	5	28	0	50
6	39	17	50	6	28	0	50
7	36	17	50	7	27	0	50
8	36	12	50	8	20	0	50
9	33	11	50	9	19	0	50
10	32	10	50	10	18	0	50
11	25	4	50				
Total	443	165	550	271	0	500	
Average	40	15	50	27	0	50	

Of the 10 healthy male controls, 4 were from within the same factory population, without known vinyl chloride exposure. Nevertheless, as long-term employment implied the possibility of some vinyl chloride exposure, albeit at low levels, we also selected 6 older controls from outside the factory environment. The average age of the controls was 27, of the subjects 40.

METHODS

The chromosome studies were performed on cultures of peripheral blood lymphocytes incubated at 37° for 65–68 h with phytohemagglutinin (Wellcome). Harvests were performed according to micromethod modifications of procedures first described

by MOORHEAD *et al.* in order to insure

In each i microscope at 1 parately so far a photographed; 1 cells examined t photographs.

Confirmed discern if a patt COURT BROWN^a such as breaks fragments, dicen monosomies, tris stable and unsta disappear from t

The system all breakage, incl

TABLE II

VINYL CHLORIDE EX BROWN^a AND BUCK

Case No.	B
1	7
2	8
3	6
4	3
5	5
6	3
7	5
8	8
9	4
10	8
11	5
Total	62
Mean($\pm$ S.D.)	5.6
Comparison	1
	0.

Control No.

1	5
2	4
3	4
4	5
5	2
6	5
7	5
8	5
9	4
10	5
Total	44.0
Mean($\pm$ S.D.)	4.4

^a B, Breaks and gaps; (monosomy, trisomy,

human data of an in-
ratory system, and of
n clastogenicity and
this new carcinogen
ok a blind study of
ers and nonexposed

ed repeated exposure
polymerization plant
as, fumes from poly-
operations. Reactor
tion of vinyl chloride
posures. Duration of
4-28 years with an
levels in the factory,
es, based on reports

by MOORHEAD *et al.*¹⁴. Media used in this study came from a single batch (Gibco) in order to insure uniform pH and other culture conditions.

In each individual studied, 50 metaphases were counted directly under the microscope at 1600 $\times$. The A, B, D, E, F, G, and Y chromosomes were evaluated separately so far as possible. Suspected aberrations and some normal metaphases were photographed; karyotypes were performed where helpful. From the total of 1050 cells examined there were 281 photographs taken and 140 karyotypes made from the photographs.

Confirmed aberrations were classified according to two systems in order to discern if a pattern of breakage might exist. The system of BUCKTON *et al.*³ and COURT BROWN⁵ has three categories of aberrations: *B*, cells with simple aberrations such as breaks and gaps; *C_u*, cells with "unstable" chromosome changes such as fragments, dicentrics, and rings; *C_s*, cells with "stable" chromosome changes such as monosomies, trisomies, deletions, and exchanges. The distinction between cells with stable and unstable aberrations is related to their tendency to either remain in or disappear from the circulation.

The system of HIRSCHHORN AND COHEN¹¹ differs in two respects. It considers all breakage, including the total number of breaks in cells with more than one aber-

TABLE II
VINYL CHLORIDE EXPOSURE AND CHROMOSOME ABERRATIONS (CLASSIFICATION SYSTEM OF COURT
BROWN³ AND BUCKTON *et al.*³)

Case No.	B^A	C_u^A	C_g^A
1	7	2	3
2	3	2	3
3	6	0	3
4	3	1	3
5	5	1	3
6	3	0	0
7	5	2	2
8	8	2	6
9	4	0	4
10	8	4	5
11	5	3	5
Total	62.0	17.0	37.0
Mean ($\pm$ S.D.)	5.64 ($\pm$ 1.91)	1.55 ($\pm$ 1.29)	3.36 ($\pm$ 1.63)
Comparison	$t = 1.84$ $0.1 > p > 0.05$	$t = 2.863$ $p = 0.01$	$t = 0.56$ $0.6 > p > 0.5$

<i>Control No.</i>			
1	5	1	3
2	4	0	7
3	4	0	1
4	5	0	3
5	2	1	6
6	5	0	3
7	5	0	3
8	5	1	2
9	4	0	0
10	5	0	1
Total	44.0	3.0	29.0
Mean ($\pm$ S.D.)	4.40 ($\pm$ 0.97)	0.30 ($\pm$ 0.48)	2.90 ($\pm$ 2.18)

^a B, Breaks and gaps; C_u, "unstable" changes (fragments, dicentrics, rings); C_s, "stable" changes (monosomy, trisomy, deletions, exchanges).

factory population, low employment levels, we also have an average age of the

peripheral blood lymphocytes (Wellcome). Harvests were first described

ration, and it gives weighted consideration to those aberrations which are the apparent result of two "hits". Therefore, complex breakage (C) such as rings, dicentrics, and exchanges is counted twice in the total, whereas simple breaks and deletions (S) are counted once.

RESULTS

Gaps and breaks were the predominant aberrations among the 1050 cells examined, as seen in the B column of Table II. Subjects have nonsignificant increases of such aberrations when compared to controls by a *t*-test for comparison of the means ($0.1 > P > 0.05$), and marginally significant increases by an *F*-ratio for comparison of the variances ($0.05 > P > 0.01$). The difference in cells with stable aberrations, or those that BUCKTON *et al.* and COURT BROWN found to persist in circulating lymphocytes^{3,5}, is also nonsignificant. In our study, stable aberrations were usually cells with random chromosome loss. However, cells with unstable aberrations were observed significantly more frequently in the cultures from exposed workers: *t*, $P = 0.01$; *F*, $0.01 > P > 0.001$.

Table III focuses on breaking events only. The total of simple breaks, including

TABLE-III

CHROMOSOME BREAK EVENTS (IN 50 CELLS/INDIVIDUAL) AND VINYL CHLORIDE EXPOSURE (SYSTEM OF HIRSCHHORN AND COHEN)

Case No.	S ^a	C ^a	Total break events (S+2C)
1	4	3	10
2	2	3	8
3	2	0	2
4	1	2	5
5	0	1	2
6	0	0	0
7	3	2	7
8	1	2	5
9	4	0	4
10	1	4	9
11	3	5	13
Total	21	22	65
Mean(±S.D.)	1.91(±1.44)	2.00(±1.67)	5.91(±3.91)
Comparison	<i>t</i> = 1.12 0.3 > <i>p</i> > 0.2	<i>t</i> = 2.80 0.02 > <i>p</i> > 0.01	<i>t</i> = 2.75 0.02 > <i>p</i> > 0.01
Control No.			
1	3	1	5
2	1	0	1
3	1	0	1
4	0	0	0
5	2	2	6
6	2	0	2
7	0	0	0
8	1	1	3
9	2	0	2
10	1	0	1
Total	13	4	21
Mean(±S.D.)	1.30(±0.95)	0.40(±0.70)	2.10(±2.02)

^a S, Single hit events (breaks, deletions); C, complex events (rings, dicentrics, exchanges).

those from multial is shown in the S column in those exposed, breaking events (t , $0.02 > P > 0.01$ total of all breaks, in these with t , 0.0

Exaggerated because of their high 3.91 (±2.47 S.D.) per individual. Significant.

DISCUSSION

There are other and it would clear experimental evidence than chromosome loss as well as subject. Within these limits vinyl chloride is a variety as determined most evident for HIRSCHHORN AND damage, shows the crease in all breaki

In this small with either the de former may never be is only a crude guess exhibited increased showed damage sin

Stating that mosomes leaves us all predictive of environmental vinyl chloride exposure with induced neoplasia studies, we will know important role to play

The second question: vinyl chloride? We feel such unwarranted. The conclusions findings in any single. However, they may exposure to unreacted experimental findings

high are the ap-
rings, dicentrics,
and deletions (S)

the 1050 cells
significant increases
comparison of the
F-ratio for com-
with stable aber-
sist in circulating
ons were usually
aberrations were
exposed workers:

breaks, including

EXPOSURE (SYSTEM

exchanges).

those from multiaberrant cells, is increased but not significantly in the subjects. This is shown in the S column. Complex breaking events in C are significantly more frequent in those exposed, with: t , $0.02 > P > 0.01$; and F , $0.01 > P > 0.001$. The total of breaking events ($S + 2C$) shows a similarly significant increase in the subjects (t , $0.02 > P > 0.01$; F , $0.01 > P > 0.001$). Not included in the charts is a combined total of all breaks, gaps, and deletions. There was a marginally significant difference in these with t , $0.05 > P > 0.02$; F was not significant.

Exaggerated secondary constrictions of the No. 9 chromosome were easily noted because of their high degree of visibility. The average subject was observed to have $3.91 (\pm 2.47 \text{ S.D.})$ and the average control had $2.00 (\pm 1.63)$ in the 50 cells examined per individual. Significance was marginal at most, $P < 0.05$ by a t -test; F not significant.

DISCUSSION

There are obvious perils in drawing strong conclusions from small samples, and it would clearly be preferable to have an age-matched control group despite experimental evidence that age is generally unrelated to chromosome changes other than chromosome loss⁵. Also, the relatively high degree of breaks and gaps in controls as well as subjects is somewhat disconcerting, although subjects do have more. Within these limitations, it is clearly indicated that chronic high level exposure to vinyl chloride is clastogenic. Much of the increased damage was of the unstable variety as defined by BUCKTON *et al.*³. The difference between cases and controls is most evident for unstable aberrations in general and fragments in particular. HIRSCHHORN AND COHEN's system, which is a better overall index of chromosome damage, shows that long-term exposure is associated with an evidently significant increase in all breaking events.

In this small sample it has not been possible to correlate the degree of damage with either the degree of exposure or with vinyl chloride disease symptoms. The former may never be possible, as the best estimate of total exposure for any individual is only a crude guess. In general, the majority of subjects selected from this factory exhibited increased breakage rates, and those with the shortest duration of exposure showed damage similar to those with the longest duration.

Stating that chronic vinyl chloride exposure almost certainly damages chromosomes leaves us with two questions. First, can chromosome damage studies be at all predictive of environmentally induced carcinogenesis? Ionizing radiation and now vinyl chloride exposures have been studied for genetic properties *after* association with induced neoplasms. If we can reverse the order of events, perhaps in animal studies, we will know with more certainty if chromosome examination has an important role to play in predicting environmental carcinogenesis.

The second question is: what kind of genetic studies should be done with vinyl chloride? We feel strongly that chromosome study of individual concerned fathers is unwarranted. The degree of damage discovered here is unlikely to yield meaningful findings in any single individual. Women are not employed in polymerization work. However, they may work in polyvinyl chloride processing industries, in which some exposure to unreacted vinyl chloride may occur. This could be important in light of experimental findings of transplacental carcinogenesis¹².

Carefully controlled examination of larger groups with vinyl chloride and polyvinyl chloride exposure are obviously needed, first to provide the larger data base necessary to confirm clastogenicity for vinyl chloride exposure, and also to evaluate dose-response relationships. Mutagenicity is being assessed in other test systems, including bacterial studies, insect and animal studies, along with *in vitro* chromosome studies.

The case for studying other genetically suspect chemicals is now stronger than ever.

REFERENCES

- 1 BENDER, M. A., AND P. C. GOOCH, Persistent chromosome aberrations in irradiated human subjects, *Radiat. Res.*, 16 (1962) 44-53.
- 2 BLOOM, A. D., AND J. H. TJIO, *In vivo* effects of diagnostic X-irradiation on human chromosomes, *New Engl. J. Med.*, 270 (1964) 1341-1344.
- 3 BUCKTON, K. E., P. A. JACOBS, W. M. COURT BROWN AND R. DOLL, A study of the chromosome damage persisting after X-ray therapy for ankylosing spondylitis, *Lancet*, ii (1962) 676-682.
- 4 CREECH, J. L., AND M. N. JOHNSON, Angiosarcoma of liver in the manufacture of vinyl chloride, *J. Occup. Med.*, 16 (1974) 150-151.
- 5 COURT BROWN, W. M., Human population cytogenetics, in A. NEUBERGER AND E. L. TATUM (Eds.), *Frontiers of Biology*, Vol. V, North-Holland, Amsterdam, 1967, pp. 1-31.
- 6 EVANS, H. J., Population cytogenetics and environmental factors, in PATRICIA A. JACOBS, W. H. PRICE AND PAMELA LAW (Eds.), *Human Population Cytogenetics*, Williams and Wilkins, Baltimore, 1970, pp. 191-216.
- 7 FORNI, A. M., A. COPPELLINI, E. PACIFICO AND E. C. VIGLIANI, Chromosome changes and their evolution in subjects with past exposure to benzene, *Arch. Environ. Health*, 23 (1971) 385-391.
- 8 FORNI, A. M., E. PACIFICO AND A. LIMONTA, Chromosome studies in workers exposed to benzene or toluene or both, *Arch. Environ. Health*, 22 (1971) 373-378.
- 9 GERMAN, J. (Ed.), *Chromosomes and Cancer*, Wiley, New York, 1974.
- 10 HEATH, C., AND J. WAGONER, *Report of a Working Group on Vinyl Chloride*, Lyon, 24-25 June, 1974, IARC No. 74/005, World Health Organization, pp. 18-19.
- 11 HIRSCHHORN, K., AND M. M. COHEN, Drug-induced chromosomal aberrations, *Ann. N.Y. Acad. Sci.*, 151 (1968) 955-987.
- 12 HOOPINGAMER, R., AND A. W. BLOOMER, Lymphocyte chromosome analysis of pesticide exposed individuals, in *7th Int. Congr. Plant Protection, Paris, 1970*, p. 772.
- 13 MALTONI, C., AND G. LEFEMINE, Carcinogenicity bioassays of vinyl chloride, I. Research plan and early results, *Environ. Res.*, 7 (1974) 387-405.
- 14 MOORHEAD, P. S., P. C. NOWELL, W. J. MELLMAN, D. M. BATTIPS AND D. A. HUNGERFORD, Chromosome preparations of leukocytes cultured from human peripheral blood, *Exp. Cell Res.*, 20 (1960) 613-616.
- 15 NICHOLSON, W. J., E. C. HAMMOND, H. SEIDMAN AND I. J. SELIKOFF, Mortality experience of a cohort of vinyl chloride-polyvinyl chloride workers, *Ann. N.Y. Acad. Sci.*, 255 (1975) 225-230.
- 16 TOUGH, I. M., AND W. M. COURT BROWN, Chromosome aberrations and exposure to ambient benzene, *Lancet*, i (1965) 684.
- 17 YODER, J., M. WATSON AND W. W. BENSON, Lymphocyte chromosome analysis of agricultural workers during extensive occupational exposure, *Mutation Res.*, 21 (1973) 335-340.