

From the desk of
H. C. GODT, JR.

JUL 29 1983

7/29/83

George!

As requested. Don't
forget, I want copies
of your slides from
the Dioxin symposium.

C03732

CONFIDENTIAL
SUBJECT TO PROTECTIVE ORDER.

HISTORICAL OVERVIEW: 2,4,5-T, Chloracne & 2,3,7,8-TCDD

- 1930s -- Chloracne, a dermatitis characterized by blackheads similar to teen-age acne but often more severe, was noted among workers dealing with chlorinated aromatic hydrocarbons (chloracnegens).
- 1941 -- 2,4-D and 2,4,5-T first synthesized.
- 1944 -- 2,4-D and 2,4,5-T reported as herbicides and marketed by Amchem Products.
- 1947 -- Effectiveness of 2,4,5-T and derivatives against woody plants resistant to 2,4-D established by Dow.
- 1948 -- 2,4,5-T registered for control of weeds on rangelands, fence rows and rights of way.
- 1949 -- Explosion in Monsanto's 2,4,5-trichlorophenol (TCP) unit at Nitro, W.Va. exposed 250 workers to reaction mass causing 122 cases of chloracne, some very severe. TCP is intermediate in 2,4,5-T manufacture.
- 1950 -- Presence of a "highly toxic impurity" in production of TCP recognized by Dow.
- 1957 -- 2,3,7,8-TCDD first implicated as causative agent of chloracne associated with TCP production. (Kimmig and Schultz, Germany)
- 1962 -- U.S. military began spraying 2,4,5-T formulations in Vietnam to take away foliage cover from the enemy.

CONFIDENTIAL

SUBJECT TO PROTECTIVE ORDER.

HISTORICAL OVERVIEW (CONT'D.)


- 1965 -- 2,3,7,8-TCDD recognized by Dow as causative agent of chloracne in TCP wastes. Alerted health authorities and other U.S. 2,4,5-T manufacturers. Shared analytical techniques.
- 1970 -- 2,3,7,8-TCDD's teratogenicity and fetotoxicity first reported in animals. U.S. military stopped spraying Agent Orange in Vietnam. U.S. Dept. of Agriculture suspended uses of 2,4,5-T.
- 1973 -- Vietnamese study linked liver cancer, abortions, and birth defects to Agent Orange spraying.
- 1976 -- Explosion at ICMESA chemical plant in Seveso, Italy released heavy concentration of 2,3,7,8-TCDD in densely populated area. Workers (156) and residents (37,000) exposed. More than 500 treated for presumed toxic symptoms; 134 cases of chloracne. Mortality rate normal to date. Event received worldwide publicity through news media.
- 1978 -- Dow announced chlorinated dioxins found in particulate matter in air emissions from various commercial and domestic combustion processes.
- 1979 -- Suit filed by Vietnam veterans against five U.S. chemical companies that produced Agent Orange. Companies, in turn, filed third party action against U.S. government for its negligent misuse of the chemical.
- 1982 -- Times Beach

CONFIDENTIAL

SUBJECT TO PROTECTIVE ORDER.

DIOXIN


Chemical Nomenclature: Dioxin = Dibenzo-p-dioxin




Media: Dioxin = 2,3,7,8-Tetrachlorodibenzo-p-dioxin
= 2,3,7,8-TCDD

CONFIDENTIAL
SUBJECT TO PROTECTIVE ORDER.


CHLORINATED DIOXINS (CDDs)



- 75 CDDs
- 22 TCDDs
- 2,3,7,8-TCDD - most toxic isomer to the male guinea pig




- CCDs 10 to 20 times more toxic than chlorinated dibenzofurans in the rabbit ear test



CONFIDENTIAL
SUBJECT TO PROTECTIVE ORDER.

STRUCTURE AND PHYSICAL PROPERTIES
OF 2,3,7,8-TCDD



Physical State:	White, crystalline solid
Molecular Weight:	322
Melting Point, °C.:	303-305
Decomposition Point, °C.:	980-1000
Solubility:	Water = 2×10^{-8} g./L Benzene = 0.57 g./L <u>o</u> -Dichlorobenzene = 1.4 g./L

CONFIDENTIAL
SUBJECT TO PROTECTIVE ORDER.


SOURCES AND FORMATION
OF DIOXINS

- Formed during manufacture of chlorophenols, intermediates in synthesis of pesticides. Thus contaminants in chlorophenol derived pesticides.
- Pyrolysis (<750°C) of chlorophenols or chlorophenol derived pesticides.
- Formed in a wide variety of commercial and domestic combustion processes.


CONFIDENTIAL
SUBJECT TO PROTECTIVE ORDER.

REQUIREMENTS FOR DIOXIN FORMATION

- Precursor must contain an o-substituted benzene ring with one of the substituents being an oxygen atom attached directly to the ring.



- The two substituents must be able to react with each other to form another compound.




- The reaction is favored by basic conditions (e.g., sodium hydroxide) and temperatures of 180-400°C.
- Presence of a metal catalyst (e.g., copper or iron powder) promotes the reaction.


CONFIDENTIAL
SUBJECT TO PROTECTIVE ORDER.

MANUFACTURE OF 2,4,5-T


Step 1




Step 2



Step 3



Step 4




CONFIDENTIAL (2,4,5-Trichlorophenoxy-
acetic acid)

SUBJECT TO PROTECTIVE ORDER.

FORMATION OF 2,3,7,8-TCDD



AS SIDE REACTION IN

MANUFACTURE OF NaTCP




CONFIDENTIAL
SUBJECT TO PROTECTIVE ORDER.

MANUFACTURE OF
PENTA AND SANTOBRITE



CONFIDENTIAL
SUBJECT TO PROTECTIVE ORDER.

FORMATION OF
CHLORINATED DIBENZO-p-DIOXINS AND
CHLORINATED DIBENZOFURANS
AS SIDE REACTIONS IN
MANUFACTURE OF PENTA




CONFIDENTIAL
SUBJECT TO PROTECTIVE ORDER.

METHODS OF DESTRUCTION
OF CHLORINATED DIOXINS

- Incineration at temperatures of 1000 to 1500° in industrial incinerators. Future: molten salt combustion or microwave plasma destruction.
- Degraded quickly by sunlight or artificial ultraviolet light (photolysis).
- Chemical destruction by ozonolysis, chlorinolysis, catalytic wet oxidation (Zimpro process), and catalytic dechlorination processes.
- Biological destruction by new genetic engineered bacteria may hold future promise.




CONFIDENTIAL
SUBJECT TO PROTECTIVE ORDER.

CHEMISTRY OF INCINERATION OF 2,3,7,8-TCDD



CONFIDENTIAL
SUBJECT TO PROTECTIVE ORDER.

CHEMISTRY OF PHOTOLYSIS OF 2,3,7,8-TCDD



CONFIDENTIAL
SUBJECT TO PROTECTIVE ORDER.