

PLAINTIFF'S EXHIBIT

IHF-397

88

I H F

1940 through 1955

#68-11451

A.S. Alden and W.M. Howell

The asbestos corn.

TITLE

ORIG. REF.

VOL.

PP.

DATE

ASSIGN. REF.

VOL.

PP.

DATE

Arch. Dermat. Syph

49

312

May '44

JUL 1944

26

183

NOV. '44

ABSTRACT

The Asbestos Corn. H.S. Alden and W.M. Howell. Arch. Dermat. and Syph., 49, 312 (May, 1944)

The author describes corns or warts peculiar to asbestos workers, first noticed by Derwitz in 1930. They have no relation to asbestos of the lungs. One characteristic is that they appear and feel as if a foreign body were present, none is ever found. It is probable, however, that one is present in the very early stages.

DATE
EXAMINED

BY

FFR

PURPOSE
OF SEARCH

ABSTRACT BLANK FOR GENERAL USE

CONT'D

MELLON INSTITUTE, PITTSBURGH, PA.

FORM 77

MELLON INSTITUTE OF INDUSTRIAL RESEARCH
PITTSBURGH, PA.

CLASSIFIED SUBJECT

#1239-12/45

NUMBER
FC

EX 752

AUTHOR: R.B. Fisher, R.L. Thorne & Van Cott
AUTHOR'S POSITION
AND ADDRESS:

TITLE: Pallgorskite, a possible asbestos substitute.

PUB. DESIGNATION: U.S. Bur. of Mines
AND SERIAL NO.: Inf. Circ. No. 7313
PLACE PUBLISHED:
AND PUBLISHER:

EDITION: DATE: Mar. '45 VOL. LIST PRICE REVIEWED IN: VOL. PAGE DATE

PAGES: 5	GRAPHIC: ILLUSTRATIONS: MAPS: COST DATA: INDEX: SUMMARY:	GENERAL: REVIEW: POPULAR: ENGINEERING: ORIGINAL: AVAILABLE AT:	THEORETICAL: MATHEMATICAL: EXPERIMENTAL: STATISTICAL:	ECONOMIC: ADMINISTRATIVE: PROPAGANDA:	EXCELLENT: GOOD: FAIR: USELESS: FOR PURPOSE:
----------	--	--	---	---------------------------------------	--

ABSTRACT: Pallgorskite is probably a variety of asbestos usually termed mountain leather. A deposit of pallgorskite has been found on an Alaskan island, and it may find a number of uses if present in sufficient quantity.

DATE EXAMINED: AT: PIR PURPOSE OF SEARCH: CONF'D
ABSTRACT BLANK FOR BOOKS, BULLETINS, ADVANCE SHEETS, ETC.

MELLON INSTITUTE OF INDUSTRIAL RESEARCH
PITTSBURGH, PA.

CLASSIFIED SUBJECT		AbB. 186 Mar'40		754		NUMBER		752	
AUTHOR		H. J. Skind							
TITLE		Clinical studies in anesthesias.							
ORIG. REFS.	VOL.	PP.	DATE	ANAL. REFS.	VOL.	PP.	DATE		
AbB. 186	41	12-21	Jun'40	J I H	22	57	Mar'40		
ABSTRACT		See Litv's, 1940 Abstract Bulletin							
DATE EXAMINED	BY	PURPOSE OF SEARCH		ABSTRACT BLANK FOR GENERAL USE					
				CONT'D.					

MELLON INSTITUTE OF INDUSTRIAL RESEARCH
PITTSBURGH, PA.

May 41 20.7.344

752

NUMBER

752

AUTHOR: Enrico C. Vicland
AUTHOR'S POSITION:
AND ADDRESS:

TITLE: Study of Asbestosis in the Manufacture of Asbestos.

PUB. DESIGNATION		AND SERIAL NO.		PLACE PUBLISHED		DATE PUBLISHED		EDITION		VOL.		PAGE		DATE	
Entg. nez. d. Propaganda per la JIH		Yor. 1940		Yor. 1940		Yor. 1940		Yor. 1940		Yor. 1940		Yor. 1940		Yor. 1940	
PAGES: 74	DISCUSSION (P.P.):	REFERENCES:	CONCLUSIONS:	INDEX:	GRAPH:	ILLUSTRATIONS:	MAPS:	COST DATA:	INDEX:	GENERAL:	REVIEW:	POPULAR:	ENGINEERING:	ORIGINAL:	AVAILABLE:

ABSTRACT: (See May Abstract Bulletin)

DATE RECEIVED:	BY:	PURPOSE OF SPACING:	CONFIDENTIALITY:

ABSTRACT BLANK FOR BOOKS, BULLETINS, ADVERTISING BOOKS, ETC.

MELLON INSTITUTE, PITTSBURGH, PA.

CLASSIFIED SUBJECT
#407-4/47

FELLOWSHIP
NUMBER
NFC

FILE NUMBER

752

AUTHOR: J.H. Skavlem and R.J. Ritterhoff

TITLE: Coexistent pulmonary asbestosis and sarcoidosis.

ORIG. REFS.	VOL.	PP.	DATE	ABSTR. REFS.	VOL.	PP.	DATE
Am.J. Path.	22	493-517	May '46	JHK&T	29	29	Mar. '47

ABSTRACT:

This is a necropsy report of a 42-year old man who died a respiratory death after 11 months of increasing severe exertional dyspnea. Previously the patient had worked for 25 years in an asbestos plant. The pathological study revealed a coexistent asbestosis and sarcoidosis of both lungs.
— J.I.H.&T.

DATE
EXAMINED:

By: FFRJJH

PURPOSE
OF SEARCH:

ABSTRACT BLANK FOR GENERAL USE

CONT'D

7353-446

FELLOWSHIP
NUMBER
FC
752

FILE NUMBER

AUTHOR: H.E. Fleischer, F.J. Viles, Jr., R.L. Gade and Philip Drinker
TITLE: A health survey of pipe covering operations in constructing naval vessels.

ORIG. REF. JHET VOL. 28 PP. 9-16 DATE Jan. '46

ABSTRACT: ABSTR. REF. VOL. PP. DATE

(1) The character of asbestos pipe covering industry on board naval vessels is such that conclusions drawn from other asbestos industries such as textiles, cannot be applied. (2) The operations of bond saw cutting, grinding, cement mixing, and installation on board ship should be equipped with exhaust ventilation to keep the total dust concentration low. (3) The incidence of asbestosis among pipe coverers in the shipyards studied was low, 0.29 per cent or three cases out of 1074. In view of the nature of shipyard pipe covering work, this low incidence is not surprising. (4) Since each of the three cases of asbestosis had worked at asbestos pipe covering in shipyards for more than 20 years, it may be concluded that such pipe covering is not a dangerous occupation — Authors' conclusions.

DATE RECEIVED: Authors' conclusions PURPOSE OF SEARCH: CONT'D

ABSTRACT BLANK FOR GENERAL USE

MILLON INSTITUTE, PITTSBURGH, PA.

AUTHOR: Luton, Champelx and Faure
TITLE: Asbestosis of the lungs.

ORIG. REFS.: Arch. d. mal. profess. 7 VOL. 299 PP. 1946 DATE 1946
ABSTR. REFS.: Occ. Med. VOL. 4 PP. 234 DATE Aug. 1947

ABSTRACT

A review of the literature reveals no previous cases of asbestosis in France, but the authors have seen several cases in a factory in which asbestos is used, and in which they make regular examinations. Details are given in one case of a man of 64, exposed to concentrations of 48 to 62 mg. of dust per cubic meter, total silica 40%, and free silica 4.3 to 14.6%. Complaints of weakness, dyspnea, coughing, and loss of weight were noted in 1943. Thoracic expansion was 2 cm. In the sputum asbestosis bodies were found but no tubercle bacilli. Roentgenography showed much more fibrosis than usual in the lower third of the lungs. During the next two years the symptoms became more severe.

-- From Occ. Med.

FORM 77

DATE EXAMINED:	BY:	PURPOSE OF SEARCH:	CONT'D.
	FR		

ABSTRACT BLANK FOR GENERAL USE

MELLON INSTITUTE, PITTSBURGH, PA.

CLASSIFIED SUBJECT
#234-2/47

FELLOWSHIP
NUMBER
MFC
FILE NUMBER
752

AUTHOR: E.J. King, J.M. Clegg and V.M. Roe

TITLE: Effect of asbestos, and of asbestos and aluminum, on lungs of rabbits.

ORIG. REFS.	VOL.	PP.	DATE	ABSTR. REFS.	VOL.	PP.	DATE
Thorax	1	188-197	Sept. 46	JAMA	153	808	Jan. 18, 1947

Abstract:

The purpose of the experiments described by King and his associates was to test whether short fiber asbestos is less damaging in the lungs of the rabbit than is long fiber asbestos and whether metallic aluminum will suppress any toxic effect produced by fiber asbestos. Four groups of animals were tested. All groups showed a foreign body reaction in the lungs. In animals in the long fiber experiments there developed a nodular reticulosis comparable with the experimental silicotic nodule. In animals in the short fiber experiment there developed a diffuse interstitial reticulosis. The addition of aluminum did not afford any benefit in any group. — J.A.M.A.

DATE EXAMINED: ST. FR:JAMA
PURPOSE OF SEARCH:
ABSTRACT BLANK FOR GENERAL USE

CONT'D

100 6-14
FORM 77

AUTHOR: L. Moro

TITLE: Histology of asbestosis.

ORIG. REFS.

VOL.

PP.

DATE

ABSTN. REFS.

VOL.

PP.

DATE

Acta Pathol. & Micro- 23 63-59 1946
biol. (Scand.)

ABSTRACT:

Two workers in an asbestos factory died under circumstances which indicated that the cause of death was their trade. When the workers of this factory were examined roentgenologically asbestosis was found in 65 per cent of the 167 workers examined. The microscopic picture of asbestosis is characterized by a reaction caused by the needles in the tissue, which becomes manifest in exudate cell infiltration, accompanied by numerous giant cells containing foreign particles and an increase of interstitial connective tissue and fibrosis. Moro is inclined to think that in the pathogenesis of asbestosis the mechanical factors are the most important, although the chemical factors may have some significance in a later phase. — J.A.M.A.

MELLON INSTITUTE, PITTSBURGH, PA.

DATE
EXAMINED:

BY:

PURPOSE
OF SEARCH:

ABSTRACT BLANK FOR GENERAL USE

CONT'D:

MELLON INSTITUTE, PITTSBURGH, PA.

CLASSIFIED SUBJECT
#227 - 3/10/47

FILE NUMBER
FC

FILE NUMBER
752

AUTHOR: Anon. Safety Rev. (U.S. Navy)

TITLE: Asbestos dust.

ORIG. REF.: Safety Rev.

VOL. 4

DATE Jan. '47

VOL. 1

DATE

ABSTRACT:

The brief article calls attention to the hazard of exposure to asbestos dust and the need of adequate localized ventilation. All working places with possible exposure should be checked for asbestos dust, and where mechanical exhaust ventilation must be supplemented by wearing of personal protective equipment, it should conform to Bureau of Mines specifications.

DATE EXAMINED:

BY: FFR

PURPOSE OF SEARCH:

ABSTRACT BLANK FOR GENERAL USE

CONF'D.

MELLON INSTITUTE, PITTSBURGH, PA.

AUTHOR: V. Wyss.

NOTE: The presence of asbestos fibers in urine of workers exposed to asbestos hazard.

ORIG. REF.:

VOL.

PP.

DATE

ASSIN. REF.

VOL.

PP.

DATE

Rass. med. Ind.
(Italian)

22

55-56

1953

Blol. Absts.

28

1580 July '54

ABSTRACT:

In urinary sediment of two asbestos workers asbestos fibers were seen; the same finding was detected in urine of two female patients suffering from asbestosis, who had left their work seven years before.

-- Blol. Absts.

DATE
RECORDED

BY

PURPOSE
OF SEARCH

ABSTRACT BLANK FOR GENERAL USE

CONT'D.

J. Champeix and J. Boulay (1996)

Electron Microscopy of Asbestos-like Bodies and the Fibers of Asbestos.

THIS MONTH	VOL.	PP.	DATE	ALBION NEWS	VOL.	PP.	DATE
Arch. Neale's Profess (Fr.)	11	607	1950	Brit. J. Ind. Med.	8	198	7/51

Asbestos bodies recovered from sputum have a central fiber of asbestos coated with a colloidal gel, which is probably protein. Removal of the enveloping gel by micromanipulation reveals some transparency of the edges of the asbestos fiber, believed to result from solution. Fibers shorter than 10 microns are not found in the bodies and it is suggested that the smaller fibers dissolve completely. On this evidence the authors suggest that pulmonary fibrosis may result from solution of asbestos rather than from some physical irritation. -- Brit. J. Ind. Med.

DATE	PURPOSE
10/1/57	OF BEACH
10/1/57	ABSTRACT PLANN FOR GENERAL USE

MILLON INSTITUTE, PITTSBURGH, PA.

CLASSIFIED SUBJECT

218 2/52

FELLOWSHIP
NUMBER
NFC 752

FILE NUMBER

AUTHOR: P. Lepine and O. Croissant

TITLE: A morphological study with the electron microscope of particles of asbestos found in the air.

ORIG. REF.: Ann. Inst. Pasteur 80 305-8 3/51 Bull. Hyg. 26 1000 10/51

Six samples of airborne dust from workplaces in Quebec where asbestos is used were examined with the electron microscope after removal of all but the most minute particles. The dust was found to contain crystalline needles, smaller in size but similar to those seen in an ordinary microscope; these needles comprised 65 to 75% of the particles. The remainder consisted of amorphous particles, mostly of serpentine. The needles had a diameter of 0.028 micron; they occurred in bundles about 0.85 micron or more in diameter. Such particles could readily gain access to the alveoli and undergo phagocytosis, with formation of the tissue lesions seen in asbestosis. Photomicrographs of the particles are shown.

-- Cond. from Bull. Hyg.

DATE
EXAMINED: 8/1

PURPOSE
OF SEARCH:
ABSTRACT BLANK FOR GENERAL USE

CONT'D.

MELLON INSTITUTE, PITTSBURGH, PA.

AUTHOR:

J. Clerens

TITLE:

Research into Pulmonary Asbestosis in Belgium

ORIG. REF.

VOL.

PT.

DATE

ABST. REF.

VOL.

PT.

DATE

Arch. Belg. med. 8 557-565 11/51 Bull. Hyg. sociale, HFR., med. du travail et ind. 26 402 4/51

ABSTRACT:

The investigation concerns the fibro-cement industry, in which asbestos is crushed and mixed with cement. Enclosed apparatus prevents most of the dust from escaping; in the most dusty locations, 20 to 40 million particles per cubic foot were found, with 50% under 5 microns. The workers have been examined periodically over a period of three years. Out of 75 men employed for seven to over thirty years, there were four doubtful cases, and only one presented x-ray shadows indicating pneumoconiosis. The man died of progressive circulatory insufficiency, and details of post-mortem findings are given.

DATE

BY

PURPOSE

OF SEARCH

ABSTRACT PLANE FOR GENERAL USE

SOURCE

MELLON INSTITUTE, PITTSBURGH, PA.

CLASSIFIED SUBJECT

0219 2/52

NUMBER
FC 752

AUTHOR: J. M. Smith, I. D. P. Wootton and E. J. King

TITLE: Experimental asbestosis in rats

ORIG. REF.	VOL.	PF.	DATE	ABST. REF.	VOL.	PF.	DATE
Thorax	6	127-36	6/51				

A study has been made of the effect of particle size on the production of lung fibrosis in rats by asbestos. Short-fiber asbestos (2.5 microns) caused only a moderate peribronchial fibrosis, whereas long-fiber asbestos (15 microns) caused well-marked interstitial fibrosis resembling that seen in the human disease. Alumina injected together with asbestos of varying particle size increased the fibrosis produced.

-- Authors' Summary

DATE EXAMINED	BY	PURPOSE OF SEARCH	CONT'D.
		ABSTRACT BLANK FOR GENERAL USE	

MELLON INSTITUTE, PITTSBURGH, PA.

CLASSIFIED SUBJECT

FELLOWSHIP
NUMBER

FILE NUMBER

#1144 1/50

NFC

753

AUTHOR: H. Ollivier, P. Morand, and R. Brun.

TITLE: Granulomata of the Skin Due to Asbestos

ORIG. REF.	VOL.	PP.	DATE	ARCH. REF.	VOL.	PP.	DATE
Arch. Malad. Profession-	516-517	1949		Bull. Hyg.	25	1482-488	May 1950
nelles (French)	10						

A man who was working with asbestos sheets wounded the palm of his hand slightly. Two weeks later, at the site of the cut, a swelling which was increasing in size appeared. There was no inflammation and the swelling was hard at the center. The small white nucleous of this core was seen under the microscope to be composed of needles of asbestos together with a mass of crystalline matter. This matter was identified as cholesterol, which formation would seem to indicate a fundamental and biochemical defensive process of the body against aggression by asbestos.

-- Condensed from Bull. Hyg.

DATE RECEIVED	BY	PURPOSE OF SEARCH	CONT'D.
		ABSTRACT PLANNED FOR GENERAL USE	

115-1 88

NUMBER
NFC 752

MELLON INSTITUTE, PITTSBURGH, PA.

AUTHOR: K. Holoman

TITLE: The Asbestosis Problem in Slovakia

ORIG. REFS.	VOL.	PP.	DATE	ACRIF. REFS.	VOL.	PP.	DATE
Pracovni Lekarstvi	2	69	1950	Arch. Ind. Hyg. & Occ. Med.	2	610	11/50

A¹ At present not a single case of asbestosis of the lungs has developed in three plants which were investigated. This is due to the short exposure to dust-containing fibres of the majority of the workers, and also to the use of raw and wet asbestos, which allows only long and less noxious fragments to be suspended in the atmosphere. Control examinations after longer periods of time will show whether certain suspected roentgenologic findings have an anatomic basis due to asbestos or are due to other causes.

-- Adaptation of Author's Summary

DATE RECEIVED	BY	PURPOSE OF SEARCH	ABSTRACT BLANK FOR SEPARATE USE	REMARKS

855

7/55

NUMBER
NFC

752

AUTHOR: F. Lecher,

TITLE: Primary cortical-cell tumor of the peritoneum in a case of asbestosis.

ORIG. REF.: German,

VOL.

PP.

DATE

ABSTR. REF.:

VOL.

PP.

DATE

Arch. Gewerbepathol. 13 382-392 (1954)

Gewerbepath.

A case of asbestosis following long exposure to asbestos is

described. In addition to typical symptoms and findings, leptomenin-

gitis was present and tubercle bacilli were found in the spinal fluid.

The patient died with cachexia. Autopsy showed tuberclosis of the

lumbar spine with bilateral psoas abscess and mixed infection in both

lungs. Asbestosis, stage two, was found in the lungs with definite fi-

brosi; there was slight asbestosis in the lymphatic glands, and asbes-

tos bodies were found in the spleen. On the peritoneum of the mesen-

tery and over the peritoneal cavity there were thickened patches and

many small nodules; a flat tumor, half the size of a hand and 1 cm. in

thickness, was attached to the peritoneum on the under side of the dia-

phragm. The results of the examination of the tumor are fully describ-

ed and illustrated. Asbestos was found in the tumor tissues by x-ray

diffraction. It is uncertain how the asbestos reached the peritoneal cav-

ity; the most probable route was thought to be by direct penetration of

the asbestos fibers from the lung through the pleura and thence through

the diaphragm.

MELLON INSTITUTE, PITTSBURGH, PA.

favour the retention of particles within the lungs. (5) Particles under 5 microns in length restricted cellular motility more than larger particles, for the reason that they were more readily phagocytosed. It would seem that wandering phagocytes are rarely able to ingest particles larger than 10 microns. Giant cells sometimes formed in tissue cultures, apparently by fusion of two or more ordinary phagocytes, and were then able to engulf larger particles or clusters of particles. (6) There was no evidence of the formation of asbestos bodies in these experiments, nor have we noted the appearance of brown granular pigment like that seen in the lungs of experimental animals.

-- Authors' summary

CLASSIFIED SUBJECT

FOLLOW-UP
NUMBER

FILE NUMBER

FC

752

AUTHOR:

T.F. Belt, J. Friedmann and E.J. King

TITLE

The effect of asbestos on tissue cultures: a comparative study with quartz and coal dust.

ORIG. REF.:

J. Path. & Bact.

VOL. PP.

59 159-164

DATE

1947

ABSTR. REF.:

VOL. PP.

DATE

ABSTRACT:

(1) Phagocytosis, even of small particles of asbestos, took place slowly in tissue cultures and a considerable portion of the asbestos dust, approximately 50 per cent, remained untouched, though within range of phagocytes throughout the experiments. Compared with coal or quartz, asbestos was less readily phagocytosed, particle size being equal. (2) Asbestos particles in small amount had little or no toxic effect on living cells in tissue culture. The growth of fibroblasts was neither retarded nor accelerated and no necrosis of cells was observed as a result of contact with asbestos particles, while fatty change appeared to occur more slowly in asbestos phagocytes than in coal phagocytes. (3) Asbestos particles underwent no visible change, and in particular no visible diminution in size, after having been ingested by phagocytes for as long as five weeks. (4) The motility of wandering phagocytes was decreased after the ingestion of asbestos particles. This effect, not necessarily of a toxic character, may be deleterious in the case of cells within the lungs, since it tends to restrict the migration of cells and so

DATE

Authors' summary

EXAMINED:

BY:

FTR

PURPOSE
OF SEARCH:

ABSTRACT BLANK FOR GENERAL USE

CONT'D. ✓

MILLON INSTITUTE, PITTSBURGH, PA.

FORM 77

MELLON INSTITUTE, PITTSBURGH, PA.

CLASSIFIED SUBJECT

FOLLOW-UP
NUMBER

FILE NUMBER

752-338

AUTHOR:

J. D. Hindley-Smith

TITLE:

Some medical problems of the asbestos cement industry.

ORIG. REFS.:

VOL.

PP.

DATE

ABSTR. REFS.:

VOL.

PP.

DATE

Brit. J. Phys. Med.

10

50

Mar.-Apr.
1947

Occ. Med.

4

469

Oct. '47

ABSTRACT:

The author encountered no incipient cases of asbestosis, and only two or three of long-standing chronic disease, in five years' experience in an asbestos cement factory, showing that ventilation and enclosure were adequate. A dermatitis occurring in some workmen was found to be caused by an abrasive constituent of the cement, other than the asbestos. Cleanliness and the use of a barrier cream were effective in preventing it.

DATE
EXAMINED:

BY

FTR

PURPOSE
OF EXAMIN:

COM'DR

ABSTRACT BLANK FOR OTHER USE

CLASSIFIED SUBJECT

FELLOWSHIP
NUMBER

FILE NUMBER

605

6/53

n/c

752

AUTHOR: F. Doernke

TITLE: Lung carcinoma caused by asbestos inhalation.

ORIG. REF.:

VOL.

PP.

DATE

ACCT. REF.:

VOL.

PP.

DATE

Med. Monatsschr.
(German)

7

77-81

1953

Chem. Absts.

4

1471-55-10-53

The relation between fibrotic changes caused by asbestos deposits in the lung and carcinoma is discussed. The lower lobe is the site of predilection. Characteristic are the multicentered developments of flat epithelioma carcinoma.

-- Chem. Absts.

MELLON INSTITUTE, PITTSBURGH, PA.

FORM 77 10

DATE
STAMPED

BY

PURPOSE
OF STAMP

ABSTRACT BLANK FOR GENERAL USE

CONT'D.

MILLON INSTITUTE, PITTSBURGH, PA.

3. *Asbestosis, its treatment, M. Giroux and M. Leduc*
Asbestosis and pulmonary tuberculosis.

ORIG. RECD. VOL. 23 1820 1947 DATE APR. 1947
 Sem. Hop. Paris (French) Brit. J. Ind. Med. 5 100 APR. 1947

ABSTRACT

The authors report from Quebec on asbestosis in the Canadian mines and on animal experiments. They disagree with Gardner's statements that pneumoconiosis due to asbestos is not an important factor in the production and progress of pulmonary tuberculosis, and that the presence of asbestos bodies in the sputum has no value in the diagnosis of asbestosis. Several cases of asbestosis are described; in one, death was due to tuberculosis and asbestosis, giving support although not proof for the authors' theory. Two cases of asbestosis with lung cancer are also described briefly. In animal experiments by Giroux, lung lesions and asbestos bodies were produced in guinea pigs but not in rabbits or dogs. In these animals the author traces the formation of asbestos bodies: fibers reaching the alveoli are rapidly covered by phagocytic cells, movement of the fibers results in microscopic bleeding from damaged alveolar walls; the hemoglobin from the liberated red cells is taken up by the phagocytes which already enclose the asbestos fibers. Subcutaneous or intratesticular injections into guinea pigs of suspensions of

DATE EXAMINED.

BY FR:Brit. J.

IND. MED.

Abstract blank for general use

CONFIDENTIAL

71045-91471

MFC 752

AUTHOR:

C. Megolius

TITLE:

Changes in the lungs in 126 cases of asbestosis observed in Finland.

ORIG. REF.:

Acta Radiologica

VOL.

28

PP.

139-152

DATE

1947

ASSIN. REF.:

JHEP

VOL.

29

PP.

91

DATE

Sept. '47

ABSTRACT:

Asbestosis of the lungs is an industrial disease not often hitherto observed in Scandinavia, only five cases have been published from Norway. In Finland, where a considerable asbestos industry has been established for a couple of decades, asbestosis of the lungs appears to occur to an appreciable extent. The author gives an account of 126 cases of asbestosis of the lungs, 94 or 75% of those cases being the mild, 23 or 18% in the moderate and 9 or 7% in the advanced stage of asbestosis. The roentgenological characteristics of the changes in the lungs, the differential diagnosis, the classification in different stages and the importance of the time of exposure are discussed in detail. The different stages of asbestosis of the lungs are illustrated by typical pictures. -- Author's summary.

Author's summary

DATE
EXAMINED:

BY: JFR

PURPOSE
OF SEARCH:

ABSTRACT BLANK FOR GENERAL USE

CONT'D:-----

MELLON INSTITUTE, PITTSBURGH, PA.

MELLON INSTITUTE, PITTSBURGH, PA.

CLASSIFIED SUBJECT

#10446-9/47

FOLLOWUP NUMBER

FILE NUMBER

AUTHOR:

P. Luton and J. Champelx

NFC

752

TITLE:

Asbestosis.

ORIG. REFS.:

VOL.

PP.

DATE

ABST. REFS.

VOL.

PP.

DATE

Arch. Mal. Prof.

7

365-378

1947

JHELT

29

91

Sept. '47

ABSTRACT:

The authors list the following symptoms of asbestosis: (1) Functional signs; (2) Radiological findings; (3) Sputum, characterized by the presence of asbestos fibers and bodies. After a presentation of several case histories and their accompanying x-ray films and sputum photographs, it is concluded that diagnosis can only be made by a combination of observations: functional, radiological, sputum. Because of the progressive nature of the disease, yearly examinations should be made with comparison of the measurements of respiratory function (spirometry, etc.) and x-rays. But functional signs must guide the physician when a change of environment is advisable, and when the worker can be said to have contracted asbestosis. -- From J.I.M.B.T.

DATE EXAMINED:

BY:

FFR:FROM JIM OF SEARCH

CONT'D

asbestos in olive oil resulted in the production of granulomatous lesions but no asbestos bodies. When, however, the suspension was injected into an area into which 1 ml. of blood had been injected immediately before, asbestos bodies were found after a month. He considers that the presence of asbestos bodies in sputum has significance, since it implies tissue damage in the lungs.

-- From Brit. J. Ind. Med.

AUTHOR: K.M. Lynch and W.M. Cannon

TITLE: Asbestosis: VI Analysis of Forty Necropsied Cases.

ORIG. REFS.:

VOL.

PP.

DATE

ABSTR. REFS.:

VOL.

PP.

DATE

Dis. Chest.

14

874-880

1948

Brit. J. Ind. Med.

6

268-277

Oct '47

MELLON INSTITUTE, PITTSBURGH, PA.

The authors report an analysis of 40 cases in which asbestosis was found at necropsy. Cases are divided into minor, medium, and advanced grades, the advance of the disease and the prominence of asbestosis bodies showing a direct relation to the duration of exposure. In 4 of 14 cases of advanced grade, pulmonary tuberculosis was the primary cause of death. Only in cases of recent exposure was recent fibrosis found, this providing support for Gardner's belief that fibrosis does not progress indefinitely after cessation of exposure. Presence of asbestosis bodies is not invariable in experimentally produced asbestosis, but is more typical of human asbestosis. Lung damage occurs before the formation of asbestosis bodies, which may represent an attempt by the tissues to segregate the particles; they persist in the lung indefinitely. Pleural fibrosis is an outstanding, though not invariable, feature of asbestosis; the reason for this is not clear as the particles do not reach the pleura. The occurrence in asbestosis of nodular fibrosis of silicotic type, which King and others have produced experimentally, was confirmed in 8 cases. Carcinoma of the lung occurred in three out of 40 cases--an incidence of 7.5 per cent, which is seven times greater

EXAMINED

BY

OF SEARCH

ABSTRACT BLANK FOR GENERAL USE

than general incidence--all in cases of medium--or advanced grade--asbestosis. This figure is in accordance with the experience of other workers, and calls for further study. Tuberculosis occurred in six cases, but it is suggested that this incidence is not significant; this is not quite convincing. Gardner stated that experimental asbestosis does not predispose to tuberculosis. (Unfortunately the occupational histories of these cases are not complete).

-- Brit. J. Ind. Med.

MELLON INSTITUTE, PITTSBURGH, PA.

MFC 752

4257 3/59

A Frenchini and G. Caneva

Title: Pulmonary Asbestosis; Anatomopathologic Study of a Case.

Orig. Refs.

Med. d. lavoro

Vol. 40

No. 161

Date: Jan-Ju, 1949

Arch. Ind. Hyg. & Occ. Med.

No. 362

Date: March 1950

Italian

A man 40 years old had worked in a factory for 22 consecutive years, applying insulating material to articles of asbestos. He never complained about respiratory disorders, and he died as the result of a streetcar accident. During the necropsy the lungs were found to be the seat of extensive asbestosis. The author's direct attention to the disproportion between the gross features described and the awareness of trouble. The results prove the importance of prophylactic and therapeutic treatment and economic assistance to asbestos workers generally, particularly to those who apply insulating material. -- Condensed from Arch. Ind. Hyg. & Occ. Med.

DATE EXAMINED

BY

PURPOSE OF EXAMIN.

COM'D BY

ABSTRACT BLANK FOR GENERAL USE

AUTHOR:

P. Iuton, J. Champetix and P. Faure

TITLE:

The significance of the presence of asbestos bodies in the sputum of asbestos workers.

ORIG. REF.:

VOL.

PP.

DATE

ASST. REF.:

VOL.

PP.

DATE

Arch. malad. profess. 8

56-50

1947

Ind. Hyg. Index

1

3-4

Nov. Dec. 1947

ABSTRACT:

33 ref., 7 photomicrographs

The authors examined 176 persons who worked or had worked with asbestos. They find that asbestos bodies appear somewhat early in the sputum of these workers, sometimes after only 20 days from the time of exposure. These fibers may persist even after cessation of exposure, but occur in almost all workers exposed to asbestos dust, whether they exhibit functional or radiographic evidence of asbestosis or not. Brown or golden yellow bodies are seen in the sputum along with uncolored fibers or spicules. The presence of these bodies, more or less scattered, indicates that the subject has inhaled asbestos dust, but it does not necessarily signify that asbestosis exists. In other words, the discovery of asbestos bodies in the sputum is only one in the ensemble of elements which permit such a diagnosis. It is true that in the majority of cases where asbestosis is predicated by reason of the intensity of functional manifestations and radiographic signs, numerous asbestos bodies are noted, massed together, and displaying a radiate formation. As such, they are identical to the clumps that are observed on histological examination of the lungs of deceased cases, in the alveoli.

-- Ind. Hyg. Index

DATE EXAMINED:

22 FEB 1948

PURPOSE OF EXAMIN:

ASSIGNMENT BLANK FOR GENERAL USE

MELLON INSTITUTE, PITTSBURGH, PA.

AUTHOR: Virginia Cameron

TITLE: The Cost of Research on Tuberculosis in the United States. Analysis of Allocations in 1947-1948.

ORIG. REFS.: Am. Rev. Tuberc.

VOL. PP.

60

393-405

DATE Oct. '49

ASSN. REFS.

VOL. PP.

DATE

1267 The Cost of Research on Tuberculosis in the United States. Analysis of Allocations in 1947-1948. Virginia Cameron. Am. Rev. Tuberc. 60, 393-405 (Oct. 1949).

The present research programs of various agencies are outlined, and their contributions to research on tuberculosis are summarized. The total cost was over six million dollars. Official health agencies account for 56.8% of the total, manufacturing firms 32.8%, and the remaining 10.4% is supported by universities, foundations, and voluntary health agencies. Most of the commercial firms are manufacturers of x-ray apparatus or pharmaceutical houses. However, other industries, principally light and heavy metal manufacturing and mining operations, made a notable contribution (\$83,550) during the period. Fifty percent of this amount was designated for research on silicosis and tuberculosis.

CLASSIFIED SUBJECT

FELLOWSHIP NUMBER

FILE NUMBER

DATE EXAMINED.

BY

PURPOSE OF SEARCH.

ABSTRACT BLANK FOR GENERAL USE

CONT'D.

MELLON INSTITUTE, PITTSBURGH, PA.

CLASSIFIED SUBJECT 867 FILE NUMBER 377 75-2
 479
 471
 FC

AUTHOR: Editorial
 TITLE: Asbestosis and Cancer of the lung.

ORIG. REF.: J.A.M.A. VOL. 140 PP. 1219-20 DATE Aug 13-49
 140 1219-20 Aug 13-49

867 Asbestosis and Cancer of the lung. Editorial.
 J.A.M.A. 140, 1219-1220 (Aug. 13, 1949).

Records of English, American and German physicians and The Annual Report of the Chief Inspector of Factories in England for 1947 show that the occurrence of cancer of the lung is related to pulmonary asbestosis. This relation is supported by the following observations: (1) The incidence rate of cancer of the lungs in asbestotic patients is 10 to 15 times as high as among the general population. (2) The male-female ratio is more nearly equal than it is for the general population, which indicates that an environmental and evidently occupational carcinogen tended to equalize the incidence rate of cancer of the lungs for both sexes. Recent experimental observations support this interpretation of clinical evidence.

DATE EXAMINED BY: PURPOSE OF SEARCH: CONT'D:
 ABSTRACT BLANK FOR GENERAL USE

MELLON INSTITUTE, PITTSBURGH, PA.

7053 71.54

NUMBER
MPC 752

AUTHOR: H. Myers
TITLE: Asbestosis

ORIG. REF.: Postgraduate Med. J. VOL. 25 PP. 631-638 DATE: Dec. 1949 ABSTR. REF.: Bull. Hyg. VOL. 25 PP. 477-478 DATE: May 1950

Asbestos and talc are the only two silicates which have been shown to produce pneumoconiosis. Gardner, in 1938, suggested, on the basis of animal experiments that the effect produced by asbestos was mechanical, since it was produced only by long fibers. A later authority made the observation that workers exposed to very short-fiber asbestos were not affected by it. Asbestosis was officially recognized as an industrial disease in 1930. The condition is thought to be due to blocking of the finer bronchioles by long fibers which seem to irritate the bronchial wall and cause fibrosis. Tuberculosis and carcinoma of the lungs may follow, as complications. Recognition of the disease, and precautions which have been taken have lessened its severity and incidence.

-- Condensed from Bull. Hyg.

DATE EXAMINED: BY: PURPOSE OF SEARCH: ABSTRACT BLANK FOR GENERAL USE: CONT'D:

CLASSIFIED SUBJECT: 254 905 FOLLOWUP NUMBER: nfc FILE NUMBER: 752

588 1/5

(L)

NFC

102

AUTHOR: P. Cartier

TITLE: A Contribution to the Study of Asbestosis.

CARD. REFS.	VOL.	PP.	DATE	ARCH. IND.	HYG. & OCC.	MED.	VOL.	PP.	DATE
Arch. Malad. Profess.	10	889-595	1949	Arch. Ind. HYG. & Occ.		Med.	3	428	4/51

The author has been in medical charge of 3,242 men employed in asbestos mines in Canada, where over 70 per cent of the world's supply of asbestos is mined. Cases of asbestosis were found only in those who had been employed for at least 14 years and exposed to air containing at least 5 million particles of asbestos fibers per cu. ft., the fibers being from 10 to 250 microns in length. The outstanding feature of this report is the mildness of any pulmonary trouble among the asbestos miners compared to that reported in the United States and Great Britain as occurring in factories where asbestos is spun and woven. No clinical symptoms of pulmonary fibrosis were present. No case of pure asbestosis was detected in those under age 60, and no physical incapacity was evident. Cases of tuberculosis were not more frequent among the miners than among the clerical staff and the rest of the community, although that incidence was high. It was difficult to ascribe any cardiac affections present to asbestosis, because of the late age at which it appeared. Postmortem findings and roentgenologic examinations indicate that asbestosis is a pathological entity; but among the miners it develops so slowly as not to shorten working life, and there is no evidence that it predisposes to tuberculosis. The conclusions reached are supported by the opinions of experts at Garmac. -- Cont. from Bull. Hyg.

1009 152 752 752

AUTHOR: R. Stoll, R. Bass, and A. A. Angrist.

TITLE: Asbestosis Associated with Bronchogenic Carcinoma.

ORIG. REF.	VOL.	PP.	DATE	ABSTR. REF.	VOL.	PP.	DATE
Arch. Intern. Med.	88	831-834	Dec. 51	Bull. Hyg.	27	444	5-52
(In German)							

ABST

Data for statistical proof of any association of asbestosis with pulmonary carcinoma are hard to collect, but three investigators have found carcinoma in 7.5 to 14.8% of cases of asbestosis. A case is reported in which a worker had been exposed to asbestos for 6 years and had refused to wear a respirator. He came under notice with weakness and a persistent cough. Clumps of malignant cells were seen in the bone marrow, and x-ray showed generalized large discrete oval shadows indicative of pulmonary metastases. After death, nodules varying to 4 cm. in diameter were found scattered throughout both lungs, with interstitial fibrosis. Large areas of necrosis existed throughout the lungs, and typical asbestos bodies were seen inside and outside the tumor areas. Metastases were found in the kidneys, brain, and liver. The opinion is suggested that the silicate in asbestos must itself be directly carcinogenic. It establishes an additional reason for preventive measures in the use of asbestos.

DATA

ABSTRACT BLANK, FOR GENERAL USE

Cond. from Bull. Hyg.

MELLON INSTITUTE, PITTSBURGH, PA.

CLASSIFIED SUBJECT

MEMORANDUM

FILE NUMBER

nfc 752

author: P. Luton, J. Champetx, and P. Faure.

title: A Case of pulmonary asbestosis, with its development followed by

X-ray and its post-mortem study.

ORIG. REF.	VOL.	PP.	DATE	ASSIN. REF.	VOL.	PP.	DATE
Arch. maladies profess	12	629-33	1951	Bull. Hyg.	27	942	10-52

The case reported is said to have been the first one of asbestosis

known to have occurred in France. The man had been exposed to high concentrations of asbestos for 11 years. Symptoms and progressive x-ray findings are described. After death the lungs showed general thickening throughout, especially of the plurae, but with no localized nodules. Many asbestos bodies were seen, isolated and in groups, together with giant cells and mineral particles. No evidence of tuberculosis was found. The dust particles could be traced not only in the lung vessels, but also in the vessels of the liver, kidneys and spleen. The action of asbestos is apparently mechanical rather than chemical.

-- Cond. from Bull. Hyg.

DATE REAMINED: _____ BY: _____ PURPOSE OF SEARCH: _____ COMMENTS: _____

ABSTRACT PLANS FOR CREDIT: (10)

MELLON INSTITUTE, PITTSBURGH, PA.

34 102 1
 NUMBER
 NTC 752

AUTHOR: J. Clerens

TITLE: Research into Pulmonary Asbestos in Jn. Belgium

ORIG. REF. Arch. Belges med. 8 557-565 11/51 Dull. 11981 26 402 4/51
 SOCIETY, N.Y.R. med. du travail et med. legale
 ABSTRACT:

The investigation concerns the fibro-cement industry, in which asbestos is crushed and mixed with cement. Enclosed apparatus prevents most of the dust from escaping; in the most dusty locations, 20 to 40 million particles per cubic foot were found, with 50% under 5 microns. The workers have been examined periodically over a period of three years. Out of 75 men employed for seven to over thirty years, there were four doubtful cases, and only one presented x-ray shadows indicating pneumoconiosis. The man died of progressive circulatory insufficiency, and details of post-mortem findings are given.

DATE EXAMINED: 3/7/51 PURPOSE OF SEARCH: 60710

MELLON INSTITUTE, PITTSBURGH, PA.

CLASSIFIED SUBJECT

FELLOWSHIP
NUMBER

FILE NUMBER

NFC

752

AUTHOR:

A. Rohme

TITLE:

Results of Periodical Examinations of Workers in an Asbestos Factory

ORIG. REF.:

VOL.

PP.

DATE

ABST. SER.

VOL.

PP.

DATE

Reitarsz z. Silkose 11 | 27/31 | 1951
Forschung (German)

Bull. Hyg.

26

511

5/51

Since radiological supervision and control measures were adopted in an asbestos factory in 1936, the dust hazard has been greatly reduced, as shown by comparing results of surveys in 1936, 1940, and 1949. A comparison of different working places shows that the risk is low in spinning and weaving, but high in preparation, carding and brake polishing. A table shows results for 14 cases from 1936 to 1949. Some progressed, others did not. Inquiries among former workers reveal little if any tendency to progression after the dust exposure has ceased. This is in contrast to silicosis. On the other hand a personal factor exists, as in silicosis, as a number of workers do not get asbestosis in spite of high dust exposures.

Progressive tuberculosis is rare, only two cases being found, both started in the parts of the lung least affected by asbestosis. One combination of asbestosis and carcinoma of the lung was found. The main result of this study is the demonstration that the hazard can be reduced by dust control.

--Cond. from Bull. Hyg. --

MELLON INSTITUTE, PITTSBURGH, PA.

AUTHOR: W. Behrens
TITLE: Experimental asbestosis.

ORIG. REF.: Schweitz, Z. allg. path. u. bakt. VOL. 14 PP. 275-97 DATE 1951 ABSTR. REF.: Bull. Hyg. VOL. 26 PP. 1181 DATE 12/51

ABSTRACT:

Pure chrysotile asbestos was injected intraperitoneally and intratracheally into mice and rats. The fibers varied from 5 to 180 microns in length. Unsuccessful attempts to secure fibers of standard length are described. The general picture is one of non-specific focal fibrosis resulting from the introduction of foreign bodies. No solution or toxic action of the asbestosis could be demonstrated. Asbestosis bodies were formed. The theories of the pathogenesis of asbestosis are discussed.

-- Cond. from Bull. Hyg.

DATE EXAMINED: 2/1 PURPOSE OF SEARCH: ABSTRACT BLANK FOR GENERAL USE CONT'D.

MELLON INSTITUTE, PITTSBURGH, PA.

CLASSIFIED SUBJECT

C174 5/52

FELLOWSHIP
NUMBER

FC

FILE NUMBER

752

AUTHOR: A. J. Lanza

TITLE: Asbestosis

Paper read before Fourth Conference of McInlyre Research Found. on Silicosis

ORIG. REF.

VOL.

PP.

DATE

ASST. REF.

VOL.

PP.

DATE

2 pp

1/1952

ABSTRACT: The action of asbestosis on the body is briefly reviewed. It is pointed out that asbestosis is regarded much more seriously in Great Britain than in Canada or the United States; the British believe it is worse than silicosis and often leads to cancer, while in America cancer resulting from asbestosis is practically non-existent.

DATE
EXAMINED

BY

PURPOSE
OF SEARCH

ABSTRACT BLANK FOR GENERAL USE

CONT'D.

354

3/63

8

n/c

752

AUTOR.

H. Bastenier, H. Denoln, A. Decoster, and R. Denoln-Reubens.

TITRE.

A clinical and pathological study of a case of pulmonayr asbestoses.

ORIG. REF.	VOL.	PP.	DATE	ASSIN. REF.	Bull. Hyg.	VOL.	PP.	DATE
Arch. belges med. sociale, hyg., med. [travail et med. legale (French)]	10	61-70	Feb. 52			27	1126	12-52

A diagnosis of asbestosis must depend upon a history of occupational exposure to asbestos dust, associated with effects on the respiratory organs. A case is reported in which a woman had worked for 25 years weaving asbestos in a factory with heavy dust exposure. The symptoms described are typical. The authors compared this case with those of uncomplicated silicosis; in the latter residual air is frequently increased, expiration is prolonged, and the maximum respiratory capacity is reduced, but hyperpnea is less, rapid breathing is not so intense, and anoxemia is never so important during effort, as in asbestosis. These differences are stressed as important when necessity may arise in determining the occupational origin of a case with obscure diagnosis.

-- Cond. from Bull. Hyg.

CONT'D.

604 63 752 nfc

AUTHOR: W. Behrens, Jr.

TITLE: The clinical picture and pathology of asbestosis.

ONIS. REFS. VOL. 45 129-40 DATE June 15, 1952 ABSTIN. REFS. Bull. Hyg. VOL. 28 192 3-53 DATE

Z. Unfallmed. u. Berufkrankh.

Abstract.

This paper presents an extensive review of literature on asbestosis including distribution and types of asbestos, its mining, treatment, and use, dust hazards, and effects of length of exposure of the dust in different concentration. Symptomatology, diagnosis, and x-ray findings in the different stages of asbestosis are described. Special mention is made of the association of asbestosis with carcinoma, which is much more frequent than that of silicosis. Post-mortem reports from the literature indicate that in 309 cases of asbestosis there were 44 lung carcinomas (14.2%), whereas in 2,204 cases of silicosis only 32 carcinomas were found (1.4%). The experience of the association of tuberculosis and asbestosis varies greatly in different countries. English investigators have recorded a high rate of tuberculous infection in asbestos workers, but it is a rare complication in America and the rate is low in Germany. The macroscopic and microscopic appearances of the lungs are described, and the nature and significance of asbestosis bodies are discussed and an extensive reference list dealing with that controversial subject is given.

-- Cond. from Bull. Hyg.

MELLON INSTITUTE, PITTSBURGH, PA.

73

1/54

1 m/c

1000

AUTHOR:

J. Rutner

TITLE:

Asbestosis bodies and pseudo-asbestosis bodies.

JOURN. REF.

VOL.

PR.

DATE

ACCESSION

VOL.

PR.

DATE

Schweiz. Z. Allgem. Pathol. u. Bakteriolog.

15

628-31

1952

Bull. Hyg.

28

719

Sept. '53

(German)

The literature on asbestosis bodies is reviewed. Similar products were found by the author in a lung which only contained carborundum and graphite; the bodies were formed around plates and needles of carborundum. The protein and iron which form the bulk of the "bodies" must come from the lung tissues and not from the mineral matter of the dust. Similar bodies could also be produced in vitro but they have not so far been found in animal experiments.

-- Bull. Hyg.

MILLON INSTITUTE, PITTSBURGH, PA.

RECEIVED
 U. S. DEPT. OF HEALTH
 DIVISION OF LABOR STANDARDS

CENT 2

CLASSIFIED SUBJECT

FOLLOWING
NUMBER

FILE NUMBER

#262

3/54

nfc

752

AUTHOR:

G. Farris

TITLE:

Asbestos warts and other dermatoses in workers with lagging materials.

ORIG. REFS.

VOL.

PP.

DATE

ASST. REFS.

VOL.

PP.

DATE

Rass. med. ind.
(Italian)

22

1-17

Jan-Feb
1953

Bull. Hyg.

28

953

Nov. '53

The author has investigated cases of warts on the hands of 29 workers with lagging materials, including asbestos cord which is wound around pipes, various asbestos-containing slabs, and mixtures of asbestos and cement to be molded to the shape of the tank or other object to be insulated. One to four warts were found in every case. They are due to asbestos fibers which penetrate into the epidermis and become lodged in the corium or true skin. The epidermis shows acanthosis (thickening of the prickle-cell layer) and hyperkeratosis (increased horny tissue) and the corium shows a granulomatous type of reaction with the appearance of giant cells in some cases. Exudative dermatitis may also be present. This condition is attributed to the combined action of the hygroscopic and alkaline cement and the irritating asbestos fibers. The gross and microscopic appearances of the lesions are illustrated.

-- Cond. from Bull. Hyg.

MELLON INSTITUTE, PITTSBURGH, PA.

FORM 77 10

DATE

EXAMINED:

BY:

PURPOSE

OF SEARCH:

ABSTRACT BLANK FOR GENERAL USE

CONT'D.

CLASSIFIED SUBJECT

FELLOWSHIP
NUMBER

FILE NUMBER

25

53

n/c

752

AUTHOR: M. A. Kovnatskil and others.

TITLE: Silicosis: etiology, pathogenesis, and clinical treatment.

ORIG. REFS.	VOL.	PP.	DATE	ABSTN. REFS.	VOL.	PP.	DATE
Gigiena i Sanit. No. 8 (Russian)	28-32	1952		Chem. Absts.	46	11400	1/25/52

ABST: A review is given of the disease caused by particles of silicates, rather than those of the type producing silicosis proper. A study of the disease caused by asbestos particles showed that the chemical and biological features are increased rate of respiration (compensatory) and in stages the level of blood saturation with oxygen is essentially normal, although changes in the myocardium are evident on the electrocardiogram. The tone of the veins is reduced, and the permeability of capillary walls suffers a decline in a considerable number of cases. This affects the oxidative processes as shown by the increase of oxidized glutathione and a decrease of total glutathione.

-- Chem. Absts.

DATE EXAMINED	BY	PURPOSE OF SEARCH	CONT'D.
		ABSTRACT PLAIN FOR GENERAL USE	

152
A. S. Ramaswamy, D. S. Venkatesh, and R. Rama Rao

152
The nature of lesions caused by asbestos and mica in experimental pneumoconiosis in guinea pigs.

ORIG. REF. 1
J. Indian Inst. Sci. 35 319-31 Oct. 1953 Bull. Hyg. 29 172 Feb. 1954

Guinea pigs were subjected to intratracheal injections of mica and asbestos dusts. Stress is laid on the development, after long periods, of lesions in other organs besides the lungs, especially in the adrenal glands. Mica dust was found to lead to some degree of nodular formation in the lungs, but the asbestos dust set up diffuse fibrosis only. These differences are portrayed in a series of photomicrographs. Mica induced hypertrophy of the adrenal cortex with extensive foamy disintegration of the cortical cells; asbestos induced lesser changes, but small areas of necrosis were occasionally found in the zona fasciculata and zona reticulata. In a theoretical discussion, the authors attempt to draw a relation between the atomic structure of mica and asbestos and the pathological lesions in the lungs and other organs. Their theory, however, leaves many points unexplained. --Cond. from Bull. Hyg.

MELLON INSTITUTE, PITTSBURGH, PA.

FORM 77 10

DATE RECEIVED BY PURPOSE OF RESEARCH CONT'D.
ASSIGNMENT BLANK FOR GENERAL USE

Author: J. Krausler and H. Seyss.

Title: Early diagnosis of asbestosis by enlarged radiographs.

ORIG. REF.	German	VOL.	PP.	DATE	ADJIN. REF.	VOL.	PP.	DATE
7. Unfalmed. u.	47	59-64	Mar 15 '54	Bull. Hyg.	29	829	8/54	
Beaufskrankh.								

After a brief description of asbestosis and the changes in the lung tissue

in the early stages, the authors describe a radiographic technique with a fine focus. Of 16 asbestos workers, with 1 to 18 year's exposure, eight showed changes in the lung tissue with the enlargement technique; two of those showed no change by normal technique, and in the other six cases the degree of change shown was minimal.

-- Cond. from Bull. Hyg.

MELLON INT

FORM 77

DATE EXAMINED:	BY:	PURPOSE OF SEARCH:
		ABSTRACT WORK FOR GENERAL USE

CONT'D.

AUTHOR, J. F. Knox and J. Beattie.

TITLE, Distribution of mineral particles and fibers in the lung after exposure to asbestos dust.

ORIG. REFS. Arch. Ind. Hyg. & Occ. Med. 10 30-36 (July '54)

PHILADELPHIA, PA.

ABSTRACT

The particle-size distributions in the Incombustible and acid-insoluble

residues from the lungs of 27 persons who had been exposed to asbestos dust were determined. The number of particles with greatest lengths in excess of 26 microns was considerable up to the eighth year after the last exposure to asbestos dust and then sharply fell. The number of particles within the size ranges of five microns and less and between five and 15 microns remained constant over a very long period in those cases which showed no sign of asbestosis or minimal fibrotic change. The cases of severe asbestosis showed a significant rise in the numbers of such particles.

It is considered that the mechanism which is responsible for the production of human asbestosis is not the same as that which is responsible for the production of asbestotic changes in experimental animals exposed to high concentrations of asbestos dust. The appearance of such changes seems to be related to the breakdown of asbestosis bodies which may liberate some fibrogenic agent.

-- Authors' summary

UNIT EXAMINED BY PURPOSE OF SEARCH

ABSTRACT BLANK FOR GENERAL USE

CLASSIFIED SUBJECT

FOLLOWUP
NUMBER

752

AUTHOR: J. F. Knox and J. Beattie.

TITLE: Mineral content of the lungs after exposure to asbestos dust.

ORIG. REF.

VOL.

PP.

DATE

ABST. REF.

VOL.

PP.

DATE

Arch. Ind. Hyg. &
Occ. Med.

10

23-29

(July '54)

ABSTRACT

The mineral content of the lung parenchyma; the lung hilar region, and the tissues of the pleural and subpleural regions of the lung has been determined in lungs from 27 workers in the asbestos-textile industry. The exposure time varied from five to 33 years, and the time interval between the last exposure to dust and death (survival time) was from less than one year to 21 years. There was an indication that the mineral material found in the lungs increased in amount as the exposure time lengthened. The amount so accumulated varied considerably from person to person. As the survival time increased, the mineral content of the lung tended to decline but again at a very variable rate. The severity of the asbestotic lesions in the lungs found at death was related not to the mineral content but to the sum of the exposure and survival times.

-- Authors' summary

FORM 77 14

DATE
EXAMINED

BY

PURPOSE
OF SEARCH

ABSTRACT BLANK FOR GENERAL USE

CONT'D

91- 1/1

NFC 752

AUTHOR: Queries and minor notes, JAMA

TITLE: Asbestosis.

ORIG. REF.: J. Am. Med. Assn. VOL. 56 PF. 1546 DATE (Dec. 18, 1944) ASSIN. REF.: VOL. PF. DATE

A 50-year-old man worked at covering pipe with asbestos insulation from 1940 to 1945. The question is the possibility that pulmonary fibrosis evident on roentgenograms in 1952 was due to asbestosis when the films had been normal in 1949. The answer is that the cause of the fibrosis must be sought among other sources than the 1940-45 occupation. Little evidence exists of progression of asbestosis after cessation of exposure to asbestos fibers. Moreover, exposure would have had to be much more severe than in the man's occupation in order to produce asbestosis, and it would have manifested itself in the 1949 roentgenogram.

MELLON INSTITUTE, PITTSBURGH, PA.

FORM 77 20

DATE EXAMINED:

BY:

PURPOSE OF SEARCH:

ABSTRACT BLANK FOR GENERAL USE

CONT'D.

55-1/2

NFC 752

AUTHOR: W. C. Hueper.

Experimental studies in metal cancerigenesis. 6. Tissue reactions in rats and rabbits after parenteral introduction of suspensions of arsenic, beryllium, or asbestos in lanolin.

CRON, R.

CRIG. REFS.	VOL.	PP.	DATE	ANIM. REFS.	VOL.	PP.	DATE
J. Natl. Cancer Inst.	15	113-129	(1954)	Chem. Absts.	48	13043	11/10/1954

55

Experimental Studies in Metal Cancerigenesis. 6. Tissue Reactions in Rats and Rabbits after Parenteral Introduction of Suspensions of Arsenic, Beryllium, or Asbestos in Lanolin. W. C. Hueper. J. Natl. Cancer Inst. 15, 113-129 (1954).

When powdered arsenic, beryllium, or asbestos in lanolin was injected into the marrow cavity of the femur, pleural cavity, or paranasal sinuses of rats, there was no development of any significant number of cancers, either at the site of deposition or remotely, causally related to the chemicals introduced. Similar negative results were obtained with lanolin alone. The sarcoma attack rate at the site of the injection for the 250 rats used stands at 0.8%, contrasting sharply with that of 30% for rats injected with nickel suspensions and 20% for rats given parenteral

FORM 77 28

DATE	EXAMINED	BY	PURPOSE	OR BRANCH	COM'D.
					(over)

ABSTRACT BLANK FOR GENERAL USE

(actions of uranium. Multiple pulmonary osteomas were found in four rats following a traumatization of the femur by the use of an electric drill. Conclusion: under proper experimental conditions the administration of chemicals into the connective tissues of rats is a reliable method for determining carcinogenic potentialities.

-- Chem. Absts.

CLASSIFIED SUBJECT

NUMBER

8.28 1156

NEC 55

MILLON INSTITUTE, PITTSBURGH, PA.

AUTHOR: E. S. Speer and C. W. Mueck.

TITLE: Local inflammatory response produced by solutions sterilized with asbestos type filters.

ORIG. REF.: Proc. Soc. Exptl. Biol. Med. 88 | 89-93 (Jan 55) | Biol. Absts. 29 | 2142 | 9/55

It was demonstrated that filtering through an asbestos type filter disc will change the local cellular responses of the organism to that solution. This reaction is due to particles and fibers which have pulled off the disc and are present in the filtrate at the time of injection. A similar eosinophilia appeared following injections of asbestos and talc, but not following silica, cotton, or paper fibers. Ground glass and mica produced only slight increases in the number of local eosinophils. It was pointed out that precautions should be taken to avoid accidental introduction of asbestos fibers from Seltz filters into serums and other biological materials intended for injection. These experiments demonstrate that a study of the local cellular responses to an injected material can lead to a better understanding of the effects of a solution upon an organism.

-> Biol. Absts.

FORM 77

DATE OF EXAMINATION: _____ PURPOSE OF SEARCH: _____
 ANALYST: _____ ABSTRACT BLANK FOR GENERAL USE: _____
 CONT'D: _____

572

755-1

FC

752

BURGH, PA.

AUTHOR: O. A. Sander.

TITLE: Asbestosis as differentiated from other pneumoconioses.

CMIO. REF. Arch. Dec. Health

VOL. 11 | 183-211 (Mar. '55)

DATE

Asbestosis as Differentiated from Other Pneumoconioses. O. A. Sander.

Difficulties in differential diagnosis are considered. All possible causes of the pathology revealed by the chest film must be considered. Poorly studied cases result in misdiagnoses which not only are embarrassing to the physician but which may cause irreversible harm to the patient. Every obscure case should be thoroughly investigated so that such unfortunate situations do not arise.

MELLON INS

FORM 77 21

DATE EXAMINED BY

PURPOSE OF SEARCH
ABSTRACT BLANK FOR GENERAL USE

CONT'D.

FILE NUMBER

ORIG. REF.	VOL.	PT.	DATE	ANSW. REF.	VOL.	PT.	DATE
Arch. Soc. Health	11	183-21	(Mar. '55)				

COMT.D.

SURON. PA.

CLASSIFIED SUBJECT

572

572

FELLOWSHIP
NUMBER

FC

FILE NUMBER

752

AUTHOR. P. Cartier.

TITLE. Some clinical observations of asbestosis in mine and mill workers.

ORIG. REF. Arch. Occ. Health

VOL. 11

PP. 183-211

DATE (Mar. '53)

ABST. REF.

VOL. PP.

DATE

Some Clinical Observations of Asbestosis in Mine and Mill Workers.
P. Cartier.

Statistics and comments are presented on 128 cases of asbestosis observed by the author over a period of eight years. In his experience, no cases of asbestosis of clinical importance have been diagnosed by the pathologist without previous detection by the roentgenologist. The wide discrepancy in the appreciation of the degree of asbestosis by different pathologists leads to confusion, especially before a compensation board. The limitations of the table are obvious, but it shows that the clinical status of 43 cases is good and that 59 employees in the early stages of asbestosis are able to work without discomfort. For the remaining 29 cases, no good correlation between degree of asbestosis and clinical status has been found.

DATE
EXAMINED

BY

PURPOSE
OF SEARCH

ABSTRACT BLANK FOR GENERAL USE

CONT'D

CLASSIFIED SUBJECT

FELLOWSHIP
NUMBER

FILE NUMBER

572

6/55

FC 752

AUTHOR: L. J. Bristol.

TITLE: Roentgenologic aspects of silicosis and asbestosis.

CMO. REF.	VOL.	PP.	DATE	ASSN. REF.	VOL.	PP.	DATE
Arch. Dec. Health	11	183-211	(Mar. '55)				

Roentgenologic Aspects of Silicosis and Asbestosis. L. J. Bristol.

Typical roentgenograms show the appearance of silicosis and of asbestosis in the three stages of each and of silicosis with tuberculosis. The functional impairment in silicosis is due to diffuse obstructive emphysema, and x-ray examination in full expiration aids in evaluating the presence or absence of this complication. On the other hand, not all cases of asbestosis have a breathing problem. Expiration films are not of much value in determining the presence or absence of respiratory impairment in these cases. Some persons with slight x-ray changes indicating asbestosis, show definite functional impairment. This situation does not occur in silicosis.

DATE
EXAMINED:PURPOSE
OF SEARCH
ABSTRACT BLANK FOR GENERAL USE

CONT'D.

CLASSIFIED SUBJECT

FELLOWSHIP
NUMBER

FILE NUMBER

572

5/55

FC

752

AUTHOR: G. W. Wright.

TITLE: Functional abnormalities of industrial pulmonary fibrosis.

CHIEF, REPT.	VOL.	PR.	DATE	ASSN. REPT.	VOL.	PR.	DATE
Arch. Oct. Health	11	183-211	(Mar. '54)				

SCRON, PA.

Functional Abnormalities of Industrial Pulmonary Fibrosis. G. W. Wright.

The author describes the function of the respiratory and cardiocirculatory apparatus in a simplified form for the benefit of lay persons, and discusses the methods of quantitating some of the physiologic aspects of the respiratory and circulatory systems. He shows how the results of these methods indicate silicosis in its various stages. The typical appearances and clinical conditions in asbestosis are then described and contrasted with those in silicosis. As in silicosis, there is a gross correlation between intensity and duration of exposure, but no evidence is found on the least exposure necessary to produce asbestosis. The physiologic abnormalities are recognizably different in the earlier stage of the two diseases, but in the terminal stage the distinction may not be entirely clear and may lead to confusion. A further difficulty in diagnosis is the frequent coexistence of the diseases. Both may be roentgenologically manifest without recognizable physical impairment.

DATE
EXAMINED.

BY

PURPOSE
OF RESEARCH
ABSTRACT BLANK FOR GENERAL USE

CONT'D.

572

5/55

FC

752

AUTHOR: A. J. Lanza.

TITLE: Recent trends in industrial health.

ORIGIN: PA.

ORIG. REF.	VOL.	PP.	DATE	ABST. REF.	VOL.	PP.	DATE
Arch. Occ. Health	11	103-211	(Mar. '55)				

Recent Trends in Industrial Health. A. J. Lanza.

This short address serves as an introduction to the scientific papers and emphasizes the need of continued progress in the study of dust diseases.

MILLON INSTITUTE

FORM 77 30

DATE EXAMINED	BY	PURPOSE	CONT'D.
		OF J. Lanza	
		ABSTRACT BLANK FOR GENERAL USE	

1113 4/524

NFC 752

AUTHOR R. Doll.

TITLE Mortality from lung cancer in asbestos workers.

CMIO DATA VOL. 12 PP. 81-86 DATE APR. 1955

ABSTRACT:

1113 Mortality from Lung Cancer in Asbestos Workers. R. Doll. Brit. J. Ind. Med. 12, 81-86 (Apr. 1955).

In necropsies of 105 persons who had been employed at an asbestos works, lung cancer was found in 18 instances. In 15 cases asbestosis was also found. These 15 workers were employed at least nine years before 1923, when dust control regulations became effective. An additional 113 men who had been exposed to asbestos dust for 20 years were followed up and their mortality was compared with that of the whole male population. Thirty-nine deaths occurred in the group whereas 15.4 were expected. The excess was entirely due to excess deaths from lung cancer (11 against 0.8 expected) and from

DATE JANUARY 1955 BY BT. PURPOSE OF SEARCHING ABSTRACT BLANK FOR GENERAL USE (OVER) CONT'D

MILLON INSTITUTE, PITTSBURGH, PA.

752

FC | 752

AUTHOR: K. M. Lynch.

TITLE: Pathology of asbestosis.

ORIG. RECD.	VOL.	PP.	DATE	JOBIN. RECD.	VOL.	PP.	DATE
Arch. Occ. Health	11	183-211	(Mar. '55)				

FROM: PA. Pathology of Asbestosis. K. M. Lynch.

The pathological appearances of asbestosis are described and illustrated. It appears that asbestos crystal fragments may traumatize living tissues by their physical qualities, and the present belief is that the larger particles do the damage, in contrast to the chemical action of silica and the harmful effects of the smaller particles.

MELLON INS

FORM 77 20

DATE EXAMINED	BY	PURPOSE OF SEARCH	CONT'D.
		ABSTRACT BLANK FOR GENERAL USE	

12-29 11/53
AUTHOR: K. W. Smith.

TITLE: Pulmonary disability in asbestos workers.

AND REFS: Arch. Ind. Health 12 98-203 Aug. '55
VOL. 12 PP. 98-203 DATE Aug. '55
VOL. 12 PP. 98-203 DATE Aug. '55

Respiratory disease experience among several thousand male and female asbestos workers in the United States and Canada is reported. Of all workers exposed to the fibers, very few develop asbestosis. This disease is insidious in onset and progresses slowly with continued exposure, causing respiratory embarrassment and cardiac failure. Expansion of the lung is difficult, and there is impaired gas transfer through the lung. Diffuse, obstructive emphysema is not common. The typical x-ray pattern

cannot be confused with other pneumoconioses. An x-ray survey of 708 employees in a milling operation showed that the majority had normal x-ray patterns, that ten or more years of exposure were necessary to produce x-ray changes, and that no cases of asbestosis were found who had worked less than 20 years in the dust. The incidence and duration of nonoccupational respiratory disease were not increased by exposure. Asbestos does not predispose to tuberculosis, nor does it aggravate an apparently healed lesion. There are several reasons for different opinions expressed concerning the relationship of asbestosis

CONF. V

and bronchogenic carcinoma. Differences in asbestos fibers were noted.
-- Cond. from author's summary

MELLON INSTITUTE, PITTSBURGH, PA.

CLASSIFIED SUBJECT

7/52 3/2

FELLOWSHIP
NUMBER

NFC

FILE NUMBER

752

AUTHOR: G. Jacob and H. Bohlig.

TITLE: Roentgenographic complications of pulmonary asbestosis.

ORIG. REF.	German.	VOL.	PF.	DATE	ASSIG. REF.	VOL.	PF.	DATE
	Fortschr. Gebiete					Am. Rev. Tuberc.	73	202
	Rontgenstrahlen	83	515-525	Oct. 1953				5/56

ABSTRACT:

Over a period of approximately twenty years, 343 cases of asbestosis, 130 in women and 213 in men, were observed in Dresden, a center for industries using asbestos. The most common complications were pleural thickening, compensatory emphysema, and nonspecific peribronchial, peribronchovascular, and bronchopneumonic inflammations. Pleural changes were found in 49 per cent of the cases, 5 per cent of which had extensive, irregularly shaped calcifications. Active pulmonary tuberculosis was present in 4.4 per cent of the cases; bronchogenic cancer was not more frequent than in the general population but, when present, was found relatively often in the lower lobe.

-- Am. Rev. Tuberc.

MILLON INSTITUTE, PITTSBURGH, PA.

FORM 77 20

DATE
EXAMINED

BY: FBI

PURPOSE
OF SEARCH:
ABSTRACT BLANK FOR GENERAL USE

CONT'D.

A Case of Pulmonary Asbestosis Accompanied by "Tuberculosis."

•

•

abstract. During the period 1933-1939 inclusive, there were 90 known deaths from asbestosis in Great Britain, of which 32 were associated with tuberculosis. This reports the case of a 44 year old man who had been an asbestos blower, supplying the material to a cleaning machine. Although supplied with a respirator, he had never worn it. He was thin, had a high temperature, emphysema, and bilateral crepitations over wide areas. Clubbing of the fingers was marked and the sputum contained both asbestosis bodies and tubercle bacilli. The roentgenogram showed widespread progressive nodular tuberculous infiltration with cavitation in all zones of the right lung fields and in the upper and mid-zones of the left lung fields.... The right dome of the diaphragm was elevated. There was no radiographic evidence of pneumoconiosis." A diagnosis of asbestosis and tuberculosis was made. The patient died within a month of spontaneous pneumothorax. Pulmonary findings showed both to have extensive fibro-casous tuberculosis with cavitation; macroscopically they were like those of pure tuberculosis. Microscopically, there was revealed caseous tuberculosis with many tubercle bacilli and typical reticular fibrosis of asbestosis with thickened pleura. There is a discussion of the diagnosis of the pure and complicated disease.

1

ABSTRACT BLANK FOR GENERAL USE

Dept 44, Abt # 711 753 753

AUTHOR: M. Nordmann and A. Sorge

TITLE: Pulmonary Cancer Produced by Asbestos Dust in Experimental Animals.

ORIG. REF. Zeltschr. f. Krebsforsch. VOL. 51, 163-182, 1941

ABSTRACT: Experimental work with mice inhaling asbestos dust over a period of from 47 to 90 days for 3-1/2 to 5 hours a day produced asbestos bodies and pronounced degrees of diffuse fibrosis after the 20th week. Proliferation of the bronchial epithelium was found, such hyperplasia encroaching on neighboring alveoli. This was followed by a metaplasia to pavement epithelium with cornification, which later seemed to be a transition to malignant neoplasia. The first well-developed carcinoma was found after 240 days. By this time only 10 of the mice had survived and 2 carcinomas were found among them. In the first series, 33 of 60 mice lived longer than 6 weeks; 14 showed abnormal growth of the epithelium, 5 metaplasia and cancer. In the second series, 28 (of 60) lived more than 6 weeks; 16 showed abnormal growth, 5 metaplasia and cornifying cancer.

DATE EXAMINED: 11/1/51

ABSTRACT BLANK FOR GENERAL USE

MELLON INSTITUTE, PITTSBURGH, PA.

MELLON INSTITUTE, PITTSBURGH, PA.

153
AUTHOR: H. H. Holleb and A. Angerist
TITLE: Bronchogenic Carcinoma in Association with Pulmonary Asbestosis. Report of Two Cases.

ORIG. REF.:
Am. J. Pathology

VOL. 18
PP. 123-125
DATE Jan. 1942

ASSN. REF.:

VOL. PP.

ABSTRACT: Two cases are reported in which men between 40 and 60 were found to have both undifferentiated asbestosis and a primary carcinoma of the lungs. A table listing 10 reported cases, collected from the literature, in which both diseases occurred, is presented. The authors conclude that the data from 2 cases, with some reports in the literature in doubt, are insufficient for a statistical evaluation, and that caution should be used in ascribing carcinoma to asbestosis. In one case the carcinoma was in a region least affected by asbestosis, but when carcinomas often occur.

DATE EXAMINED:

BY: FFR-JIH

PURPOSE OF SEARCH:

ABSTRACT BLANK FOR GENERAL USE

CONFID.

5/43-#480

752

FELLOWSHIP
NUMBER

FILE NUMBER

-753

AUTHOR: J. Kuehn

TITLE: Electron Microscopy on Asbestos Dust and on Lungs in Asbestosis.

ORIG. REF.	VOL.	PP.	DATE	ANAL. REF.	VOL.	PP.	DATE
Arch. f. Gowerbe-	10	473	April 6,				
path.			1942				

The author reports some microscopic studies of asbestos and of lungs of asbestosis victims for the purpose of determining what variety of asbestos is responsible for asbestosis. Serpentine (chrysotile) asbestos is the form generally used, and it was supposed that it would be found in lungs with asbestosis. Sundius and Bygden disclosed, however, that only hornblende asbestos is detectable in the lungs, suggesting that serpentine dust is decomposed in the tissues. The author used the electron microscope to study the form, size and structure of asbestos, both in free dust and as found in the lungs. It is usually easy to distinguish the two forms under the electron microscope, the serpentine variety being generally curved, clear and transparent, and hornblende rigid

MELLON INSTITUTE, PITTSBURGH, PA.

and greater in diameter. However, examination of the particles from 3 lungs did not indicate clearly what variety was present. The photomicroscopic determination of the refraction exponent, suggested by Sundius and Bygden, was then employed. That method revealed that the large asbestos needles isolated from the lungs were of the hornblende type. It is concluded that serpentine asbestos is decomposed in the lungs and is the essential cause of fibrosis. When asbestos lungs are macerated with nitric acid and hydrogen peroxide, both free asbestos needles and asbestos bodies are obtained. Hornblende asbestos shows no signs of decomposition by the organism, either in the free needles or on the asbestos bodies.

MEILLON INSTITUTE, PITTSBURGH, PA.

AUTHOR: A. Bolme

TITLE: Examination of workers in an asbestos factory.

ORIG. FILE: Arch. f. Gererobachth. 11 435-452 1942
 VOL. 11 435-452
 DATE: Nov. 10, 1942

ABSTRACT:

This is a record of clinical and radiological observations made in an asbestos factory during the years 1936-1937 and the year 1940. The factory handled

raw asbestos and manufactured a variety of asbestos products. It employed up to 500 workers. Of 132 persons examined, 29 per cent showed radiological evidence of asbestosis. Correlating these findings with exposure, it is stated that there were in workers with less than three years of exposure 5 per cent of cases, in those with 3-5 years of exposure 12 per cent, in 5-10 years 36 per cent, and in over 10 years 79 per cent. Five cases of pulmonary tuberculosis were noted. The clinical histories of these are recorded.—

DATE: _____ BY: _____ PURPOSE: _____ CONT'D: _____
 RESEARCH: _____ OF RESEARCH: _____

ABSTRACT BLANK FOR GENERAL USE

MELLON INSTITUTE, PITTSBURGH, PA.

Sept 12 - 22 # 834

752

752

AUTHOR. L. U. Gardner

TITLE. Chrysotile Asbestos as an Indicator of Subtle Differences in Animal Issues.

ORIG. REPR. Am. Rev. of Tub.

VOL. 45

PP. 762

DATE June, 1942

VOL.

PP.

DATE

ABSTRACT. In the course of experiments upon the action of asbestos fibres upon living tissue it has been observed that this irritant is not like finely divided quartz, which provokes specific cellular responses in any organ of all species of animals investigated (tadpoles, fish, chickens, mice, rats, guinea pigs, rabbits, cats, dogs, monkeys and man). Asbestos fibres are acted upon to produce the "bodies," described by Cooke, only in the lungs of a few species like man, guinea pigs and white mice. They do not form in other organs of these species nor have they been observed in the lungs of white rats or dogs; in rabbits' and cats' lungs there are occasional atypical ones. Fibrous tissue reaction is also limited to the lungs, but none at all has developed in those of mice and dogs. The subtle differences in tissues responsible for such variation have not been identified but they may be of significance to the general problem of host resistance to other irritants like tubercle bacilli.

Authors summary)

DATE RECEIVED

BY FRH

PURPOSE OF SEARCH

CONFIDENTIAL

ABSTRACT BLANK FOR GENERAL USE

MELLON INSTITUTE, PITTSBURGH, PA.

AUTHOR: A. Welz

TITLE: Further Observations on Occupational Carcinoma in Asbestos Workers.

ORIG. REF.	VOL.	PP.	DATE	ABSTR. REF.	VOL.	PP.	DATE
Arch. f. Gewerbehyg.	11	536-560	Nov. 10, 1942				
U. Gewerbehyg.							

ABSTRACT: Abstract filed under 471

DATE EXAMINED:

BY: FRL-JIH

PURPOSE OF SEARCH:

CONT'D:

ABSTRACT BLANK FOR GENERAL USE

CLASSIFIED SUBJECT

FELLOWSHIP
NUMBER

FILE NUMBER

7/43-#7121

589 152

152

AUTHOR: Illinois

TITLE: Wartime Operations Emphasize Industrial Hygiene Problems of Asbestos

Industry

ORIG. REFS:

VOL.

PP.

DATE

ABSTR. REFS.

VOL.

PP.

DATE

Illinois Labor Bull

3

10, 11

Feb. 28, 1943

ABSTRACT:

The control of asbestosis must be an engineering control because there are no medical means to detect the disease until much harm has been done. Engineering control consists of two important principles: (1) Adequate exhaust; 2. Good housekeeping. By experiment it has been estimated that if the air contains less than 5,000,000 particles of asbestos per cubic foot there is small chance of the workers developing asbestosis. The article lists the procedures in the manufacture of rigid and of textile asbestos products and shows that further protective measures are still needed in most plants. —John C. Timbille.

MELLON INSTITUTE, PITTSBURGH, PA.

FORM 10-10-31

DATE EXAMINED:

BY: FJR-JIH

PURPOSE OF SEARCH:

ABSTRACT BLANK FOR GENERAL USE

CONFIDENTIAL

MELLON INSTITUTE, PITTSBURGH, PA.

258 3/44 / 753

FELLOWSHIP NUMBER 753

AUTHOR: R. Desmeules and others

TITLE: Asbestosis and Pulmonary Tuberculosis

ORIG. REF.: VOL. 8 PP. 217 DATE Mar. 1943

Leval. Med. (French)

ABSTRACT:

Asbestosis is an occupational disease caused by the inhalation of asbestos dust. It is diagnosed by the presence of asbestos bodies in the sputum, and is characterized by the transformation of normal lung tissue into fibrous tissue. Symptoms appear usually five to seven years after the beginning of exposure. Both asbestosis and tuberculosis were seen in 8 workers exposed to asbestos dust for 20 years. In three of those cases asbestosis was masked by tuberculosis.

DATE EXAMINED: BY: JHR

PURPOSE OF SEARCH:

CONT'D.

ABSTRACT BLANK FOR GENERAL USE

2921 5/14
AUTHOR: L. LOUSSEAU

753

NUMBER

753

TITLE: PULMONARY ASBESTOSIS

ORIG. REF.:

VOL. 9

PP. 233

DATE: MAY. 1943

VOL.

PP.

DATE

Level Med.

(French)

ABSTRACT:

The radiological picture of pulmonary asbestosis is not characteristic; it is usually similar to that of tubercular fibrosis. The diagnosis is, therefore, often made only after asbestos bodies have been found in the sputum. Asbestos fibers may be seen in the sputa of individuals who do not suffer from asbestosis; asbestos bodies are due to a pathologic tissue reaction. Microscopical examination of the lungs of patients who have died from an intercurrent disease may show asbestosis, which had not been found on X-ray examination. ~~Asbestos bodies have been found in the sputum of patients who have died from an intercurrent disease.~~

and gastric examinations were negative for tubercle bacilli. Asbestos bodies were found in the sputum.

~~Asbestos bodies have been found in the sputum of patients who have died from an intercurrent disease.~~ The case history of a forty-two year old man with asbestosis without tuberculosis is given. The patient had been working in the asbestos industry for eighteen years. For two years the patient had been suffering from increasing cough, expectoration and dyspnea. Roentgenological examination showed mottling of both lungs. Sputa

MELLON INSTITUTE, PITTSBURGH, PA.

DATE: BY: FFM-JIH

PURPOSE OF SEARCH:

CONT'D.

ABSTRACT BLANK FOR GENERAL USE

FELLOWSHIP NUMBER

FILE NUMBER

290 244 ✓ 100 1/53

AUTHOR: M. GLEUX

TITLE: Asbestos Bodies

MELLON INSTITUTE, PITTSBURGH, PA.

FORM 77 10M 3-43

DATE EXAMINED:	ORIGINAL NO.	VOL.	PP.	DATE	ABSTRACT	VOL.	PP.	DATE
	Level Med.	0	239	Aug. 1943				
	(French)							

ABSTRACT:

The "curious bodies" found in the sputum of patients suffering from pulmonary asbestosis usually measure 24 to 39 μ by 12 to 21 μ but can be as large as 250 μ . They consist of a central fiber of asbestos which is covered by a golden-yellow substance containing iron. Rabbits, guinea pigs and one dog were exposed to inhalation of asbestos dust. Asbestosis did not develop in the rabbits nor in the dog. The guinea pigs showed, after several weeks, macrophages with asbestos fibers in their alveoli. Erythrocytes appeared in the alveoli, and hemophilin derivatives containing iron could be found on asbestos fibers two to three months after the beginning of the inhalation of the asbestos dust. When a suspension of asbestos in olive oil was injected subcutaneously or into the testicles of guinea pigs no asbestos bodies were formed. However, if first one cc. of blood was injected and then the asbestos dust through the same needle, asbestos bodies could be found one month later. The presence of asbestos bodies in the sputum is believed to be a definite sign of pulmonary asbestosis.—Am. Rev. Tuberc.

by: FRI-JIH

PURPOSE OF SEARCH:

ABSTRACT BLANK FOR GENERAL USE

CONT'D.

II. Descentules and II. Gironx
asbestosis and pulmonary tuberculosis

ORIG. PERG.	VOL.	PP.	DATE	ADDM. REF.	VOL.	PP.	DATE
Loyal, died. (French)	8	227	Mar. 1943				

The case history of a 44 year old man is given who had worked for 20 years in an asbestos mill. After 10 years he had developed productive cough which increased gradually. Shortly before he was admitted to the hospital he started to run fever and had pains in the chest. The clinical diagnosis of ulcers-fibrosis tuberculosis and asbestosis was confirmed by the autopsy. Diffuse peribronchial and perivascular sclerosis was also seen. It is believed that the first symptoms of asbestosis appeared ten years before the symptoms of tuberculosis, but the matter could not be decided radiologically and pathologically.

DATE: _____
EXAMINED: _____

NY. **PRIN**

**PURPOSE
OF SEARCH**

ABSTRACT BLANK: FOR EXPERIAL USE

CONT'D

ASBESTOSIS and pulmonary carcinoma.

ORIG. REF.: Deut. med. Wochenschr. (German) VOL. 67, PP. 575-576, DATE Aug. 6, 1943, ABSTR. REF.: JHEP VOL. 26, PP. 183, DATE Nov. 44

ABSTRACT:

Asbestosis and Pulmonary Carcinoma. H.W. Wedler. Deut. med. Wochenschr., 69, 575-576 (Aug. 6, 1943). German.

The author has collected the published autopsy records on asbestosis from various countries and finds that there have been 14 instances of malignant disease of the lungs and pleura in 92 post-mortem examinations (16 percent). This is in excess of the proportion of lung carcinoma in autopsies generally (2-6 percent). Carcinoma as a complication of asbestosis occurred most frequently in males between 35 and 41 and was generally in the part of the lung most affected with asbestosis. Development of the lesion was slow but corresponded with the length and intensity of the exposure (12-42 years). There was often a long interval between the cessation of exposure and the development of the cancer, which, in the author's view, must now be regarded as an occupational cancer. -- Bull. Hyg.

FFR:Bull Hyg.

DATE EXAMINED:

BY:

PURPOSE OF SEARCH:

CONT'D:

ABSTRACT BLANK FOR GENERAL USE

MELLON INSTITUTE, PITTSBURGH, PA.

FORM 77

102 0-24

MELLON INSTITUTE, PITTSBURGH, PA.

AUTHOR:

F. Homburger

TITLE: Coincidence of Primary Carcinoma of the Lungs and Pulmonary Asbestosis

ORIG. REF.:

VOL.

PT.

DATE

ABSTR. REF.:

VOL.

PT.

DATE

ART. J. Path.

19

797-807

Sept. 1943

ABSTRACT:

In the medical literature there are now at least 19 known cases of asbestosis associated with primary pulmonary carcinoma. In 4137 autopsies at the pathology laboratory at Yale University School of Medicine, there were 45 cases of pulmonary carcinoma in the years from 1918 to 1938. Asbestosis was diagnosed 8 times, all in 17 times. Pulmonary carcinoma was found in 4 of the 8 cases of asbestosis and twice in the 17 silicotic cases.

Despite the high correlation of the foreign body asbestosis in the lungs with primary pulmonary carcinoma, this is a small series of cases and the value of these data is limited. Three cases are given in detail. The diagnosis of pulmonary asbestosis was diagnosed according to the following criteria: 1) fibrosis of the lung and 2) presence of asbestosis bodies. —G. J. Path. L.

DATE EXAMINED:

BY: FER-JIH

PURPOSE OF SEARCH:

ABSTRACT BLANK FOR GENERAL USE

CONT'D

4436-3/45

NUMBER
NFC 752

AUTHOR: M. Cloux

TITLE: Experimental Asbestosis: Pathogenomic Value of Asbestos Bodies

ORIG. REF.: Vol. 8, 239-50, 1945
DATE: 1945
ABST. REF.: C.A.
VOL. 39, 366, Jan. 29, 1945

ABSTRACT:

Experiments with animals indicate that asbestos bodies form only after intense tissue reaction and with an appreciable deposit of iron furnished by hemoglobin. The presence of asbestos bodies in the sputum or in fluid obtained by pulmonary puncture indicates pulmonary asbestosis but not the degree of pulmonary sclerosis since the degree of fibrosis may vary. — C.A.

DATE RECEIVED: C.A.:GMR

PURPOSE OF SEARCH:

ABSTRACT BLANK FOR GENERAL USE

CONT'D.

AUTHOR: F. I. JOSEPH
AUTHOR'S POSITION:
AND ADDRESS:

Title: Asbestos Bodying.

MELLON INSTITUTE OF INDUSTRIAL RESEARCH
PITTSBURGH, PA.

FORM 78 10-1-33

PAGES	ILLUSTRATIONS	GENERAL	THEORETICAL	ECONOMIC	TECHNICAL
DISCUSSION (if any)	MAPS	REVIEW	MATHEMATICAL	ADMINISTRATIVE	GOOD
REFERENCES	ECST DATA	POPULAR	EXPERIMENTAL	PROPAGANDA	PAIR
CONCLUSIONS	AUTHOR INDEX	ENGINEERING	STATISTICAL		USELESS FOR PURPOSE
TABLES	SUBJECT INDEX	ORIGINAL AVAILABLE AT:			
GRAPHS		CALL NUMBER			

ABSTRACT: This paper gives a detailed study of the optical and histological nature of asbestos bodies with excellent photomicrographs. Chrysotile-asbestos was identified in a lung deposit. By crystal-lattice studies it was shown that the hydrated magnesium silicate is dissolved by the acid body fluids, leaving a shell of silicic acid. Then form of the needles and their double refracting power remains unchanged. Asbestos bodies form as a result of protein absorption by the needles and subsequent conglutination as a gel. These beliefs are confirmed by ex-

DATE RECEIVED: BY: PURPOSE OF REVIEW: COMMENTS:

ABSTRACT BLANK FOR ALL PERIODICALS AND REGULARLY ISSUED PROCEEDINGS.

pariment with asbestos and albumen. The bodies are stained brown by granular streaks of ferric oxide.

AUTHOR: H. Bastenier, H. Denolin, A. de Coster, and M. Englent.

TITLE: A study of respiration in pulmonary asbestosis.

ORIG. REF.: French. = VOL. 16 PR. 546-553 DATE 1955 ABSTR. REF.: Bull. Hyg. VOL. 31 PR. 544 DATE 5/56

In order to test the respiratory value of lungs affected with asbestosis nine cases were closely studied. The seven men and two women had been exposed to asbestos-dust, 1,000 to 2,000 particles per cu. m., for 10 to 30 years. The patients displayed definite symptoms of asbestosis with troublesome cough and dyspnea, and all showed x-ray shadows like ground glass extending over both lungs. No tubercle bacilli were found in any case. A definite reduction was found in vital capacity, but the maximum expiratory volume was satisfactory. Breathing, however, was accelerated by hyperventilation. At rest breathing was sufficient and satisfactory, but on exertion dyspnea appeared, varying considerably according to the pathological lesions in the lungs. These functional changes were in no way different from those seen in other cases of pneumoconiosis. Varying emphysema was found, but no special strala was thrown upon the heart.

-- Cond. from Bull. Hyg.

MELLON INSTITUTE, PITTSBURGH, PA.

FORM 77 10M 7-57

DATE EXAMINED: ST. FFR PURPOSE OF SEARCH: ABSTRACT BLANK FOR GENERAL USE

RELATIONSHIP: FILE NUMBER

CONT'D

MELLON INSTITUTE, PITTSBURGH, PA.

CLASSIFIED SUBJECT

FELLOWSHIP
NUMBER

FILE NUMBER

NFC

753

AUTHOR. G. Jacob and H. Bohlig.

TITLE. Incidence and characteristics of lung cancer in asbestos workers.

ORIG. REF.	German.	VOL.	PP.	DATE	ABST. REF.	VOL.	PP.	DATE
Arch. Gewerbehyg.	14	10-28	(1955)	Bull. Hyg.		30	1080	12/55
Gewerbehyg.								

This is a study of the incidence of lung cancer among asbestos workers in Dresden. The international literature is based exclusively on the results of autopsies; the incidence among asbestos workers appears to vary between 12 and 17%. In Dresden, however, from research into cases at the clinic and among asbestos workers, it appears to be rare, it being estimated that only about one-fourth of these workers have an effective exposure to the dust. In cooperation with the radiological and radium institutes the authors reviewed the cases of asbestosis reported since the middle of the 1930s. These numbered 339 in all stages; they are classified according to stage and sex. Siliacosis was found in eight males. In five females and ten males (4.4% of the total) active open tuberculosis was found. Four fatal cases of lung cancer in asbestos workers, which occurred in Dresden up to 1954, are described. A long discussion of the literature on cancer among asbestos workers includes an examination of the characteristics of the disease; these are no longer convincing evidence of a high incidence of cancer in asbestosis. In the future, cases of cancer of the lung in asbestosis should not be published individually but should always be considered in the framework of the whole asbestos industry in the region.

-- Cond. from Bull. Hyg.

DATE EXAMINED	BY	PURPOSE OF SEARCH	CONT'D.
	ACB		
ABSTRACT PLANNED FOR GENERAL USE			

MELLON INSTITUTE OF INDUSTRIAL RESEARCH
PITTSBURGH, PA.

CLASSIFIED SUBJECT

Abst. 460 June '40

752

NUMBER

752

AUTHOR, E. C. Vigilant

TITLE Two fatal cases of pulmonary asbestosis.

DATE RECEIVED	VOL.	PT.	DATE	ABSTN. RECEIVED	VOL.	PT.	DATE
Mass. Men. Indust.	11	26-52	'40	J. I. H.	22	128	June '40

ABSTRACT: This first of these cases concerned a woman who worked for thirty years in a factory using asbestos in the manufacture of cord. The second woman worked for seven years in an asbestos factory. Her susceptibility to colds increased while she was working and she suffered all her life from bronchitis and asthma. She died at the age of fifty, with symptoms of a lung disease, nearly thirty years after having stopped work. The autopsy showed interstitial sclerosis of the lungs and asbestosis bodies.

DATE
RECEIVED

BY F. S. J. - J. H.

PURPOSE
OF STATEMENT

CONT'D.

ABSTRACT BLANK FOR GENERAL USE

MILLON INSTITUTE OF INDUSTRIAL RESEARCH
PITTSBURGH, PA.

CLASSIFIED SUBJECT

Abst. 503 July '40

752

FELLOWSHIP
NUMBER

FILE NUMBER

152

AUTHOR: L. Roethel, H. Kempf and H. K. Fedler

TITLE: Investigations on pulmonary function in asbestosis.

ORIG. REF.

VOL.

PP.

DATE

ABST. REF.

P.

VOL.

PP.

DATE

Leut. Arch. F. Klin. Med.

185 55

3/7 '40

J A M A

114

2513

6/23 '40

ABSTRACT: According to these authors, the relatively slight clinical symptoms of asbestosis are frequently in contrast to serious impairment of function. A difference between the x-ray appearance and the clinical picture is also not uncommon. It is difficult to determine the degree of functional impairment. The authors studied the oxygen consumption in nineteen cases of asbestosis. They found considerable conformity between the spirographic data and the severity of the clinical symptoms. This method permits an estimation of the pathologic changes and discloses the extent of pulmonary insufficiency.

DATE
EXAMINED:

BY: I S M - JAMA

PURPOSE
OF SEARCH:

ABSTRACT BLANK FOR GENERAL USE

CONT'D.

MILLON INSTITUTE OF INDUSTRIAL RESEARCH
PITTSBURGH, PA.

CLASSIFIED SOURCE

753

135

FELLOWSHIP
NUMBER

FILE NUMBER

Author: J. L. H. H. H.

Title: The Dust Retaining Ability of the Nose in Asbestosis

ORIG. REF.	VOL.	PR.	DATE	ASSN. REF.	VOL.	PR.	DATE
Arch. f. Gewerbepath.	10	309-35	1940				

Abstract. The author examined the dust-retaining ability of the nose in 238 asbestos workers, using Lehmann's method. No constant relation between this ability and tendency to asbestosis could be discovered. The dust-retention capacity was below 40% in several men who showed asbestotic lung changes, but many others remained healthy in spite of a similar low capacity. The two men with severe asbestosis had a good dust retaining capacity. The author concludes that this test has no essential value and may be abandoned in pre-employment examinations of asbestos workers.

DATE EXAMINED.	BY	PURPOSE OF STATEMENT	CONT'D.
	PTL-JIH		

ABSTRACT BLANK FOR GENERAL USE

MELLON INSTITUTE OF INDUSTRIAL RESEARCH
PITTSBURGH, PA.

CLASSIFIED SUBJECT

752

APR 41, Also #2611

NAME. The Struggle Against Asbestosis Today

ORIG. REF. Arbelschutz

VOL. 191-3

DATE 1940

ASSN. REF.

VOL.

PP.

DATE

ABSTRACT. The paper is a brief description of asbestosis, rather than an account of the struggle against it. The author states that no relationship was found between nasal dust filtering ability and degree of asbestosis.

DATE RECEIVED.

BY

FER-JIH

PURPOSE OF SEARCH.

CONF'D.

ABSTRACT BLANKS FOR GENERAL USE

FELLOWSHIP NUMBER

FILE NUMBER

752

MILLON INSTITUTE OF INDUSTRIAL RESEARCH
PITTSBURGH, PA.

CLASSIFIED SUBJECT

Ab9.732,10/40

752

FELLOWSHIP
NUMBER

752

FILE NUMBER

AUTHOR: G. Motturn and E. Faglano

TITLE: Anatomy, Pathology and Pathogenesis of Pulmonary Asbestosis

ORIG. REF.	VOL.	PT.	DATE	ABST. REF.	VOL.	PT.	DATE
Rass. med. Indust.	11	233-284	1940				

ABSTRACT: These are the histopathological findings in two cases of asbestosis, both in women. There is also a discussion of the subject of asbestosis bodies.

DATE
EXAMINED

BY

(JH)

PURPOSE
OF SEARCH

ABSTRACT SLANT FOR ANNUAL VOL

CONT'D.

MELLON INSTITUTE OF INDUSTRIAL RESEARCH
PITTSBURGH, PA.

CLASSIFIED SUBJECT

752

FELLOWSHIP
NUMBER

FILE NUMBER

Apr '41, also # 213

AUTHOR: Anonymous

TITLE: Methods of Avoiding the Dust Danger in Asbestos Plants, Valid from

August 1, 1940.

ORIG. REFS:
Arbolschutz

VOL. 263-5
DATE 1940

ABST. REFS.
VOL. PP. DATE

ABSTRACT: The new regulations concerning exhaust systems, established by the manufacturers, insurance association and German factory inspectors, are described. The efficiency of the systems must be tested by a representative of the insurance association.

DATE
EXAMINED:

FFR-JIN

PURPOSE
OF SEARCH:

ABSTRACT BLANK FOR GENERAL USE

CONT'D.

MILLON INSTITUTE OF INDUSTRIAL RESEARCH
PITTSBURGH, PA.

CLASSIFIED SUBJECT

752

753

APR 4, 1941
AUTHOR: Brachmann

TITLE: Asbestosis in Grinders and Drillers of Brake Bands.

ORIG. REFS.

VOL.

PP.

DATE

ABST. REFS.

VOL.

PP.

DATE

Arbeitschutz

172-4

1940

ABSTRACT: Brake bands are made of sheets of asbestos, or are pressed out of asbestos waste products and a filler, impregnated with artificial resin. Afterwards the bands and plates are ground and drilled. In Germany all workers in asbestos plants must be examined annually. Clinical and x-ray examination of the grinders, drillers and men who file individual pieces has shown a few light cases of asbestosis after 2 years' work, and in nearly all such workers after 5 years employment.

DATE
RECEIVED

BY:

FTR-JIH

PURPOSE
OF SEARCH

ABSTRACT BLANK FOR GENERAL USE

CONF'D.

MELLON INSTITUTE OF INDUSTRIAL RESEARCH
PITTSBURGH, PA.

CLASSIFIED SUBJECT

752

FELLOWSHIP
NUMBER

FILE NUMBER

752

AUTHOR, A. Wolff

TITLE, Asbestosis.

ORIG. REF.

Nord.Hyg.Tid.

Z.O.U.

VOL. 21

1-33

27/193-4

DATE

1940

ASSN. REF.

C. A.

VOL. 35

3357

DATE

5/20/41

Abstract. The author describes three cases of asbestosis which occurred in Norway in factories where asbestosis was used as packing material for acid containers. In one case asbestos particles were present in the expectoration. The disease has not hitherto been observed in Scandinavia.

DATE
EXAMINED

BY

FTR-CA

PURPOSE
OF SEARCH

ABSTRACT BLANK FOR GENERAL USE

CONFIDENTIAL

CLASSIFIED SUBJECT

Feb 43 182 / 708

FELLOWSHIP
NUMBER

FILE NUMBER

752

AUTHOR: Dr. J. Lindbach and H. W. Hedler

TITLE: Occupational Cancer Among Asbestos Workers.

ORIG. REFS.

VOL.

PP.

DATE

ASST. REFS.

VOL.

PP.

DATE

Virchow Arch. f. 307 387-409 1941

Orth. Anat.

ABSTRACT: A 61 year old worker who spent 18 years in an asbestos plant contracted asbestosis in the earlier years of occupation there. Two years after third stage asbestosis had been recognized radiologically, a sharply defined, intensive, nearly homogeneous shadow about two inches in diameter appeared above the right diaphragm. Development of a large cavity three months later led to the diagnosis of cancer, which was confirmed on post mortem examination, which also disclosed asbestosis, nodules scattered irregularly throughout the lungs. The cancer was a squamous cell carcinoma. From this case and a study of the literature the author concludes that asbestosis creates a local disposition to cancer, one factor of which is mechanical irritation from hornblende asbestos and the other chemical irritation from silicic acid derived from the easily soluble serpentine asbestos.

DATE
RECEIVED

BY: PER-JIH

PURPOSE
OF SERVICE

ABSTRACT BLANK FOR GENERAL USE

MARION LINDBACH, PITTSBURGH, PA.

Oct 11, 1941
AUTHOR: J. Kühn

ULTRAMICROSCOPIC EXAMINATION OF ASBESTOS DUST AND ASBESTOTIC LUNGS.

ORIG. REFS. VOL. 10, 473-485, 1941
ARCH. F. Gewerbe- path.
DATE AREA, REFS. VOL. DATE

ABSTRACT: Interesting pictures of different kinds of asbestos and asbestos dust from lung tissue are presented. Magnifications are from 10,800 to 29,000 times. Very small needle-shaped particles up to 6 millimicrons in diameter and 400 millimicrons in length have been found in the lungs. These particles seem to be hornblende -- serpentine seems to be dissolved. Filters of respirators do not exclude particles less than 0.2 micron and hence cannot prevent these tiny particles from reaching the lungs.

DATE EXAMINED BY FSN-JIH
PURPOSE OF SEARCH
ABSTRACT BLANK FOR GENERAL USE

CONF'D.

MELLON INSTITUTE, PITTSBURGH, PA.

FORM 77

13M 4-

MELLON INSTITUTE, PITTSBURGH, PA.

CLASSIFIED SUBJECT		752		FELLOWSHIP NUMBER		752	
AUTHOR: P. Filman							
TITLE: Pulmonary Asbestosis							
ORIG. REF.	VOL.	PP.	DATE	ASSIN. REF.	VOL.	PP.	DATE
Proc. Roy. Soc. Med.	34	557	July, 1941				
ABSTRACT: This woman worked for a period of one year, thirty-one years ago sewing asbestos mattresses. Her sputum still contains asbestosis bodies. She has had dyspnoea, lassitude, cough and loss of weight. There is impairment at the base of both lungs, crackling rales and diffuse basal fibrosis, with some mid-zone involvement.							
DATE EXAMINED:	BY: FCM-JIH	PURPOSE OF SEARCH:		CONF'D.			