

MCA

MANUFACTURING CHEMISTS ASSOCIATION CHEMICALS  
RESEARCH

1825 CONNECTICUT AVENUE, N.W., WASHINGTON, D.C. 20009

APR 19 1979

TELEPHONE: (202) 328-4200

TELEX: 89617 (MCA WSH)

FK \_\_\_\_\_

OCK \_\_\_\_\_

AJLu \_\_\_\_\_

DJR \_\_\_\_\_

CMS \_\_\_\_\_

WRBe \_\_\_\_\_

ATCL \_\_\_\_\_

\_\_\_\_\_

LM \_\_\_\_\_

April 16, 1979

TO: Vinyl Chloride Technical Panel Members Sponsoring  
University of Louisville Research

Gentlemen:

Enclosed is your copy of the annual report covering 1978-79 research completed under the MCA agreement with the University of Louisville entitled "Research Techniques and Methods for Detection and Prevention of Carcinogenesis in Industrial Workers".

To avoid possible confusion in subsequent references to this document by University of Louisville personnel, be apprised of the possibility that they might refer to it as the "Second Annual Report from the University of Louisville Chemical Monomer Group".

Currently, it is anticipated that research being performed under the 1979-80 agreement will be covered in a report to issue during April or early May, 1980.

Sincerely,



J. T. Seawell  
Project Administrator  
Vinyl Chloride Research

JTS:pmt  
Enclosure

VVC 000004839



University of Louisville  
Health Sciences Center

School of Medicine  
Department of Medicine  
Division of Digestive Diseases and Nutrition

Louisville, Ky. 40232  
P.O. Box 35260  
Walnut & Preston Streets

April 6, 1979

Mr. Joseph T. Seawell  
Program Manager  
Manufacturing Chemists Association  
1825 Connecticut Avenue, N.W.  
Washington, D.C. 20009

RE: Annual Report for the Manufacturing Chemists Association's Agreement  
with the University of Louisville for the 1978-79 Fiscal Year.

Dear Mr. Seawell:

The following describes what has been completed during the second year of the Manufacturing Chemists Association's agreement with the University of Louisville entitled, "Research Techniques and Methods for Detection and Prevention of Carcinogenesis in Industrial Workers". Progress for each technical proposal will be reported separately.

Technical Proposal A -- Immunological Systems for the Detection of Vinyl Chloride and Other Chemical Injury. H. P. Fortwengler

During the second year, Fortwengler's laboratory has continued evaluating three independent aspects of the immune system of industrial workers to determine whether cancer can be detected earlier or to identify those at high risk.

Part I. Evaluation of Immunocompetence of Humans Chronically Exposed to Vinyl Chloride.

It has been demonstrated that lymphoid cells (T cells) can be cytotoxic to human tumor cells and are often found decreased or poorly functioning in cancer patients resulting in various degrees of immunodepression. The purpose of this study was to determine the immunocompetence of individuals that have undergone prolonged exposure to VC monomer and have developed liver lesions which, in some instances, are thought to presage the development of the cancer angiosarcoma.

The scientific literature is replete with demonstrations of immunodepression in cancer patients at various stages of disease. There is very little information, however, concerning

VVC 000004840

immunoevaluation prior to diagnosis of frank malignancy. We used immunological assays that have demonstrated usefulness in indicating immunodepression in cancer patients. These tests include the enumeration of lymphocytes and lymphocyte function assays. (For details, see report for fiscal year 1976-77).

These tests have been used to evaluate the immunocompetence of individuals with possible pre-malignant lesions or other liver disease as demonstrated by biopsy. A comparison of the employees with demonstrated disease versus those employees with no clinical or laboratory evidence of disease showed no immunological difference between the two groups. A comparison of VC workers having high VC exposure (those with lifetime exposure above the plant median VC exposure) with individuals having low exposure (below the plant median) demonstrated a slight pattern of immunodepression when lymphocytes were stimulated by PHA and Con-A. However, further evaluation of this data, even after additional immunological parameters were examined, indicated there is no statistical significance between the high and low exposure groups (1).

At this time, our preliminary interpretation of these results is that there is no residual immunological depression as a result of chronic exposure to increased levels of VC as determined by the standard battery of immunological tests.

#### Part II. HLA Frequencies in V.C. Workers.

An increased incidence of certain HLA types has been shown by several reports to be associated with susceptibility to various diseases. The first occupational disease-HLA correlation may have already been found: HLA-B27 antigen has been found more often in workers suspected of having the occupational disease asbestosis than among a control population.

The microdroplet lymphocyte cytotoxicity test has been used in our laboratory to HLA type approximately 429 individuals from the Louisville vinyl chloride polymerization plant. As individuals do not change their genetic complement of HLA antigens, these determinations need to be done only once and do not need to be repeated periodically as in various other biochemical assays.

Tissue typing for 11 separate HLA-A antigens and 16 HLA-B antigens and their possible increased association with angiosarcoma or other chemically-related diseases is about two-thirds completed.

Mr. Joseph T. Seawell  
April 6, 1979  
Page 3

Our compiled data is being compared with the frequencies obtained by two other large HLA studies. Frequencies of the healthy individuals studied by Scott et al., 1977, and the World Health Organization compare favorably to the frequencies found in the study here at the University of Louisville.

A pattern of potential differences from normal frequencies has been seen in VC workers found to have liver disease. Although definitive statistical analysis will not be done until the study is completed, differences are being seen in the frequencies of antigens A9, A13, A15, and B17. Whether these particular antigens may be used to identify individuals susceptible to chemical injury will have to await completion of the study.

### Part III. A Search for Evidence of a VC-Induced Tumor Antigen.

New antigens arise on tumors formed as a response to carcinogens. Their presence on methylcholanthrene-induced sarcomas was discovered by Foley in 1953. This discovery in mice was verified and extended by Prehn and Main in 1957 to conclude that there were antigens peculiar to and specific for tumor tissue. Subsequently, evidence for tumor antigens was found in humans by the Hellstroms, Vankey, Halliday and Maluish, Thompson and others. The majority of the evidence suggested that the tumor antigens found were distinctive for each histological type of tumor.

Since the body mounts an immune reaction to cancer, a cell-mediated test system (lymphocyte transformation) was used to search for evidence of specific immune reactions. Lymphocytes (the cells responsible for immunity) from the individual tested were isolated by centrifugation over Ficoll-Hypaque. These cells were then grown in the presence of a liver reagent prepared from either a normal individual or an individual who had angiosarcoma. A three-times increased incorporation of  $H^3$ -thymidine into stimulated cultures as compared to unstimulated cultures is a positive reaction.

A comparison of the responses between vinyl chloride plant workers and normal non-chemical plant workers indicated that there were many people in the general population having reactivity to tissue antigens irrespective of whether these antigens were from liver angiosarcoma or normal liver. Non-tumor specific reactions of this type may be due to sensitization by "natural" means, injections of human or animal substances, transfusions, etc.

VVC 000004842

Vinyl chloride workers were classified according to known VC exposure and tested for reactivity to reagents prepared from normal liver or angiosarcoma liver. Lymphocyte responses from all individuals tested having low VC exposure (below plant median exposure) were compared to responses from all individuals having high exposure. No quantitative statistical differences were noted between the reactivities of the two groups.

When viewed qualitatively, concomitant reactions by an individual's lymphocytes to both normal liver and angiosarcoma liver cannot be interpreted. Angiosarcoma reagent contains both normal and tumor antigen. Therefore, only those remaining individuals with reactions to either normal liver or angiosarcoma liver alone were compared further. The composition of that group of individuals reacting to the angiosarcoma liver reagent was striking -- the only reactions obtained against this tumor preparation were from individuals with high VC exposure. That is, the individuals reacting to the tumor antigen reagent were all from the high risk group.

These results must be interpreted with caution because of the small number of individuals having reactions to only tumor antigen. We are now in the process of testing control subjects in replicate to determine if test and operator variations can be further minimized. Repeat determinations of selected exposed individuals will, as a consequence of reduced variation, take on greater statistical meaning.

In addition to testing for evidence of reactions to an "angiosarcoma antigen", we have found that the tumor cells contain antigenic coagulation Factor VIII. Factor VIII has been found in tumor tissue from three individuals having had angiosarcoma. Hoyer's finding in 1973 that only endothelial cells contain Factor VIII, when combined with our findings, indicate that the specific lining cell which becomes aberrant during the course of angiosarcoma is the endothelial cell (2).

Technical Proposal B -- Biochemical Enzymatic Systems for the Detection of Vinyl Chloride and Other Chemical Injury and Cancer Development in Industrial Workers. J. T. Du

#### Animal Studies

Animal studies are continuing to be conducted to determine which biochemical changes occur early in the course of vinyl chloride exposure and how the determination of these biochemical changes can be used to a) detect early injury, b) to demonstrate a level of exposure with no biological effect.

VVC 000004843

The previous long-term experiment in Du's laboratory identified those metabolic pathways which best reflected vinyl chloride handling. The second phase of this study, extending exposure to over 250 hours, has verified the original findings and illustrated adaptation of the detoxifying mechanisms of rat liver with prolonged exposure at high levels (3).

Our second vinyl chloride exposure study exposed rats to 28,000 ppm VC for 70, 140 and 210 hours in durations of 2, 4 and 6 weeks. We examined the rat liver's ability to oxidize vinyl chloride by study of the microsomal P-450 enzyme system (mixed function oxidase) and detoxification mainly by conjugation via glutathione (GSH) and glutathione transferase. The results showed an elevation of glutathione reductase in the liver (the enzyme which regenerates reduced glutathione from its oxidized form), followed by an elevation of the concentration of reduced glutathione. This was later followed by an elevation of the detoxifying enzymes, glutathione epoxide-S-transferase (GEST) and glutathione aralkyl-S-transferase (GAST). These results suggest that the rats had the capacity to induce detoxifying enzymes as well as to maintain high glutathione concentrations to strengthen detoxification capability. These biochemical changes were evident while the conventional clinical liver tests showed no abnormalities (4).

Also, our finding a higher level of glutathione, and glutathione reductase after vinyl chloride exposure, may be an early adaptive biochemical mechanism which precedes the precancerous alteration. This is similar to Fiala's finding that administration of hepatocarcinogens to rats led to an increase in the concentration of glutathione in the liver and that the concentration remained high until the development of hyperplastic nodules (J. Natl. Cancer Institute, 57:591-598, 1976). Further, there was a decrease of P-450 concentration in the livers of rats exposed to vinyl chloride. The decreasing P-450 would help the animal to produce less toxic metabolites, and may reflect another means of biological adaptation and help understand why the liver cell does not become malignant.

The present working hypothesis to explain why the primary liver cell does not develop tumors in the adult animal is that it has the ability to adequately detoxify the carcinogenic metabolites of vinyl chloride. The adjacent sinusoidal lining cells most likely develop the tumor (angiosarcoma) because of their decreased ability to detoxify the metabolites. Further studies with isolated hepatocytes and endothelial lining cells and animal exposure studies will be performed to elucidate the mechanisms.

### Isolation Studies

Some preliminary studies of liver cell isolation have been started. Hepatocytes were isolated by collagenase perfusion and endothelial and Kupffer cells by Pronase digestion. We have successfully obtained cells of good viability. Newer techniques using ultracentrifugation have become available which will help further improve our yield and start the next phase of our studies.

Hopefully, the use of isolated various liver cells to determine their ability to oxidize and detoxify chemicals may be used to determine biological threshold limits in more realistic fashion.

### Technical Proposal C -- Glycosaminoglycan Changes in Earlier Detection of Fibrotic Injury and Hepatic Cancer. C. E. Kupchella

Glycosaminoglycans (GAGs) are involved in wound healing and scar formation (fibrosis). Certain GAGs are elevated in malignant tumors including hepatic tumors, and it has been postulated that GAGs may be important determinants of tumor cell properties. A number of laboratories, including Kupchella's, have established that urinary GAGs may serve as markers in the pathogenesis of chemical injury, fibrosis, and cancer. Although urinary GAG analyses have long been used clinically to detect and diagnose genetically-determined metabolic disorders of GAG metabolism, a systematic evaluation of urinary GAG patterns in the detection and diagnosis of acute and/or chronic necrotic and/or fibrotic liver injury -- or cancer -- has never been made.

The objective of this proposal is to determine the usefulness of urinary and tissue glycosaminoglycan measurements in the detection of chemically-induced liver injury.

Kupchella has reported a number of findings, (references 5-11), some of which are described in previously submitted MCA reports. A summary of the pertinent results follows. Abstracts previously not submitted are provided in the attached appendix.

1. Human hepatic angiosarcoma and fibrotic liver diseases are accompanied by elevated tissue GAGs. (5)
2. The GAGs in the angiosarcoma tumor tissue are different from those in fibrotic tissue adjacent to the tumor. (5)

3. Angiosarcoma and hepatoma patients have characteristic urinary GAG patterns -- patterns not found in normal controls. (6)
4. Heparin sulfate (a type of GAG) is elevated in hepatic tissue undergoing experimentally-induced fibrosis and heparin sulfate is elevated in the urine of experimental animals. (7)
5. Heparin sulfate and hyaluronic acid levels -- but not heparin -- are 3-4 times higher in experimentally transplanted hepatomas than in normal liver and, urinary excretion reflects both the tumor GAG composition and the size of tumors. (8)
6. Livers of animals bearing metastasizing hepatoma (5123tc) have 10-fold greater concentrations of a non-sulfated, neutral, uronic, acid-positive material than is found in the livers of animals bearing two other, non-metastasizing hepatomas. (8)
7. Hepatic necrosis is accompanied by significant tissue GAG elevations, but hepatic regeneration is not. (9)
8. Gross (non-fractionated) urinary GAG determinations give a better indication of liver disease than ultrasound analysis. However, modifications in GAG analysis must be evaluated further as to specificity and sensitivity. (10)
9. Exacting urinary GAG analysis (fractionated) is potentially able to differentiate active from inactive liver disease. (11)

PRACTICAL SIGNIFICANCE:

These studies address the needs for:

- a. Useful screening tests for liver injury and for active versus inactive disease -- urine tests of the type to be evaluated obviously fit the ideal of being non-invasive and having zero morbidity/mortality and not requiring "time off".
- b. Methods of therapeutic intervention, i.e., the elucidation of the role of the GAGs in the pathogenesis of fibrotic liver disease may well lead to the identification of strategies by which fibrogenesis can be blocked and/or reversed.

- c. Tests to identify individuals at risk of chemical injury -- if we are able to find characteristic CAG changes reflective of chronic alcohol injury or other similar injury, we would have a way of identifying individuals with liver impairment in the screening of job applicants.

Technical Proposal D -- Histological Systems of Detection. C.H. Tamburro, R. Schrodtt

Scar tissue (fibrosis) is a common early finding associated with chemical injury to the liver as well as other organs. Vinyl chloride and other chemicals have been shown to produce mild injury undetectable by standard biochemical means but reflected by increase in fibrosis associated with prolonged exposure.

Drs. Schrodtt and Tamburro have been developing a means of analyzing light microscopic sections of liver tissue obtained from vinyl chloride workers to determine the feasibility of quantitating the amount of scar tissue in these individuals related to their exposure. Sinusoidal cell size and collagen deposits within the sinusoidal Space of Disse have been determined by utilizing a relatively newly developed Hewlett-Packard 9864-A digitizer and a 9815-A micro computer. With this equipment, Schrodtt and Tamburro have been able to quantitate the areas of trichrome stainable collagen (fibrosis) within biopsy samples. These morphometric analyses have been done on randomly selected fields from biopsies obtained during medical evaluation in vinyl chloride workers. There are now some 110 biopsies approximately 50 of which have been reviewed. Morphometric analysis, however, has had to be delayed since there was no known standards of normal human collagen content known for human adults.

A study has been begun to determine the normal distribution and content of collagen at varying ages in normal individuals without history of chemical, viral or medical disease or injury of the liver. This study is one-third complete. Preliminary review of the data suggests that there may be an increase in the collagen deposition (fibrosis) in the normal human liver associated with age. If this holds true upon analysis after completion of the study, age corrected standards will have been established so that the data obtained from the vinyl chloride exposed human biopsies may be accurately interpreted (12).

Resumption of the morphometric analysis of the vinyl chloride exposed liver biopsies will be resumed upon completion of the normal control study.

VVC 000004847



Technical Proposal E -- Chemical Systems of Detection of Toxicity of Vinyl Chloride. J. L. Wong

Wong's current study of vinyl chloride toxicity/carcinogenicity has been concerned with (1) the chemistry of vinyl chloride metabolism, i.e., the structure of the intermediates and their reaction with cytoplasmic chemicals, and (2) the putative actions of the primary metabolites on nuclear materials. Two putative metabolites, chlorooxirane (COR) and chloroacetaldehyde (CAA), have been shown to react in different ways with sulfhydryls (detoxification study) as well as with nucleic acid constituents (mutagenesis and carcinogenesis study). Radiolabeled derivatives are being prepared to observe disposition and conversions in animals and isolated liver cells.

Results and Discussion

The detoxification studies are summarized in Table I. The central question is how are the primary metabolites 1 and 2

TABLE I. DETOXIFICATION OF VINYL CHLORIDE

|                                                                                       |                                                             |                                               |                                                                |  |
|---------------------------------------------------------------------------------------|-------------------------------------------------------------|-----------------------------------------------|----------------------------------------------------------------|--|
| PRIMARY METABOLITES                                                                   | COR <sup>1</sup><br>↓                                       |                                               | CAA <sup>2</sup><br>↓                                          |  |
| INTERMEDIARY METABOLITES                                                              | S-ACETALDEHYDE <sup>3</sup><br>THIAZENES <sup>5</sup> , --- |                                               | HEMITHIOACETAL <sup>4</sup><br>THIAZENES <sup>5</sup>          |  |
| URINARY METABOLITES                                                                   | S-ACETIC ACID <sup>6</sup>                                  |                                               | S-ETHYL ALCOHOL <sup>7</sup><br>CHLOROACETIC ACID <sup>8</sup> |  |
| 1  | 2 $\text{H}-\text{C}(=\text{O})-\text{CH}_2\text{Cl}$       | 3 $\text{RS}-\text{CH}_2\text{CHO}$           | 4 $\text{RS}-\text{CH}(\text{OH})-\text{CH}_2\text{Cl}$        |  |
| 5  | 6 $\text{RS}-\text{CH}_2\text{CO}_2\text{H}$                | 7 $\text{RS}-\text{CH}_2\text{CH}_2\text{OH}$ | 8 $\text{Cl}-\text{CH}_2\text{CO}_2\text{H}$                   |  |

(RS FROM 3,4-DICHLOROBENZENETHIOL, N-ACETYLCYSTEINE  
R' = ACETYL, H)


detoxified. Do they yield the same products or different ones? We have studied their reaction with sulfhydryl compounds 3,4-dichlorobenzenethiol and N-acetylcysteine. The benzenethiol was used by the Stockholm group in a preliminary study to detect the formation of CAA and COR from VC. The cysteine derivative is a cellular sulfhydryl component as well as a close analog of

glutathione. In the case of COR and benzenethiol, the sulfur-conjugation product S-acetaldehyde, compound 3, is formed. However, CAA and benzenethiol forms the hemithioacetal, compound 4. These two reactions are distinctly different. The formation of hemithioacetal is reversible but that of the S-acetaldehyde is not. With N-acetylcysteine, both COR and CAA yield the same cyclic condensation product, a dihydrothiazinecarboxylic acid, compound 5, in aqueous solution. This thiazene is a multi-step reaction product, formed much faster with COR than with CAA. It is plausible that 5 is the origin of the identified urinary metabolites: S-acetic acid, compound 6, and S-ethyl-alcohol, compound 7, and may itself be present in the urine. Furthermore, the COR reaction when titrated with hydroxide to maintain pH 7 yields a new product, the structure of which is yet undetermined. These results and continuation study will enable us to undertake a more comprehensive detection study of all the vinyl chloride detoxification products in biological specimens.

The detection study of the putative action of vinyl chloride is summarized in Table II. Our hypothesis is that such action comes from the modification of the nucleic acid materials by the

TABLE II. PUTATIVE ACTION OF VINYL CHLORIDE. REACTION  
 WITH NUCLEIC ACID BASES

|     |                                                                                                                                |
|-----|--------------------------------------------------------------------------------------------------------------------------------|
| CAA | ETHENO—C <sup>1</sup> , ETHENO—A <sup>2</sup><br>L—ETHENO—G <sup>3</sup> , A—ETHENO—G <sup>4</sup> , HEMIACETAL—G <sup>5</sup> |
| COR | G—7—ACETALDEHYDE <sup>6</sup> + ...                                                                                            |



primary metabolites COR and CAA. Although CAA is long known to react with nucleic acid bases such as cytosine and adenine to form the etheno derivatives, compounds 1 and 2, very little is known about the reaction of CAA on the most reactive base guanine.

Mr. Joseph T. Seawell  
April 6, 1979  
Page 11

By using a battery of modern analytical tools such as HPLC, GC-MS and FT-NMR, we have found that the guanine base in various forms, as the nucleoside, nucleotide and polyguanylic acid, gives rise to two tricyclic ethenoguanines, the linear etheno compound 3 and the angular etheno compound 4, and a third product which is possibly an intermediate, compound 5. Their ratios change with time and pH. It is worthy of note that some of them exhibit fluorescence properties which would allow direct detection in a cell nucleus. The reaction of COR with the guanine base is more tricky due to the instability of COR in aqueous medium. A multitude of products are formed which we have found to be different from those of the CAA reaction. One major product is tentatively identified as compound 6. It is important to note that this is the first indication that COR and CAA show different molecular events in their putative action.

#### Significance

Broadly speaking, this study will lead to early detection and prevention of industrial cancers. Our chemical methodologies (synthesis, structure, and analysis), applied as an integral part of the multidisciplinary approach, will elucidate specific molecular events in the effects of vinyl monomers on industrial workers. This information will form a rational basis for safe use of chemicals and design of preventive measures. Our molecular studies also provide the opportunity to develop useful marker(s) in the form of metabolites in the pathogenesis of chemical injury.

Technical Proposal F -- Assays for the Carcinogenic Potential of Industrial Chemicals Utilizing Prokaryotic and Eukaryotic Systems. U.N. Streips

In the second year of funding, Streip's laboratory has primarily developed and expanded testing capabilities relative to industrial chemicals. Thus, two new screening tests have been implemented: the Comptest and the III test which are indicators for SOS repair function. SOS repair is induced in bacteria following massive insult to DNA. This repair disregards normal DNA sequences and actually results in mutations. SOS repair is postulated to participate in the evolution of a neoplastic cell following chemical damage.

The Comptest examines SOS induction in the bacterium Bacillus subtilis and the III test determines the inhibition of interferon induction in mammalian cell lines (also a suggested SOS function). Since SOS repair is extremely error prone, we postulate that these tests will be specific for carcinogens, not just act as mutagen

VVC 000004850

screens. These tests are described in an upcoming publication (14) and are summarized in Table III.

Table III  
COMPOSITE MUTAGENICITY SPECTRUM OF CHEMICAL MONOMERS<sup>a</sup>

| Chemicals               | Salmonella | Comptest | "Repair Assay" | Forward Mutation | III test |
|-------------------------|------------|----------|----------------|------------------|----------|
| Chloroacetaldehyde      | +          | ND       | +              | +                | ND       |
| Styrene oxide           | +          | ND       | -              | ±                | +        |
| Methyl methanesulfonate | +          | +        | +              | +                | +        |
| Ethyl methanesulfonate  | +          | -        | +              | +                | -        |

<sup>a</sup>Results are expressed as (+) positive in the assay performed; (-) negative in the assay performed; and (ND) not determined. Styrene oxide was weakly reactive in the forward mutation test and must be considered to have borderline activity in this assay (+).

As can be seen in Table III, the Comptest and the III test discriminate between ethylmethanesulfonate (EMS) and methylmethanesulfonate (MMS), both potent mutagens but only MMS is carcinogenic. We are currently testing chloroacetaldehyde and styrene oxide in both of these tests. We will, in the third year, be able to expand these new tests to be a part of our complete battery of rapid screens for the assay of a wide spectrum of industrial chemicals. In addition, our recent results should be applicable to industrial screening laboratories and result in better overall monitoring of environmental hazards.

At this time chloroacetaldehyde has been highly positive (15) in all assays tried and must be considered to be the active metabolite of vinyl chloride. Styrene oxide shows variable activity indicating it may have a different route of attack to cells than most other active chemicals. Since it is strongly positive in the III test, we will have to postulate that styrene oxide may be carcinogenic. This finding is being tested by the Comptest, and styrene oxide will be examined in whole animal systems for carcinogenesis.

Mr. Joseph T. Seawell  
April 6, 1979  
Page 13

Technical Proposal G -- Tissue Antigens and Antibodies in the Detection of Vinyl Chloride Injury. E. Espinosa

Espinosa's finding of an antigen missing in VC-related liver angiosarcoma (detailed in the previous annual report) stimulated further studies of antigenic deletion in chemically-induced hepatomas and in cultured human liver carcinoma cells.

Some of the work in the past year has centered on the characterization of these liver tissue antigens in normal and chemically-induced diseased states. Two liver antigens found to be absent in the fast growing and undifferentiated chemically-induced Morris hepatoma 7777 were characterized and partially isolated. In studies of their occurrence in other tissues, one of these antigens (Antigen I) was shown to be present in kidney and spleen in addition to liver. The second antigen (Antigen II) was detected only in liver. Antigen II was found unrelated to liver-specific F-antigen, differing in a number of properties and in immunologic reactivity.

In studies of their subcellular distribution in normal liver, Antigen I appeared localized in cytosol (54%) and mitochondrial (38%) fractions. Antigen II was about equally distributed in cytosol, mitochondria and nuclei fractions with little amounts in microsomes. Antigen I has a electrophoretic mobility in immuno-electrophoresis close to that of serum gamma-globulins and Antigen II to that of serum alpha-globulins. The two antigens were completely inactivated with Pronase indicating that both antigens are proteins or protein associated. Both antigens were relatively thermolabile; they were partially inactivated following incubation at 56°C and completely inactivated at higher temperatures. Both antigens were completely inactivated when incubated in pH buffer lower than 3.5. In Sephadex-G200 gel filtration, Antigen I behaved like a protein of approximately 51,000 Daltons, using as standards serum albumin, ovalbumin, chymotrypsinogen and ribonuclease. The molecular size of Antigen II (determined on a Bio-gel A5m column) was approximately 240,000 Daltons, using aldolase, catalase and ferritin as markers.

The two antigens were found in the more differentiated and slower growing hepatomas 5123tc and 9618A at about the same concentration as normal liver. The fact that hepatoma 7777 is the fastest growing and least differentiated of the tumors studied suggests a possible functional relationship between the absent antigens and these properties (16). These antigenic deletions may be used as indicators in the early detection of liver tumors and in the evaluation of the rate of growth, histologic differentiation and metastatic properties of such hepatomas.

Mr. Joseph T. Seawell  
April 6, 1979  
Page 14

Another liver constituent which may serve as a sensitive indicator of chemically-induced liver tumors, liver-specific F-antigen, was studied. In studies on the behaviour of this antigen in Morris hepatomas, it was found that different types of these chemically-induced tumors have quite different levels of F-antigen.

F-antigen appeared to be absent in the fast growing hepatoma 7777. In the slow growing hepatoma 9618A, the concentration was very low ranging from less than 2% to 10% of the normal liver concentration. The medium growing hepatoma, 5123tc, highly metastatic, had about twice the concentration as normal liver. F-antigen of hepatoma 5123tc and of normal liver were found localized in the cytosol subcellular fraction and were determined to be immunologically identical and to have equivalent electrophoretic mobility and molecular weight.

Since the antigen was undetectable in the fast growing hepatoma and undetectable or very low in the slow hepatoma, the level of F-antigen does not appear to correlate with the rate of growth of these tumors. A possible relationship between metastatic properties and F-antigen is now being considered because the hepatoma with the increased concentration of F-antigen was by far the most highly metastatic. This may prove useful in treatment of tumors (17).

In studies on cultured human liver carcinoma cells (after establishing optimal conditions required for the maintenance in serum free media of PLC/PRF/5 human liver carcinoma cells) it was determined that these hepatoma cells, similar to the experimental Morris hepatoma 7777, are deficient in liver-specific F-antigen. Nevertheless, these cells, like normal liver cells, produce serum albumin, fibrinogen, transferrin, alpha-1 antitrypsin and alpha-2 macroglobulin (18). These data add further support to the clinical observation that tissue antigens are more useful for treatment and follow-up care than screening and early detection, and that antigenic deletions may prove useful in early screening.

This completes the second annual report from the University of Louisville Chemical Monomer Research Group. If there is need for any further information or clarification, please contact me.

Sincerely yours,

  
Carlo H. Tamburro, M.D.  
Professor of Medicine  
Chief, Division of Digestive  
Diseases and Nutrition

CHT:vb

VVC 000004853

# CITED REFERENCES

1. Fortwengler, H.P., Dever, M.E., Tamburro, C.H., and Espinosa, E. Lymphocyte Transformation Tests in Vinyl Chloride (VC) Workers. Federation Proceedings, 37:362, 1978.
2. Fortwengler, H.P., Jones, D., Tamburro, C.H., Espinosa, E. Factor VIII Content as Evidence for Endothelial Origin of Vinyl Chloride Associated Liver Angiosarcoma (VCA). Federation Proceedings, 38:999, 1979
3. Du, J.T. and Tamburro, C.H. Decreased Glucose-6-phosphatase Activity in Liver in Vinyl Chloride Exposed Rats. Proc. Amer. Fed. of Biological Chemists, 35:1422, 1976.
4. Du, J.T., and Tamburro, C.H. Elevated Glutathione Content, Glutathione-S-Transferase and Glutathione Reductase in Liver of Rats Exposed to Vinyl Chloride. Federation Proceedings, 37:1545, 1978.
5. Kupchella, C.E. and Tamburro, C.H., 1978. Urinary and Tissue Glycosaminoglycan Patterns in Angiosarcoma and Other Vinyl Chloride Exposure Associated Liver Injury. In: Detection and Prevention of Cancer, H.E. Neiburgs, Ed., Part 1, Vol. 1, Marcel Dekker, Inc., New York.
6. Curran, K.L., Kupchella, C.E. and Tamburro, C.H., 1977. Urinary Glycosaminoglycan Patterns in Angiosarcoma of the Liver. Cancer 40:3050-3053.
7. Kupchella, C.E., Jarvis, J.O., Curran, K.L. and Tamburro, C.H., 1977. Tissue and Urinary Glycosaminoglycans (GAGs) changes in Hepatic Fibrosis. Presented at the meeting of the American Association for the Study of Liver Disease. Chicago, IL., November 1, 1977. Gastroenterology 73(5):1229.
8. Kupchella, C.E., Drake, E., Curran, K.L., Kennedy, J. and Tamburro, C.H., 1979. Tissue and Urinary Glycosaminoglycans in Transplantable Hepatomas. Gastroenterology (Abstract in Press).
9. Kupchella, C.E., Secskas, E., Kennedy, J. and Espinosa, E., 1979. Glycosaminoglycan Changes Associated with Hepatic Tumors: The Contributions of Regeneration and Necrosis. To be presented at the Annual (National) Meeting of the American Federation for Clinical Research, Washington, D.C., May 7, 1979. Clinical Research (Abstract in Press).
10. Greenberg, R.A. and Tamburro, C.H. (with C.E. Kupchella, et al.) 1978. Early Detection of Disease in Individuals Exposed to Vinyl Chloride. Presented at the 1978 Annual Meeting of the American Public Health Association, San Diego, CA.

11. Curran, K.L., Kupchella, C.E., Sandoz, J. and Tamburro, C.H., 1979. Urinary Glycosaminoglycan Patterns in Human Hepatic Angiosarcoma, Hepatoma and in Workers at Risk for Angiosarcoma. Gastroenterology (Abstract in Press).
12. Barrows, G.H., Joyce, M.J., Schrodt, G.R., Greenberg, R.A., Tamburro, C.H., 1979. Computer-Assisted Morphological Quantitation of Collagen in Human Liver Biopsies. Laboratory Investigations 40:3.
13. Elmore, J.D., Wong, J.L., Laumbach, A.D. and Streips, U., 1976. Vinyl Chloride Mutagenesis by the Metabolites Chlorooxirane and Chloroacetaldehyde Monomer Hydrate. Biochem. Biophys. Acta 442:405.
14. Streips, U.N., Laumbach, A.D. and Yasbin, A.B. In Microbial Testers for Chemical Carcinogenesis, I.C. Felkner, (Ed.) Marcel Dekker, N.Y., N.Y. in press.
15. Laumbach, A.D., Streips, U.N. and Wong, J.L., 1978. Chloroacetaldehyde Induced Damage to Bacillus subtilis. Abs. Ann. Mtg. Amer. Soc. Microbiol, p. 125, H 128.
16. Espinosa, E., Caple, S., Kupchella, C. and Chia, S., 1979. Two Liver Antigens Undetectable in a Fast Growing Line of Transplanted Hepatomas. Federation Proceedings 38:1069.
17. Espinosa, E., Chia, S., Caple, S. and Kupchella, C., 1979. Liver-specific F-antigen in Transplantable Hepatomas Having Different Growth Rates. Federation Proceedings 38:1069.
18. Johnston, P.B., Espinosa, E., Chia, S. and Caple, S. Properties of 14 Week Maintenance Cultures of PLC/PRF/5 Cells. Abst. of 30th Mtg. Tissue Culture Association, In Vitro in press.

VGM Sumner Univ of Louisville - 4/6/76

Crech - 10 arjos in Linnell plot

4 found by testing procedure - 29 after fibrosis

1100 room for 1000 employees -

Makle - Histology

How remained all family, Canadian, per George

Dispersing - 80-81% competent in wide spread disease

## Spitzer - Acute Effects of Vom

Trying to relate exposure to response

Tomkins: Should always use constant questions in order to compare results for ~~the~~ Std. forms

Naumville COL. Cinn ~~Pa.~~ Mortality Study (males)  
Laumville 20 hie Pa.

VVC 000004856

*Herbert CDC Atlanta Surveillance*

45 from paleologists 37 from bones from r = 24/5

from 160 cases  
103 cases confirmed by Pathology 

Per Hyster.  
65% of cases by local hospital or confirmed  
76% 40-79 yrs.

1% to 88 yrs.

1 - 10-19 22 40-49

1 - 20-29

5 - 30-39

M vs F 59% 40-79 yrs M

No evidence of race; AI, Ethnic lead.

Reidentified analyses re PVC plant & UGM.

1.4 cases of angio / 10mm perph.

Calculated ~~total~~ 5 cases in area red plants.

Found 3 cases - no exposure to UGM, Bz, asbestos.

ESA calculated 5 case misperception.

9 cases of angio in printers, found was, 8 were not.

Checking farmer:

VVC 000004857

16 in white, 10 confirmed (1 thorotrast, 9 unknown).

found 4 farmers, & ∴ May have faint to involvement.

Bu list shi

Nelson

Radiography

PM Session

1. Clinical Screening

~~False~~  
False  
Granting

Data Collection & Interpretation

Need to get Concor data on computer program. CLW needs assistance.

Next to put everything possible in

personal data

organizing job

work history  
set of chemicals (19/1000 PBF)  
set of job functions (14)  
includes apparatus

VVC 000004858

Chemicals used

acrylic acid, acrylic acid, methacrylate,  $C_3H_7$   $n$   $C_8$   $n$  MA

acryl,  $C_2H_5$ , acrylate, Diphent A BP

capryl chloride, chlorinated solvents, chloromethyls

medical  
data  
CCl<sub>4</sub>, C<sub>2</sub>H<sub>5</sub>, C<sub>2</sub>H<sub>5</sub>Cl, EPC

acrylate - ethyl methacrylate, methyl methacrylate, 2 EH  
n-butyl.

diethyl malate,  $HgCl_2$ ,  $CA_3OH$ , phenol,  
toluene, xylene, VDC, VAC, PVC dust

Derived threshold limits from several 1000 to  
Kumt 1 ppm. Now divide into 6  
levels wide open.

Forbes - Have derived no exposure levels since  
1939. Did not refer to TLVs when  
first set up.

Level 0 - no exposure

1 - min. exp. to level <sup>not handled</sup> by int.

2 - min. exp. incl. work around chem.

3 - within area, subject to his supervision

4 - within area where level hi;

5 - intimate contact (reactor cleaner)

slurry handle, etc.

Spider program comparable to BFR, diff guidelines

VVC 000004859


3 Employee Response & Problems  
Modalities

4/7/76 AM

Research Studies on Chemical Carcinogens



Metabolic Derivatives of UCM - John Wong

Studied in two systems



VVC 000004860


Mon-ay.



Brodmann / Bepflanzung Ost (1976) Wang  
St. Leipzig

*Mammox hydrate principal carrier form*





Oxidation traps

Conjugation (C<sub>18</sub>N<sub>2</sub> or electron trap)

study V<sub>id</sub>, C, E, A as traps  
Carotenes or melanophylls

VVC 000004862

Deconjugation of N<sub>2</sub> retention medium  
exponentially

DMS - nonconjugation <sup>pH 7.5</sup> Ethylene glycol

Body fluid of experimental animals after exposure to  
then water. Metabolism pathway  
via NADH flavin transform

Strep - Screening - Analysis

Bacillus Subtilis

Dot Key effect  
DNA repair  
screen

165 Min

11CR-9 (JB01-200)

FB-13

MC-1

Salmonella Typhimurium Tests

TA 1535

TA 100

TA 1537

TA 1538

TA 98

VVC 000004863

4 nitrophenyl-oxides

Repair Assay: strat. / unit / year

fills  
proper  
sat. with  
mitogen

strat. / repair

24 hr incubation

Summary of Mitogen Testers

Serial

VGM  $\pm$  NR

$Cl_2C=CCl_2$  + ++

$ClCH_2CO$  ++ NR+ +

epidichloroprit + NR ++

chloroform ++ ++

minimally  
added to  
first small  
stage

all forms of  $ClCH_2CO$  mitogen-

in some stages

epi-chloroform the last

VVC 000004864

deamination repair.  
this type most error prone of all DNA repair

epi deamination doesn't give same type  
of mutations as the others.

Specifically affect deamination repair.  
MUT - really destroyed

cells may cause window movement repair.  
could cause cross-linked DNA.

Reversion of amino acids to bases  
Chromosomes allowed for complete replication  
cycle. cells - even better at killing  
non-replicating chromosomes.

Does not need active cell to do cross-linking.

What do cells do with DNA?

CAT & GP1 don't physically damage DNA  
in vitro

VVC 000004865

~~look~~  
treat CBA with particles  
in vivo

DNA is being damaged in the  
DNA.

Mutagen - ~~Particle~~ - <sup>bound</sup> to surface membrane.  
damaged -

Way to see what induction <sup>mutagen</sup> numbers are  
Then look for what effect, then how to solve  
problem.

Hoffman - Sulfonamide metabolism &  
Carcinogenesis

VVC 000004866

Involvement -

glutathione. Detoxification - accomplished by conjugation  
with -SH groups, PAH epoxides, VMA metabolites

2. Metabolism analog, ethionine, is a carcinogen
3. Metabolism adrenergic stress response. highly  
repressed & repressed in tumor function in

in comparison to neonatal liver.

comparison to neonatal adults.

protein  $\rightarrow$  methionine  $\rightarrow$  cysteine  $\rightarrow$  glutathione  
50-100 nmol/mole  
90% go to cysteine, 10% to protein synthesis.

S-adenosyl-L-methionine


CH<sub>3</sub> transfer important for 4 major DNA transfer.

Diethylamine, flavone - very potent, <sup>coronin</sup>

Nucleoly - deactivate enzyme.

SAM - inactivated by Cl<sub>2</sub>, CO

1/2 molecule. Dithionite, 4 moles S<sub>2</sub>  
did not prevent inactivation of SAM.



herbicide, not started yet.

Julie Alu - Eugene assay of Experiments Carcinogen

Bile acid clearance test -

Chromatography of Conjugate of Cholic Acid

New Eng J of Med (1975) Hoffman

|          |               |             |
|----------|---------------|-------------|
| 49 ppm   | 15 hr         | 754         |
| uris     | 58 hr         | 67.1        |
| pus      | 2.7           | 3.8         |
| epid co  | 9.8           | 14.0        |
| lyst v m | <u>      </u> | <u>0.02</u> |
|          | 70.5          | 84.92       |

New Metabolism Wash (above)

Residue is in tissues (75% 13%)

VVC 000004868



DNA derangement.

Kupffer cell -

Hepato/Hepatocyte Birkhead exploration

Now - see decrease of glutathione in liver.

Experiments in BFB vials??

up to 20,000 ppm exposure 4-8 hr.

Have checked S.M.P. - 12 at all

levels of exposure - found no abnormalities.

up to 137 hr. Checked also for  
marker enzymes -

total of glutathione levels in liver -

VVC 000004870

Found change in glutathione reductase,  
increased with increased exposure  
liver.  $\rightarrow$  glucose-6-phosphate - decreased.

enzyme.

See effect also glucose-6-phosphate  
dehydrogenase.

Fortmeyer - Immunology

25 BFG oncoprotein typed - with & without  
leucine deacetylase

Tumor destruction by cells, lymphocytes

Transplants or viruses suppressed.

Of all oncoproteins, only those  
treated with chemotherapy developed  
metastases.

Cellulose is neutral

Cancer tests - Immunofluorescence

tumors but tumor antigens

Leukocytes can recognize antigens.

VVC 000004871


Empirical findings  
 WBC adheres to glass  
 Lymphatics cause WBC to fall off glass.  
 L.A.T.

add lymphocytes from suspected patient.  
 add antigen from carcinomas with same type.  
 Showed to good predictive test.

Sigdestad  $\rightarrow$  Partial Hepatectomy Model -  
 No partial ~~hepatectomy~~ <sup>hepatectomy</sup>, increase 40-60% response  
 of carcinomas. Normal liver, few cells at work.  
 Proliferous state when remainder of it all go  
 almost simultaneously to DNA make.

1 in 20-30, on in mytotic (rest) 3% in DNA make.

- On rats - remove 65% of liver, regenerate  
 max in 6-12 days.



VVC 000004872

CCl<sub>4</sub>, hepatitis, chronic alcohol - all will trigger  
response.

Partial hepatectomy + VEGF could be informative.

need  
proof/hold.

1. Determine angles in normal rat
2. Determine angles in partial hepatectomy + VEGF
3. Determine angles in " " + CCl<sub>4</sub> CMO.

VEGF maybe stimulating proliferation.

study cell kinetics after partial hepatectomy.

Chromosome aberration experiments.

↔ Possibility radiation damaged liver which has  
repaired etc could cause repair with hepatocarcinoma.

Kupffer cells → Exploitation.

Neovascularization of angio

Study by etiology of angio.

VVC 000004873

glycosaminoglycan  
glycosaminoglycan? gag  
amino

protein-carbohydrate enzymes -

fibrosis can cause rises & falls in gag levels.

- Seems to contradict if something important

a lack of        suggests in urine analysis.

       in case of VCM exposure (non-argus)

<sup>Argus</sup> Esplenoma - Antigens not present in normal liver.

Search for antigenic changes

new antigens

antigens absent

- Data supports presence of antigen  
present in argus cases not in n. liver.

Abnormal (liver)

live  
cell changes, not found in circulating blood,  
viral hepatitis and alcoholism do have  
circulating antigens.

VVC 000004874

Sheet ~~2~~ Electron Micrographic Studies.

VVC 000004875