

*Fisheries Research Board of Canada Biological Station
St. Andrews, N. B.*

Methods and Materials

Five *Gammarus* were added to each beaker which was maintained at 5 ± 0.5 C and the survival time (± 6 hours) of each animal recorded. Bioassays were continued for 30 days and solutions changed weekly.

Apart from some deaths caused by cannibalism, no mortality occurred in sea water alone or Corexit 7664 in sea water in concentrations up to 1.9 g/l. Colloidal solutions of Aroclor 1254 solubilized in Corexit 7664 (1:19 W/W) or emulsions (1:10, 1:1, and 1:0.1 W/W) in sea water were lethal at concentrations as low as 0.01 mg/l and 0.1 mg/l respectively. Observations suggested that moulting or freshly moulted animals were particularly vulnerable. Plotting the survival time of *G. oceanicus* against concentration (Fig. 1) showed a "critical point" of response below which there was, and above which there was no time/concentration effect. The lethal threshold is between 0.001 and 0.01 mg/l for colloidal solution and between 0.01 and 0.1 mg/l for emulsions of Aroclor 1254 solubilized in Corexit 7664 in sea water. After exposure to Aroclor 1254 in sea water for periods greater than 150 hours some *Gammarus* which died were found to have severely necrosed branchiae. Necrosis was indicated by a

cnyls
panicus

igate
owing
stem
of the
lor

Corexit
dis-
red by
ss beak-
Con-
ter and

aker
vival
assays
nged

annibal-
e or
p to
solu-
ons
lethal
mg/l
ulting
vulner-
zicus
tical
d above
The
g/l for
ng/l for
exit
r 1254
ers some
ely ne-
a




Figure 1. Median and range of survival time of *G. oceanicus* in Aroclor 1254:Corexit 7664 in sea water.

chitinous wound callus which isolated the affected brownish tissue. A less extensive, sublethal necrosis was found in some animals down to 0.001 mg/l of colloidal PCB in sea water. A sublethal branchial oedema was attributed to the effect of Corexit 7664 in sea water effective at a concentration down to 0.19 mg/l.

Other data (2) show that the highest concentration of Corexit 7664 used (1.9 g/l) to solubilize Aroclor 1254 is 15 times lower than the 48-hr LC50, and thus no mortality was caused directly by the emulsifier. Nevertheless it is possible that a synergistic effect occurred between the emulsifier and PCB.

References

1. ZITKO, V. Bull Environ. Contamination and Toxicol.
2. WILDISH, D. J. Fisheries Research Board of Canada, MS Report No. 1084. Polychlorinated biphenyls in sea water: bioassay of *Gammarus oceanicus* (May 1970).

This
out at Mar
tensive ef
quality up

Fish Avoid

Many
centrarch
areas when
important
the fish c
fish type
reaction a
pearance c
tant to th
this time
dealing wi
fish life
informatio
would be p

The f
general fi
fish, shin
abundance
bass and w
population
tion cause
ditions mo
diverse sa
thirty yea
carp and g
migratory
Hudson lis
species cor

In ord
glass tank
to that des
are fine pl
connected t