Monsanto

FROM (NAME & LOCATION) R. E. Keller - R&D Laboratories - South Second Street

DATE June 30, 1970

SUBJECT CHLORINATED DIBENZOFURANS - PCB

REFERENCE

TO : W. B. Papageorge
General Offices

cc H. S. Bergen - GO
W. R. Richard - GO
E. P. Wheeler - GO
M. W. Dietrich - Res. 2
E. M. Emery - Res. 2

The attached letter and manuscript have been received from Dr. J. G. Vos at the University of Utrecht. It is in reply to our request for details of their work on the identification of chlorinated dibenzofurans in PCB. Please note that this manuscript has been sent on a confidential basis and it is not for general distribution.

We will attempt to provide chlorinated naphthalenes requested for gas chromatography work by Vos. Bill Richard thinks we might be able to obtain these commercially. Dr. Emery is asked by copy of this memo to determine if we have any of these compounds in SRC.

Drs. Dietrich and Emery are asked by copy of this memo to please review the manuscript and determine the difficulty of applying this technique to our Aroclor products. Assuming this is within our capabilities with a reasonable amount of effort, we will look at Aroclor products used at Bio-Test for feeding studies and compare results with those for current production grade materials.

R. E. Keller

db
Attachments - 2

IM 10 REV 11 69

VETERINAIRE FARMACOLOGIE
EN TOXICOLOGIE
DER RIJKSUNIVERSITEIT
TE UTRECHT

DIRECTEUR: Prof. H. VAN GENDEREN

No.

Bericht op:

Betreffende:

Bijlagen: +

UTRECHT, June 24, 1970.
Biltstraat 172
Telefoon No. 715544

Dr. R.E. Keller,
Applied Sciences Section
Monsanto Company,
1700 South Second Street,
ST. LOUIS, Missouri 63177.
USA

Dear Dr. Keller,

Enclosed you will find a photocopy of the manuscript concerning the identification of chlorinated dibenzofurans in PCB. We use fractionation GC/mass spectrometry techniques; the details are given in the manuscript which is not for general distribution since it is not yet accepted by the Editors of the Journal of Food and Cosmetics Toxicology.

During the visit of you and your colleagues at our Institute we mentioned a.o. severe skin lesions (hyperkeratosis) in cattle. The floor of the stable in which the animals were housed was covered with second-hand conveyor belts, covered with a new anti-slip layer. Gas chromatographic analyses of a sample of the rubber mat and of the liver of a severely affected cow revealed a peak pattern which was very indicative for PCB, as we also mentioned during your visit. Mass spectrometric analyses of the rubber revealed compounds with molecular weights of probably penta- and hexachloronaphthalene, well known for their hyperkeratotic action in cattle.

Chlorinated naphthalenes are not for sale and since they are very important for us as reference compounds in gas chromatography, we would like to ask you to mediate for the supply of chlorinated naphthalene mixtures and perhaps a pure sample of octa chloronaphtalene (perchloronaphthalene). Dr. Koeman told me that he will send you a total-body sample of a cormorant containing a high concentration of PCB's, for a comparison of the quantitative aspects of the analytical methods.

Sincerely yours,

J.G. Vos.

Institute of Veterinary
Pharmacology and Toxicology,
University of Utrecht,
Biltstraat 172,
UTRECHT,
The Netherlands

MONS 201245

Confidential

Identification and Toxicological Evaluation of Chlorinated Dibenzofurans and Chlorinated Naphthalene in Two Commercial Polychlorinated Biphenyls.

J.G. Vos.
Institute of Veterinary Pathology,
Biltstraat 172, Utrecht, The Netherlands.
J.H. Koeman and M.L. van der Maas.
Institute of Veterinary Pharmacology and Toxicology,
Biltstraat 172, Utrecht, The Netherlands.
M.C. ten Noever de Brauw.
R.H. de Vos.
Central Institute for Nutrition and Food Research,
TNO, Zeist, The Netherlands.

Running title:

Toxic impurities in technical PCB preparations.

This write-up was sent on a confidential basis to R. E. Keller on 6-24-70. It is not for general distribution and should not be reproduced.

RE Keller

MONS 201246

Abstract.

In a previous study (Vos & Koeman, 1970) a significant difference in toxicity was found between 3 commercial PCB preparations; high mortality, liver necrosis and chick oedema-like lesions occurred in 2 PCB's.

In the present study it was found, by means of column chromatography and gas chromatography, that in the 25% diethylether fractions of these 2 PCB's polar compounds are present. A chicken embryo assay proved the high toxicity of this fraction. Subsequent mass spectrometric and microcoulometric analyses revealed the identity of these compounds; a.o. tetra- and pentachlorodibenzofurans. The significance of these impurities are discussed. It is concluded that evaluation of animalexperiments with PCB's is extremely difficult due to the presence of chlorinated dibenzofurans in some commercial PCB's.

MONS 201247

Introduction.

Polychlorinated biphenyls (PCB) have been identified in tissues of fish and wildlife (Jensen, 1966; Holmes, Simmons & Tatton, 1967; Holden & Marsden, 1967; Koeman, Ockhup, Veen, Brouwer, Booth, Zwart, van de Broek & van Genderen, 1967; Risebrough, Riecke, Foskall, Herman & Kirven, 1968; Koeman, ten Noever de Brauw & de Vos, 1969; Jensen, Johnels, Olsson & Otterlind, 1969), and have been recently been implicated in the mortality of sea birds in Britain (Bourne & Head, 1969).

Commercial PCB preparations are extremely stable and only slightly soluble in water. They are used as lubricants, as heat transfer media, in protective coatings for wood most metal and concrete, and in many other applications. They are compatible with paints, synthetic resins, varnishes and waxes, and improve such properties of these products as their chemical resistance, water resistance, adhesiveness and flexibility (Anonymus, 1960; Hubbard, 1964). However, it has not yet been established which of the materials treated with PCB has contributed to the present environmental contamination.

Experiments with rats and guinea pigs have shown that the ingestion of PCB may give rise to liver injury (Miller, 1944). Hydropericardium, occasionally accompanied by abdominal oedema, was found in chicks (McCune, Savage & O'Dell, 1962; Flick, O'Dell & Childs, 1965; Vos & Koeman, 1970) and Japanese quail (Koeman et al., 1969) after ingestion of PCB of American (Aroclor) and French (Phenoclor) origin, respectively. A marked resemblance between the gas chromatograms and mass spectra in mass spectrometer analyses of 3 commercial PCB preparations (Phenoclor DP6, Clophen A60, and Aroclor 1260) was found (Koeman et al., 1969).
The occurrence of "chick oedema" like lesions in birds fed PCB (Koeman et al., 1969; McCune et al., 1962; Flick et al. 1965) was

MONS 201248

the reason for a recent comparative toxicological test with the 3 mentioned PCB preparations in chickens (Vos & Koeman, 1970). In this study a significant difference in toxicity between the 3 tested compounds was found; Clophen A60 and Phenoclor DP6 showing the highest, Aroclor 1260 showing the lowest mortality. Subcutaneous and abdominal oedema, and centrolobular liver necrosis was recorded only in those chicks fed with Clophen and Phenoclor. Hydropericardium was recorded in nearly all chicks fed with Clophen and Phenoclor and only occasionally in chicks fed with Aroclor. Chemical porphyria was found as a general PCB effect.

In the present study, the sequel to that mentioned above, we describe the identification and evaluation of compounds responsible for the difference in toxicity between the 3 commercial PCB preparations.

Experimental.

The PCB preparations, which contain an average of 60% chlorine, were obtained from Prodelec, France (Phenoclor DP6), Bayer, Germany (Clophen A60, Lot no. 912434), and Monsanto, U.S.A. (Aroclor 1260, Lot no. Ak-3).

The screening method, used for detection of chick oedema factor (Anonymus, 1960a) was modified and used to isolate from the PCB preparations, compound(s) responsible for differences in toxicity. The clean-up step of this procedure was omitted. In the column chromatograph; 8g. activated florisil (column ϕ 10mm) was used instead of 15g. activated alumina. For fractionation redist. hexane was used as elution solvent instead of petroleum ether. Two mg PCB was dissolved in 2 ml hexane and fractionated on the florisil column. This column was eluted with hexane until nearly all PCB's were eluted (tested by gas chromatography). The 2nd fraction (5% diethylether in hexane) was also tested until no more PCB could be detected. The final fraction (fraction 3), eluted with 25% diethylether (reagent grade) in hexane, was evaporated to 1 ml. An aliquot of this concentrate was injected into a 20%-1B Varian Aerograph gas chromatograph with electron capture detection. The Pyrex glass column (5 feet x 1/8 inch), filled with 10% DC 200 on Gaschrom Q (80-100 mesh) was operated at 200°C with nitrogen as carrier gas (about 50 ml/min.). The same procedure was used to fractionate aliquots of about 550 mg of the PCB preparations in order to obtain sufficient quantities for a chicken embryo assay and for mass spectrometer analyses. The PCB was dissolved in 200 ml hexane. The column (ϕ 32 mm) was filled with 180 g activated florisil. The more polar compounds were eluted in the final (3rd) fraction and, after evaporation, tested by gas chromatography.

A chicken embryo assay (Verrett, Harliac & McLaughlin, 1964) was used to determine the toxicity of the 3 PCB preparations.

(doses 5... µg/egg, dissolved and kept at 37°C in ethanol. Group 1 = Aroclor, group 2 = Phenoclor and group 3 = Clophen) and of the 3rd fraction of Clophen (dose: 3rd fraction obtained from 35, 3.5, and 0.35 and 0.035 µg. Clophen in groups 4, 5, 6 and 7 respectively). The samples, after dissolving in 0.05 ml ethanol were injected into the air cell of fresh White Leghorn eggs prior to incubation. The livers of all hatched chicks were removed for histological examination, fixed in 10% buffered formalin and processed in a conventional manner. Moreover, cryostat sections were studied in a fluorescence microscope.

Mass spectrometry was used to identify compounds present in the polar 3rd fractions. This analysis was carried out using a gas chromatograph-mass spectrometer combination. For this purpose a CH4 mass spectrometer (Varian Mat. G.m.b.H.) was coupled to a Varian Aerograph oven, type A550B, with a Pyrex glass column (5 feet x 1/8 inch) filled with 10% DC200 on Gaschrom Q (80-100 mesh). Particular attention was paid to the design of the interface between the gas chromatograph and the mass spectrometer in order to obtain the best operating conditions (ten Noever de Brauw & Brunde, 1967). High resolution mass spectra were taken on a double focus spectrometer type SH_1 (Varian Mat. G.m.b.H.). Exact mass measurements were done with the peak matching method (Measurements were also done from spectra which were recorded on photographic plates processed by Varian Mat. on a Leitz comparator DA/11).

To obtain information about the quantities, microcoulometric analyses of the 3rd fractions were carried out. The equipment used was a Dohrmann microcoulometer, coupled to a Microtek MT 220 gas chromatograph, with a glass column (6 feet x ¼ inch). The same 10% DC 200 column packing was used. The column effluents were oxidized in a Dohrmann combustion furnace and the HCl, produced by chlorinated compounds, was detected and quantitatively measured in a P-300-S titration cell.

Results

The resulting gas chromatogram of fraction 3, obtained with the small column (Ø 10mm) revealed the presence of polar compounds in Phenoclor and Clophen (fig.1). These were not found in the Aroclor extract. The same results were found with fraction 3 obtained with the large column (Ø 32 mm). The 3rd fraction 3, obtained of Clophen was subsequently used in the chicken embryo assay. This assay (table 1) confirmed the differences in toxicity between the 3 tested PCB preparations found in the previous study (Vos & Koeman, 1970). Mortality in group 1 (Aroclor) did not differ significantly (X^2 test) from mortality in group 9 (control). Mortality in group 4 and 5 is dose related; early mortality in the former, scattered mortality in the latter group. Corresponding mortality in group 3 (3.5 mg Clophen) and group 6 (3rd fraction from 3.5 mg Clophen) indicates strongly that the polar residue in the latter determines the toxicity of the commercial PCB. Dissection of all fresh embryos and of the one-day-old chicks revealed no abnormalities except for one embryo of group 3 which showed a severe subcutaneous oedema. Oedema and beak deformities are described after yolk sac injection of 25 µg and 10 mg Aroclor 1242, resulting in respectively 100% and 95% embryonic mortality (McLaughlin, Murliac, Verrett, Hutchler & Fitzhugh, 1963) M.L. stain liver sections revealed no morphologic alterations. In the cryostat section of the liver of the only hatched weak chicken of group 3, a red fluorescence in some lobules was seen in the u.v. microscope. This fluorescence is indicative of the porphyrin biosynthesis disturbance found in the prior study (Vos & Koeman, 1970).

Mass spectrometric analyses revealed identical chlorinat compounds in fraction 3 from Phenoclor and Clophen in contrast wit Aroclor. The results are given in table 2. Mass spectra of compoun with mass number 304 and 330, obtained from peak 1 and 4 (fig.1) are given in fig.2 and 3. From exact mass measurements it was foun that the formulas of the molecule peaks 304 and 330 were C_{12}

MONS 201252

$C_{12}H_4O^{35}Cl_4$ and $C_{12}H_3O^{35}Cl_5$. The formulas of the fragment ions 241 and 275 are $C_{11}H_4{}^{35}Cl_3$ and $C_{11}H_3{}^{35}Cl_4$. From this data it can be concluded that the parent ion has a preferential fragmentation for the loss of a single ClCO unit. Mass spectra were compared with the spectrum of a compound isolated from toxic fat (Wootton & Courchene, 1964). Subsequent study, using X-ray diffraction, has revealed that the identity of one of the toxic factors was 1, 2, 3, 7, 8, 9-hexachlorodibenzo-p-dioxin (Wootton & Courchene 1964). The mass-spectrum of this compound has a similar fragmentation pattern i.e. loss of successive 2 mass units 63 suggests a loss of 2 ClCO units.

Microcoulometric analyses of the 3rd fraction of Clophen and Phenoclor revealed chromatograms with the same peak pattern as in fig.1. The "peak" of the tetrachlorodibenzofuran (mass number 304) is too masked by the proximity of the hexachloronaphthalene peak (peak no.2) to make quantitative calculations possible. For similar reason peak no.4 in the chromatogram, found to be a mixture of heptachlorobiphenyl and pentachlorodibenzofuran, could not be measured directly. However, when assuming that the whole peak is produced by the dibenzofuran compound alone, a certain maximum level can be indicated. This level was found to be 5 ppm pentachlorodibenzofuran in Clophen and 20 ppm in Phenoclor. However the minimum level could be indicated with the estimated lower limit of detection of the mass spectrometric equipment. Under the conditions used this level was 1 ppm.

Discussion.

Mass spectrometric results have not conclusively proved the identity of the chlorinated dibenzofurans. But synthesis of these compounds, necessary for comparison with the mass spectra obtained experimentally, was found extremely dangerous; a single oral dose of 0.5-1.0 mg/kg tri- and tetrachlorodibenzofuran caused severe and often lethal liver necrosis in rabbits (Bauer, Schulz & Spiegelberg, 1961; Hofman, 1958). Application on the ear of a rabbit resulted in chloracne (Bauer et al., 1961). The related compound 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin (following the description of Chemical Abstracts), which may appear in the synthesis of 2, 4, 5-trichlorphenol by hydrolysis (under pressure and with the presence of sodium hydroxide and methanol) of 1, 2, 4, 5-tetrachlorobenzene (Bauer et al., 1961), usually caused a lethal liver necrosis in the rabbit after a single oral dose of 0.05-0.1 mg/kg. Ear application, also in a 10 times lower dose compared with chlorinated dibenzofuran, resulted in chloracne (Bauer et al., 1961; Linn Jones & Krinck, 1962). 1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin, which can arise in the pyrolysis of 2,3,4,6-tetrachlorophenol (Higginbotham, Huang, Firestone, Verrett, Ress & Campbell, 1960), has been isolated and identified as one of the toxic substances of the Chick Oedema Factor (Wootton & Courehone, 1964; Anonymus, 1968b). The relationship between the toxic nature of PCB and the Chick Oedema Factor has already been discussed (Flick et al., 1965).

Air cell injection in chicken eggs of 0.05 ug hexachlorodibenzo-p-dioxin resulted in 100% embryonic mortality (Higginbotham et al., 1969). The experiment using an injection of the 3rd fraction obtained from 35 mg Clophen into the air cell resulted in 100% mortality (table 1). It can be calculated that the maximum

dose to 0.2 ug pentachlorodibenzofuran (using a value of max. 5 ppm). This confirms the order of toxicity discussed above.

The combination of toxicological, pathological and chemical analytical data makes it almost certain that the compounds of mass numbers 304 and 338 are respectively tetra- and pentachlorodibenzofuran. The origin of these toxic factors in Clophen A60 and Phenoclor DP6 can possibly be explained by differences in manufacture. Particularly in the distillation procedure of crude PCB, in which sodium hydroxide can be used (Fay & Richards, 1947). PCB reacts with sodiumhydroxide at elevated temperatures to yield phenolic compounds, e.g. PCB is saponified into polychlorohydroxybiphenyls by means of sodium hydroxide in a polyhydric alcohol medium (Anonymus, 1957). A further loss of hydrochloric acid could produce dibenzofuran derivatives.

Chlorinated naphthalenes are far less toxic than chlorinated dibenzofurans (Hofmann, 1958). Therefore the latter may be assumed to determine the toxicity of fraction 3.

The occurrence of hydropericardium in chick fed with Arocolor (Flick et al., 1965), may indicate the presence of a very small quantity toxic factor.

From this study it is concluded that evaluation of PCB residues in wildlife animals, based on effects and residues in test animals, is extremely difficult due to differences in toxicity between commercial PCB preparations.

Work is in progress to detect and identify chlorinated dibenzofuran from wildlife specimens.

MONS 201255

Acknowledgements.

We acknowledge the contribution of Mr. E.W.A. Kampordijk and Mr. F.P. Nijkamp, and the technical assistance of Mr. J.M. Penni

This investigation was supported by a grant from the Central Organization for Applied Scientific Research (T.N.O.) which made it possible to accomplish the mass spectra analyses.

MONS 201256

References

Anonymus, (1957). Method for producing phenolic substances through the treatment of chlorinated diphenyls and polyphenyls and substances obtained thereby. Brit. Pat. 779, 221, July 17 (to Procil).

Anonymus, (1960). Aroclor Plasticizers. Monsanto Company Tech. Bull. no. PL-506.

Anonymus, (1960a). Changes in Methods, 26. Oils, fats and waxes. J. Ass. Offic. Anal. Chem. 51, 489-490.

Anonymus, (1965b). The chick edema factor. Nutr. Rev. 26, 28-30.

Bauer, H., Schulz, K.H. & Spiegelberg, U. (1961). Berufliche Vergiftungen bei der Herstellung von Chlorphenol-Verbindingen. Arch. Gewerbepath. Gewerbehyg. 18, 530-555.

Bourne, B. & Mead, C. (1969). Seabird slaughter. B.T.O. News, ed. by Keith Clark for the British trust for Ornithology, no. 36.

Fay, J.W.J. & Richards, J.H. (1947). Impregnants used in German paper capacitors. Office Tech. Serv. PB rept. 75059, BIOS Final Rept. 893.

Flick, D.F., O'Dell, R.G. & Childs, V.A. (1965). Studies of the chick edema disease. 3. Similarity of symptoms produced by feeding chlorinated biphenyl. Poultry Sci. 44, 1460-1465.

Higginbotham, G.R., Huang, A., Firestone, D., Verrett, J., Ress, J. & Campbell, A.D. (1968). Chemical and toxicological evaluations of isolated and synthetic chlorodderivatives of dibenzo-p-dioxin Nature, 220, 702-703.

Hofmann, H.Th. (1958). Neuere Erfahrungen mit hochtoxischen Chlorkohlenwasserstoffen. Experimentelle Pathologie Pharmakolog 232, 220-230.

Holden, A.V. & Marsden, K. (1967). Organochlorine pesticides in seals and porpoises. Nature 216, 1274-1276.

MONS 201257

hydrocarbons in British wildlife. Nature, 216, 227-229.

Hubbard, H.L. (1964). Chlorinated biphenyl and related compounds. Encyclopedia of Chemical Technology, ed. by Eller Kirk, R. & Othmer, D.F., 2nd rev. ed., 5, 289-297.

Jensen, S. (1966). Report of a new chemical hazard. New Sci. 32, 612

Jensen, S. Johnels, A.G., Olsson, M. & Otterlind, G. (1969). DDT and PCB in marine animals from Swedish waters. Nature, 224, 247-250.

Koeman, J.H., Oekamp, A.A.G., Veen, J., Brouwer, E., Rooth, J., Zwart, P., v.d. Brock, E. & van Genderen, H.,(1967). Insecticides as a factor in the mortality of the sandwich tern (Sterna sandvicensis. Mededel.Rijksfac.Landbouwwetenschappen Gent, 32, 841-854.

Koeman, J.H., ten Noever de Brauw, M.C. & de Vos, R.H. (1969). Chlorinated biphenyls in fish, mussels and birds from the river Rhine and the Netherlands coastal area. Nature, 221, 1126-1128.

Linn Jones, E. & Krizek, H. (1962). A technic for testing acnegenic potency in rabbits, applied to the potent acnegen, 2,3,7,8,-tetrachlorodibenzo-p-dioxin. J. Invest. Dermatol. 39, 511-517.

McCune, E.L., Savage, J.E. & O'Dell, B.L. (1962). Hydropericardium and ascites in chicks fed a chlorinated hydrocarbon. Poultry Sci. 41, 295-299.

McLaughlin, J., Marliac, J., Verrett, M.J., Hutchler, M.K. & Fitzhugh, O.G., (1963). The injection of chemicals into the yolk sac of fertile eggs prior to incubation as a toxicity test. Toxicol. Appl. Pharmacol., 5, 760-771.

Miller, J.W. (1944). Pathologic changes in animals exposed to a commercial chlorinated diphenyl. Pub. Health Rep., 59, 1085-1093

MONS 201258

Risebrough, R.W., Riechc, P., Peakall, D.B., Herman, S.G. & Kirven, M.N. (1968). Polychlorinated biphenyls in the global ecosystem. Nature, 220, 1098-1102.

Ten Hoover de Braun, M.C. & Brunnée, C.Z. (1967). Ergebnisse mit einer optimierten Kopplung von Gas Chromatograph und Massenspektrometer. Zeitschr. f. Anal. Chem. 229, 321-335.

Verrett, M.J., Marline, J. & McLaughlin, J. (1964). Use of the chicken embryo in the assay of aflatoxin toxicity. J. Ass. Offic. Agr. Chem. 47, 1003,-1006.

Vos, J.G. & Koeman, J.H. (1970). Comparative toxicological study with polychlorinated biphenyls in chickens with special reference to porphyria, edema formation, liver necrosis, and tissue residues. Toxicol. Appl. Pharmacol. In press.

Wootton, J.C. & Courchene, W.L. (1964). A contribution to the knowledge of the structure of two hydropericardium-producing factors from a toxic fat. Agr. Food. Chem. 12,-94-99.

MONS 201259

Table 1. Chicken Embryo Assay of Chlorinated Biphenyls Aroclor 1260, Phenoclor DP 6 and Clophen A60 and the 25 per cent Diethylether Fraction of Clophen A60.

Group	Compound	Dose/egg[a]	Number	Number of Infertile Eggs	Per cent Hatch of Fertile Eggs	Remarks
1	Aroclor 1260	3.5 mg	15	0	80	
2	Phenoclor DP 6	3.5 mg	15	0	0	Most embryos died during firs half of incubation period.
3	Clophen A 60	3.5 mg	20	1	5	Most embryos died during firs half of incubation period.
4	Clophen A 60 3rd fraction of	35 mg	15	0	0	All embryos died during firs four days of incubation peri
5	Clophen A 60 "	3.5 mg	15	0	7	Most embryos died during firs half of incubation period.
6	Clophen A 60 "	0.35 mg	15	2	92	
7	Clophen A 60 "	0.035mg	15	2	100	
8	Ethanol		20	2	94	
9	Control		20	0	90	

[a]The samples, dissolved in 0.05 ml ethanol, were injected into the aircell of fresh eggs before incubation.

Table 2. Retentions and Collected Mass Spectrometric Data of the Peaks in fig.1.

Peak no.	Relative retention[a]	Mass number(s)[b], and number of chlorine atoms/molecule	Identity of the compound(s).
1	1.40	304(4Cl)	tetrachlorodibenzofuran
2	1,53	332(6Cl)	hexachloronaphthalene
3	1.74	358(6Cl)	hexachlorobiphenyl
		392(7Cl)	heptachlorobiphenyl
4	2.42	338(5Cl)	pentachlorodibenzofuran
		392(7Cl)	heptachlorobipheryl
5	3.48	366(7Cl)	heptachloronaphthalene

[a]Retention related to dieldrin = 1

[b]Mass number calculations based on Cl = 35

MONS 201261