

ARITZ

Br. J. Cancer (1973) 28, 173

MESOTHELIOMATA IN RATS AFTER INOCULATION WITH ASBESTOS AND OTHER MATERIALS

J. C. WAGNER, G. BERRY AND V. TIMBRELL

From the Medical Research Council's Pneumoconiosis Unit, Llandough Hospital, Penarth, Glamorgan

Received 21 March 1973. Accepted 10 April 1973

Summary.—Four experiments in which SPF Wistar rats were inoculated intrapleurally with asbestos or other materials are described. Mesotheliomata were observed in a considerable proportion of animals with all the samples of asbestos used and with a sample of brucite. A few were produced with synthetic aluminium silicate fibres and single ones with barium sulphate, glass powder and aluminium oxide. The risk of developing a mesothelioma at a given time after injection was approximately proportional to the dose. Of the UICC standard reference samples, crocidolite was the most carcinogenic and removal of the oils by benzene extraction did not alter the carcinogenicity of these samples. Chemical properties also seem unlikely to be the main factor producing mesotheliomata but the results support the hypothesis that the finer fibres are the more carcinogenic, and this is additional to the known aerodynamic advantage which the finer fibres have in penetrating to the periphery of the lung.

We report here the results of 4 experiments in which asbestos and other test materials were administered to rats by intrapleural inoculation. These experiments were planned to obtain more information on the carcinogenic effect of asbestos and other materials than could be obtained from our original 2 experiments (Wagner and Berry, 1969). Preliminary results of some of the present experiments were given by Wagner, Berry and Timbrell (1970), Wagner (1970, 1972). In this paper the complete results are given, with emphasis on the light they throw on the aetiology of mesotheliomata, taking into account the oils and waxes present in asbestos, other chemical characteristics and the physical characteristics.

MATERIALS AND METHODS

In all 4 experiments specific pathogen-free (SPF) rats, of the Wistar strain were used. These rats had been bred at the Unit from stocks given to us by Imperial Chemical

Industries, Pharmaceutical Division at Alderley Edge, Cheshire in 1964 and 1968.

The following materials were used:

1. *SFA chrysotile*.—A super fine sample obtained from a Canadian mine, and produced by water sedimentation separation from grade 7, the most fully milled commercial product.

2. *Crocidolite*.—Prepared from virgin fibre from a mine in the North West Cape. Both (1) and (2) were from the same samples as used in the earlier experiments (Wagner and Berry, 1969).

3. *UICC Standard reference samples*.—Samples of amosite, anthophyllite, Canadian chrysotile, Rhodesian chrysotile and crocidolite (Timbrell, Gilson and Webster, 1968) prepared following recommendations of l'Union Internationale Contre le Cancer (UICC).

4. *Benzene-extracted UICC Standard reference samples*.—Samples of (3) which had been repeatedly extracted for 64 hours by hot benzene using a Soxhlet apparatus to remove oils and other benzene-soluble substances. After extraction the benzene was

1 rats per dose, and
March 1966.

SFA chrysotiles

Materials were 7 of the
samples, SFA chrysotile.
The dose was 20 mg per
rats for each Canadian
chrysotile and 48 controls
during December 1966.

Canadian samples and Canadian

used were the 5 (100
both in the normal and
S Canadian chrysotile
and barium sulphate and
dose was 20 mg per rat
for each of the Canadian
each of the other treat-
ment took place between
February 1968.

Inoculation dusts

ected were ceramic fibre,
wider, aluminium oxide,
also the second sample of
e dose was 20 mg per rat
to 36 rats per treatment
of animals it was not
36 to all treatments and
tion fatalities could not
on took place in June

ment animals were allo-
cations. The age of the
was about 6 weeks for
13 and 13 weeks for
4. In Experiments 1 and
1 numbers of male and
Experiment 2 there were
males as males, and in
there were twice as many

al materials were made up
physiological saline with a
1 mg/ml for Experiments
Experiment 1 the con-
h that the required dose
in 0.4 ml of suspension,
thetized with ether and a
a two-way tap was then
right axilla at the level of
One arm of the two-way
a capillary manometer.

which gave a negative reading when the
needle reached the pleural cavity. Details of
the method of inoculation were given by
Wagner and Berry (1969). Following injec-
tion the rats were caged in fours isolated in a
special unit. They were fed on a proprietary
brand of autoclaved cubes, and water *ad
libitum*. Each rat was allowed to live until it
died or appeared to be distressed and a full
necropsy examination was carried out, except
for a few which had been cannibalized.

The results have been analysed using the
model given by Pike (1966) and shown to be
valid for experiments of this type (Berry and
Wagner, 1969). Fuller details are given in
in the appendix, and it need only be noted that
this method of analysis allows a constant c to
be estimated for each of the treatment groups
and that this constant, which we will refer to
as the "carcinogenicity factor", serves as a
single index summarizing the mesothelioma
experience of each group. It combines the
information on the proportion of animals
which developed a mesothelioma with the
times after inoculation at which the meso-
theliomata occurred. Also, since the method
of estimation eliminates mortality due to
other causes, chance variations in natural
mortality between different treatment groups
do not affect the treatment comparisons, nor
do systematic differences in natural mortality
between different experiments, such as that
resulting from the animals in Experiments 2
and 4 being older than those in Experiments
1 and 3, affect comparisons between experi-
ments.

Where significance levels are quoted they
are usually based on the *chi-squared* approxi-
mation to a likelihood-ratio test.

RESULTS

There were 13 rats for which histo-

logical examination was not possible,
leaving 1112 rats included in the results.

The predominant finding was that a
high proportion of most asbestos treated
groups developed mesotheliomata and the
results will be given mainly in terms of the
number of mesotheliomata and the time
when they occurred. Some details of
the results are given in Tables I-IV. A
total of 386 mesotheliomata occurred but
the histological features of these tumours
will not be described here as there is
nothing to add to the features described
for the original experiments (Wagner and
Berry, 1969). Also, in the presentation
and analysis of the results, no account has
been taken of the sex of the rats. The
original experiments show males and
females equally likely to develop a meso-
thelioma, and the present experiments
confirm this.

Experiment 1

There is a relationship between the
number of mesotheliomata and the dose
for both SFA chrysotile and crocidolite.
This implies that the carcinogenicity is
related to dose (d) and we considered this
relationship in the form of the carcino-
genic factor being proportional to a
power of dose, i.e. $c = bd^p$ where b and p
are constants. The power p was estimated
separately for each dust, giving 0.73 for
chrysotile and 0.96 for crocidolite. Be-
cause of the small number of mesothe-
liomata, however, these estimates are not
very precise and the approximate 95%
limits are 0.3-1.3 and 0.2-1.9 for chry-
sotile and crocidolite respectively. There

TABLE I.—Experiment 1 Results

		Number of rats with histology	Number with a mesothelioma	Survived time (days) of first mesothelioma	Mean survival (days)
SFA chrysotile	0.5 mg	12	1	512	754
SFA chrysotile	1 mg	11	3	615	729
SFA chrysotile	2 mg	12	5	425	664
SFA chrysotile	4 mg	12	4	470	762
SFA chrysotile	8 mg	12	8	496	692
Crocidolite	0.5 mg	11	1	992	809
Crocidolite	1 mg	12	0	-	760
Crocidolite	2 mg	12	3	562	777
Crocidolite	4 mg	13	2	917	819
Crocidolite	8 mg	11	5	799	689

TABLE II.—*Experiment 2 Results*

	Number of rats with histology	Number with a mesothelioma	Survival time (days) of first mesothelioma	Mean survival (day)
Canadian chrysotile A	16	8	416	642
Canadian chrysotile B	16	10	416	594
Canadian chrysotile C	16	5	188	702
Canadian chrysotile D	16	10	461	624
Canadian chrysotile E	16	7	405	573
Canadian chrysotile F	16	10	421	619
Canadian chrysotile H	16	4	384	602
SFA chrysotile	32	22	376	553
Control	48	0	—	728

TABLE III.—*Experiment 3 Results*

Material	Number of rats with histology	Number with a mesothelioma	Survival time (days) of first mesothelioma	Mean survival (days)
<i>UICC samples</i>				
Amosite	32	12	377	716
Amosite benzene-extracted	32	11	590	718
Anthophyllite	32	8	498	761
Anthophyllite benzene-extracted	32	14	533	728
Chrysotile (Canadian)	32	10	541	747
Chrysotile (Canadian) benzene-extracted	32	9	632	753
Chrysotile (Rhodesian)	31	7	502	693
Chrysotile (Rhodesian) benzene-extracted	32	5	659	686
Crocidolite	32	19	586	682
Crocidolite benzene-extracted	30	19	468	657
Canadian chrysotile A	24	14	488	712
Canadian chrysotile B	22	9	437	636
Canadian chrysotile C	24	9	460	717
Canadian chrysotile D	23	12	534	669
Canadian chrysotile E	21	9	489	660
Canadian chrysotile F	21	13	484	675
Canadian chrysotile G	23	16	576	659
Canadian chrysotile H	23	14	429	663
Brucite	32	18	502	680
Barium sulphate	30	1	436	783
Saline control	32	0	—	818

TABLE IV.—*Experiment 4 Results*

	Number of rats with histology	Number with a mesothelioma	Survival time (days) of first mesothelioma	Mean survival (days)
Ceramic fibre	31	3	743	736
Fibreglass	35	0	—	774
Glass powder	36	1	516	751
Aluminium oxide	35	1	646	710
SFA chrysotile	36	23	325	568
SFA chrysotile (2nd sample)	32	21	382	639

are some theoretical grounds for choosing p to be an integer and therefore p was taken as unity. The values of the carcinogenicity factor adjusted to a dose of 20 mg were then 4.10×10^{-9} for chrysotile and 1.70×10^{-9} for crocidolite.

Experiments 2, 3 and 4

Comparing first the effects of the separate Canadian samples (Table V), there is considerable variation between Experiments 2 and 3 and this is mainly because of the small number of animals in

TABLE V.—*Experiments 2 and 3*

Material	Number of rats with histology	Number with a mesothelioma	Survival time (days) of first mesothelioma	Mean survival (day)
Canadian chrysotile	16	8	416	642
Canadian chrysotile	16	10	416	594
Canadian chrysotile	16	5	188	702
Canadian chrysotile	16	10	461	624
Canadian chrysotile	16	7	405	573
Canadian chrysotile	16	10	421	619
Canadian chrysotile	16	4	384	602
SFA chrysotile	32	22	376	553
SFA chrysotile (2nd sample)	32	21	382	639
Brucite	32	18	502	680
Barium sulphate	30	1	436	783
Ceramic fibre	31	3	743	736
Glass powder	36	1	516	751
Aluminium oxide	35	1	646	710

each group. Table V shows that the results of Experiment 2 are not significantly different from the results of Experiment 3 and that the lowest carcinogenicity factor was obtained from this group of samples. The results of Experiment 4 are also not significantly different from the results of Experiment 3 and the results of Experiment 4 are also not significantly different from the results of Experiment 2. The correlation of the results of the three experiments is therefore very good. The correlation of the results of the three experiments is therefore very good. The correlation of the results of the three experiments is therefore very good.

Turning now to the results of the three experiments, the results of Experiment 2 are not significantly different from the results of Experiment 3 and the results of Experiment 4 are also not significantly different from the results of Experiment 3 and the results of Experiment 4 are also not significantly different from the results of Experiment 2.

TABLE V.—*Estimates of Carcinogenicity Factor ($\times 10^9$) for Experiments 2, 3 and 4*

Time of first tumour	Mean survival (days)		Experiment		
			2	3	4
642		Canadian chrysotile A	1.25	1.20	—
594		Canadian chrysotile B	2.23	1.16	—
702		Canadian chrysotile C	0.68	0.67	—
624		Canadian chrysotile D	2.39	1.23	—
573		Canadian chrysotile E	1.89	0.97	—
619		Canadian chrysotile F	1.90	1.40	—
602		Canadian chrysotile G	—	2.09	—
553		Canadian chrysotile H	1.10	1.84	—
728		SFA chrysotile	4.72	—	2.85
		SFA chrysotile (2nd sample)	—	—	2.28
		Brucite	—	1.21	—
		Barium sulphate	—	0.04	—
		Ceramic fibre	—	—	0.16
		Glass powder	—	—	0.04
		Aluminium oxide	—	—	0.05

Time of first tumour

377	716
590	718
498	761
533	728
541	747
632	753
502	693
659	686
586	682
498	687
488	712
437	636
430	717
534	669
—	660
—	675
—	659
429	663
502	680
436	783
—	818

Time of first tumour	Mean survival (days)
1	736
2	774
3	751
4	710
5	568
6	639

Table 4
the effects of the
samples (Table 1).
ble variation bet
1.3 and this is m
ll number of animals

each group. There was overall about 30% more carcinogenicity in Experiment 2 than in Experiment 3 but the difference is not significant ($P > 0.1$). Sample C has the lowest carcinogenicity in each experiment and overall is significantly the least carcinogenic ($P < 0.05$) but apart from this no differences between the samples were detected. These samples have been analysed for certain metals (Holmes, Morgan and Sandalls, 1971; Morgan and Calley, 1973) and in Table VI the results of these analyses are shown, together with the carcinogenicity factor obtained by combining the 2 experiments. The correlation coefficients between the carcinogenicity factor and the different metals are -0.13 for iron, -0.58 for chromium, 0.04 for cobalt, -0.02 for nickel, -0.39 for scandium and -0.04 for manganese. None of these is significant and it is reasonably clear from examination of the chemical properties of sample C that the low carcinogenicity of this sample is not because of a low content of any of these metals.

Turning now to the UICC reference samples (Table VII), there are no significant differences between the normal and benzene-extracted samples, and overall 58 mesotheliomata occurred with the benzene-extracted samples and 56 with the untreated samples. Crocidolite was the most carcinogenic sample with the

others in order amosite, anthophyllite, Canadian chrysotile and Rhodesian chrysotile. The difference between the two samples of chrysotile is not significant. Holmes *et al.* (1971) also carried out chemical analyses on the UICC reference samples. There were very large differences between the samples in the amount of the different metals present and it is clear that these bear no relation to the carcinogenicity.

The sample of brucite proved as carcinogenic as the Canadian samples of chrysotile (Table V). Non-asbestos materials which produced the occasional mesothelioma were ceramic fibre, barium sulphate, glass powder and aluminium oxide.

The second sample of SFA chrysotile proved similar in carcinogenic effect to the original sample.

DISCUSSION

The application of the test materials by intrapleural inoculation may be criticized as unrealistic in comparison with human exposure, about which the animal experiments are intended to provide relevant information. Nevertheless, this type of experiment has an important part to play. With an inhalation experiment, which provides a realistic route of entry of the test material, there are 2 factors involved. First, the penetration of dust through the airways and alveoli will differ with different samples of dust (Timbrell, 1965). The second factor is the effect of the dust, given that it has reached the pleura. In inoculation experiments only the second factor is relevant and hence these experiments are simpler to interpret. This makes intrapleural inoculation a more suitable method for the investigation of questions such as whether extraction of the oils alters the carcinogenicity of an asbestos sample. The two types of experiment supplement one another and we will be reporting separately on 2 experiments in which rats were exposed to dust clouds of the UICC reference samples.

The varying dose experiment gave results which indicated that the risk of developing a mesothelioma at a given time after injection was proportional to the dose. This form of dose relationship was also found by Pike and Doll (1965) for lung cancer and smoking in man whereas Lee and O'Neill (1971) showed that after repeated applications of benzo(a)pyrene to the backs of mice the incidence rate of tumours was proportional to the square of the dose.

The carcinogenicity of the SFA chrysotile sample was similar in Experiments 1 and 2, after adjusting the former to a dose of 20 mg, and in Experiment 4 was lower but not significantly so. In all these 3 experiments the carcinogenicity of the SFA chrysotile was significantly greater than in the earlier experiment (Wagner and Berry, 1969) when the estimate of the carcinogenicity factor was 1.68×10^{-9} . The crocidolite was also more carcinogenic in Experiment 1 than in the earlier experiment ($c = 1.16 \times 10^{-9}$) but not significantly so. These differences could be the result of a change in susceptibility of the rats or of a change in the dust during storage.

The suggestion that natural oils and waxes (Harington, 1962), or contaminating oils from the preparation of the fibre (Harington and Roe, 1965; Roe, Walters and Harington, 1966) or from plastic storage bags (Commins and Gibbs, 1969) might contribute to the carcinogenicity of asbestos receives no support from our present experiments, which is in agreement with our original experiments (Wagner and Berry, 1969) when removal of the oils from the crocidolite sample resulted in no detectable change in carcinogenicity.

Harington and Roe (1965) also advanced the possibility that the presence of trace metals might be relevant to the carcinogenicity of asbestos. In our experiments with the Canadian samples the carcinogenicity was not related to the content of iron, chromium, cobalt, nickel, scandium or manganese. Also, the fact

that all the types of asbestos, having very different chemical compositions, produce mesotheliomata makes it unlikely that the carcinogenicity of asbestos could be due to chemical properties.

Our experiments offer some evidence that the development of mesotheliomata is associated with the presence of fine fibrous material within the pleural cavity. First, UICC Canadian chrysotile is a mixture of batches of material from 8 Canadian mines, and the separate samples used were taken from the same batches (Timbrell and Rendall, 1971). The main difference in the subsequent preparation of the material was that the separate Canadian samples were ground more finely than the composite UICC sample. Comparing Tables V and VII, the carcinogenicities of all the separate Canadian samples were greater than that of the UICC Canadian chrysotile. Also, the samples of SFA chrysotile were from mine D. These were superfine samples and resulted in a very high carcinogenicity. It should be noted that of the Canadian samples the one with the lowest carcinogenicity (C) in both experiments was from a mine in British Columbia whilst the others were from 7 mines in the Quebec area. However, sample C could not be distinguished from the other samples by its size distribution.

A full quantitative analysis of our experimental results will only be possible when techniques are available for complete size characterization of the experimental materials, both before injection and present in the lungs at postmortem. Such techniques to determine the mass and the diameter and length distributions of the particles are being developed. However, even with the characterization methods at present available, a relationship emerges between the observed incidence of mesotheliomata and the physical factors.

A further factor that must be taken into account is the tendency of chrysotile fibres to fragment longitudinally into fibrils in lung fluids, the degree of frag-

TABLE VI

Sample	Carcinogenicity Factor
A	
B	
C	
D	
E	
F	
G	
H	
I	
J	
K	
L	
M	
N	
O	
P	
Q	
R	
S	
T	
U	
V	
W	
X	
Y	
Z	

TABLE VII.—*Ex*
Factor $\times 10^9$
Samples in *Ex*

A	crocidolite
B	chrysotile
C	chrysotile (Canadian)
D	chrysotile (Rhodesian)
E	chrysotile

mentation and fibres and fibrils the precise physical conditions. Any on the other hand fibre-diameter should appear to retain Pooley and Wagner.

To illustrate electron micrographs are presented of their cross-sections give short consideration to those that are less and also greater. The non-chrysotile considered first. In for UICC crocidolite (Fig. 1), ceramic fibre (Fig. 2) it is of a significant decreasing carcinogenicity. The glass fibre, fibres but the material is $< 5 \mu\text{m}$. Or

TABLE VI.—*Carcinogenicity and Chemical Analysis of Canadian Samples—Experiments 2 and 3 Combined*

Sample	Carcinogenicity factor ($\times 10^3$)	Iron (%)	Chromium parts/10 ⁶	Cobalt parts/10 ⁶	Nickel parts/10 ⁶	Scandium parts/10 ⁶	Manganese parts/10 ⁶
G	2.37	1.9	380	53	1400	7.5	420
D	1.70	4.0	930	78	1550	6.6	600
F	1.40	2.9	730	78	1900	7.7	450
B	1.59	3.4	780	110	2400	4.1	580
H	1.54	3.2	480	42	550	5.6	530
E	1.33	3.2	435	57	895	5.4	610
A	1.22	4.8	515	63	1150	5.0	540
C	0.67	2.0	1200	60	1800	12.0	420

TABLE VII.—*Estimates of Carcinogenicity Factor ($\times 10^3$) for UICC Reference Samples in Experiment 3*

	Normal form	Benzene-extracted form
Amosite	0.66	0.66
Anthophyllite	0.33	0.77
Chrysotile (Canadian)	0.50	0.42
Chrysotile (Rhodesian)	0.44	0.31
Crocidolite	1.45	1.87

mentation and hence the number of fibres and fibrils produced depending on the precise physical and physiological conditions. Amphibole types of asbestos, on the other hand, have characteristic fibre-diameter distributions which they appear to retain in lung tissue (Timbrell, Pooley and Wagner, 1970).

To illustrate this relationship, the electron micrographs of some of the materials are presented in decreasing order of their carcinogenicity (Fig. 1-8). For reasons given by Timbrell (1973), we shall consider as "significant" fibres those that are less than $0.5 \mu\text{m}$ in diameter and also greater than $10 \mu\text{m}$ in length. The non-chrysotile materials will be considered first. In the electron micrographs for UICC crocidolite (Fig. 2), UICC amosite (Fig. 4), UICC anthophyllite (Fig. 5), ceramic fibre (Fig. 7) and glass fibre (Fig. 8) it is evident that the number of "significant" fibres decreases with decreasing carcinogenicity of the materials. The glass fibre, for instance, contains long fibres but the majority of these are thicker than $0.5 \mu\text{m}$. On the other hand, whereas

a high proportion by weight of the brucite consists of large fibres, there are also present a number of very fine long fibrils. It is difficult to compare chrysotile samples (Fig. 1 and 6) with other types of material on this basis. But even so, the relative positions of the chrysotiles in the classification by carcinogenicity appears to correspond with the number of "significant" fibres present. For example, although the SFA chrysotile (Fig. 1) contains a high proportion by weight of non-fibrous particles, even before injection the fibres were in a highly dispersed state. The enormous number of fibres that complete fragmentation of chrysotile can produce will be clear from the illustration that a single fibre may fragment into 1000 fibrils. The fact that this SFA sample was the most carcinogenic of all the materials used corresponds to its highly dispersed state and its high content of "significant" fibres.

The above theory has been examined further using the results of Stanton and Wrench (1972). Their experiments were similar to ours and they used 17 samples, including several materials (UICC samples and glass fibres) after partial pulverization. They analysed their results by discounting submicroscopic fibrils and converting all longer fibres into microfibrils of standard size ($1.25 \times 3.75 \mu\text{m}$) on the assumption that fragmentation of both glass and asbestos occurred *in vivo*. The numbers of microfibrils were then compared with the carcinogenicity of the materials. At first sight their results seem to conflict with our findings. But,

os, having by compositions, produce it unlikely that the bestos could be due

s. offer some evidence it of mesotheliomata the presence of fine in the pleural cavity, lian chrysotile is a of material from s

t the separate samples on the same batches dl. 1971). The main bsequent preparation is that the se) de

were ground a-

posite UICC sample, and VII, the carcino-

e separate Canadian er than that of the

hrysotile. Also, the

osotile were from mine

aperfine samples and

gh carcinogenicity. It

hat of the Canadian

th the lowest carcino-

periments was from

umbia whilst the

mines in the Quebec

ample C could not be

the other sample why...

ative analysis of our

s will only be possible

re available for com-

rization of the experi-

both before injection

lungs at postmortem.

o determine the mass

nd length distributions

are being developed.

th the characterization

it available, a relation-

liomata and the physical

or that must be taken

tendency of chry

longitudinally into

ds, the degree of frag-



FIG. 1.—Electronmicrograph of SPA chrysotile ($\times 1875$).



FIG. 2.—Electronmicrograph of UIC crocidolite ($\times 1875$).

10 μ m



FIG. 1.—Electromicrograph of SFA dry-salt ($\times 1875$).



FIG. 2.—Electromicrograph of UIV crocidolite ($\times 1875$).



FIG. 3.—Electromicrograph of UIV amosite ($\times 1875$).

FIG. 4.—Electromicrograph of UIV amosite ($\times 1875$).



FIG. 3. Electron micrograph of UICV anthophyllite ($\times 1875$).

FIG. 6. Electron micrograph of UICV Canadian chrysotile ($\times 1875$).



FIG. 5.—Electron micrograph of UICC anthophyllite (1875).

FIG. 6.—Electron micrograph of UICC Canadian chrysotile (1875).



FIG. 7.—Electron micrograph of ceceane fibre (1875).

FIG. 8.—Electron micrograph of glass fibre (1875).

when their materials were assessed in the manner used for our own samples, the order obtained by classification of the materials according to the number of "significant" fibres was in good agreement with the reported order of carcinogenicity. The smaller number of mesotheliomata observed from pulverized materials did not correlate with the estimated total number of particles that this pulverization would produce, but did correlate with the estimated number of "significant" fibres.

Stanton and Wrench (1972) also produced mesotheliomata with very fine fibreglass. Mesotheliomata were not produced by our sample of fibreglass but they were by synthetic aluminium silicate fibre, which was finer than our glass fibre (Fig. 7 and 8). Again, the apparent contradiction is explicable in terms of physical characterizations. The 2 samples of fibreglass can only be compared using light microscope size data; in our sample 55% of the fibres had diameters exceeding $2.5 \mu\text{m}$ compared with less than 10% of Stanton and Wrench's sample.

Although a direct association between physical factors and the development of mesotheliomata has not been demonstrated, these characteristics appear to be the relevant properties. If the finer fibres are the more carcinogenic when applied to the pleura then, since the finer fibres are also able to penetrate to the pleura more easily after inhalation, these 2 factors would combine together to give the finer fibres more relative importance than even the aerodynamic differences would suggest.

Clearly, the experiments described in this paper have involved a large amount of daily effort over a number of years and we are grateful to all our colleagues who have contributed to this. We are also grateful to Dr B. T. Commins of the MRC Air Pollution Unit who prepared the benzene-extracted samples and the barium sulphate sample.

REFERENCES

- BERRY, G. & WAGNER, J. C. (1969) The Application of a Mathematical Model Describing the Times of Occurrence of Mesotheliomas in Rats following Inoculation with Asbestos. *Br. J. Cancer*, **23**, 582.
- COMMINS, B. T. & GIBBS, G. W. (1969) Contaminating Organic Material in Asbestos. *Br. J. Cancer*, **23**, 358.
- HARINGTON, J. S. (1962) Occurrence of Oils Containing 3:4-Benzopyrene and Related Substances in Asbestos. *Nature, Lond.*, **193**, 43.
- HARINGTON, J. S. & ROE, F. J. C. (1965) Studies of Carcinogenesis of Asbestos Fibres and their Natural Oils. *Ann. N.Y. Acad. Sci.*, **132**, 439.
- HOLMES, A., MORGAN, A. & SANDALLS, F. J. (1971) Determination of Iron, Chromium, Cobalt, Nickel and Scandium in Asbestos by Neutron Activation Analysis. *Am. industr. Hyg. Ass. J.*, **32**, 281.
- LEE, P. N. & O'NEILL, J. A. (1971) The Effect of Time and Dose Applied on Tumour Incidence Rate in Benzopyrene Skin Painting Experiments. *Br. J. Cancer*, **25**, 759.
- MORGAN, A. & CRALLEY, L. J. (1973) Chemical Characteristics of Asbestos and Associated Trace Elements. In *Proc. Conf. Biological Effects of Asbestos*, Lyon, 2-5 October, 1972. In print.
- PIKE, M. C. (1966) A Method of Analysis of a Certain Class of Experiments in Carcinogenesis. *Biometrics*, **22**, 142.
- PIKE, M. C. & DOLL, R. (1965) Age at Onset of Lung Cancer: Significance in Relation to Effect of Smoking. *Lancet*, **i**, 665.
- ROE, F. J. C., WALTERS, M. A. & HARINGTON, J. S. (1966) Tumour Initiation by Natural and Contaminating Asbestos Oils. *Int. J. Cancer*, **1**, 491.
- STANTON, M. F. & WRENCH, C. (1972) Mechanisms of Mesothelioma Induction with Asbestos and Fibrous Glass. *J. natn. Cancer Inst.*, **48**, 797.
- TIMBRELL, V. (1965) The Inhalation of Fibrous Dusts. *Ann. N.Y. Acad. Sci.*, **132**, 255.
- TIMBRELL, V. (1973) Physical Factors as Aetiological Mechanisms. In *Proc. Conf. Biological Effects of Asbestos*, Lyon, 2-5 October, 1972. In print.
- TIMBRELL, V., GILSON, J. C. & WEBSTER, L. (1968) UICC Standard Reference Samples of Asbestos. *Int. J. Cancer*, **3**, 406.
- TIMBRELL, V., POOLEY, F. & WAGNER, J. C. (1970) Characteristics of Respirable Asbestos Fibres. In *Pneumoconiosis. Proc. Internat. Conf. Johannesburg, 1969*. Ed. H. A. Shapiro. Cape Town: Oxford University Press. p. 120.
- TIMBRELL, V. & RENDALL, R. E. G. (1971) Preparation of the UICC Standard Reference Samples of Asbestos. *Powder Technol.*, **5**, 279.
- WAGNER, J. C. (1970) The Pathogenesis of Tumours following the Intrapleural Injection of Asbestos and Silica. In *Morphology of Experimental Respiratory Carcinogenesis. Proc. Conf. Gathburg 13-18 May 1970*. Ed. P. Nettesheim, M. G. Hanna Jr. and J. W. Deatherage Jr. U.S. Atomic Energy Commission Symposium Series No. 21. p. 347.
- WAGNER, J. C. (1972) The Significance of Asbestos in Tissue. In *Recent Results in Cancer Research*, Vol. 39. *Current Problems in the Epidemiology of Cancer and Lymphomas*. Ed. E. Grundmann and H. Tulinius. Berlin: Springer-Verlag. p. 37.
- WAGNER, J. C. & BERRY, G. (1969) Mesotheliomas following the intrapleural injection of asbestos. *Cancer*, **23**, 567.
- WAGNER, J. C. & BERRY, G. (1969) Mesotheliomas in Rats following the intrapleural injection of Asbestos. *Proc. Internat. Conf. Johannesburg, 1969*. Ed. H. A. Shapiro. Cape Town: Oxford University Press. p. 216.

APP

The results have been taken following the model given by Berry and Wagner to be valid for experimental specific death rate of a mesothelioma taken as $ck(t-t_0)^{k-1}$ constants. These estimated separately each experiment but the estimates make the original experiment 417 mesotheliomata could be taken as 3 that in the lapse period mesotheliomata occurred; in particular amosite was found longer than that crocidolite but the mesothelioma occurred only 398 days. For reported here, although the lapse period of different treatments, but due to the imprecision and it is no strong evidence there is no evidence being dependent on the analysis the best based on all our these are $m = 270$ with all the data that very precise; for example

- (1969) The Appearance of Mesotheliomas in Rats following Inoculation with Asbestos. *Br. J. Cancer*, 23, 582.
- G. W. (1969) Continuous Asbestos. *Br. J. Cancer*, 23, 587.
- WAGNER, J. C., BERRY, G. & TIMBRELL, V. (1970) Mesotheliomas in Rats Following the Intra-pleural Inoculation of Asbestos. In *Pneumoconiosis. Proc. Internat. Conf. Johannesburg, 1969*. Ed. H. A. Shapiro. Cape Town: Oxford University Press. p. 216.
- Y. L. J. (1973) Asbestos and Associated Lung Cancer. *Conf. Biological Effects of Asbestos*, October, 1972. In print.
- Method of Analysis of a Certain Type of Carcinogenesis. *Int. J. Cancer*, 1973.
- (1965) Age at Onset of Lung Cancer in Relation to Effect of Asbestos. *Int. J. Cancer*, 1, 491.
- M. A. & HARRINGTON, J. S. (1972) Mechanism of Action of Asbestos and Crocidolite. *Cancer Inst.*, 48, 797.
- Inhalation of Fibrous Asbestos. *Int. J. Cancer*, 1972.
- Physical Factors as Aetiological in Lung Cancer. *Conf. Biological Effects of Asbestos*, October, 1972. In print.
- J. C. & WEBSTER, J. (1969) Reference Samples of Asbestos. *Int. J. Cancer*, 1, 129.
- F. & WAGNER, J. C. (1970) Respirable Asbestos Fibres. *Proc. Internat. Conf. Johannesburg, 1969*. Ed. H. A. Shapiro. Cape Town: Oxford University Press. p. 129.
- ALL, R. E. G. (1971) Preparation of Standard Reference Samples of Asbestos. *Int. J. Cancer*, 5, 279.
- The Pathogenesis of Tumours following Intra-pleural Injection of Asbestos. *Proc. Conf. Guelph, 1970*. Ed. P. Nettesheim. and J. W. Deatherage Jr. New York: Commission Symposium. p. 47.
- The Significance of Asbestos in Lung Cancer. *Br. J. Cancer*, 1970. Ed. E. G. W. Berlin: Springer-Verlag. p. 17.
- KEY, G. (1969) Mesotheliomas in Rats following Inoculation with Asbestos. *Br. J. Cancer*, 23, 587.
- WAGNER, J. C., BERRY, G. & TIMBRELL, V. (1970) Mesotheliomas in Rats Following the Intra-pleural Inoculation of Asbestos. In *Pneumoconiosis. Proc. Internat. Conf. Johannesburg, 1969*. Ed. H. A. Shapiro. Cape Town: Oxford University Press. p. 216.

APPENDIX

The results have been analysed using the model given by Pike (1966) and shown to be valid for experiments of this type (Berry and Wagner 1969). The age-specific death rate of animals dying with a mesothelioma t days after injection is taken as $ck(t-w)^{k-1}$ where c , k and w are constants. These 3 constants could be estimated separately for each treatment of each experiment but correlations between the estimates make them imprecise. In the original experiments, with a total of 417 mesotheliomata, it was shown that k could be taken as 3 for all treatments but that w , the lapse period before any mesotheliomata occurred, varied with treatment; in particular the lapse period for amosite was found to be about 200 days longer than that for chrysotile and crocidolite but there was an isolated mesothelioma occurring with amosite after only 398 days. For the experiments being reported here, although the estimates of the lapse period vary widely over the different treatments, this wide range could be due to the imprecision of the estimates, and there is no strong evidence that it is invalid to use a common value. Also, there is no evidence of the lapse period being dependent on dose. Therefore, in the analysis the best estimates of k and w based on all our evidence were used; these are $w = 270$ and $k = 3.25$. Even with all the data these estimates are not very precise; for example the pairs (300,

2.9) and (220, 3.8) would be acceptable, as also would the estimates (250, 3.0) which we used in our preliminary reports. However, these alternatives all lead to similar conclusions.

The constant c was estimated for each of the treatment groups and this constant, which we will refer to as the "carcinogenicity factor", serves as a single index summarizing the mesothelioma experience of each treatment group. It combines the information on the proportion of animals which developed a mesothelioma with the times after inoculation at which the mesotheliomata occurred. Also, the method of estimation eliminates mortality due to other causes, so that neither chance nor systematic variations in natural mortality between different groups will affect comparisons between these groups.

As an example of the elimination of natural mortality, the proportion of rats developing mesotheliomata after injection with SFA chrysotile in Experiments 2 and 4 (69% and 64%) were similar to the proportion (65%) in SPF rats in our original experiment (Wagner and Berry, 1969). However, the rats in Experiments 2 and 4 were injected at 13 weeks of age opposed to 6 weeks in our original experiment, and also the natural mortality was less in our original experiment, even after allowing for this age difference. Hence the carcinogenicity of the chrysotile was least in our original experiment and this is reflected in the values of the carcinogenicity factor which were 1.7×10^{-9} in the original experiment, 4.7×10^{-9} in Experiment 2 and 2.9×10^{-9} in Experiment 4 (Table V). The non-significant difference between the values for Experiments 2 and 4 was revealed after eliminating the chance lower natural mortality in the group used in the latter experiment.