

FILE NAME: Metropolitan Life (ML)

DATE: 1930

DOC#: ML211

DOCUMENT DESCRIPTION: Unpublished Report of the Physical Examinations of Asbestos by Pedley

Jandwer 1370007 -

REPORT OF THE PHYSICAL EXAMINATIONS
EXAMINATIONS OF ASBESTOS WORKERS IN Asbestos
AND THETFORD MINES, QUEBEC

~~1/2~~ The report herewith presented deals with the examination of 141 employees of the Canadian Johns-Manville Co., Asbestos, Que. and 54 employees of the Asbestos Corporation of Canada, the Keasby-Mattison Co. and Johnson's Mine, all of Thetford Mines, Que. 40 of the men of the Canadian Johns-Manville Co. were examined by Dr. D. M. Stevenson and Dr. Joseph Wyatt, who performed the physical examinations and X-ray examinations respectively. The other examinations were all performed by me, except the X-ray examinations. Actually, Drs. Stevenson and Wyatt examined 57 men; but since 17 of these men were re-examined by me I am including them in my list. Drs. Stevenson and Wyatt carried on their work in November 1929. My examinations were done in July 1930 in Asbestos, and in September 1930 in Thetford Mines.

The Asbestos and Thetford groups are very dissimilar with respect to age distribution and length of service in the asbestos industry, but racially and economically the two groups are much alike.

Table I gives the age distribution of the men and Table II the distribution according to length of service in the asbestos industry.

REPORT OF THE PHYSICAL TEAM / FBI AND U.S. DEPT. OF HEALTH
EXAMINATIONS OF ASBESTOS EXPOSURE IN A
AND THETFORD MINES, QUEBEC.

The report herewith presented deals with the examination of 141 employees of the Canadian John-McNeill Co., Asbestos, Que. and 54 employees of the Asbestos Corporation of Canada, the Thetford-Mattison Co. and John-McNeill's Mine, all of Thetford Mines, Que. 40 of the men of the Canadian John-McNeill Co. were examined by Dr. W. H. Stevenson and Dr. Joseph Hyatt, who performed the physical examinations and X-ray examinations respectively. The other examinations were all performed by me, except the X-ray examinations. Actually, Drs. Stevenson and Hyatt examined 57 men; but since 17 of these men were re-examined by me I am including them in my list. Drs. Stevenson and Hyatt carried on their work in November 1929. The examinations were done in July 1930 in Asbestos, and in September 1930 in Thetford Mines.

The Asbestos and Thetford groups are very dissimilar with respect to age distribution and length of service in the asbestos industry, but racially and economically the two groups are much alike.

Table I gives the age distribution of the men and Table II the distribution according to length of service in the asbestos industry.

REPORT ON THE PHYSICAL AND X-RAY EXAMINATIONS OF ASBESTOS MINERS IN THE
AND THETFORD MILLS, UTAH.

The report herewith presented deals with the examination of 141 employees of the Canadian Johna-Johnville Co., Arbeston, and 54 employees of the American Corporation of Canada, the Thetford-Mattinson Co. and Johna-Johnville Mine, all of Thetford Mines, Que. 40 of the men of the Canadian Johna-Johnville Co. were examined by Dr. W. H. Stevenson and Dr. Joseph Hyatt, who performed the physical examinations and X-ray examinations respectively. The other examinations were all performed by me, except the X-ray examinations. Actually, Drs. Stevenson and Hyatt examined 57 men; but since 17 of these men were re-examined by me I am including them in my list. Drs. Stevenson and Hyatt carried on their work in November 1929. The examinations were done in July 1930 in Arbeston, and in September 1930 in Thetford Mines.

The Arbeston and Thetford groups are very dissimilar with respect to age distribution and length of service in the asbestos industry, but racially and economically the two groups are much alike.

Table I gives the age distribution of the men, and Table II the distribution according to length of service in the asbestos industry.

TABLE 3.

AGE DISTRIBUTION OF ACIDWATER MONITORS

<u>AGE GROUP</u>	<u>Amherst</u>		<u>Thetford Falls</u>	
	<u>Number</u>	<u>Percentage</u>	<u>Number</u>	<u>Percentage</u>
15 - 19	3	2.1		
20 - 24	24	17.0	2	3.7
25 - 29	24	17.0	5	9.3
30 - 34	26	18.5	6	11.1
35 - 39	25	17.7	4	7.4
40 - 44	18	12.8	15	27.8
45 - 49	14	10.0	7	12.9
50 - 54	3	2.1	6	11.1
55 - 59	1	0.7	4	7.4
60 - 64	1	0.7	6	11.1
65 - 69	1	0.7	1	1.9
70 - 74	1	0.7		
<u>TOTAL</u>	<u>141</u>	<u>100.00</u>	<u>54</u>	<u>100.0</u>

T A B L E I I

LENGTH OF SERVICE IN ASBESTOS INDUSTRY

Years of Service.	Asbestos		Thetford Mines	
	Number	Percentage	Number	Percentage
0 - 4	44	31.2	1	1.85
5 - 9	65	46.1	8	14.8
10 - 14	11	7.8	8	14.8
15 - 19	10	7.1	16	29.6
20 - 24	6	4.3	9	16.7
25 - 29	2	1.4	9	16.7
30 - 34	2	1.4	1	1.85
35 - 39	1	.7		
40 - 44			2	3.7
T O T A L	141	100.0	54	100.0

It will be noted that while 68 percent of the men in Thetford Mines were over 40 years of age only 28 percent of the men in Asbestos were over 40. And further with respect to length of service 83.4 percent of the men examined in Thetford Mines had worked in the asbestos industry for 10 years or more, but in Asbestos only 22 percent had worked for more than ten years.

Table III shows the disposition of the men in various parts of the industry. Since the numbers are

small broad occupational groupings are used. In Asbestos the occupations are divided into 3 groups - mill, pit and factory; in Thetford Mines only the first two occupational groups are used, since no asbestos manufacturing is carried on there. All the men of the Thetford Mines group could be so classified, but 5 men were excluded from the Asbestos group since they did not appear to be exposed to an unusual amount of dust.

T A B L E III

LENGTH OF SERVICE IN DIFFERENT OCCUPATIONS IN ASBESTOS INDUSTRY

Years of service	ASBESTOS			THETFORD MINES	
	Mill	Drilling pit	Factory	Mill	Drilling pit
0-4	14	7	24	1	
5-9	36	8	16	5	5
10-14	6	6		7	1
15-19	7	3		12	4
20-24	3	1		6	1
25-29	1	1		6	1
30-34	1	1		1	
35-39	1				
40-44					2
TOTAL	69	27	40	40	14

The past occupational history in industries other than the asbestos industry was significant only in 5 cases. Of these one had been a grain inspector

for 6 years and so exposed to very dusty conditions, one had been an underground copper miner for 18 years, one a copper miner for 2 years, a fourth had worked in a cement plant for 9 years, and the fifth had operated a drill in an iron mine. This last individual showed X-ray evidence of pneumoconiosis, the others did not have pneumoconiosis.

In Table IV are given observations of weight in relation to height by age. The average weights are taken from the figures compiled by the Association of Life Insurance Medical Directors and the Actuarial Society of America.

T A B L E IV.

WEIGHTS AND HEIGHTS ACCORDING TO AGE

	Pounds Under average				Pounds over average			
	20-29	10-19	5-9	-4 to 4	5-9	10-19	20-29	Age
AMERICAN	No. 21	23	19	24	16	19	9	4
	15.6	20.0	15.6	17.1	11.4	15.6	6.4	2.9
EUROPEAN	No. 9	11	8	9	8	6	7	1
	16.7	20.4	14.8	16.7	5.6	11.1	12.9	1.9
FROM 10,000 male Industrial workers	No. 15	18	11	21	9	14	10	5

The group of 10,000 male industrial workers referred to in the above table was studied by the United States Public Health Service (see Public Health Bulletin

No. 162). It will be noted that there is very little distinction between this large group and the group of asbestos workers so far as degrees of overweight or underweight are concerned.

In the consideration of individual defects some interesting facts have been found. In this study only the 101 men at Asbestos and the 54 men at Thetford Mines are considered, since the records of the men examined by Dr. Stevenson are not strictly comparable with mine.

TEETH:-

Dividing the men into two groups according to whether the teeth did or did not need attention, of the Asbestos group 49.5% required attention, and of the Thetford Mines group 51.8 %.

HEARING:-

Acuity of hearing was tested by means of a watch. (My own hearing, which has been tested by an audiometer, was considered normal, and used as a standard) The watch test is admittedly very rough. In noisy occupations impaired hearing is frequently found in the older employees. Of the 101 men at Asbestos 10 showed definite hearing loss. This is not an excess proportion since an

group of adults will show a similar picture. In Thetford Mines 28 of the 54 men suffered from well marked hearing loss. This is a very high percentage, and comparable to what may be found in weaving sheds and boiler shops. All these men, except two, had been exposed to loud noise in their work for 10 years or more, and it is reasonable to suppose that noise played a part in their hearing disability.

The number of men with infected TONSILLS was 27 in Asbestos and 13 in Thetford Mines, and almost identical ratio to the total in both groups.

Defects of the THYROID were rare. 4 men in Asbestos and 1 in Thetford Mines had enlarged thyroids.

BLOOD PRESSURE:-

Readings were recorded in all men over 40 years of age. Except in exceptional circumstances the blood pressure was not taken in the younger men. Consequently although the groups examined in Thetford Mines was smaller than that in Asbestos more blood pressure readings were taken there. 2 men in Asbestos had blood pressures of over 150 and 2 blood pressures under 100 m.m. of mercury. 8 men in Thetford Mines had blood pressures of over 150 and 2 blood pressures under 100.

The incidence of HERNIA is of interest from the compensation standpoint. In Asbestos all new employees must pass a physical examination before employment. There were 4 cases of inguinal hernia discovered among the 101 men examined there. In addition to these 5 men had been operated upon for this condition.

In Thetford Mines physical examination is not required before employment. There were 5 cases of inguinal hernia among the 54 men examined and 2 cases which had been operated upon.

Some interest has arisen in connection with the occurrence of the so-called ASBESTOS CORNS. These are usually pin head sized elevations of the skin, somewhat like minute warts. None of these were detected in the Asbestos men probably due to my ignorance of their appearance. 8 of the Thetford Mines men had these growths on their hands and several others reported having had them at one time or another.

L U N G S

Of the 101 men examined by me at Asbestos only 4 suffered from definite first stage pneumoconiosis. These will not be considered on account of the doubtful diagnosis.

Of the 40 men examined by Drs. Stevenson and Wyatt 14 were diagnosed as first stage pneumoconiosis (asbestosis 1) (I am including the 17 men examined by Drs. Stevenson and Wyatt and subsequently re-examined by me in my group of 101). The much lower incidence of pneumoconiosis in my group of men may seem surprising, but it must be remembered that the X-ray negatives were not read by the same men. In my group the X-ray diagnoses were made by Dr. Merivether of Pitcher, Oklahoma, and in the other group the interpretations were by Drs. Wyatt and ~~Stevenson~~ ^{Merivether}. Further the ~~group~~ ^{men} examined by Drs. Stevenson and Wyatt were, in general, older than those in my group.

Of the 54 men examined at Thetford Mines 24 were diagnosed by X-ray as suffering from pneumoconiosis (asbestosis), of these 4 were diagnosed as probably first stage asbestosis, 8 as early stage, 9 as definite first stage and 3 as second stage.

The pertinent facts regarding the definite cases of asbestosis are given in tables V and VI.

Table V

Case No.	Diagnosis	Age	Occupational history	Symptoms	Signs
<u>Asbestosis Group.</u>					
66	A.1	52	Sherman cyclone 10 years	cough, dyspnea, loss of strength	deficient expansion, crepitations left apex.
71	A.1	47	general operator waakt foramen 7 years	cough, dyspnea, frequent colds	poor expansion, percussion dullness left chest, crepitations all over both lungs.
72	A.1	42	loading cars in mill 3 foramen 23 years	no symptoms	no abnormal signs.
73	A.1	50	driller 7 years	no symptoms	no abnormal signs.
74	A.1	41	millwright 31 years	dyspnea, cough	no abnormal signs
75	A.1	7	motor attendant for 11 years	no symptoms	faint moist rales 2nd rib in front.
76	A.1	33	millwright 16 years	no symptoms	no abnormal signs
77	A.1	50	labourer watchman in mill 7 years	no symptoms	slight dullness right base, left chest on right more than right. breath sounds at right base
78	A.1	47	labourer to floor mill, 7 years	no symptoms	no abnormal signs
79	A.1	47	in mill 16 years	no symptoms	expansion slightly limited
80	A.1	42	labourer in mill 27 years	no symptoms	expansion limited
81	A.1	41	mine (work not stated) 17 years	no symptoms	no abnormal signs
82	A.1	41	driller 4 years	no symptoms	no abnormal signs
83	A.1	41	driller 25 years	no symptoms	no abnormal signs
84	A.1	60	driller 4 years in foramen 23 years	no symptoms	abundant vocal fremitus
85	A.1	37	driller 7 years trackman 4 years	no symptoms	bronchial rales left side
86	A.1	33	pipe coverer (factory) 4 years	no symptoms	no abnormal signs.

Table VI

No.	Initial	Age	Occupational history	Symptoms	Signs.
Thetford Union Group					
1	A.1	47	filling bags 6 years oilier 12 years foreman 4 years	no symptoms	corroded aortic aorta & at both bases behind.
3	A.1	43	driller 23 years	no symptoms	no abnormal signs
6	A.1 (probably)	39	oilier in mill 3 yrs. filling bags 4 yrs.	dyspnea	no abnormal signs
11	A.1	37	millwright 27 years	dyspnea	no abnormal signs
12	A.2 (early)	43	sweeper 12 years	cough, ex- pectoration, loss of weight	corroded crevices left a. ex behind.
13	A.1	51	millwright 21 years	cough, ex- pectoration, dyspnea, loss of weight	dry rales over rt. chest behind, ex-pulsion deficient
17	A.1 (probably)	47	millwright 21 years	cough	occasional dry rales at left base
18	A.1	40	driller 11 years	no symptoms	no abnormal signs
19	A.1 (early)	43	driller 17 years	history of pleurisy, sister died of t.b.c., dyspnea	no abnormal signs
21	A.1 (early)	37	millwright 14 years	dyspnea	no abnormal signs
22	A.1 (early)	41	driller 40 years	no symptoms	waxy crevices right base.
26	A.1 (early)	33	driller 6 years general work in pit 4 years	no symptoms	no abnormal signs
27	A.1	44	filling bags 1 year last coal 1 year	history of leucopneumonia	no abnormal signs
31	A.1 (early)	33	in mill 3 years in tunnel feeding crusher 3 years	cough, dyspnea, suspected t.b.c. in l. l.	no abnormal signs
32	A.1 (early)	33	crusher 11 years sweeper in mill 4 yrs. oilier in mill 3 years	no symptoms	no abnormal signs
33	A.1	46	loading boxes in pit 27 years, feeding the mill 3 years	sister died of of t.b.c. 11 yrs.	showing of fine crevices both lungs behind from vertebra down; crevices note diminished
34	A.1 (early)	33	millwright 6 years foreman 26 years	dyspnea	no abnormal signs
4	A.1	40	feeding crusher 10 yr clearing in mill 15 yrs.	dyspnea	crevices in rt. axilla & l. base behind,
41	A.1 (probably)	36	feeding crusher in mill 6 years, driver 1 year, foreman in mill 3 years	no symptoms	no abnormal signs

Table VI (continued)

No.	Diagnosis	Age	Occupational history	Symptoms	Signs
-----	-----------	-----	----------------------	----------	-------

Thetford Mill Group. (cont'd)

63	A.I. (early)	34	loading boxes in pit 7 yrs., driller 6 yrs.	no symptoms	no abnormal signs.
71	A.I. (probably)	31	operator in mill 11 yrs., millwright 11 years	no symptoms	pericardium large & prominent over right lung, small & creditable over left. Frontal view, over left u. or lower front. view.
73	A.I.	35	otter in mill 10 yrs. foreman mill 7 yrs.	non-diab. t.b.c. daughter has tuberculosis.	examination poor, very creditable in r. side, creditable in l. u. or lower front. view.
74	A.I.	41	mill repairs 11 yrs. mill foreman 11 yrs.	cough, dyspnea	examination poor, pericardium poor at 2. lower. no signs.
75	A.I. (late)	44	Foreman mill 17 yrs. millwright 7 yrs.	no symptoms	no abnormal signs.

It will be noted that of the 14 men examined at Ankerston 17 were diagnosed as first stage asbestosis, while of the 34 examined at Thetford Mines 21 were diagnosed as first stage asbestosis and 3 as second stage asbestosis. This is not to be interpreted as indicating an excess hazard in Thetford Mines. The increased incidence of pneumoconiosis in Thetford Mines is simply a reflection of the different distribution with respect to age and length of service. Reference to tables 1 and 2 will recall the great differences in these two things.

In the great majority of cases the length of service in dusty work was more than 5 years as indicated in the following table:

length of service	number of cases	number of men examined
0 - 4 yrs.	2	32
5 - 9	2	75
10 - 14	7	10
15 & over	13	23
Total	44	140

From this table one would conclude that asbestosis is not likely to develop with an exposure of less than 5 years, but that with exposure of more than 15 years there is a great possibility of its developing. It is, however, of interest to remember that many men with long periods of exposure do not develop any signs of asbestosis.

As to occupations of 41 drillers 13 were cases of asbestosis, of 1 mill workers 27 were cases of asbestosis and of 40 workers in the factory only 1 case showed x-ray evidence of the disease. Evidently there is little difference in the hazard in mill workers and drillers. The mill workers appear to be exposed to far greater dust concentrations than the drillers and it is rather difficult to understand the high incidence of asbestosis among the drillers. However, it may or may not be of interest to note the

Tuberculosis did not play an important part in the study. Even if the cases of anemias appeared to be complicated with this disease. In fact, the incidence of definite tuberculosis in the two groups of individuals was very low. There were several suspected cases on clinical examination which were not confirmed by x-ray and several suspected cases by x-ray which showed no real evidence in the way of signs and symptoms. Only one case in the group can be considered as actively tuberculous.

It was difficult to obtain sputum from most of the men since very few of them were coughing. About 17 sputa were examined for tubercle bacilli and anthea bodies. No tubercle bacilli were found but one case of anthea bodies. This case was diagnosed by x-ray as tuberculosis (first stage) but is not included in the number of tuberculosis cases since the x-ray film was not satisfactory.

Conversation with physicians and mine managers in the anthracite region indicated that no hazard to health was suggested in connection with work in the anthracite mining and milling industry.

In general it may be said that the hazard of ^{hazard} ^{pneumoconiosis} ~~anthracite~~ in the anthracite mining and milling industry has been demonstrated, but evidently the disease does not incapacitate to any great degree and, therefore, it is not particularly associated with tuberculosis.

Frank G. Bailey
1930