

Mortality among Workers Exposed to Coal Tar Pitch Volatiles and Welding Emissions: An Exercise in Epidemiologic Triage

MICHAEL SILVERSTEIN, MD, MPH, NEIL MAIZLISH, PhD, ROBERT PARK, MS, AND FRANKLIN MIRER, PhD

Abstract: The United Automobile Workers International Union has established a system of epidemiologic triage to evaluate patterns of mortality among groups of union members. In response to worker concerns, the Union examined mortality at a metal stamping plant, using a method which linked pension records with the State of Michigan computerized death registry. The observed proportion of malignant neoplasms was nearly twice that expected (95% Confidence Limits 1.36, 2.62). Two- to five-fold excess proportional mortality from cancer of the digestive organs, lung cancer, and leukemia accounted for most of the overall excess. Strong associations were found between lung and digestive organ cancer and

employment as maintenance welders or millwrights in the plant (odds ratios > 10). High levels of six polycyclic aromatic hydrocarbons with mutagenic and carcinogenic properties were found during hot coal tar application to wood block floors, work conducted by the high-risk groups. These levels were substantially reduced following the purchase of new tar pots. The example demonstrates that epidemiologic tools can play a valuable role in occupational health decision making, but care must be taken to avoid mechanical reliance on quantitative testing and to acknowledge the important role of social and political value judgments in the establishment of responsible public policy. (*Am J Public Health* 1985; 75:1283-1287.)

Introduction

This paper describes recent union-based efforts to develop and use epidemiologic surveillance techniques as routine, accessible tools in the day-to-day conduct of a health and safety program.

The United Automobile Workers International Union (UAW) represents 1.2 million workers in more than 2,500 workplaces throughout the United States and Canada. These workers face an enormous range of potential health hazards on jobs ranging from paint formulation to salt mining. Automobile manufacture, employing one-half of UAW members, is itself a collection of many distinctly different types of work.

The UAW has had an active health and safety program for many years, utilizing occupational health professionals and local union health and safety representatives to identify workplace hazards and to develop technical and political strategies for their control. Stimulated by recent intense membership concern with apparent excess cancer mortality in various workplaces, it became apparent that the health and safety program needed a new dimension—one which could evaluate patterns of illness, injury, and death that might not be apparent to even a skilled observer with the training to recognize known hazardous materials and conditions.

A practical system of epidemiologic triage was required so that threshold decisions about health and safety strategy could be made on firm grounds. For example, with several workplaces reporting apparent clusters of cancer deaths, a prompt choice among three courses of action is necessary: 1) a priority program to reduce worker exposures to known or likely hazards through engineering or other controls; 2) systematic research to explore suspected health effects supplemented with modest, interim protective measures; 3) reassurance to workers that work-related health problems have not been demonstrated and are unlikely to have occurred and that little additional action is needed. It is not

reasonable to expect that any epidemiologic method can determine unequivocally which of these activities is correct for a given workplace. Moreover, many epidemiologic techniques are too slow and cumbersome to have short-term decision-making value in the workplace. However, it has been possible to adapt standard epidemiologic methods to

serve as public health tools in the workplace, so that quantitative data can be used in choosing among alternatives at a given workplace and in determining priorities among different locations according to relative urgency. One such method, addressing mortality, has been established in Michigan by making use of the State's computerized death registry. Three steps are involved:

- First, when a potential problem is identified, local Union and international Union records are searched to produce a listing of persons at risk, or known to have died, with names and social security numbers. The full cohort of interest, all those who worked in the plant over a defined period, is usually not accessible or cannot be generated quickly. However, an inventory of available records (e.g., seniority lists, dues cards, application records, and sometimes death records) usually provides a fairly complete specification of those at risk. The fact that such a list might have gaps and sampling inconsistencies is not a deterrent in this triage process, as long as there is reasonable certainty that major confounding and selection bias have been avoided. This can often be assured by using a subgroup for which nearly complete and unbiased ascertainment is possible, such as all those hourly employees with at least 10 years of credited pension service.

- Second, names and social security numbers on the list of deceased or at-risk persons are matched against computerized mortality files such as those obtained from the State of Michigan Department of Public Health. The Michigan files contain all in-state deaths since 1970 with underlying cause coded by the International Classification of Disease (ICD) classification revision in effect at the year of death. As a useful but not essential adjunct, out-of-state deaths are identified by searches of Social Security Administration (SSA) records.

- Third, the resulting deaths identified, with underlying cause and demographic data, are used as input to the Monson program (with revisions) to generate standardized proportional mortality ratios (SPMR).¹

From the United Automobile Workers, Health and Safety Department. Address reprint requests to Michael Silverstein, MD, MPH, UAW Health and Safety Department, 8000 E. Jefferson, Detroit, MI 48214. Dr. Maizlish is currently with the Occupational and Environmental Health Unit, University of California, Davis. This paper, submitted to the *Journal* October 15, 1984, was revised and accepted for publication May 8, 1985.

Editor's Note: See also related editorial p 1259 this issue.

© 1985 American Journal of Public Health 0090-0036/85\$1.50

As an example, we describe the course of a mortality investigation at a metal stamping plant where maintenance welders and millwrights were exposed to coal tar and welding emissions.

Methods

Development of Study

The study site was a metal stamping plant located near Kalamazoo, Michigan. The plant, which began operations in 1966, has employed on average approximately 3,000 UAW members. The plant operations have included metal stamping, spot welding, tool and die making, metal machining, and pattern making.

The Local Union became concerned in 1980 about a possible high number of cancer deaths among its members. The Local Union Health and Safety Representative began to compile a list of deceased members along with information about their causes of death and jobs, which he was able to abstract from information available at the plant. The UAW had not yet developed most of the epidemiologic tools described in this paper. However, manual calculations of proportional mortality were done and did not reveal excesses of cause-specific mortality or associations between mortality and specific work assignments.

By 1982, the number of known decedents was 58 and the Union had developed the computerized capacity for PMR analysis.¹ Twenty-two cancer deaths among White men were observed, compared with eight expected based upon national rates. The Union decided to continue the investigation on a high priority basis.

A site visit was conducted by International Union staff to gain insight into the nature of chemical exposures potentially associated with various jobs. It was observed that the floors in this plant consist of wood blocks which are set and protected with hot coal tar in an operation that closely resembles roof tarring. Further, employees reported that: 1) millwrights have been responsible for laying the floor tar and have been exposed to smoke and vapors from this work; 2) the millwrights at this plant also work closely with the maintenance welders who weld or flame cut floor-bolts at floor level where tar and wood blocks sometimes ignite and generate smoky emissions; and 3) the wood blocks have been treated with wood preservatives, and have absorbed solvent and oil spills.

Historically, maintenance welding and floor tarring work had been done with a minimum of local exhaust ventilation or respiratory protection. Routine quantitative measurements of these exposures had not been made, but in 1979 personal and area air samples taken by the employer during floor tarring revealed coal tar pitch volatile concentrations in excess of 200 µg/cubic meter TWA (8-hour time weighted average) with peak exposures much higher.

This additional information about the chemical environment prompted a decision to expand the mortality evaluation and, simultaneously, to press for reduced exposure to tar emissions. The outcome of this more thorough review is presented below.

Study Group

The target population for the investigation was all deceased local union members ever-employed at the stamping plant with a date of death between January 1, 1966 and December 31, 1982. Sources of data included local union lists of deceased workers, union pension records, and employer life insurance records. These sources routinely received

reports of deaths or copies of death certificates of active workers and retirees. Additionally, the local union newsletter circulation file was used to identify those who left employment before vesting or left employment with vested pension rights but might have died before retirement. This file was matched to SSA records of deceased persons using Social Security number as the match key.

Ninety-eight (98) deaths were reported from all sources including eight uniquely ascertained from SSA records. Our judgment is that the group of 98 decedents is a virtually complete ascertainment of hourly workers who died while active or retired; a small number of deaths among workers of deferred vested status, or those terminated with less than 10 years seniority may have been missed.

Death certificates (or, in two cases, abstracted records) were obtained for 94 of the 98 deaths (96 per cent). For 62 of the Michigan deaths, underlying cause of death was determined by record-linkage of the death file to the computerized Michigan death certificate registry using Social Security numbers for matching. The remaining deaths, which included out-of-state deaths and those which occurred before the State computerized its files in 1970, were coded according to the 8th Revision of the ICD by the investigators.

The final study group included 91 White males, 2 Black males, and 1 White female.

Work Histories

Personnel records were obtained from the employer for 83 of the study group. Following site visits, discussions with the plant management and the Local Union, and a review of the available industrial hygiene reports, decedents grouped into two exposure categories: 1) maintenance welders and millwrights; and 2) all other job classifications. This classification scheme was based on the judgment that maintenance welders and millwrights are most likely to be exposed to coal tar pitch volatiles.

Pre-plant employment as a welder or millwright was coded on an ever-never basis.

Statistical Analyses

SPMRs, standardized for age, sex, race, and calendar year of death, and case-control studies within the decedent population were carried out. Standardized Proportional Mortality Ratios (SPMR) for 54 causes of death were calculated for the 91 White males with US White males as the reference population, updated to include National Center for Health Statistics (NCHS) mortality data through 1978. SPMRs were also calculated for restricted subgroups based on employment duration and prior occupational history as a welder or millwright.

Case-control analyses were designed to examine the relationship between cancer mortality and specific jobs which share, among other things, hot tar operations or welding and flame cutting near tarred surfaces. Cancer cases were defined by underlying cause of death and grouped according to specific ICD codes: lung, 150-159; digestive organs, 150-159; leukemia, 204-207; and all malignant neoplasms, 140-209. Controls were defined as all non-cancer deaths. Nonmalignant respiratory disease deaths were therefore included as controls in the case-control analysis of lung cancer, a decision made to conserve statistical power despite the possibility that shared etiologic factors might result in a reduced ability to demonstrate associations. "Exposed" was defined as ever-maintenance welder or ever-millwright and "non-exposed" were all other job classifications.

Cases and controls were drawn from the 83 decedents

MORTALITY AMONG STAMPING PLANT WORKERS: TRIAGE

TABLE 1—Standardized Proportional Mortality, White Males, 1968–82, by Cause of Death

Cause of Death	Observed	Expected	SPMR	95% C.L.
All Cancer	28	1482	189	1.36, 2.62
CA of Lung	10	4.75	220	1.18, 3.76
CA Digestive Organs	7	3.10	226	1.12, 4.56
CA of Testis	2	0.41	485	—
Leukemia	4	0.94	428	1.75, 10.45
All Cardiovascular	22	26.79	82	0.59, 1.13
All Nonmalignant Lung	0	2.96	0	—
All Nonmalignant Digestive	2	5.22	25	—
All External Causes	37	33.28	111	0.90, 1.38
All Other	2	7.92	25	—
TOTAL	91	91		

2—Characteristics of Cases and Controls

Characteristics	Cases n = 27	controls n = 56
Age at Death, Mean (SD)	47 (11)	38 (12)
Age at Death, Range	22–66	20–68
Employment Duration, Mean (SD)	9 (5)	7 (4)
Former Welders/Millwrights, Number (%)	7 (26)	4 (7)

with complete work histories: 80 White males, 2 Black males, and 1 White female. Tobacco smoking patterns were not known. Stratification was used to control potential confounding by age and pre-plant occupation.

Results

The observed number of malignant neoplasms (Table 1) is nearly twice that expected ($PMR = 28/14.82 = 189$, 95% C.L. 1.36, 2.62). Two-fold to five-fold excess proportional mortality from cancer of the digestive organs, lung cancer, and leukemia accounts for most of the overall cancer excess. Essentially the same excesses were found in the subgroups with five or more years seniority or with no pre-plant employment as a welder or millwright (data not shown).

Table 2 gives age and occupational characteristics of cases and controls. On average, both cancer cases and controls died at relatively young ages and short employment duration. Because cases tended to be older at death than controls and were overrepresented by former welders or millwrights, case-control analyses were stratified by age and pre-plant occupation.

Table 3 presents age-stratified odds ratios for the association between cancer mortality and employment as maintenance welders and millwrights. The mean age in the 40+ age stratum was 50.3 (49.4 among nine welder/millwrights and 50.6 among 34 non-welder/millwrights) and in the <40 age stratum was 30.9 (33.8 among four welder/millwrights and 3.6 among 36 non-welder/millwrights). Odds ratios of more than 10 were found overall and in the 40+ age strata. Similarly, the odds ratios for lung cancer and digestive organ cancers were highly elevated in the 40+ strata (Table 4).

Table 5 presents cancer mortality and welder or millwright classification stratifying on prior welding or millwright occupation. While odds ratios are elevated in both strata, the odds ratio was much greater and statistically more stable in the subgroup without prior welding or millwright occupations. Odds ratios for lung cancer or cancer of digestive organs followed a similar pattern.

Further, case-control analysis simultaneously stratifying

TABLE 3—Association between All Cancer Mortality and Welder/Millwright Classification

Classification	All Ages Cases	Controls
Welder/Millwright	10	3
Other	17	53
OR = 10.4; 95% CL: 2.6–42.2		
<40 Years		
Welder/Millwright	2	2
Other	5	31
OR = 6.2; 95% CL: 0.7–54.6		
40+ Years		
Welder/Millwright	8	1
Other	12	22
OR = 14.7; 95% CL: 1.6–131.6		

TABLE 4—Association between Cancer Mortality, Two Sites, and Welder/Millwright Classification, Age 40+ Years at Death

classification	Lung Cancer Cases	Controls
Welder/Millwright	3	1
Other	5	22
OR = 13.2; 95% CL: 1.1–154.9		
All Digestive Cancers		
Welder/Millwright	3	1
Other	4	22
OR = 16.5; 95% CL: 1.1–162.2		

TABLE 5—Association between All Cancer Mortality and Job Classification by Prior Occupation

Classification	Former Welders or Millwrights Cases	controls
Welder/Millwright	6	2
Other	1	2
OR = 6.0; 95% CL: 0.3–107.0		
Not Former Welders or Millwrights		
Welder/Millwright	4	1
Other	16	51
OR = 12.8; 95% CL: 1.3–122.5		

on age, employment duration, and prior occupation also revealed similar associations between cancer mortality and millwright or maintenance welder classification, but statistical power was diminished, as expected with decreasing cell sizes.

Discussion

A union-based epidemiologic surveillance program proved to be an effective and efficient guide to occupational health action in a metal stamping plant. Use of a mortality surveillance system designed for routine use revealed excess proportional mortality from cancer of the lung, digestive organs, and leukemia. This lent credibility to local union concerns and led the union to assign this situation very high

TABLE 6—Stamping Plant Exposure Data* Before and After Installation of New Tar Pots

Chemical	Old Tar Pot: Vent	OM Tar Pot: Floor	New Tar Pot: Floor
Phenanthrene	500	50	0.88
Anthracene	200	20	0.55
Acridine	10	1	ND**
Pyrene	100	10	0.16
Chrysene	20	1	ND
Benz(a)pyrene	1	ND	ND

*All values in mgcu m

Short-term Air samples

**Non-Detectable

priority for further investigation and to undertake interim efforts to control suspect chemical exposures. Subsequent case-control analysis documented associations between mortality from lung cancer and digestive cancers and work as maintenance welders and millwrights which involved probable exposure to coal tar pitch volatiles and welding emissions. This served as the basis for a further decision to seek more vigorous in-plant environmental intervention.

Following discussions of the findings the employer took short-term source and area samples during tarring which showed substantial levels of six specific polycyclic aromatic hydrocarbons with known carcinogenic or mutagenic properties (Table 6). Agreement was reached to take steps to reduce exposures of maintenance welders and millwrights to the suspect emissions. The employer purchased new, thermostatically controlled electric tar pots and agreed to obtain more than 20 portable ventilation units, construct several exhausted welding booths, purchase additional respirators, and initiate a search for non-coal tar floor coating substitutes. A series of educational meetings was held with more than 200 affected workers in the plant. At the urging of the union, the employer sent a bulletin to its other locations asking that floor tarring operations be evaluated and that steps be taken either to reduce emissions or find substitute materials. Follow-up area air sampling during tarring was conducted after the installation of the new tar pots and showed large reductions in airborne polycyclic aromatic hydrocarbons (Table 6).

We believe that these findings are sufficiently plausible to warrant an alert to the public health community of a probable, but previously unrecognized, hazard. The observed associations are large; feasible control measures are readily available; and the findings have considerable biologic plausibility.²⁻⁴ Exposure of roofers and coke oven workers to coal tar pitch volatiles containing polycyclic aromatic hydrocarbons has been associated with excess multiple-site cancer mortality.^{5,6} The group experiencing excess mortality in this study uses similar materials under analogous conditions,⁴ and exposures to specific mutagens and carcinogens have been documented (Table 6).

Efforts should be made to replicate the findings at other locations where similar work is done in such a manner that the combined and independent effects of coal tar and welding exposures can be evaluated. In our judgment, concurrent efforts should be made to notify representatives in potentially affected workplaces of the possible hazards and to encourage actions to reduce worker exposures to chemical emissions during floor tarring operations as the prudent response to the surveillance findings.

Mortality surveillance enabled the union to distinguish this plant situation from others which, on the surface, might

have appeared equally urgent. For example, in another location there were 54 cancer deaths in 260 total deaths among White men. More than 60 malignancies were expected. There were slight excesses of brain and lymphopoietic cancers but a review of job histories revealed no clustering and did not direct attention to any specific chemical exposures. The mortality file has been updated yearly but no other intervention has been made.

Some advocates for an increased role for epidemiology in public health have argued that quantitative decision rules can be used by occupational and environmental health policy makers as virtually sufficient and complete guides to action. Dinman and Sussman, for example, have proposed, "a set of criteria for acceptance or rejection of any published study."⁷ These criteria would be used to generate a summary score for studies in a given area of policy concern. The score would be used to ensure that "the public decision-making process optimally utilizes data in an organized, systematic fashion." Although they concede that "these scaled systems should not be considered as providing a mechanical method for arriving at judgment without thought," their single example seems to rely upon just such mechanical application. Dinman and Sussman calculate scores for five published studies of mortality among workers exposed to coal tar emissions in several industries. They average the scores for two studies of workers exposed to emissions in the early stages of coal tar production (i.e., coke oven workers) and for three studies of workers exposed to emissions after coal tar has been further distilled to produce coal tar pitch (i.e., roofers and aluminum reduction workers). They direct attention to an "information/option nonogram," a scale along which the epidemiologic scores correspond to public policy options. Use of this scale directs that strong action is necessary to "restrict manufacture" in coke ovens and that the appropriate measures for roofers and aluminum reduction workers fall somewhere between "no action" and "simple educational measures." Dinman and Sussman conclude that "the fumes of pitch are indeed less carcinogenic than tar if, indeed, they are carcinogenic at all."

Lanes and Poole have critiqued the Dinman/Sussman scheme and conclude that, with respect to their coal tar: coal pitch assessment, "It is difficult to imagine a less secure evaluative foundation for such a strong and sweeping statement about causality."⁸ We echo their critical judgment sharing the belief that the fault lies not merely with the specific scoring system but with the very notion that public health policy decisions can be reduced to any set of epidemiologic rules and that the "formulation is best left to epidemiologists."⁷

Our concern with such reductionist systems is that even the most rigorous are essentially deceptive. Their apparent objectivity invariably masks a set of value judgments even bit as politically charged as those expressed by the most rhetorical of lay policy makers. The statistical concepts upon which scientific decision rules are based are inherently judgmental. For example, the choice of a statistically significant level expresses the willingness of an investigator to risk a conclusion which might be falsely positive. At a level of $p = .05$ the investigator is unwilling to conclude that a finding is significant unless there is less than a 5 per cent chance of being wrong. This embodies a value judgment about the level of certainty necessary for action, one which is more stringently conservative than we apply to most decisions of daily living.

While we advocate the increased input of quantitative

ACKNOWLEDGMENTS

This work would not have been done without the efforts and determination of Jack Kelly and Roger Snell of UAW Local Union 488, Kalamazoo, Michigan. This research was supported, in part, by NIOSH Contract #210-81-5104. Some of these data were originally presented at the American Public Health Association Annual Meeting, Dallas, November 7, 1983.

REFERENCES

1. Monson R: Analysis of relative survival and proportional mortality. *Comput Biomed Res* 1974; 7:325-332.
2. Lindstedt G, Sollengren J: Polycyclic aromatic hydrocarbons in the occupational environment. *Scand J Work Environ Health* 1982; 8:1-19.
3. Bingham E, Trosset R, Warshawsky D: Carcinogenic potential of petroleum hydrocarbons—a critical review. *J Environ Pathol Toxicol* 1980; 3:483-563.
4. Cooke M, Dennis A, Fisher G (eds): *Polyaromatic Hydrocarbons*. 6th International Symposium. Columbus, Ohio: Battelle Press, 1982.
5. Redmond C, et al: Long-term mortality study of steelworkers: VI. mortality from malignant neoplasms among coke oven workers. *JOM* 1972; 14:53-61.
6. Hammond C, et al: Inhalation of benzo(a)pyrene in man. *Ann NY Acad Sci* 1976; 271:116-124.
7. Dinman B, Sussman N: Uncertainty, risk and the role of epidemiology in public policy development. *JOM* 1983; 25:511-516.
8. Lanes S, Poole C: Truth in packaging?: the unwrapping of epidemiologic research. *JOM* 1984; 26:571-574.

A New Publication: The Journal of Rural Health

The Journal of Rural Health, a new periodical which began publication in January 1985, serves as a medium for communication among health scientists and professionals in practice, education, and research settings. The principal goal of the journal is to advance professional practice, research, theory development, and public policy related to rural health.

According to its sponsoring organizations—the American Rural Health Association, and the National Rural Health Care Association—"For some time there has been a cadre of rural health professionals who have not had an adequate communication channel for their views, ideas, activities, and scholarly research results. *The Journal of Rural Health* will fill that void and provide further identity to the field of rural health. This identity is essential if the world around us is to become sensitized to the vast differences which exist between rural and urban America, and to evaluate the health care consequences of those differences."

The peer-reviewed journal will be organized around six sections, to appear routinely in the journal when a sufficient number of manuscripts have been reviewed and accepted for publication to produce a meaningful contribution to the literature. The six sections will be:

- Education for Rural Health Professionals
- Rural Health Policy
- Rural Health Practice Applications
- Current Research in Rural Health
- Book Reviews
- Research Reviews

Contributed manuscripts are welcome and should be sent directly to the journal editor, Raymond T. Coward, PhD, Center for Rural Studies, University of Vermont, 448 Waterman, Burlington, VT 05405-0160. Instructions for manuscript submission appeared in the January 1985 issue of *The Journal of Rural Health*, or can be obtained by writing to the editor. Subscription information is also available from the editor.

Book Reviews—To be considered as a potential book reviewer, to suggest books that might be critiqued, or to reply to a particular review, write to: Dr. Ben F. Banahan, III, Health Services Research Division, Research Institute of Pharmaceutical Sciences, University of Mississippi, University, MS 38677.

Mortality Among Workers Employed in Petroleum Refining and Petrochemical Plants

Terry L. Thomas, M.S.; Pierre Decoufle, Sc.D.; and Rafael Moure-Eraso, M.S.

The cause-specific mortality experience of 3,705 members of the Oil, Chemical and Atomic Workers International Union was examined to determine if there were unusual patterns of fatal disease that may be indicative of hazardous agents in the work environment. Deaths among active Union members that were reported by locals in Texas between 1947 and 1977 were identified through membership records, and proportionate mortality was analyzed in several broad industrial categories. PMRs for cancers of the liver and biliary passages, pancreas, lung and skin were elevated among refinery and petrochemical plant workers; however, risks did not increase with length of membership. Increased relative frequencies of stomach cancer, cancer of the brain, leukemia and multiple myeloma were confined to white males in the same category who had been Union members for 70 or more years. Excess deaths from stomach cancer and brain cancer were found among white male members employed at one specific oil refinery and petrochemical plant. Observed numbers of deaths from cancer of the stomach were greater than expected among whites and nonwhites, and an elevated PMR for lung cancer among nonwhites was found at an additional plant. Findings suggest that workers in this industry may be at increased risk of certain cancers and indicate areas for further investigation.

Several studies have indicated the need for further evaluation of cancer risks among persons employed in petroleum refineries and petrochemical plants. In a recent study of cancer mortality by county, high rates of

cancers of the lung, nasal cavity and sinuses, and skin were found among male residents of counties where the petroleum industry is heavily concentrated.¹ Early studies indicated increased risk of scrotal cancer among wax pressmen in a refinery² and cancer of the larynx and nasal sinuses among workers engaged in the manufacture of isopropyl alcohol.³ Although no increased cancer risks have been found in more recent studies of U.S. refinery workers,^{4,5} a Canadian study has reported elevated risks of esophageal, stomach, and lung cancer among workers exposed to crude petroleum and its products.⁶

The opportunity to evaluate cancer risks among a large group of workers employed in petroleum refineries, petrochemical plants and related industries arose when officials of the Oil, Chemical, and Atomic Workers International Union (OCAW) invited the National Cancer Institute to review data sources within the Union that might be useful for an epidemiologic study. The present report examines the cause-specific mortality experience of Union members to determine if there was an unusual pattern indicative of hazardous agents in the work environment. The purpose of this study is to point out areas for further investigation.

Materials and Methods

In considering what kind of study design might be possible, a retrospective cohort study was deemed not feasible because of limitations in the Union's record-keeping system. Since records of deaths among active Union members are maintained by the Membership Department, we decided to examine the relative frequency of specific causes of death as a preliminary means of obtaining an estimate of site-specific cancer risks in various broad industrial categories within the petroleum refining, petrochemical, and related industries.

Because of the heavy concentration of the petroleum industry in the state of Texas, and the large OCAW

¹ In the Occupational Studies Section, Environmental Epidemiology Branch, National Cancer Institute, 3C07 Landow Building, Bethesda, MD 20205. Dr. Decoufle is affiliated with the School of Health Related Professions, University of Arizona, Tucson, Ariz., and Mr. Moure-Eraso is with the Health and Safety Office, Oil, Chemical, and Atomic Workers International Union, Denver, Colo.

Table 6. — Mortality for Selected Cancer Sites in White Male Texas OCAW Members in Category 2 by Length of Union Membership.

site (8th Revision, ICD)	Length of Membership											
	< 10 Years (N = 578)							10 + Years (N = 1144)				
	O ^a	E ^b	PMR	E ^c	PCMR	E ^d	TPMR	O ^a	E ^b	PMR	E ^c	TPMR
Liver and biliary passages (155 156)	5	2.4	208	2.6	190	2.7	184	7	4.6	152	6.8	103
Pancreas (157)	15	4.4	3.38 ^e	5.3	2.81 ^e	5.2	2.39 ^e	14	12.5	1.12	167	0.84
Lung (162, 163)	35	21.7	1.61 ^e	24.3	1.44 ^e	25.9	1.35	95	74.3	1.28 ^e	91.1	1.04
Skin (172 173)	6	2.3	2.56 ^e	2.7	2.23	3.4	179	8	4.9	165	6.2	129

^aO = Observed deaths

^bE = Expected deaths based on proportionate mortality for U.S. white males

^cE = Expected deaths based on proportionate cancer mortality for U.S. white males

^dE = Expected deaths based on proportionate mortality for Texas white males

^eStatistically significant at the 0.05 level

relative frequencies of specific causes of death, excesses for one or more causes will automatically force others to be in deficit. In addition, whenever multiple comparisons are made, a certain number of differences will be statistically significant based on chance alone. Despite its limitations, a PMR analysis is useful as a preliminary, hypothesis-generating step.

Results

Distribution of study subjects by industrial category is shown in Table 1. Most of the decedents were classified into Category 2. Only 55 decedents worked for companies that could not be classified.

Since the study group is comprised entirely of workers who were active Union members at the time of death, less than 1% were age 65 or older; however, over half of the 3,105 decedents were 50 or older (Table 2).

Table 3 indicates that 62% of the whites and 57% of the nonwhites died 10 or more years after initiation into the Union. A substantial number (> 25%) were members for 20 or more years prior to death.

PMRs for major causes of death are shown in Table 4, by race, for each industrial category that had at least 100 deaths. Among whites, the relative frequency of cancer deaths is elevated only in Category 2. Ratios for heart disease are elevated in both racial groups and in each industrial category. Observed numbers of deaths from respiratory and digestive disease are lower than expected in each category and exceptionally low among whites in Category 4. An increased relative frequency of motor vehicle accidents is seen among whites in Categories 1 and 3 and among nonwhites. Non-motor vehicle accidental deaths are elevated in each category except 3.

Site-specific cancer mortality is shown in Table 5 for each of the industrial categories in which there were at least 100 total deaths. Most interesting are results for the petroleum refinery and petrochemical plant category (Category 2). Among whites, observed numbers of deaths are significantly greater than expected for cancers of the digestive and respiratory systems, skin, and brain. A non-significant excess of respiratory cancer deaths appears in the crude oil and gas extraction category (Category 1). No other site/category combination is associated with unusual mortality, although the numbers of deaths are small in many instances. Observed numbers of deaths from digestive and respiratory cancer are greater than ex-

pected for nonwhites. Further analyses of cancer risks are limited to decedents employed in petroleum refined and petrochemical plants (Category 2).

If occupational exposure contributes to cancer risk, we generally expect risk to increase with increasing duration of exposure, or, in this case, with length of membership. Specific cancer sites that were associated with increased relative frequencies in Category 2 were examined with respect to length of membership. Not all cancer sites with elevated PMRs in the petroleum refinery and petrochemical category show a positive gradient. Table 6 shows cancer sites for which risk among members in the category does not increase with length of membership. The elevated PMR for cancer of the pancreas is confined to members initiated less than 10 years prior to death. Observed numbers of deaths from cancers of the liver, lung, and skin are greater than expected regardless of length of time between initiation and death. PCMRs computed for each of these sites show essentially the same pattern for the < 10 year membership group, however, ratios are smaller for decedents who were members for 10 or more years. Results using Texas white males as a comparison group vary little from the expected values based on the total U.S. males for each site except skin. TPMRs for skin cancer are much lower than PMRs but remain elevated.

Table 7 shows cancer sites for which risk increases with length of membership for white males in the petroleum refinery and petrochemical plant category. An increased relative frequency of stomach cancer is confined to members who joined the Union 20 or more years prior to death. There were four deaths from multiple myeloma in the same length-of-membership group and only 1.3 were expected. Deaths from leukemia and from cancers of the brain and central nervous system seem to cluster among males who were Union members for 10 or more years. Although an increased relative frequency of kidney cancer deaths is not present in the total group, there is a significant excess of deaths among members who were initiated at least 20 years prior to death, with seven deaths observed and 3.3 expected. Expected numbers of deaths based on proportionate cancer mortality are higher than those based on proportionate mortality, however, the same trends are present.

Table 8 shows cancer sites contributing to the excess shown in Table 5 for nonwhite males. Increased relative

Table 4. — Mortality Experience of Texas OCAW Members by Race and Industrial Category, 1947-1977.

Underlying Cause of Death (19th Revision, ICD)	White Males - Industrial Category										Nonwhite Males - Industrial Category			
	All Categories		1		2		3		4		All Categories		2	
	n	PMR	n	PMR	n	PMR	n	PMR	n	PMR	n	PMR	n	PMR
Cancer (140-209)	541	463.1	1.14	46	50.1	0.92	394	313.6	1.26	145	33.4	1.03	29	16.1
Circulatory disease (390-458)	1247	1180.0	1.06	131	134.1	0.98	839	801.9	1.05	73	64.2	1.14	183	172.1
Arteriosclerotic heart disease (410-413)	1029	845.6	1.22	111	93.2	1.19	687	575.7	1.19	61	46.8	1.30	114	76.4
Vascular lesions of CNS (430-438)	85	120.3	0.71	7	14.3	0.49	62	81.7	0.76	6	12.4	0.48	36	48.2
Respiratory disease (460-519)	45	110.6	0.41	9	12.3	0.73	27	74.8	0.36	1	6.0	0.17	15	30.3
Digestive disease (520-577)	92	155.6	0.59	11	16.4	0.67	68	104.8	0.65	2	8.5	0.23	19	29.2
Accidents, suicide, homicide (800-998)	499	414.7	1.20	66	38.7	1.71	302	268.1	1.13	31	22.0	1.41	134	97.8
Motor vehicle accidents (810-827)	161	143.0	1.13	28	13.2	2.12	81	90.7	0.89	8	7.8	1.02	45	23.3
Other Accidents (800-807, 830-949)	219	144.1	1.52	26	14.1	1.84	145	93.9	1.54	20	21.3	0.94	41	30.0
Suicide (950-959)	91	94.5	0.96	8	8.9	0.90	59	62.2	0.95	14	13.5	1.03	8	5.5
All other causes	139	238.9	0.58	16	27.5	0.58	92	158.8	0.58	17	28.0	0.61	38	84.8
All causes	12563	2563.0	1.00	1279	279.0	1.00	11722	1722.0	1.00	304	304.0	1.00	542	542.0

* 0 = Observed deaths

†E = Expected deaths based on proportionate mortality for U.S. males

‡Statistically significant at the 0.01 level

Table 5. — Mortality for Selected Cancer Sites in Texas OCAW Members by Race and Industrial Category.

Site (8th Revision, ICD)	White Males - Industrial Category										Nonwhite Males - Industrial Category			
	All Categories		1		2		3		4		All Categories		2	
	n	PMR	n	PMR	n	PMR	n	PMR	n	PMR	n	PMR	n	PMR
Digestive organs & peritoneum (150-159)	140	131.3	1.07	9	15.3	0.59	111	89.3	1.24	9	1.37	0.66	24	27.6
Respiratory system (160-163)	188	130.8	1.25	21	15.6	1.35	134	102.6	1.31	18	17.5	1.02	33	24.9
Skin (172, 173)	19	0.8	1.76	1	1.0	—	14	7.2	1.94	3	14	—	2	0.5
Genitourinary organs (185-189)	42	4.2	1.04	3	4.6	—	28	27.2	1.03	6	4.5	1.34	7	8.5
Brain & central nervous system (191, 192)	31	21.3	1.46	2	2.2	—	25	14.2	1.76	4	2.5	—	1	1.4
Lymphatic & hematopoietic tissue (200-209)	63	54.4	1.16	4	5.5	0.72	44	36.3	1.21	8	1.13	—	7	6.3

* 0 = Observed deaths

†E = Expected deaths based on proportionate mortality for U.S. males

‡Statistically significant at the 0.05 level

— Not calculated

Table 6. — Mortality for Selected Cancer Sites in White Male Texas OCAW Members in Category 2 by Length of Union Membership.

Site (8th Revision, ICD)	Length of Membership													
	< 10 Years (N = 578)							10 + Years (N = 1144)						
	O*	E†	PMR	E‡	PCMR	E§	TPMR	O*	E†	PMR	E‡	PCMR	E§	TPMR
Liver and biliary passages (155, 156)	5	2.4	2.08	2.6	1.90	2.7	1.4	7	4.6	152	6.3	103	5.7	1.22
Pancreas (157)	15	4.4	3.38†	5.3	2.81†	5.2	2.89†	14	12.5	112	16.7	0.34	12.4	1.13
Lung (162, 163)	35	21.7	1.61†	24.3	1.44†	25.9	1.35	95	74.3	1.28†	91.1	1.04	76.8	1.24†
Skin (172, 173)	6	2.3	2.56†	2.7	2.23	3.4	1.79	8	4.9	165	6.2	129	6.3	1.26

*O = Observed deaths

†E = Expected deaths based on proportionate mortality for U.S. white males

‡E = Expected deaths based on proportionate cancer mortality for U.S. white males

§E = Expected deaths based on proportionate mortality for Texas white males

†Statistically significant at the 0.05 level

relative frequencies of specific causes of death, excesses for one or more causes will automatically force others to be in deficit. In addition, whenever multiple comparisons are made, a certain number of differences will be statistically significant based on chance alone. Despite its limitations, a PMR analysis is useful as a preliminary, hypothesis-generating step.

Results

Distribution of study subjects by industrial category is shown in Table 1. Most of the decedents were classified into Category 2. Only 55 decedents worked for companies that could not be classified.

Since the study group is comprised entirely of workers who were active Union members at the time of death, less than 1% were age 65 or older; however, over half of the 3,105 decedents were 50 or older (Table 2).

Table 3 indicates that 62% of the whites and 57% of the nonwhites died 10 or more years after initiation into the Union. A substantial number (> 25%) were members for 20 or more years prior to death.

PMRs for major causes of death are shown in Table 4, by race, for each industrial category that had at least 100 deaths. Among whites, the relative frequency of cancer deaths is elevated only in Category 2. Ratios for heart disease are elevated in both racial groups and in each industrial category. Observed numbers of deaths from respiratory and digestive disease are lower than expected in each category and exceptionally low among whites in Category 4. An increased relative frequency of motor vehicle accidents is seen among whites in Categories 1 and 3 and among nonwhites. Non-motor vehicle accidental deaths are elevated in each category except 3.

Site-specific cancer mortality is shown in Table 5 for each of the industrial categories in which there were at least 100 total deaths. Most interesting are results for the petroleum refinery and petrochemical plant category (Category 2). Among whites, observed numbers of deaths are significantly greater than expected for cancers of the digestive and respiratory systems, skin, and brain. A non-significant excess of respiratory cancer deaths appears in the crude oil and gas extraction category (Category 1). No other site/category combination is associated with unusual mortality, although the numbers of deaths are small in many instances. Observed numbers of deaths from digestive and respiratory cancer are greater than ex-

pected for nonwhites. Further analyses of cancer risks are limited to decedents employed in petroleum refineries and petrochemical plants (Category 2).

If occupational exposure contributes to cancer risk, we generally expect risk to increase with increasing duration of exposure, or, in this case, with length of membership. Specific cancer sites that were associated with increased relative frequencies in Category 2 were examined with respect to length of membership. Not all cancer sites with elevated PMRs in the petroleum refinery and petrochemical category show a positive gradient. Table 6 shows cancer sites for which risk among members in this category does not increase with length of membership. The elevated PMR for cancer of the pancreas is confined to members initiated less than 10 years prior to death. Observed numbers of deaths from cancers of the liver, lung, and skin are greater than expected regardless of length of time between initiation and death. PCMRs computed for each of these sites show essentially the same pattern for the < 10 year membership group, however, ratios are smaller for decedents who were members for 10 or more years. Results using Texas white males as a comparison group vary little from the expected values based on the total U.S. males for each site except skin. TPMRs for skin cancer are much lower than PMRs but remain elevated.

Table 7 shows cancer sites for which risk increases with length of membership for white males in the petroleum refinery and petrochemical plant category. An increased relative frequency of stomach cancer is confined to members who joined the Union 20 or more years prior to death. There were four deaths from multiple myeloma in the same length-of-membership group and only 1.3 were expected. Deaths from leukemia and from cancers of the brain and central nervous system seem to cluster among males who were Union members for 10 or more years. Although an increased relative frequency of kidney cancer deaths is not present in the total group, there is a significant excess of deaths among members who were initiated at least 20 years prior to death, with seven deaths observed and 3.3 expected. Expected numbers of deaths based on proportionate cancer mortality are higher than those based on proportionate mortality, however, the same trends are present.

Table 8 shows cancer sites contributing to the excesses shown in Table 5 for nonwhite males. Increased relative

Significantly increased relative frequencies of brain (5 observed; 1.5 expected) and of lung cancer deaths (22 observed; 11.7 expected) were seen among 198 whites who worked at Plant C, which functions as a refinery and products terminal. Although numbers were very small, we examined mortality patterns by length of membership and found that the brain cancers clustered among males who were members for less than 10 years (all of the decedents in this group with brain tumors were members for one year) while the lung cancer excess was confined to workers with at least 10 years of membership. No excesses of specific cancers were seen among members who worked in the other two plants that were examined separately.

PMRs for some causes of death may appear higher than they should because of lack of information on members who retired or left the Union for other reasons. No conclusions can be drawn with respect to exposure to specific agents, however, categories of industrial processes and individual plants associated with higher than expected risks can be identified. In spite of limitations imposed by the "healthy worker effect" and the lack of exposure information, the finding that risks of stomach

Table 7. — Mortality for Selected Cancer Sites in White Male Texas OCAW Members in Category 2 by Length of Union Membership.

Site (8th Revision, ICD)	Length of Membership											
	< 10 Years (N = 578)				10-19 Years (N = 609)				20+ Years (N = 535)			
	O*	E†	PMR	ES	O*	E†	PMR	ES	O*	E†	PMR	ES
Stomach (151)	7	7.1	0.98	8.6	9	7.2	1.26	9.2	14	5.2	2.69	8.1
Kidney (189)	1	2.3	—	2.8	4	3.1	—	3.9	7	3.3	2.14	4.2
Brain & central nervous system (191, 192)	7	4.7	1.49	5.5	12	5.2	2.30	6.6	6	4.3	1.40	5.6
Leukemia (204-207)	6	5.0	1.19	5.8	8	5.0	1.61	6.3	7	3.9	1.77	5.6
Multiple myeloma (203)	0	1.0	—	1.2	2	1.3	—	1.6	4	1.3	—	1.8

* = Observed deaths
 † = Expected deaths based on proportionate mortality for U.S. white males
 ‡ = Expected deaths based on proportionate cancer mortality for U.S. white males
 § = Expected deaths based on proportionate mortality for Texas white males
 ¶ Statistically significant at the 0.05 level
 — Not calculated

Table 8. — Mortality for Selected Cancer Sites in Nonwhite Male Texas OCAW Members in Category 2 by Length of Union Membership.

Site (8th Revision, ICD)	Length of Membership													
	< 10 Years (N = 150)							10 + Years (N = 259)						
	O*	E†	PMR	E‡	PCMR	E§	TPMR	O*	E†	PMR	E‡	PCMR	E§	TPMR
Stomach (151)	5	2.1	2.42¶	2.7	1.87	2.2	2.31	12	4.3	2.80¶	6.2	1.92¶	4.6	2.59¶
Lung (162, 163)	7	3.8	1.82	4.7	1.48	4.8	1.46	19	14.1	1.35	16.6	1.14	12.3	1.55

*O = Observed deaths

†E = Expected deaths based on proportionate mortality for U.S. nonwhite males

‡E = Expected deaths based on proportionate cancer mortality for U.S. nonwhite males

§E = Expected deaths based on proportionate mortality for Texas nonwhite males

¶Statistically significant at the 0.05 level

Table 9. — Mortality for Selected Cancer Sites in White Male OCAW Members Employed at Plant A by Length of Union Membership.

Site (8th Revision, ICD)	Length of Membership								
	< 10 Years (N = 107)			10-19 Years (N = 177)			20 + Years (N = 137)		
	O*	E†	PMR	O	E	PMR	O	E	PMR
Stomach (151)	3	1.4	—	1	2.2	—	5	1.3	3.76‡
Brain & central nervous system (191, 192)	0	0.8	—	5	1.5	3.32‡	4	1.1	—

*O = Observed deaths

†E = Expected deaths based on proportionate mortality for U.S. white males

‡Statistically significant at the 0.05 level

— Not calculated

Site (8th Revision, ICD)	White Males Length of Membership						Nonwhite males		
	< 10 Yrs (N = 111)			10 + Years (N = 191)			(N = 111)		
	O*	E†	PMR	O	E	PMR	O	E	PMR
Stomach (151)	1	1.7	—	6	2.0	2.97‡	7	2.1	3.40‡
Lung (162, 163)	5	4.5	1.11	13	12.1	1.08	13	4.4	2.94‡
Skin (172, 173)	1	0.4	—	3	0.8	—	0	0.1	—
Kidney (189)	1	0.5	—	3	1.0	—	0	0.3	—
Leukemia (204-207)	3	0.9	—	3	1.5	—	0	0.4	—

cancer, cancer of the brain, leukemia, and multiple myeloma increase with longer Union membership among workers in the refinery and petrochemical category strengthens the possibility of an association with occupational factors. Furthermore, essentially the same relationships were seen in a proportionate cancer mortality analysis.

These findings are consistent with reports of an excess of stomach cancer deaths among Canadian petroleum refinery workers,¹⁴ and of an elevated risk of leukemia associated with exposure to benzene and perhaps other petrochemicals.^{15, 16} Studies of vinyl chloride workers have shown an increased relative frequency of brain tumors;^{15, 16} however, no association between employment in refineries and petrochemical plants and excess

mortality from brain tumors has previously been reported.

Although no positive gradient in risk with length of Union membership was detected for cancers of the pancreas, liver, lung, and skin, further evaluation of these sites should be made. PMR analysis using Texas males as a comparison group does not alter the conclusions; therefore, these findings cannot be explained by high Texas rates. The lack of a gradient raises the possibility that factors other than occupational exposure may be operating; however, occupational factors cannot be completely ruled out because some cancers may have shorter latent periods, depending upon the nature of the exposure.

In addition, this study has highlighted three specific plants in which Union members appear to have increased

risks of certain cancers — of brain cancer at Plants A and C, of stomach cancer at Plants A and B, and of lung cancer at Plants B and C. Further evaluations of mortality among workers at these plants will be made.

Despite limitations in data and in methodology, results of the present study suggest that workers in this industry are at increased risk of certain cancers. Further studies are necessary to support and clarify these findings.

The authors wish to thank Drs. I. F. Fraumeni, Jr., R. Hoover, A. Blair, J. Wairath and Ms S. Krekel for support and guidance, S. Mayfield, T. Prante, M. Rogers, and D. Watts for technical assistance, and M. Glasser for manuscript preparation.

References

1. Blot WJ, Brinton LA, Fraumeni JF, and Stone BJ: Cancer mortality in U.S. counties with petroleum industries. *Science* **198**:51-53, 1977
2. Hendricks NV, Linden NJ, Berry CM, et al: Cancer of the scrotum in wax pressmen A.M.A. *Arch Ind Health* **19**:524-529, 1959
3. Weil CS, Smyth HF, and Nale TW: Quest for a suspected industrial carcinogen A.M.A. *Arch Ind Hyg* **5**:535-547, 1952.
4. Baird VC: Effects of atmospheric contamination on cancer mortality in petroleum refinery employees. *J Occup Med* **9**:415-420, 1967
5. Tabershaw/Cooper Associates: A mortality study of petroleum refinery workers. Project OH-1 prepared for the American Petroleum Institute, Washington, D.C., 1974
6. Wade L: Observations on skin cancer among refinery workers *Arch Environ Health* **6**:730-735, 1973

7. Hanis NM, Stavsky KM, and Fowler JL: Cancer mortality in oil refinery workers. *J Occup Med* **21**:167-174, 1979.

8. Considine DM (Ed.): *Chemical and Process Technology Encyclopedia*. New York: McGraw-Hill, 1974.

9. International Classification of Diseases (Eighth Revision). Geneva: WHO, 1973

10. Monson RR: Analysis of relative survival and proportional mortality. *Comput Biomed Res* **7**:325-332, 1974.

11. Mantel N and Haenszel W: Statistical aspects of the analysis of data from retrospective studies of disease. *J Natl Cancer Inst* **22**:719-748, 1959.

12. McMichael AJ, Havnes SC, and Trygler HA: Observations on the evaluation of occupational mortality data. *J Occup Med* **17**:128-131, 1975.

13. Vigliani EC and Saita C: Benzene and leukemia. *N Engl J Med* **271**:872-876, 1964

14. Infante PF, Wagoner JK, Rinsky RA, and Young RI: Leukemia in benzene workers. *Lancet* **276**:78, 1977.

15. McMichael AJ, Spirtas R, Kupper LL, and Gamble IF: Solvent exposure and leukemia among rubber workers: An epidemiologic study. *J Occup Med* **17**:234-239, 1975.

16. Monson RR and Peters JM: Proportional mortality among vinyl chloride workers. *Lancet* **2**:397-398, 1974

17. Tabershaw IR and Caffey WR: Mortality study of workers in the manufacture of vinyl chloride and its polymers. *J Occup Med* **16**:509-518, 1974.

18. Waxweiler RJ, Stringer W, Wagoner JK, et al: Neoplastic risk among workers exposed to vinyl chloride. *Ann NY Acad Sci* **271**:40-48, 1976.

Meaning of Work

For many persons, work is a vital part of the process of coping with life stress. This has two aspects: first, without work, the potential for boredom and meaninglessness is immeasurably increased. Work is often the primary means by which a person feels useful in life and through which significance and personal identity are established. The young adult must find a useful and meaningful place in society, and only in this way does a man or woman develop a firm sense of identity.

Second, work is often a form of coping and a refuge. Consider, for example, how people in grief regard work. Many report that in the process of mourning and reorienting their lives after a major loss, work is not a burden, but the best refuge against constant high levels of distress and depression. One speaks here of "burying one's self in work." Thus, work may provide a psychological haven against problems that otherwise would be insurmountable, or against loneliness and depression. For such people not to work is to be deprived of the only viable means of coping with non-work-related stress.

— From *Work Stress* by Alan A. McLean. Addison-Wesley Publishing Company, Reading, Massachusetts, 1979