

X

ff Lit.
Dandinger v. UCC

IARC Monographs on the Evaluation of the Carcinogenic Risk
of Chemicals to Humans (1979), Volume 19

377

Background
material

VINYL CHLORIDE, POLYVINYL CHLORIDE and
VINYL CHLORIDE-VINYL ACETATE COPOLYMERS

Vinyl chloride

This substance was considered by a previous IARC Working Group, in June 1974 (IARC, 1974). Since that time new data have become available, and these have been incorporated into the monograph and taken into account in the present evaluation.

A literature compilation (Warren *et al.*, 1978) and a review (Milby, 1977) are available.

1. Chemical and Physical Data


1.1 Synonyms and trade names

Chem. Abstr. Services Reg. No.: 75-01-4

Chem. Abstr. Name: Chloroethene

Chloroethylene; monochloroethylene; VC; VCM; Vinyl C monomer

1.2 Structural and molecular formulae and molecular weight



$\text{C}_2\text{H}_3\text{Cl}$

Mol. wt: 62.5

1.3 Chemical and physical properties of the pure substance

From Weast (1976), unless otherwise specified

(a) Description: Colourless gas (Windholz, 1976)

(b) Boiling-point: -13.37°C

(c) Melting-point: -153.8°C

(d) Density: d_4^{20} 0.9106; vapour density, 2.2 (air = 1) (Anon., 1972)

UCC
060466

- (e) Refractive index: n_D^{20} 1.3700
- (f) Spectroscopy data: Infra-red, nuclear magnetic resonance and mass spectral data have been tabulated (Grasselli & Ritchey, 1975).
- (g) Solubility: Slightly soluble in water (0.11 g/100 g at 25°C) (Hardie, 1964); soluble in ethanol; very soluble in ether, carbon tetrachloride and benzene
- (h) Volatility: Vapour pressure is 2530 mm at 20°C (Hardie, 1964)
- (i) Stability: Flash-point, -78°C (closed cup) (Hardie, 1964); polymerizes in light or in the presence of a catalyst (Windholz, 1976); on combustion it degrades to hydrogen chloride, carbon monoxide, carbon dioxide and traces of phosgene (O'Mara *et al.*, 1971)
- (j) Reactivity: On treatment with strong alkalis at high temperatures it loses hydrogen chloride (Miller, 1969).
- (k) Conversion factor: 1 ppm in air = 2.6 mg/m³

1.4 Technical products and impurities

Vinyl chloride is generally supplied as a liquid under pressure. Usually no inhibitor is added when it is to be shipped within the US. A typical analysis of a commercial US grade is as follows: water, 50 mg/kg (ppm); nonvolatile residue, 5 mg/kg (ppm); acetaldehyde, <1 mg/kg (ppm); acetylene, <1 mg/kg (ppm); iron, 0.1 mg/kg (ppm); hydrogen chloride, <0.1 mg/kg (ppm); and hydrogen peroxide, 0.01 mg/kg (ppm).

In Japan, commercial vinyl chloride meets the following specifications: purity, 99.9% min; water, 200 mg/kg (ppm) max; hydrogen chloride, 1 mg/kg (ppm) max; iron, 1 mg/kg (ppm) max; and evaporation residue, 50 mg/kg (ppm) max. Chlorinated hydrocarbons may be present as impurities.

2. Production, Use, Occurrence and Analysis

2.1 Production and use

(a) Production,

The first synthesis of vinyl chloride appears to have been made in 1835 (Regnault, 1835). Addition of hydrogen chloride to acetylene,

formerly
halogena
in the l
ethylene
dichlori
hydrogen

Vin-
years (U
producti
US impor
1976 (US
150 mill
Canada (i

Tota
in the fo
Republic
(690), Th
and the l
kg.

In J
In 1976,
chloride;

(b)

Ab u
was for ti
The remain
duction o:
in the pro
vinylidene

Th i
plastic pi
in consume
For detail
chloride-v

Hardi.
gerant, as
production
phonamides
being used

Limits
propellant
(US Enviror

UCC
060467

formerly the most important route of synthesis, has been displaced by the halogenation of ethylene; over 95% of the vinyl chloride monomer produced in the US and Japan in 1976 was made from ethylene. In this process, ethylene is reacted with hydrogen chloride and oxygen to give ethylene dichloride, which is subsequently cracked to produce vinyl chloride and hydrogen chloride.

Vinyl chloride has been produced commercially in the US for over fifty years (US Tariff Commission, 1928). In 1976, nine companies reported the production of 2580 million kg (US International Trade Commission, 1977). US imports have been negligible; exports amounted to 291 million kg in 1976 (US Department of Commerce, 1977), and in 1977, exports were about 150 million kg to the following countries (% of total): Brazil (28), Canada (8), Colombia (11), Mexico (14), Norway (12) and Yugoslavia (14).

Total western European production in 1976 amounted to 3925 million kg, in the following countries (millions of kg): Belgium (490), the Federal Republic of Germany (990), Finland (25), France (620), Greece (25), Italy (690), The Netherlands (340), Spain (190), Sweden (95), Switzerland (30) and the UK (430). Exports from western Europe in that year were 44 million kg.

In Japan, commercial production of vinyl chloride began prior to 1946. In 1976, eighteen companies produced a total of 1281 million kg vinyl chloride; 115 million kg were exported.

(b) Use

About 96% of the 2274 million kg vinyl chloride used in the US in 1976 was for the production of vinyl chloride homopolymer and copolymer resins. The remainder was used (essentially by one company internally) in the production of methyl chloroform and as a comonomer with vinylidene chloride in the production of resins. For a detailed description of the uses of vinylidene chloride-vinyl chloride copolymers, see p. 450.

The largest use for polyvinyl chloride resins is in the production of plastic piping and conduit. Other important uses are in floor coverings, in consumer goods, in electrical applications and in transport applications. For detailed descriptions of the uses of polyvinyl chloride and vinyl chloride-vinyl acetate copolymers, see pp. 406 and 414.

Hardie (1964) reported that vinyl chloride has been used as a refrigerant, as an extraction solvent for heat-sensitive materials and in the production of chloroacetaldehyde (an intermediate in the synthesis of sulphonamides); however no evidence was found that vinyl chloride is presently being used for these purposes.

Limited quantities of vinyl chloride were used in the US as an aerosol propellant, but in 1974 it was banned from use in pesticide aerosol products (US Environmental Protection Agency, 1974a), in self-pressurized household

containers, and as an ingredient of drug and cosmetic products (US Consumer Product Safety Commission, 1974a,b).

Vinyl chloride was used in western Europe in 1977 in the production of polyvinyl chloride (95%) and for other uses, including the production of methyl chloroform (5%).

In Japan in 1976, vinyl chloride was used in the production of polyvinyl chloride (92-94%) and for other uses, such as in copolymers (6-8%).

The US Occupational Safety and Health Administration's health standards for exposure to air contaminants require that an employee's exposure to vinyl chloride not exceed an eight-hour time-weighted average of 2.6 mg/m^3 (1 ppm) in the workplace air in any eight-hour work shift of a forty-hour work week. During any work shift an employee's exposure may not exceed a ceiling concentration limit of 13 mg/m^3 (5 ppm), averaged over any period of 15 minutes or less (US Occupational Safety and Health Administration, 1974).

The work environment hygiene standards for exposure to vinyl chloride in various countries, in terms of time-weighted averages (8-hr) and ceiling concentrations (10- or 15-min), were as follows in 1977: Canada, 10 ppm (8-hr) and 25 ppm (15-min); Finland, 5 ppm (8-hr) and 10 ppm (10-min); Italy, 50 ppm (8-hr), although this is expected to change to 25 ppm (8-hr); Japan, expected to be 10 ppm; The Netherlands, 10 ppm (8-hr); Norway, 1 ppm (8-hr) and 5 ppm (15-min); Sweden, 1 ppm (8-hr) and 5 ppm (15-min); USSR, 12 ppm (Bertram, 1977). In France, the standards were reported to be 5 ppm for 1 week, with a ceiling concentration of 15 ppm, in already existing factories, and 1 ppm and 5 ppm, respectively, for new factories; in Spain, no limits; in Denmark, 1 ppm (8-hr); in Belgium, 5 ppm (1 week), ceiling 15 ppm; in the Federal Republic of Germany, the same as for France in existing factories, and 2 ppm (1 year) and ceiling 15 ppm (1-hr) for new factories; in the UK, 10 ppm (8-hr), ceiling 30 ppm max; and in Switzerland, 10 ppm (1 week, 8-hr day for 5 days) (Thomas, 1977).

In August 1977, the proposed European value was 3 ppm over one year for existing and future plants, with an 'alarm-value' of 15 ppm (Commission of the European Communities, 1977a).

The US Environmental Protection Agency has proposed new rules to reduce the national emission standard for vinyl chloride from 10 ppm to 5 ppm in order to reduce vinyl chloride emissions by one-half within 3 years of the actual rulemaking. This would result in hourly emissions (based on new average-sized plants) of 5.1 kg from an ethylene dichloride-vinyl chloride plant (instead of 10.3 kg); 9 kg from a dispersion process polyvinyl chloride plant (instead of 17.5 kg); and 13.5 kg from a suspension process polyvinyl chloride resin plant (instead of 16 kg) (US Environmental Protection Agency, 1977).

In
is limit
0.3 mg/

Th
1 mg/kg
packagi
(Commis
level o
Republi
(Thomas

2.2 Oc

VI:

Th
and pol
(US Env

(a

The
prior to
scraping
hands di
Between
found in
Air conc
chloride
100-800
(Filato
et al.,
1976).
(Bol'she
0.35 ppm
mated th
chloride

On
mer to
(50-250
the US
chloride
vinyl ch
Barnhart
 mg/m^3 (C
ride pla

In the Federal Republic of Germany, the emission in the environment is limited to 3 kg/hr/source or 150 mg/m³/source, with a ground level of 0.3 mg/m³ (99% confidence) in inhabited areas (Thomas, 1977).

The Commission of the European Communities has adopted a level of 1 mg/kg (1 ppm) as the amount of vinyl chloride which can be present in packaging and 0.01 mg/kg (ppm) in foodstuffs packed in polyvinyl chloride (Commission of the European Communities, 1977b, 1978). A maximum migration level of 0.05 mg/kg has been adopted in Belgium, Denmark, the Federal Republic of Germany, France, Italy, The Netherlands, Spain and Sweden (Thomas, 1977).

2.2 Occurrence

Vinyl chloride is not known to occur as a natural product.

The occurrence of vinyl chloride in ambient air near vinyl chloride and polyvinyl chloride plants, in water, and in food has been reviewed (US Environmental Protection Agency, 1975a,b).

(a) Occupational exposure

The air concentration of vinyl chloride in a polymerization reactor prior to ventilation is of the order of 7800 mg/m³ (3000 ppm); during the scraping procedure, 130-260 mg/m³ (50-100 ppm); and that close to the hands during scraping, 1560-2600 mg/m³ (600-1000 ppm) (Cook *et al.*, 1971). Between 1950 and 1959, concentrations up to 10.4 g/m³ (4000 ppm) were found in one factory near the polymerization reactors (Ott *et al.*, 1975). Air concentrations of vinyl chloride in working places in polyvinyl chloride-producing factories have been reported variously to range from 100-800 mg/m³ (40-312 ppm), with peaks up to 87.3 g/m³ (33 500 ppm) (Filatova & Gronsberg, 1957); from 112-556 mg/m³ (43-214 ppm) (Angheliescu *et al.*, 1969); and >195 mg/m³ (>75 ppm) in a Yugoslav plant (Orusev *et al.*, 1976). In a Russian synthetic leather plant, <113.6 mg/m³ (44 ppm) (Bol'shakov, 1969) and in three UK cable factories, 0.4-0.9 mg/m³ (0.15-0.35 ppm) (Murdoch & Hammond, 1977) were detected. In 1974, it was estimated that 20 000 US workers, past and present, had been exposed to vinyl chloride in manufacturing plants (Heath *et al.*, 1975).

On a time-weighted average, the concentration of vinyl chloride monomer to which coagulator operators are exposed ranges from 130-650 mg/m³ (50-250 ppm) (Baretta *et al.*, 1969). However, in a more recent survey for the US National Institute for Occupational Safety and Health of three vinyl chloride plants it was reported that the time-weighted average exposure to vinyl chloride ranged from 0.2-70 mg/m³ (0.07-27 ppm) (Milby, 1977). Barnhart *et al.* (1975) found <0.03-15 mg/m³ (<0.01-5.89 ppm), 0.03-220 mg/m³ (0.01-84.77 ppm) and 0.05-57 mg/m³ (0.02-21.8 ppm) in 3 vinyl chloride plants in the US.

In 1974, it was reported that polyvinyl chloride leaving certain manufacturing plants may have contained 200-400 mg/kg (ppm) vinyl chloride monomer; on delivery to the customer, this level was about 250 mg/kg (ppm); and after processing, levels of 0.5-20 mg/kg (ppm) were reached, depending on the method of fabrication (Anon., 1974). Wilkinson *et al.* (1964) found 100 mg/kg (ppm) residual vinyl chloride monomer in polyvinyl chloride dispersions. However, new processing methods leave as little as 1-2 mg/kg (ppm) residual vinyl chloride in vinyl chloride resins (US Food and Drug Administration, 1975). Residual vinyl chloride in commercial food grade resins has been reduced by processing and stripping techniques to 115 µg/kg (ppb) for resin, less than 0.048 µg/kg (ppb) for compound and less than 0.043 µg/kg (ppb) for sheet polyvinyl chloride (Saggese *et al.*, 1976). Industrial grade polyvinyl chloride-coated films used for food packaging were found to contain 5-71 µg/kg (ppb) of monomer (Gilbert *et al.*, 1975) and plastic bottles up to 7.9 mg/kg (ppm) (Breder *et al.*, 1975).

(b) Air

It has been estimated that prior to 1975 vinyl chloride emissions from US polyvinyl chloride plants amounted to 110 million kg/year (US Environmental Protection Agency, 1975b) and that the average concentration of vinyl chloride in air around these plants was 44 µg/m³ (17 ppb) (US Environmental Protection Agency, 1976). Vinyl chloride has been determined in the air in the Houston, Texas, area (where an estimated 40% of the US production capacity is located) in concentrations of 8 µg/m³-3.2 mg/m³ (3.1-1250 ppb) (Gordon & Meeks, 1977) and in the ambient air near two vinyl chloride plants in the Long Beach, California, area in concentrations of 0.26-8.8 mg/m³ (0.1-3.4 ppm) (National Field Investigations Center, 1974). It has also been detected in the air in Delaware City, Delaware, in maximum concentrations of 3.9 mg/m³ (1.5 ppm) with a mean of 2 mg/m³ (0.8 ppm) (Lillian *et al.*, 1975).

(c) Water

Vinyl chloride has been detected in effluent discharged by chemical and latex manufacturing plants and in raw water in the US (Shackelford & Keith, 1976). The highest concentration of vinyl chloride detected in finished drinking-water in the US was 10.0 µg/l (Safe Drinking Water Committee, 1977; US Environmental Protection Agency, 1975a).

In 1974, it was estimated that about 12.3 kg/day vinyl chloride were discharged in the waste-water effluent from 2 vinyl chloride plants in the Long Beach, California, area (National Field Investigations Center, 1974).

(d) Food

In May 1973, a branch of the US Treasury Department banned the use of polyvinyl chloride for the packaging of alcoholic beverages (Anon., 1973a), as a result of studies reported by the US Food and Drug Administration indicating that up to 20 mg/kg (ppm) vinyl chloride monomer were present in

alcohol
chloride
mg/kg (levels

It
(ppm) (in butt
et al.,
chloride

(e
Vin
concentr
another
16 new c

Vin
and list
marijuan

2.3 Ana

A c
methods
sphere,
is avail
vinyl ch.
1975b).
water, at

A ga
National
chloride
5-litre a
1977).

A ga
mission o
foodstuff
into cont

An o
of German
It is bas
by gas ch
tion and
Normen Au

alcoholic beverages packaged in this material (Anon., 1973b). Vinyl chloride has been found in a variety of alcoholic drinks at levels of 0-2.1 mg/kg (ppm) (Williams, 1976a,b; Williams & Miles, 1975) and in vinegars at levels of up to 9.4 mg/kg (ppm) (Williams & Miles, 1975).

It has been found in edible oils, in concentrations of 0.05-14.8 mg/kg (ppm) (Roesli *et al.*, 1975; Williams, 1976a; Williams & Miles, 1975), and in butter and margarine, in concentrations of 0.05 mg/kg (ppm) (Fuchs *et al.*, 1975), when these products were packaged and stored in polyvinyl chloride containers.

(e) Other

Vinyl chloride has been found in 2/7 new automobile interiors in concentrations of 1-3 mg/m³ (0.4-1.2 ppm) (Hedley *et al.*, 1976). In another study (Going, 1976), no concentrations above 10 ppb were found in 16 new or used automobiles or in 4 new or old mobile homes.

Vinyl chloride has been detected in domestic and foreign cigarettes and little cigars, in concentrations of 5.6-27 ng/cigarette, and in a marijuana cigarette at a level of 5.4 ng/cigarette (Hoffmann *et al.*, 1976).

2.3 Analysis

A comprehensive critical review, containing over 100 references, of methods of sampling and analysis of vinyl chloride in the workplace atmosphere, ambient air, water, food, cigarette smoke and polyvinyl chloride is available (Egan *et al.*, 1979). Methods of collection and analysis of vinyl chloride have also been reviewed (US Environmental Protection Agency, 1975b). A review of methods used to determine vinyl chloride in air, water, and water piping is available (Laramy, 1977).

A gas chromatographic method of analysis has been accepted by the US National Institute for Occupational Safety and Health for determining vinyl chloride in the workplace atmosphere, in the range of 0.008-5.2 mg/m³ in a 5-litre air sample (National Institute for Occupational Safety and Health, 1977).

A gas chromatographic analytical method has been proposed by the Commission of the European Communities for determining vinyl chloride in foodstuffs and in vinyl chloride polymers and copolymers intended to come into contact with food (Commission of the European Communities, 1977b).

An official analytical method has been drafted in the Federal Republic of Germany for determining residual vinyl chloride in polyvinyl chloride. It is based on treatment of the polymer with *N,N*-dimethylacetamide, followed by gas chromatographic analysis of the solution with flame-ionization detection and has a limit of detection of 0.5 mg/kg (ppm) (Deutsche Industrie Normen Ausschuss, 1977).

3. Biological Data Relevant to the Evaluation of Carcinogenic Risk to Humans

3.1 Carcinogenicity studies in animals^{1,2}

(a) Oral administration

Rat: Groups of 40 male and 40 female 13-week-old Sprague-Dawley rats received gastric intubations of 0, 3.33, 16.65 or 50 mg/kg bw vinyl chloride dissolved in olive oil 4-5 times/week for 52 weeks. After 85 weeks from the initial treatment, 35, 39, 32 and 23 animals were still alive. At 120 weeks, 9 liver angiosarcomas, 2 Zymbal gland carcinomas and 3 nephroblastomas occurred in rats administered the 16.65 mg/kg bw dose; and 16 liver angiosarcomas, 2 nephroblastomas, 1 Zymbal gland carcinoma, and 1 thymic and 1 intra-abdominal angiosarcoma were found in the 50 mg/kg bw group. One intra-abdominal angiosarcoma was seen in the low-dose group, and 1 Zymbal gland tumour occurred in the control group (Maltoni, 1977a; Maltoni *et al.*, 1975).

(b) Inhalation and/or intratracheal administration

Mouse: Groups of 30 male and 30 female 11-week-old Swiss mice were exposed to concentrations of 130-26 000 mg/m³ (50, 250, 500, 2500, 6000, or 10 000 ppm) vinyl chloride in air for 4 hours/day on 5 days/week for 30 weeks. A total of 344 mice (176 males and 168 females) died within 61 weeks. At 81 weeks (end of experiment), 176 animals (3.5, 57, 66, 57, 70 and 70% in the different groups, respectively) had adenomas and/or adenocarcinomas of the lung, 60 animals (33, 32, 24, 30, 28 and 47%, respectively) had mammary adenocarcinomas and 47 animals (2, 19, 19, 20, 5 and 16%, respectively) had angiosarcomas of the liver. Except for lung tumours, which were not increased in the group treated with 50 ppm, a significantly higher number of neoplasms occurred in all treated groups. In 80 male and 70 female untreated controls, 8 pulmonary tumours and 3 lymphomas were observed (Maltoni, 1977; Maltoni *et al.*, 1974).

¹The Working Group was aware of studies in progress to assess the carcinogenicity of vinyl chloride in rats by administration in the drinking-water and by administration in the diet, and of complete but unpublished studies by inhalation in rats (IARC, 1978a).

²In all his experiments, Maltoni used vinyl chloride that contained the following impurities (mg/kg): water, 100; acetic aldehyde, 5; acetylene, 2; allene, 5; butane, 8; 1,3-butadiene, 10; chloroprene (see also, p. 131), 10; diacetylene, 4; vinyl acetylene, 10; propene, 3; methyl chloride, 100.

Gr
were ex
in air
were ob
treated
liver =
(histol
A dose-
no tumo
study w
1975).

Ex
gr

2
25
Co

¹F

Tw
albino
chlorid
exposed
ppm gro
minated
ding pu
and a m
for 26
had ang
had an
angiosa
adenoca
reticul

Groups of 100 male and 100 female CDI Swiss/ChR mice (age unspecified) were exposed to 130, 520 or 6500 mg/m³ (50, 200 or 2500 ppm) vinyl chloride in air (purity unspecified) for 7 hours/day on 5 days/week for 9 months and were observed for an additional 9 months. After 8 months' exposure, 49 treated animals died with tumours. A total of 42 pulmonary adenomas, 41 liver angiosarcomas and 11 mammary gland adenocarcinomas were observed (histological evaluation was carried out on grossly visible tumours only). A dose-related carcinogenic effect was evident (see Table I). At 8 months, no tumours were observed in 200 controls (100 females and 100 males). The study was still in progress at the time of reporting (Keplinger *et al.*, 1975).

Table I

Incidence of tumours in mice exposed to vinyl chloride
(purity unspecified) for 8 months¹

Exposed groups	No. of mice with tumours at death			Type and location of tumour		
				Adenomas	Angiosarcomas	Adenocarcinomas
	Male	Female	Total	lung	liver	mammary gland
50 ppm	1	3	4	2	2	2
200 ppm	3	12	15	12	11	3
2500 ppm	6	24	30	28	28	6
Control	0	0	0	0	0	0

¹From Keplinger *et al.* (1975), preliminary results

Two groups each of 12 male and 12 female 3-month-old NMRI outbred albino mice were exposed to either 130 or 1300 mg/m³ (50 or 500 ppm) vinyl chloride in air for 6 hours/day on 5 days/week. The 500 ppm group was exposed for 26 weeks only (due to the poor condition of the mice); the 50 ppm group was exposed for 52 weeks, at which time the experiment was terminated. In the low-dose group, 18/24 animals had developed tumours, including pulmonary adenomas in 13/24, angiosarcomas at various sites in 15/24 and a mammary carcinoma in 1 mouse. Inhalation of 500 ppm vinyl chloride for 26 weeks induced pulmonary adenomas in all mice; in addition, 8 mice had angiosarcomas, mammary adenocarcinomas were found in 4 animals, 1 mouse had an angiosarcoma of the liver, 1 an adenoma of the kidney and 1 an angiosarcoma of brown fat. In the control group, 3/48 had tumours: 1 adenocarcinoma of the mammary gland, 1 dysgerminoma of the ovary and 1 reticulum-cell sarcoma of the spleen (Holmberg *et al.*, 1976).

Groups of 36 male and 36 female 2-month-old albino CD mice were exposed to 130, 650 and 2600 mg/m³ (50, 250 and 1000 ppm) vinyl chloride (99.8% pure) in air for 6 hours/day on 5 days/week for 52 weeks; at that time, 70, 52, 46 and 38 animals were still alive, respectively. A total of 12, 22 and 48 mice developed lung adenomas in the exposed groups, respectively; 1 lung adenoma was found in untreated control animals. In addition, angiosarcomas of the liver developed in 3, 23 and 31 treated mice, respectively, and angiosarcomas in other organs in 7, 5 and 9 mice. Mammary gland tumours were found in 9, 3 and 13 mice, respectively; most of these tumours metastasized to the lungs (Lee *et al.*, 1977, 1978).

Rat: A group of 26 male 3-month-old Ar/IRE Wistar rats were exposed to an atmospheric concentration of 3% v/v (equivalent to 78 g/m³ or 30 000 ppm) commercial grade vinyl chloride (99% pure) for 4 hours/day on 5 days/week for 12 months; the experiment was terminated at 54 weeks. Skin tumours developed in the submaxillary parotid region in all 17 surviving rats (14 epidermoid carcinomas, 2 mucoepidermoid carcinomas, 1 papilloma); in addition, lung tumours developed in 7 rats and osteochondromas in 5. No tumours were observed in 25 untreated controls killed at an unstated time (Viola *et al.*, 1971) [Maltoni & Lefemine (1974) examined slides from this experiment and concluded that the skin tumours were Zymbal gland tumours and that the lung tumours were metastases from these].

Groups of 30 male and 30 female 21-week-old Sprague-Dawley rats were exposed by inhalation to 130-26 000 mg/m³ (50, 250, 500, 2500, 6000 or 10 000 ppm) vinyl chloride in air for 4 hours/day on 5 days/week for 17 weeks. At 86 weeks, 18, 15, 37, 33, 16 and 9 animals were still alive in the 6 groups. At 155 weeks (end of experiment), carcinomas of the Zymbal gland were found in 0, 1, 1, 3, 6 and 7 animals, respectively (1 in untreated controls), nephroblastomas in 1, 2, 0, 2, 1 and 1 animals, liver angiosarcomas in 0, 0, 1, 1, 1 and 0 animals and angiosarcomas at other sites in 2, 0, 1, 2, 1 and 1 (1 in controls). Brain neuroblastomas were seen in 0, 0, 0, 2, 2 and 6 animals, respectively (Maltoni, 1977a; Maltoni *et al.*, 1974).

Groups of 64-96 13-week-old Sprague-Dawley rats were also treated for 52 weeks with the above concentrations of vinyl chloride in air. The following tumours developed in various organs by the end of the experiment, at 135 weeks: carcinomas of the Zymbal gland in 29/239 rats at the 4 highest dose levels, and nephroblastomas (26/257 rats) and angiosarcomas of the liver (47/357 rats) in all the treated groups (the numbers of rats given were those alive at 26 weeks); a total of 14 angiosarcomas was observed in organs other than the liver. The tumours of the liver and kidney metastasized to other organs. No such tumours were observed in 58 untreated controls alive at 26 weeks (see Table II). In a group of 60 17-week-old Sprague-Dawley rats treated with 78 g/m³ (30 000 ppm) vinyl chloride in air for 4 hours/day on 5 days/week for 43 weeks, 30 (50%) developed Zymbal gland carcinomas, 13, liver angiosarcomas and 1, a lung angiosarcoma within the 61 weeks of observation (Maltoni *et al.*, 1974).

Mice were
vinyl chloride
weeks; at that
ly. A total
groups, res-
imals. In
31 treated
5 and 9 mice.
tively; most
(1978).

were exposed
/m³ or 30 000
ay on 5 days/
. Skin
7 surviving
papilloma);
omas in 5.
n unstated
slides from
al gland

ey re
6000 or
ek for 17
11 alive in
the Zymbal
(1 in un-
als, liver
at other
omas were
7a; Maltoni

reated
air. The
experiment,
the 4
psarcomas
ts of rats
ss was
er and
ved in 58
of 60 17-
vinyl
(50%)
a lung
(1974).

Table II¹

Incidence of tumours in Sprague-Dawley rats exposed to vinyl chloride
for 4 hours/day on 5 days/week for 52 weeks and surviving up to 130 weeks

Concentration of vinyl chloride (ppm)	Total no. of animals at start	No. of animals alive at 26 weeks	No. of Zymbal gland tumours	No. of nephro- blastomas	No. of angio- sarcomas of the liver	No. of angio- sarcomas at other sites	No. of brain neuro- blastomas	No. of other tumours
10 000	69	61	16	5	9	3	7	11
6000	72	60	7	4	13	3	3	10
2500	74	59	2	6	13	3	5	7
500	67	59	4	4	7	2	0	8
250	67	59	0	6	4	2	0	7
50	64	59	0	1	1	1	0	10
Controls	68	58	0	0	0	0	0	10

¹From Maltoni *et al.* (1974)

Wistar rats were also exposed by inhalation to 130-26 000 mg/m³ (50, 250, 500, 2500, 6000 and 10 000 ppm) vinyl chloride in air for 52 weeks. After 136 weeks of observation, 1 Zymbal gland carcinoma was found, whereas at a comparable time the Sprague-Dawley rats had developed 29 such tumours. In the Wistar rats, 1 nephroblastoma, 8 liver angiosarcomas and 1 brain neuroblastoma were found in the 10 000 ppm group; 3 nephroblastomas, 2 liver angiosarcomas and 1 brain neuroblastoma in the 6000 ppm group; 3 liver angiosarcomas in the 2500 ppm group; and 1 nephroblastoma and 4 liver angiosarcomas in the 500 ppm group (Maltoni, 1977a; Maltoni *et al.*, 1974).

The effect of length of treatment by inhalation of vinyl chloride on the incidence of liver angiosarcomas was investigated. Groups of 60-120 Sprague-Dawley rats were given either 15.6 or 26 g/m³ (6000 or 10 000 ppm) vinyl chloride in air for 4 hours/day on 5 days/week for 5, 17 or 52 weeks. The experiment was terminated at 155 weeks. Liver angiosarcomas developed in 13 (22%) and 9 (15%) of the 6000 and 10 000 ppm groups exposed for 52 weeks; 1 (0.6%) liver angiosarcoma was found in a rat exposed to 6000 ppm for 17 weeks and none in the 10 000 ppm group. No such tumours were induced in rats treated for 5 weeks (Maltoni, 1977b) [No information was given about tumours occurring at other sites].

The influence of age on the incidence of liver tumours was examined in Sprague-Dawley rats exposed to 15.6 or 26 g/m³ (6000 or 10 000 ppm) vinyl chloride in air for 4 hours/day on 5 days/week for 5 weeks, starting at the age of 13 weeks (120 rats/group) or 1 day (43 and 46 rats). The animals were observed for 135 weeks. One hepatoma was reported in the older rats treated with 10 000 ppm. In the newborn rats, 10 angiosarcomas and 13 hepatomas were observed in the 6000 ppm group, and 10 angiosarcomas and 15 hepatomas were found in rats treated with 10 000 ppm. No liver tumours were reported in the 249 untreated rats (Maltoni, 1977b).

Groups of 36 male and 36 female 2-month-old CD rats were exposed to 0, 130, 650 and 2600 mg/m³ (0, 50, 250 and 1000 ppm) vinyl chloride in air (99.8% pure) for 6 hours/day on 5 days/week for 12 months, at which time the surviving animals (72, 70, 58 and 51) were killed. In rats treated with 250 and 1000 ppm, liver angiosarcomas occurred in 12 and 22 and lung angiosarcomas developed in 3 and 13 (Lee *et al.*, 1977).

In a report of a study in progress, 4 groups of 80 Sprague-Dawley male rats received either 5% ethanol in the drinking-water or drinking-water only for 4 weeks prior to beginning inhalation of 1560 mg/m³ (600 ppm) vinyl chloride for 4 hours/day on 5 days/week for 12 months or air; ethanol-water was given until death or sacrifice. After 60 weeks from the first exposure to vinyl chloride, 55 rats had died or had been sacrificed; liver tumours were found in 21/28 (75%) in the vinyl chloride-ethanol group and 5/13 (38%) in the vinyl chloride only group (Radike *et al.*, 1977).

Hamster: Groups of 32-35 male 11-week-old golden hamsters were exposed by inhalation to 130-26 000 mg/m³ (50, 250, 500, 2500, 6000 and 10 000 ppm)

vinyl ch
weeks fr
At 109 w
treated
epitheli
in 2/70
with 50,
and 35 f
and 2 an
1974).

Rab
week for
Between 9
cinomas v
months of
inadequac

(c)

A gr
given sin
oil. One
[The Work
not given

(d)

Group
received s
oil, 2, 3
coma were
times and

(e)

Prena
given 15.6
day by inh
17/32 and
(end of exp
offspring;
cinoma; ar
and 1, a ne
a Zymbal gl

vinyl chloride in air for 4 hours/day on 5 days/week for 30 weeks. At 48 weeks from the initial treatment, 60/198 treated animals were still alive. At 109 weeks (end of treatment), 2 liver angiosarcomas were seen in hamsters treated with 500 ppm and 1 in those treated with 6000 ppm. Skin trichoe-
 epitheliomas developed in 22 treated hamsters (in 1-6 animals/group) and in 2/70 controls. Two animals treated with 6000 ppm and 1 each treated with 50, 2500 and 10 000 ppm developed melanomas. In addition, 6 lymphomas and 35 forestomach papillomas and acanthomas were found in treated animals, and 2 and 2, respectively, in controls (Maltoni, 1977a; Maltoni *et al.*, 1974).

Rabbit: A group of 40 rabbits were exposed for 4 hours/day on 5 days/week for 12 months to air containing 26 g/m³ (10 000 ppm) vinyl chloride. Between 9-15 months of exposure, 12 skin acanthomas and 6 lung adenocarcinomas were seen. No similar tumours occurred in 20 controls after 15 months of observation (Caputo *et al.*, 1974) [The Working Group noted the inadequacy of reporting].

(c) Subcutaneous and/or intramuscular administration

A group of 75 male and female 21-week-old Sprague-Dawley rats were given single s.c. injections of 4.25 mg/animal vinyl chloride in 1 ml olive oil. One nephroblastoma occurred in the treated animals (Maltoni, 1977a) [The Working Group noted that survival times and period of observation were not given].

(d) Intraperitoneal administration

Groups of 30 male and 30 female 13-week-old Sprague-Dawley rats received single injections of 4.25 mg/animal vinyl chloride in 1 ml olive oil, 2, 3 or 4 times in 2 months. One nephroblastoma and 1 s.c. angiosarcoma were found (Maltoni, 1977a) [The Working Group noted that survival times and period of observation were not given].

(e) Other experimental systems

Prenatal exposure: Two groups of 30 female Sprague-Dawley rats were given 15.6-26 g/m³ (6000 or 10 000 ppm) vinyl chloride in air for 4 hours/day by inhalation from the 12-18th day of pregnancy. Of the offspring, 17/32 and 23/54 had died by the 95th week after birth. After 143 weeks (end of experiment), 1 s.c. angiosarcoma was observed in each group of offspring; 1 animal exposed *in utero* to 6000 ppm had a Zymbal gland carcinoma; and 3 animals exposed to 10 000 ppm had Zymbal gland carcinomas, and 1, a nephroblastoma. One female rat treated with 10 000 ppm developed a Zymbal gland carcinoma (Maltoni, 1974; Maltoni, 1977a).

3.2 Other relevant biological data(a) Experimental systemsToxic effects

The 2-hour LC_{50} of vinyl chloride for mice was 294 g/m^3 (113 000 ppm); for rats, 390 g/m^3 (150 000 ppm); for guinea-pigs, 595 g/m^3 (230 000 ppm); and for rabbits, 295 g/m^3 (113 000 ppm). Vinyl chloride gas had a narcotic effect on experimental animals, the most sensitive species being mice, followed by rats, guinea-pigs and rabbits. The death of animals was preceded by excitement, contractions and convulsions, accelerated respiration, followed by respiratory failure. Rabbits and guinea-pigs had more accentuated muscular contractions and convulsions than mice and rats. Microscopically, congestion of the internal organs with more intense damage to the lungs, liver and kidneys were found (Prodan *et al.*, 1975a).

The hepatotoxicity of vinyl chloride has been shown to be increased after administration of cytochrome P-450 inducers such as phenobarbital, Aroclor 1254 and hexachlorobenzene (Ivanetich *et al.*, 1977; Reynolds *et al.*, 1975a,b). The extent of liver damage has been measured by the release of alanine α -ketoglutarate, glutamic oxalacetic and glutamic pyruvic transaminases (Reynolds *et al.*, 1975b, 1976) and of sorbitol dehydrogenase (Conolly & Jaeger, 1977) into the serum.

A single 6-hour inhalation exposure to 130 g/m^3 vinyl chloride (50 000 ppm) produced acute liver injury in male Sprague-Dawley rats pretreated with phenobarbital or Aroclor 1254. The degree of injury, as indicated by elevation of serum levels of enzymes derived from the liver, correlated with the magnitude of induction of cytochrome P-450 and morphological changes in the endoplasmic reticulum (Reynolds *et al.*, 1975a,b, 1976). Similar findings were reported in phenobarbital-pretreated male Holtzman rats (Jaeger *et al.*, 1974) and in phenobarbital-treated male Charles River CD-1 rats that received 10 daily exposures for 6 hours/day to 35 g/m^3 (13 500 ppm) vinyl chloride in air (Drew *et al.*, 1975).

Cytochrome P-450 concentration decreased during *in vivo* exposure or during *in vitro* incubation of liver homogenate from phenobarbital or 3-methylcholanthrene-induced rats (Ivanetich *et al.*, 1977; Reynolds *et al.*, 1975c).

In an abstract, it was reported that male rats pretreated by gavage with Aroclor 1254 for 3 consecutive days and exposed on day 4 by inhalation to 62.5 g/m^3 (24 000 ppm) vinyl chloride for 4 hours showed significant elevations of serum alanine- α -ketoglutarate transaminase and severe degeneration and necrosis of the liver (Conolly *et al.*, 1977). Overnight fasting, which depletes hepatic glutathione, of Aroclor-pretreated male Holtzman rats before exposure to 26 g/m^3 (10 000 ppm) for 4 hours significantly increased the hepatotoxic effects, as measured by sorbitol dehydrogenase levels in the serum (Conolly & Jaeger, 1977).

Sim
ppm viny.
injury in
vinyl chl
cantly ex
in fed ra

Expe
2 hours/c
histopath
and lungs
denoting
Large qua
by vinyl

Mice
vinyl chl
some acut
renal cor
lost weig
for 12 mo

The
exposed to
a function
1976a).

Viny
oxide and
(Border &

Embry

Pregn
(50 and 50
rats to 13
and New Ze
6-18 of ge
to 15% eth
of several
received v
1975).

Absor

The
and review
Henschler,
1975; Mala
1976; Plug
1976b,c).

UCC
060479

Simultaneous exposure to 1.75 g/m³ (671 ppm) vinyl chloride with 200 ppm vinylidene chloride prevented vinylidene chloride-induced hepatic injury in fasted male rats. However, pre-exposure to concentrations of vinyl chloride which depleted hepatic glutathione concentrations significantly enhanced early acute hepatotoxic response to vinylidene chloride in fed rats (Jaeger *et al.*, 1975a,b).

Exposure of guinea-pigs to 260 g/m³ (100 000 ppm) vinyl chloride for 2 hours/day for 3 months resulted in marked growth disturbances and intense histopathological and histochemical lesions in the liver, kidneys, spleen and lungs. Interruption of the exposure resulted in a regenerative effect, denoting a certain degree of reversibility of the hepatorenal lesions. Large quantities of vitamin C reduced the gravity of the lesions caused by vinyl chloride (Prodan *et al.*, 1975b).

Mice were exposed to 130, 650 or 2600 mg/m³ (50, 250 or 1000 ppm) vinyl chloride for 6 hours/day on 5 days/week. The highest dose caused some acute deaths with toxic hepatitis and marked tubular necrosis in the renal cortex. From the 6th month of treatment, all mice became lethargic, lost weight quickly and died. Only a few mice exposed to 50 ppm survived for 12 months (Lee *et al.*, 1977).

The non-protein, free SH-groups of the liver are depleted in rats exposed to 390-5200 mg/m³ (150-2000 ppm) vinyl chloride for 1-7 hours, as a function both of concentration and duration of exposure (Watanabe *et al.*, 1976a).

Vinyl chloride and two of its presumed metabolites, chloroethylene oxide and chloroacetaldehyde, depressed DNA synthesis in rat liver *in vivo* (Border & Webster, 1977).

Embryotoxicity and teratogenicity

Pregnant CF-1 mice were exposed by inhalation to 130 and 1300 mg/m³ (50 and 500 ppm) vinyl chloride on days 6-15 of gestation, Sprague-Dawley rats to 1300 and 6500 mg/m³ (500 and 2500 ppm) on days 6-15 of gestation, and New Zealand rabbits to 1300 and 6500 mg/m³ (500 and 2500 ppm) on days 6-18 of gestation, for 7 hours/day, with or without simultaneous exposure to 15% ethanol in the drinking-water. A significantly increased incidence of several skeletal anomalies was observed in offspring of mice that received vinyl chloride plus ethanol (John *et al.*, 1977; Schwetz *et al.*, 1975).

Absorption, distribution, excretion and metabolism

The *in vivo* and *in vitro* metabolism of vinyl chloride has been studied and reviewed (Antweiler, 1976; Bartsch & Montesano, 1975; Bonse & Henschler, 1976; Green & Hathway, 1975, 1977; Haley, 1975; Hefner *et al.*, 1975; Malaveille *et al.*, 1975; Müller & Norpoth, 1975; Müller *et al.*, 1976; Plugge & Safe, 1977; Watanabe & Gehring, 1976; Watanabe *et al.*, 1976b,c).

Low concentrations (130 mg/m³, 50 ppm, for 65 min) of vinyl chloride are readily metabolized in rats exposed by inhalation and are converted into polar metabolites, which are predominantly excreted in the urine; a very small amount is expired in air as unchanged vinyl chloride (Hefner *et al.*, 1975).

Following exposure of male rats by inhalation to 26 mg/m³ (10 ppm) ¹⁴C-vinyl chloride for 6 hours, urinary ¹⁴C-activity and expired vinyl chloride comprised 68 and 2%, respectively, of the recovered radioactivity; after exposure to 2600 mg/m³ (1000 ppm) ¹⁴C-vinyl chloride, the proportion of the radioactivity in the urine was lower and that expired as vinyl chloride higher, representing 56 and 12%, respectively. The pattern of pulmonary elimination of 10 and 1000 ppm vinyl chloride *per se* was described by apparently similar first-order kinetics, with half-lives of 20.4 and 22.4 min, respectively; the half-lives for the initial phase of excretion of ¹⁴C-radioactivity in the urine were 4.6 and 4.1 hours, respectively. ¹⁴C-Radioactivity recovered from the carcass after 72 hours was 14 and 15%, respectively; no vinyl chloride *per se* was found in tissues. The proportions of 3 urinary metabolites, *N*-acetyl-*S*-(2-hydroxyethyl)cysteine, thiodiglycolic acid (thiodiacetic acid) and an unidentified metabolite, were not markedly influenced by the level of exposure (Watanabe *et al.*, 1976b).

Following single oral administration of 0.05, 1 or 100 mg/kg bw ¹⁴C-vinyl chloride to male rats, excretion in the urine was 59, 68 and 11%, respectively; the ¹⁴CO₂ in expired air accounted for 9, 13 and 3%, respectively; pulmonary elimination of unchanged vinyl chloride represented only 1-3% of the lower dose levels and 67% of the higher level. The pulmonary clearance of 0.05 and 1 mg/kg bw doses of vinyl chloride was monophasic, with half-lives of 53.3 and 57.8 min, respectively; it was biphasic after administration of 100 mg/kg bw, with half-lives of 14.4 and 40.8 min for the fast and slow phases, respectively. The percentages of the doses left in the carcass after 72 hours were 10, 11 and 2% of the 0.05, 1 and 100 mg/kg doses, respectively. Two of 3 urinary metabolites were identified as *N*-acetyl-*S*-(2-hydroxyethyl)cysteine and thiodiglycolic acid; their proportions were not influenced by dose (Watanabe *et al.*, 1976c). It has been suggested that the metabolism of vinyl chloride in rats following oral and inhalation exposure is a saturable process (Watanabe *et al.*, 1976b,c).

The kinetic parameters and half-lives for the elimination of vinyl chloride from rats after inhalation and i.v. administration have also been reported by Withey (1976).

When rats were exposed to initial concentrations of less than 260 mg/m³ (100 ppm) [1,2-¹⁴C]-vinyl chloride, about 40% of that inspired was absorbed by the lung. Highest radioactivity levels were observed in the liver and kidney immediately after exposure. Most of the radioactive metabolites were excreted rapidly, largely by the kidneys: the radioactivity in the urine amounted to 70% within 24 hours. Some metabolites, however, remained in tissues (mostly in spleen, liver, kidneys) even 48 hours after exposure (Bolt *et al.*, 1976). Metabolites that were not excreted in urine were partly

excreted
Green &

Vin
to chloro
dehyde
in vivo.
chloride
homogena
internec
et al.,
of the 1
formed w
et al.,
of vinyl
acetalde

Met
chrome P
Salmon,
3-bromop
vinyl ch
oxide, w
(Barbin
1975).
thione S-
which is
1977) (F
which is
methyl gl
thiodigly

Chlo
bolites,
N-Acetyl-
1977; Wa
vinylcyst
after ora
ethyl)cys
ethyl)cys
chloride
obtained
taldehyde.
tion of th
animals gi
or chloroa
Hathway, 1

Follo
Hathway, 1
acid were


excreted *via* faeces and partly *via* expiration of $^{14}\text{CO}_2$ (Bolt *et al.*, 1976; Green & Hathway, 1975).

Vinyl chloride is metabolized by microsomal mixed-function oxidases to chloroethylene oxide, which can rearrange spontaneously to chloroacetaldehyde (Fig. 1). Although there is no direct evidence for this pathway *in vivo*, the following data are consistent with this hypothesis. Vinyl chloride in the presence of a mouse liver microsomal fraction, rat liver homogenate, an NADPH generating system and oxygen yielded an alkylating intermediate which reacted with either 3,4-dichlorobenzenethiol (Green *et al.*, 1974) or with 4-(4-nitrobenzyl)pyridine. The absorption spectra of the latter adduct was identical to those obtained with the product formed with synthetic chloroethylene oxide (Barbin *et al.*, 1975; Bartsch *et al.*, 1976). These studies indicate that the primary *in vitro* metabolite of vinyl chloride is chloroethylene oxide, which can rearrange to chloroacetaldehyde.

Metabolism of vinyl chloride occurs predominantly through the cytochrome P-450 system (Ivanetich *et al.*, 1977; Reynolds *et al.*, 1975c; Salmon, 1976). Inhibitors of microsomal mixed-function oxidases, such as 3-bromophenyl-4(5)-imidazole or 6-nitro-1,2,3-benzothiadiazole, reduced vinyl chloride metabolism *in vivo* (Bolt *et al.*, 1976). Chloroethylene oxide, with a half-life of 1.6 min in aqueous solution at neutrality (Barbin *et al.*, 1975), rearranges to chloroacetaldehyde (Bonse *et al.*, 1975). Chloroacetaldehyde combines directly or enzymatically *via* glutathione *S*-transferase with glutathione to form *S*-formylmethylglutathione, which is excreted as *N*-acetyl-*S*-(2-hydroxyethyl)cysteine (Green & Hathway, 1977) (Fig. 1). Chloroacetaldehyde can be oxidized to chloroacetic acid, which is either excreted as such or bound to glutathione to form *S*-carboxymethyl glutathione, which upon further enzymic degradation is excreted as thiodiglycolic acid (thiodiacetic acid) (Plugge & Safe, 1977).

Chloroacetic acid was metabolized in rats to two major urinary metabolites, *S*-carboxymethylcysteine and thiodiacetic acid (Yllner, 1971). *N*-Acetyl-*S*-(2-hydroxyethyl)cysteine (a major metabolite) (Green & Hathway, 1977; Watanabe *et al.*, 1976b,c), *S*-(carboxymethyl)cysteine and *N*-acetyl-*S*-vinylcysteine have been shown to be metabolites of vinyl chloride in rats after oral administration (Green & Hathway, 1977) and *N*-acetyl-*S*-(2-hydroxyethyl)cysteine after inhalation (Watanabe *et al.*, 1976b); *S*-(2-chloroethyl)cysteine was also identified after oral administration of vinyl chloride to rats (Green & Hathway, 1975). As thiodiglycolic acid was obtained as a common metabolite in rats dosed separately with chloroacetaldehyde, chloroacetic acid or *S*-(carboxymethyl)cysteine, the identification of the same *S*-containing metabolite from vinyl chloride-treated animals gives further support to the hypothesis that chloroethylene oxide or chloroacetaldehyde are formed and react with glutathione (Green & Hathway, 1977).

Following oral administration of ^{14}C -vinyl chloride, $^{14}\text{CO}_2$ (Green & Hathway, 1975; Watanabe *et al.*, 1976c), ^{14}C -labelled urea and glutamic acid were identified as minor metabolites (Green & Hathway, 1975).

Figure 1^a

^a Adapted from Plugge & Safe (1977)

^b Detected *in vivo*

^c Detected *in vitro*

^d GSH = glutathione

the ch depend et al. is inv the pri in vit; 1977).

Cf (Barric in the in vitr chloroa 1972). that ha or with 1977). cytidin DNA tre fed 250

The v re ide and pr t chloride

Mut

The Montesan

Vin substitu in the p 1976; B McCann e 1975; G. specimen vinyl ch tion syst supernat 1975; Mc

Vin in the Sc but produ forward m in Saccha m use liv mediated

UCC
060483

In vitro binding of ^{14}C -vinyl chloride was shown to be dependent on the thiol content of proteins (Bolt & Filser, 1977), and binding was dependent on the presence of NADPH, oxygen and microsomal enzymes (Kappus *et al.*, 1976). It has been suggested that an epoxide of vinyl chloride is involved in the covalent binding reaction (Kappus *et al.*, 1975). In the presence of a rat liver microsomal system, vinyl chloride binds to RNA *in vitro* (Kappus *et al.*, 1975) and to RNA and DNA *in vivo* (Laib & Bolt, 1977).

Chloroacetaldehyde reacts with adenosine to give 1, N^6 -ethenoadenosine (Barrio *et al.*, 1972). Chloroethylene oxide and vinyl chloride, incubated in the presence of a mouse liver-microsomal preparation with adenosine *in vitro*, produced the same product (Barbin *et al.*, 1975). Reaction of chloroacetaldehyde with cytidine gives 3, N^4 -ethenocytidine (Barrio *et al.*, 1972). 1, N^6 -Ethenoadenosine was isolated after hydrolysis of polyadenosine that had been incubated with rat liver microsomes and ^{14}C -vinyl chloride or with liver RNA of rats treated with ^{14}C -vinyl chloride (Laib & Bolt, 1977). The corresponding etheno-derivatives of deoxyadenosine and deoxycytidine were identified in hydrolysis products obtained from calf thymus DNA treated with chloroacetaldehyde *in vitro* and from liver DNA of rats fed 250 mg/l vinyl chloride in their drinking-water (Green & Hathway, 1978).

The 2-hydroxyethyl derivatives of guanine, cysteine and histidine were identified after chemical reduction of the hydrolysis products of DNA and proteins isolated from the livers of mice treated with ^{14}C -vinyl chloride (Osterman-Golkar *et al.*, 1977).

Mutagenicity and other short-term tests

The mutagenicity of vinyl chloride has been reviewed by Bartsch & Montesano (1975), Bartsch *et al.* (1976) and Fishbein (1976).

Vinyl chloride vapour induced reverse mutations of the base-pair substitution type in *Salmonella typhimurium* G46, TA1530, TA1535 and TA100 in the presence of a 9000 x g supernatant from rat liver (Andrews *et al.*, 1976; Bartsch *et al.*, 1975; Garro *et al.*, 1976; Malaveille *et al.*, 1975; McCann *et al.*, 1975; Rannug *et al.*, 1974), mouse liver (Bartsch *et al.*, 1975; Garro *et al.*, 1976; Malaveille *et al.*, 1975) and human liver biopsy specimens (Bartsch *et al.*, 1975, 1979; Malaveille *et al.*, 1975). Although vinyl chloride also induced mutations in the absence of a metabolic activation system, a much higher mutagenic response was observed when a 9000 x g supernatant from liver was added (Andrews *et al.*, 1976; Bartsch *et al.*, 1975; McCann *et al.*, 1975).

Vinyl chloride in aqueous or methanolic solution was not mutagenic in the *Salmonella* test system (Bartsch *et al.*, 1975; Rannug *et al.*, 1974) but produced reverse mutations in *Escherichia coli* K12 (Greim *et al.*, 1975), forward mutations in *Schizosaccharomyces pombe* and mitotic gene conversions in *Saccharomyces cerevisiae* in the presence of a 9000 x g supernatant from mouse liver. Forward mutations in *S. pombe* were also induced in the host-mediated assay in mice (Loprieno *et al.*, 1976, 1977).

Vinyl chloride as vapour or as ethanol solution was not mutagenic in *Neurospora crassa* in the presence or absence of a metabolic activation system (Drozdzowicz & Huang, 1977).

In inhalation experiments in *Drosophila melanogaster*, vinyl chloride was mutagenic in the recessive lethal test (Magnusson & Ramel, 1976; Verburgt & Vogel, 1977) but not mutagenic in tests for dominant lethals, translocations and sex-chromosome loss (Verburgt & Vogel, 1977).

No dominant lethals were observed in male CD-1 mice after exposure by inhalation to 7.8, 26, or 78 g/m³ (3000, 10 000 or 30 000 ppm) vinyl chloride in air for 6 hours/day for 5 days (Anderson *et al.*, 1976, 1977).

Exposure to vinyl chloride vapour in the presence of a 15 000 x g supernatant from phenobarbital-pretreated rat liver induced forward mutations in V79 Chinese hamster cells in terms of 8-azaguanine and ouabain resistance (Drevon *et al.*, 1977).

The mutagenicity of several possible metabolites of vinyl chloride has also been examined. Chloroethylene oxide was the strongest mutagen among those tested in *S. typhimurium* TA1530 and TA1535 (Bartsch *et al.*, 1975; Malaveille *et al.*, 1975; Rannug *et al.*, 1976), *E. coli* (Hussain & Osterman-Golkar, 1976), *S. pombe* (Loprieno *et al.*, 1977), *S. cerevisiae* (Loprieno *et al.*, 1977) and V79 Chinese hamster cells (Huberman *et al.*, 1975). Chloroacetaldehyde was mutagenic in *S. typhimurium* TA1535, TA1530 and TA100 (Bartsch *et al.*, 1975; Malaveille *et al.*, 1975; McCann *et al.*, 1975; Rannug *et al.*, 1976) and V79 Chinese hamster cells (Huberman *et al.*, 1975). Chloroethanol was a weak mutagen in *S. typhimurium* TA1530, TA1535 and TA100 (Bartsch *et al.*, 1976; Malaveille *et al.*, 1975; McCann *et al.*, 1975; Rannug *et al.*, 1976; Rosenkranz *et al.*, 1974). Chloroacetic acid was not mutagenic in *S. typhimurium* TA1530, TA100 or TA1535 (Bartsch *et al.*, 1975; Malaveille *et al.*, 1975; McCann *et al.*, 1975; Rannug *et al.*, 1976).

1,2-Dichloroethane, a possible by-product of vinyl chloride production and a main component of waste products from vinyl chloride industries (EDC-tar), was mutagenic in *S. typhimurium* TA1535 (Rannug & Ramel, 1977) and in *S. typhimurium* TA100 (McCann *et al.*, 1975) in the presence or absence of a liver microsomal metabolic activation system.

(b) Humans

Toxic effects

Exposure to vinyl chloride is associated with multiple systemic disorders, including a sclerotic syndrome, acro-osteolysis (sometimes associated with a Raynaud-like symptomatology), thrombocytopenia and liver damage, consisting of parenchymal damage, fibrosis of the liver capsule, periportal fibrosis associated with hepatomegaly, and splenomegaly (Lange *et al.*, 1974a; Thomas *et al.*, 1975).

Ex-
factory
disease
removed
after t

Not
found in
chloride
but fibr
premalig

Of
present:
due to p
(William

Exp
and live
vision f

Ele
200 poly
populati
was foun
symptoms
immunolo
in 2/30
order (W

A g
were exa
of disor
and sple
anaemia
syndrome
spontane
incidence
levels d
spastic
suggestiv
lial sys

Redu
vinyl chi
lence of
ting that
(Miller

Examination of 70 workers from a single polyvinyl chloride-producing factory showed a high frequency of signs and symptoms of vinyl chloride disease. Although skin and bone changes may disappear when the patient is removed from contact with vinyl chloride, the thrombocytopenia persists after termination of exposure (Veltman *et al.*, 1975).

Non-cirrhotic portal fibrosis with associated portal hypertension was found in 7 patients who had been involved in the production of vinyl chloride monomer for 4-15 years. An angiosarcoma developed in one patient, but fibrosis was a more common lesion and was considered to be probably not premalignant (Smith *et al.*, 1976a).

Of 487 workers involved in polyvinyl chloride production, two cases presenting with thrombocytopenia were found to have portal hypertension due to periportal fibrosis, with oesophageal varices and splenomegaly (Williams *et al.*, 1975, 1976).

Exposure to vinyl chloride was not only associated with circulatory and liver dysfunction and skin and bone disorders, but also deafness, vision failure and giddiness (Jühe & Lange, 1972).

Elevated carcino-embryonic antigen levels have been found in 48% of 200 polyvinyl chloride workers, as compared with 9% of a normal healthy population (Pagé *et al.*, 1976). No evidence of an auto-immune disorder was found in 13 patients employed in polyvinyl chloride production who had symptoms of 'vinyl chloride disease' (Lange *et al.*, 1974a); however, immunological data from 19/28 patients with vinyl chloride disease and in 2/30 workers exposed to vinyl chloride suggested an immune complex disorder (Ward *et al.*, 1976).

A group of 168 workers (114 from one factory and 54 from another) were examined medically at various times during 1962-1969. Manifestations of disorders of the nervous system were recorded commonly; hepatomegaly and splenomegaly occurred in 30% and 6% of workers; and some cases of anaemia and leucopenia were also observed. The incidence of Raynaud's syndrome fell from 6% in 1962 to 2.9% in 1966; this phenomenon cleared spontaneously upon removal of the subjects from exposure: these different incidence figures were associated with a 22-fold decrease in vinyl chloride levels during the period of the study. A much higher percentage of vaso-spastic changes was found in the two groups (66 and 55%, respectively), suggesting that vinyl chloride acts as an irritant in the reticuloendothelial system to produce reactive splenic enlargement (Suciu *et al.*, 1975).

Reduced pulmonary function has been observed in workers exposed to vinyl chloride (Gamble *et al.*, 1976; Miller *et al.*, 1975). The prevalence of this impairment was similar in smokers and nonsmokers, suggesting that occupational or other environmental factors were operative (Miller *et al.*, 1975).

Embryotoxicity and teratogenicity

A significant excess of foetal deaths was reported in women whose husbands were exposed to vinyl chloride: 15.8%, or 23, foetal deaths in 139 pregnancies, as compared with 8.8% (24/273) in the age-adjusted control group. This excess of foetal deaths was shown not to be a function of chronic abortions, i.e., the association was maintained after excluding pregnancies of women who had had more than 2 abortions (Infante *et al.*, 1976a). The significance of this study was questioned because data collection methods were not specified and there was no statistical treatment of the data (Paddle, 1976). Subsequently, the data collection methods were described, showing that there had been no interviewer-responder bias, and details of statistical analyses were specified (Infante *et al.*, 1976b).

In a registry-based study, Infante (1976) reported that an excess of central nervous system defects, of deformities of the upper alimentary and genital tracts, and of clubfoot has been observed in stillborn and live children in 3 cities in Ohio in which vinyl chloride polymerization plants are located.

In hospital-based studies in newborns in Painesville (Ohio), where there are two polyvinyl chloride plants, and in Kanawha county (West Virginia), where there is one plant, excesses of anencephaly and spina bifida were reported, but no association was made with vinyl chloride (Edmonds, 1977; Edmonds *et al.*, 1975, 1978).

Mutagenicity and other short-term tests

Chromosome aberrations were found in workers occupationally exposed to vinyl chloride in the US (Ducatman *et al.*, 1975; Heath *et al.*, 1977), Sweden (Funes-Cravioto *et al.*, 1975), the UK (Purchase *et al.*, 1975), Belgium (Léonard *et al.*, 1977), Hungary (Szentesi *et al.*, 1976) and Norway (Hansteen *et al.*, 1976). These aberrations were in most cases fragments, dicentrics and rings, and breaks and gaps.

3.3 Case reports and epidemiological studies¹

In 1974, more than 40 years after the introduction of vinyl chloride into industry, Creech & Johnson (1974) first reported an association of exposure to this chemical with cancer in man. Three cases of liver angiosarcoma were reported in men who were employed in the manufacture of polyvinyl chloride resins (one had cleaned reactor vessels) in a single vinyl chloride polymerization plant in the US.

¹The Working Group was aware of a study in progress on the occupational and community carcinogenic risk of vinyl chloride (IARC, 1978b).

E
matic
and an
countr
Czecho
et al.
Roche
Romani
Harry,
1974a;
of 64
(Spirt:

Nc
toxic r
review
zation
of live
cluded
detect
organs

Tr
one yea
compare
system,
of the
those n
vinyl c
for 152
of expo

In
plants
due to
were 11
1974).
grounds
related
failed
to reti:

The
each of
5 years
seniorit
a 2.3-fc
3 were c

By reviewing medical records and pathological material and by systematic medical screening, the association between exposure to vinyl chloride and angiosarcoma of the liver has been reported from a number of other countries: Canada (Delorme & Thériault, 1978; Noria *et al.*, 1976); Czechoslovakia (Lloyd, 1975); the Federal Republic of Germany (Lange *et al.*, 1974b, 1975); France (Couderc *et al.*, 1976; Ravier *et al.*, 1975; Roche *et al.*, 1978); Italy (Maltoni, 1974); Norway (Lloyd, 1975); Romania (Lloyd, 1975); Sweden (Byrén & Holmberg, 1975); the UK (Lee & Harry, 1974; Smith *et al.*, 1976b); the US (Block, 1974; Falk *et al.*, 1974a; Makk *et al.*, 1974); and Yugoslavia (Šarić *et al.*, 1976). A review of 64 reported cases in various countries as of October 1977 is available (Spirtas & Kaminski, 1978).

No history of acro-osteolysis and no evidence of exposure to hepatotoxic materials other than vinyl chloride was reported in a clinical review of 7 cases of liver angiosarcoma among US vinyl chloride polymerization workers (Heath *et al.*, 1975). In a pathological evaluation of cases of liver angiosarcoma among exposed US vinyl chloride workers, it was concluded that these tumours were often multicentric: angiosarcomas were also detected in the wall of the duodenum, in the heart and kidney, and in other organs (Thomas & Popper, 1975).

The cancer risk among a cohort of males in the US who had at least one year of occupational exposure to vinyl chloride was studied. When compared with the US male population, an excess of cancer of the digestive system, of the liver (primarily angiosarcoma), of the respiratory system, of the brain and of unknown sites, as well as lymphomas was observed in those members of the study cohort with the greatest estimated exposure to vinyl chloride (Tabershaw & Gaffey, 1974) [Vital status was undetermined for 15% of the study cohort, and only 50% had 15 or more years since onset of exposure to vinyl chloride].

In a proportional-mortality analysis of 161 deceased workers in two US plants producing and polymerizing vinyl chloride, a 50% excess of deaths due to all cancers was reported. Sites of cancer with the greatest excess were liver and biliary tract, brain, digestive tract and lung (Monson *et al.*, 1974). Falk *et al.* (1974b) questioned the authors' conclusion, on the grounds that not all deaths studied were among workers in activities directly related to vinyl chloride production or polymerization and that the study failed to include deaths among workers who had terminated employment prior to retirement or death.

The cancer mortality experience of 257 US workers (255 were traced), each of whom had been occupationally exposed to vinyl chloride for at least 5 years and observed after 10 years from onset was studied using union seniority and company employment records. Among 24 deaths from all causes, a 2.3-fold excess was observed in deaths from cancer; of the 24 deaths, 3 were due to haemangiosarcoma of the liver (Nicholson *et al.*, 1975).

No excess of total or cause-specific mortality was reported in a study of 2100 male workers in the UK exposed to vinyl chloride for periods of up to 27 years; in addition, the authors reported a decreasing risk of mortality with increasing duration of exposure to vinyl chloride (Duck *et al.*, 1975). Wagoner *et al.* (1976) challenged the conclusions of the study on the grounds of analytical shortcomings. After reanalysing the data, the authors (Duck & Carter, 1976) reported an increased risk of cancer of the digestive system 15 years after initial exposure to vinyl chloride.

The cancer mortality of 594 US workers exposed occupationally to vinyl chloride and to lesser amounts of vinylidene chloride (see p. 439) and other compounds (such as methyl methacrylate, see p. 187, and acrylonitrile, see p. 73) was studied. Although no angiosarcomas were found (no deaths due to any liver cancer), an excess of all malignancies combined was reported among those workers classified as having been highly exposed to vinyl chloride when compared with all other exposure categories. However, the number of workers in the lower exposure categories who were exposed for more than 10 years was small, resulting in part from the fact that workers first took jobs in the dry end of the polymerization process where exposures to vinyl chloride were low; many employees who remained with the units and established seniority would subsequently have moved to the higher exposure areas (Ott *et al.*, 1975).

The incidence of abnormal sputum cytology among workers in the vinyl chloride-polyvinyl chloride industry in Italy was much higher than expected, even when compared with a population of heavy smokers who did not work in chemical industries (Maltoni, 1976).

Waxweiler *et al.* (1976) studied the cancer mortality experience of 1294 individuals with 5 or more years of employment and 10 years since onset of employment in departments or jobs with direct exposure to vinyl chloride at 1 of 4 vinyl chloride-polyvinyl chloride production plants in the US. When compared with the US white male population, an excess of cancer was found in four organ systems: brain and central nervous system, respiratory system, hepatic system, and lymphatic and haematopoietic systems. This excess of organ-specific cancer was restricted to those workers with 15 or more years since onset of vinyl chloride exposure. For all malignant neoplasms combined, the standard mortality ratio was 184; for the brain and central nervous system, 498; for the respiratory system, 194; for the hepatic system, 1606; and for the lymphatic and haematopoietic system, 176. Of 14 histologically confirmed cases of biliary and liver cancer among workers from these 4 plants, 11 were angiosarcoma of the liver. Of 10 cases of brain cancer, 9 were classified histologically as glioblastoma multiforme, a cell type of brain cancer reported to be unusual in the US. Of the 14 cases of primary lung cancer, 5 were large-cell undifferentiated and 3 were adenocarcinoma.

In 771 workers employed in a Swedish vinyl chloride-polyvinyl chloride plant since its start in the early 1940's, a 4- to 5-fold excess of cancer of the liver and pancreas was found. Although the risks of cancer of the

brain and
signific:

When
currently
sure to v
those inc
initial e
(Fox & Co

Fox
males who
chloride
from liver
was thought
evidence
Relatively
even in ca
follow-up
effect of

A stu
in the pro
Republic o
an excess
phatic org
with durati

The re
of age and
facilities.
was higher
leukaemia
(1976).

The in
period in a
polymerized
were includ
was found b
et al., 1976

brain and of the lung were also increased, they were not statistically significant (Byrén *et al.*, 1976).

Whereas no excess mortality from lung cancer was demonstrated among currently employed individuals 15 years after initial occupational exposure to vinyl chloride, a 56% excess of lung cancer was observed among those individuals who had terminated employment less than 15 years since initial exposure; this excess was found for each duration of exposure (Fox & Collier, 1976).

Fox & Collier (1977) investigated the cancer mortality among 7561 males who were exposed to vinyl chloride in the manufacture of polyvinyl chloride in the UK at some time between 1940 and 1974. An excess mortality from liver cancer was reported for each group of workers, whether exposure was thought to be high, medium or low; however, the authors reported no evidence of an excess mortality from cancers other than of the liver. Relatively few subjects had long-term exposure to vinyl chloride, and even in cases in which men had completed 20 years of employment, the follow-up period was judged to be too short to evaluate the carcinogenic effect of vinyl chloride.

A study was reported of cancer mortality among 7021 males employed in the production and polymerization of vinyl chloride in the Federal Republic of Germany. When compared with the national male population, an excess of cancers was found for 4 organs: liver, brain, lung and lymphatic organs. This excess of organ-specific cancer was shown to increase with duration of exposure (von Reinl *et al.*, 1977).

The risk of cancer mortality was investigated among residents 45-years of age and older in three US communities with vinyl chloride polymerization facilities. Among males, the death rate from central nervous system cancer was higher than that for the state as a whole. No excess mortality from leukaemia and aleukaemic leukaemia or from lymphoma was found (Infante, 1976).

The incidence of liver and lung cancer was studied for a 4-year period in a city in Yugoslavia with a factory in which vinyl chloride was polymerized and polyvinyl chloride processed. Polyvinyl chloride workers were included in the study. Except for liver angiosarcoma, no association was found between cancer incidence and place of work or residence (Šarić *et al.*, 1976).

Polyvinyl chloride1. Chemical and Physical Data1.1 Synonyms and trade names

Chem. Abstr. Services Reg. No.: 9002-86-2

Chem. Abstr. Name: Chloroethene homopolymer

Atactic poly(vinyl chloride); chloroethylene polymer; poly(chloroethylene); poly(vinyl chloride); poly(vinylchloride); polyvinylchloride; PVC; vinyl chloride homopolymer; vinyl chloride polymer



Airex; AL 30; AL 31; Armodour; Aron compound HW; Astralon;
 Bakelite OYNV; Bakelite QSAH 7; Bakelite QYAC 10; Bakelite QYJV 1;
 Bakelite QYOH-1; Bakelite QYSJ; Bakelite QYSL 7; Bakelite QYTO 7;
 Bakelite UCA 3310; Benvic; Blacar 1716; Bolatron 6200; Bonloid;
 Breon; Breon 107; Breon 113; Breon 121; Breon 125/10; Breon 151;
 Breon 4121; Breon 111EP; Breon 112EP; Breon P 130/1; Breon S 110/10;
 Breon S 125/12; C 65; Caliplast; Carina S 70-01; Carina S 70-71;
 Chemosol; Chlorostop; Cobex (polymer); Contizell; Corvic 55/9;
 Corvic 65/50; Corvic 206573; Corvic 20560600; Corvic 20650600;
 Corvic C 65/02; Corvic D 55/9; Corvic D 57/15; Corvic D 57/17;
 Corvic D 60/11; Corvic D 65; Corvic D 65/02; Corvic D 65/8;
 Corvic D 75/10; Corvic D 6518; Corvic H 55/34; Corvic P 65/50;
 Corvic P 65/54; Corvic P 65/55; Corvic R 65/81; Corvic S 46/70;
 Dacovin; Dacovin 2082; Danuvil 70; Darvic 110; Darvis Clear 025;
 Daycell; Decelith H; Denka Vinyl SS 80; Denka Vinyl SS-Y; Diamond
 Shamrock 40; Diamond Shamrock 71; Diamond Shamrock 450; Diamond
 Shamrock 7602; DN 4; DN 5; Dorlyl; Durofol P; Dynadur; E 62;
 E 66; Ekavyl SD 2; Ekavyl SDF 58; Ekavyl SK64; Ekavyl SK66;
 E 66P; 103EP8; E-PVC; Escambia 2160; Escambia 2200; Europhan;
 Exon 605; Exon 640; Exon 654; Exon 965; Exon 9269; Exon 9290;
 Exon 9269A; FC 4648; Flocor; Genotherm; Genotherm N; Genotherm
 UG 200; Geon 51; Geon 59; Geon 72; Geon 101; Geon 103; Geon
 110X233; Geon 120X241; Geon 121; Geon 124; Geon 126; Geon 128;
 Geon 131; Geon 151; Geon 85542; Geon 101EP; Geon 102EP; Geon

Geor
 Geor
 Hi-
 Host
 PVP
 P;
 Kane
 Kohi
 ES;
 GB 1
 Lucc
 Lucc
 Marv
 Marv
 Movl
 Nipe
 081;
 Norv
 Norv
 Opal
 650;
 Pant
 Pevl
 Pevl
 D 10
 S 50
 Poroc
 PVKh
 QYSA;
 (poly
 540;
 Solvi
 Solvi
 Sumil

Geon 103EP; Geon 103EP8; Geon 103EPF7; Geon 135J; Geon 121L;
 Geon Latex 151; Gutttagena; Halvic 223; Halvic 229; HC 325;
 Hi-S Film No. 111L; Hishirex 502; Hishirex 502Z; Hispavic 229;
 Hostalit; Hostalit E; Hostalit P 7078; Hostalit PVP 3475; Hostalit
 PVP 5470; Hostalit S; Hostalit S 4070; HX-M; Igelite F; Igelite
 P; Improved Wilt Pruf; Kanevinyl PSH 10; Kanevinyl PSL 81;
 Kanevinyl S 1001; Kanevinyl S 1007; Khs 010; KhSE 3; Klegecell;
 Kohiner R 687; KR 800; Kureha S 901; L 5; Lak Kh SL; Lonza 380
 ES; Lonza G; Lucoflex; Lucovyl BB 800; Lucovyl BB 8010; Lucovyl
 GB 1150; Lucovyl GB 9550; Lucovyl GS 1200; Lucovyl GS 8001;
 Lucovyl PB 1302; Lucovyl PE; Lucovyl PE 1100; Lucovyl PE 1290;
 Lucovyl PE 1311; Lucovyl PE 1355; Lucovyl RB 8010; Lutofan;
 Marvinol; Marvinol; Marvinol 14; Marvinol 53; Marvinol 57;
 Marvinol 7000; Marvinol VR 50; Marvinol VR 53; Mirrex MCFD 1025;
 Movinyl 100; Mowilith F; Myraform; NIKA-TEMP; Nikavinyl SG 700;
 Nipeon A 21; Nipol 576; Nipolit CM 081; Nipolit SK; Nipolit SK
 081; Nipolit SL 082; Nipolit SM 092; Nipolit SV 13081; Norvinyl;
 Norvinyl P 2; Norvinyl P10; Norvinyl S 1-70; Norvinyl S 1-80;
 Norvinyl S 3-68; Novon 712; Ongrovil S 165; Ongrovil S 470;
 Opalon; Opalon 410; Opalon 440; Opalon 610; Opalon 630; Opalon
 650; Opalon 660; Opalon R 7611; Ortodur; P 400 (vinyl polymer);
 Pantasote R 873; Pattina V 82; Pevikon D 61; Pevikon KL 2;
 Pevikon PE 709; Pevikon PE 712; Pevikon PS 690; Pevikon R 23;
 Pevikon R 25; Pevikon R 45; Pevikon R 341; Pevikon S 602; Pliovic
 D 100X; Pliovic DB 80V; Pliovic K 906; Pliovic K 90E; Pliovic
 S 50; POK 60; Polivinit; Polwinit; Polyco 2622; Polytherm;
 Porodur; Prototype III Soft; PVKhL 4; PVKhS 60; PVKh-S 60;
 PVKh-S 65; PVKh-S 63Zh; QSAH 7; QSAN 7; Quirvil; Quirvil 278;
 QYSA; Ravinil R 100/65D; Rucon B 20; Ryurene S 800B; S 61; S 65
 (polymer); S 70; S 901; Scon 5300; Sicron; Sicron 530; Sicron
 540; Sicron 548; Sicron 548FM; SKhV 71; S-Lon; SM 200; Solvic;
 Solvic 223; Solvic 229; Solvic 239; Solvic 334; Solvic 340;
 Solvic 406; SP 60; SP 60 (chlorocarbon); SR11; Sumilit EXA 13;
 Sumilit PXA 13; Sumilit PX-A; Sumilit PX-N; Sumilit PXNH;

Sumilit PX-NL; Sumilit SX 11; Sumilit SX 13; Sumilit S X-D;
 Sumilit SX 7G; Sumilit VS 9200; Sumitomo PX 11; SV 55; SX 11;
 SX 7G; SX 8T; Takilon; Technopor; Tenneco 1742; TK 1000;
 Tocryl C 440; Trovidur; Trovidur N; Trovithern HTL; TS 1100;
 U 1; U 1 (polymer); Ultron; VA 15; VC 100; VC 410; Veron P
 130/1; Vestolit B 7021; Vestolit CH; Vestolit S 60; Vestolit S
 6554; Vestolit S 6857; Vestolit S 7054; Vestolit S 7554;
 Vestolit S 8054; Vinika KR 600; Vinika KR 800; Vinika 37M;
 Vinika 35R; Vinikulon; Viniplast; Viniplen P 73; Viniplen P 74;
 Viniplen P 73E; Viniplen P 73EM; Vinnol E 75; Vinnol H 100/65;
 Vinnol H 100/70; Vinnol H 60d; Vinnol H 70D; Vinnol H 100/70d;
 Vinnol H 75F; Vinnol P 70; Vinnol P 70E; Vinnol P 100/70e;
 Vinnol Y; Vinoflex; Vinoflex P 313; Vinylchlon 4000LL; Vinylite
 QYJV; Viplast RA/F; Volgovinyl E 62; Volgovinyl E 62P; Volgovinyl
 E 66P; VSKh-S; Vygen 85; Vygen 110; Vygen 120; Vygen 313;
 Welvic G 2/5; Welvic PRIO 953; Welvic PRO 686; Welvic R 7/622;
 Welvic RI 7/316; Welvic RIO 715; Wilt Pruf; Winidur; X-AB;
 Yugovinyl

1.2 Structural and molecular formulae and molecular weight



Mol. wt: 60 000-150 000 (average)

1.3 Chemical and physical properties of the polymer

From Windholz (1976), unless otherwise specified

- (a) Description: White or colourless granules (Hawley, 1971)
- (b) Density: 1.406
- (c) Refractive index: n 1.54

1.4 Te

A
with va
the spe

PV
tially
mer made
The com

Mo:
at a le
us d are
material
also use

PVC
(PVC res
PVC are
form is

For
samples,

(d) Solubility: Solvents for unmodified polyvinyl chloride (PVC) of high molecular weight are: cyclohexanone, methyl cyclohexanone, dimethyl formamide, nitrobenzene, tetrahydrofuran, isophorone and mesityl oxide. Solvents for lower polymers are: di-n-propyl ketone, methyl amyl ketone, methyl isobutyl ketone, acetonylacetone, methyl ethyl ketone, dioxane and methylene chloride.

(e) Stability: PVC is unstable to heat and light in the absence of added stabilizers. Thermal decomposition products can include ethylene, benzene, toluene (Eckardt & Hindin, 1973), 1,3,5-trichlorobenzene (Tsuge, 1969) and naphthalene (Dyer & Esch, 1976).

1.4 Technical products and impurities

A wide variety of vinyl chloride homo- and copolymers are available, with varying properties designed for specific applications. Consequently, the specifications vary widely.

PVC resins for the production of rigid plastics are processed essentially without plasticizer: the polymer may be a homopolymer or a copolymer made with low levels of comonomer such as vinyl acetate or ethylene. The comonomers are used to aid in the processing of the resulting polymer.

Most of the flexible and semirigid PVC plastics contain plasticizers at a level of 10-100% of the resin weight. The plasticizers most commonly used are dialkyl phthalates (e.g., dioctyl phthalate). Other compounding materials (such as pigments, fillers and light- and heat-stabilizers) are also used.

PVC dispersion or paste resins are used in the form of plastisols (PVC resin dispersed in plasticizer). In Europe, significant amounts of PVC are used in the form of latexes; in the US very little of the latex form is used.

For concentrations of unreacted vinyl chloride monomer in various PVC samples, see p. 382.

2. Production, Use, Occurrence and Analysis2.1 Production and use(a) Production

A method for the synthesis of PVC was reported in 1872 (Baumann, 1872). Commercial homopolymers of vinyl chloride were introduced in 1933 (Darby & Sears, 1968). Vinyl chloride polymer is currently produced by one of four processes: suspension, emulsion, bulk or solution polymerization. In the US in 1976, over 80% of homopolymers and copolymers were produced by the suspension polymerization process.

In 1976, 22 US companies reported production of 2065 million kg PVC resins of all types (US International Trade Commission, 1977); approximately 230 million kg of the total were copolymers. US exports of uncompounded PVC resins in 1977 were about 72 million kg, and exports of compounded PVC (excluding any additives in compounded products) were approximately 39 million kg. These exports went primarily to Belgium (13%), Brazil (12%), Canada (21%), Iran (6%), New Zealand (7%) and Venezuela (13%). US imports of PVC resins are negligible (approximately 1% of domestic production).

Total western Europe production in 1976 amounted to 3745 million kg. The major producers were the following countries (production in millions of kg): Austria (40), Belgium (200), the Federal Republic of Germany (1020), Finland (35), France (615), Greece (25), Italy (675), The Netherlands (290), Norway (55), Portugal (15), Spain (215), Switzerland (30), Sweden (110) and the UK (415). Exports from western Europe in that year were 1050 million kg and imports 690 million kg.

PVC was first produced commercially in Japan prior to 1946. In 1976, nineteen companies produced a total of 1044 million kg; 121 million kg were exported, and 15 million kg imported.

(b) Use

Use of PVC resins in the US in 1976 was as follows: building and construction industries (49%), consumer goods (15%), electrical applications (8%), packaging (9%) and transportation (7%), with miscellaneous uses accounting for the remainder. Since 1968, the major uses have been in the building and construction industries, in consumer goods, packaging and electrical wire insulation.

In building and construction, PVC resins are used in piping and conduits (including water pipes), in flooring, in windows and other rigid structures, in pipe fittings, in sidings and as swimming-pool liners. They are used in such consumer products as upholstery, wall coverings, garden hoses and appliances and also in gramophone records, stationery supplies, footwear, toys, outerwear and sporting goods. Electrical

applicat
uses of
liners ar
beverages
into the
covers, i

In
the US in
tubing ar
blood sto
parenters

The
and fitti
(10%), ar
and other

In
sheet (14
other, in

The
component
contact w
(3) paper
rigid and
may not e
effect (U

An e
tion of P

2.2 Occu

PVC

It h
PVC may h
occurred
effluents
Agency, 1
productio

The
and by in
1974b,c).

applications consist primarily of wire and cable insulation. The major uses of PVC in packaging are in plasticized film, bottles and bottle-cap liners and gaskets; however, in the US, its use for packaging of alcoholic beverages has been banned because of migration of vinyl chloride monomer into the alcohol. Major uses in transport include upholstery and seat covers, automotive tops and automotive floor mats.

In 1975, 27 million kg PVC were used in Europe and 14 million kg in the US in plastic materials for medical applications, including external tubing and catheters; in sheet form for splints; and in shunts, balloons, blood storage bags, cannulae, surgical drapes and packaging containers for parenteral substances (Halpern & Karo, 1977).

The 1977 western Europe use pattern for PVC was as follows: piping and fittings (24%), rigid and flexible films (20%), profiles (12%), cable (10%), artificial leather cloth (8%), bottles (7%), gramophone records (3%) and other uses (16%).

In Japan in 1976, it was used as follows: piping (31%), film (17%), sheet (14%), extrusions and artificial leather (12%), wire/cable (11%) and other, including flooring (15%).

The US Food and Drug Administration permits the use of PVC as a component of the following products when they are intended for use in contact with food: (1) adhesives; (2) resinous and polymeric coatings; (3) paper and paperboard (in contact with dry food only); and (4) semi-rigid and rigid acrylic and modified acrylic plastics. The amount present may not exceed that which is reasonably required to produce the intended effect (US Food and Drug Administration, 1977).

An estimated 700 000 to 2 million workers are employed in the production of PVC in the USA (Infante, 1977).

2.2 Occurrence

PVC is not known to occur as a natural product.

It has been estimated that prior to 1975 more than 22.7 million kg PVC may have been discharged into the environment in the US. These losses occurred as particulates in air emissions, suspended solids in water effluents, and components of solid wastes (US Environmental Protection Agency, 1974b). Workers in the USSR have been exposed to PVC dust in the production of block polymer (Filatova *et al.*, 1974).

The disposal of PVC by ocean dumping, as solid waste, as landfilling and by incineration has been reviewed (US Environmental Protection Agency, 1974b,c).

2.3 Analysis

No information was available to the Working Group on methods for determining PVC residues in foods or other parts of the environment.

A rapid and simple way of identifying fifteen packaging films, including PVC, has been described in which the films were treated with ten different solvents and the solubility and physical appearance of the film at room temperature and at the boiling-points of the solvents were noted (Van Gieson, 1969).

Plastics, including PVC, have been identified by measurement of the pH of an aqueous solution of the pyrolysis products, followed by thin-layer chromatography, and by reaction of PVC with pyridine (Braun & Nixdorf, 1972).

PVC has been separated from plasticizers, stabilizers, modifiers, fillers, pigments and other inorganic additives by gel permeation chromatography or sequential solvent extraction. The resulting additive-free fraction has been analysed by chlorine analysis, infra-red spectroscopy, molecular-weight determination by viscosity measurements and thermal gravimetry and differential thermal analysis (Brookman, 1974).

Pyrolysis-gas chromatography has been proposed to identify PVC in polymers used in medical applications (Nematollahi *et al.*, 1970), in three commercial polymers (in this case, pyrolysis-gas chromatography was used in combination with thermogravimetric analysis and differential thermal analysis) (Boettner *et al.*, 1969), and among some chlorine-containing polymers (Tsuge *et al.*, 1969). Pyrolysis-gas chromatography has also been used to identify polymers, including PVC, among 37 commercial polymers (Okumoto & Takeuchi, 1972), in plastics and rubbers (Fischer, 1967) and in adhesives used in building (in this case, pyrolysis-gas chromatography was used in combination with thermogravimetry, differential thermogravimetry and differential thermal analysis) (Rona, 1971).

Thin-layer chromatography of pyrolysis products has been used to identify high polymers, including PVC (Pastuska, 1969).

Thermogravimetry and infra-red spectral analysis have been used to determine PVC in floor tiles (Powell, 1974).

3.1 Carcinogenicity

Subcutaneous

Rat:
abdominal
some additive
appearance
(38.6%) mal-
the site of
similar but
risk. No
implant of
of 0.03 mm
a similar g
the report
were never

Groups
into the ab-
received an
animals were
rats were k
were found
developed in

A group
(of unstated
animals were
after the im-
fibrosarcoma

In rats
films or per
(Kogan & Tug-

¹The Wor-
carcinogenic
administratio-

UCC
060497

3. Biological Data Relevant to the Evaluation of Carcinogenic Risk to Humans¹

3.1 Carcinogenicity studies in animals¹

Subcutaneous or intraperitoneal administration

Rat: A group of 45 adult Wistar rats were given s.c. implants in the abdominal wall of squares or discs of a commercial PVC known to contain some additives; the implants were 0.04 mm thick and 15 mm wide. At the appearance of the first tumour, 44 animals were still alive. Seventeen (38.6%) malignant tumours (fibrosarcomas and 1 liposarcoma) developed at the site of film implantation, with a latent period of 189-727 days; a similar but perforated film did not produce local tumours in 27 rats at risk. No local tumours were found in a group of 50 rats given a s.c. implant of cotton (Oppenheimer *et al.*, 1952, 1955). With a pure PVC film of 0.03 mm thickness, 4 malignant tumours were obtained after 533 days in a similar group (Oppenheimer *et al.*, 1955) [The Working Group noted that the reporting of this experiment was preliminary and that final results were never forthcoming].

Groups of 35 (male and female) Wistar rats were given a s.c. implant into the abdomen of PVC film 4 x 5 x 0.16 mm. A group of 25 control rats received an implant of glass of similar size. After 300 days, 30 and 20 animals were still alive in the two groups, respectively; all surviving rats were killed 800 days after implantation. One sarcoma and one fibroma were found after 580 days in the PVC-treated rats, whereas no local tumours developed in the control group (Russell *et al.*, 1959).

A group of 80 outbred albino rats (sex unspecified) received implants (of unstated size) of PVC film by laparotomy to surround the kidney. The animals were sacrificed at 3, 10, 15, 30, 90, 195, 285, 300 and 380 days after the implantation. Of rats that survived 285-375 days, 6/16 developed fibrosarcomas at the site of implantation (Raikhlin & Kogan, 1961).

In rats implanted in the kidney with either PVC capsules, whole PVC films or perforated PVC films, 5/16, 2/5 and 1/5 sarcomas were observed (Kogan & Tugarinova, 1959).

¹The Working Group was aware of a study in progress to assess the carcinogenicity of PVC powder by inhalation exposure and intrapleural administration to rats (IARC, 1978a).

3.2 Other relevant biological data

(a) Experimental systems

Rats and guinea-pigs exposed continuously to PVC dust for 24 hours/day for periods varying from 2-7 months were found to have extensive lung damage (Frangia *et al.*, 1974).

The growth of granulation tissue around PVC particles implanted into the muscles of rabbits has been reported (Guess & Stetson, 1968). Severe fibroblastic reactions were found in rats (Calnan, 1963) and dogs (Harrison *et al.*, 1957) which received PVC sponge implants into the s.c. tissue of the anterior abdominal wall.

Ingested and rectally absorbed PVC particles (5-110 μ m) were found to be transported by both the lymphatic and the portal system from the intestinal wall of rats, guinea-pigs, rabbits, chickens, dogs and pigs (Volkheimer, 1975).

In rats, inhalation of fumes from heated PVC produced interstitial oedema, as well as focal bronchial and intra-alveolar haemorrhage in the lungs of some animals (Cornish & Abar, 1969).

No data on the embryotoxicity, teratogenicity, metabolism or mutagenicity of this compound were available to the Working Group.

(b) Humans

Workers exposed to PVC dust during the manufacture of articles made from PVC showed alterations in the respiratory organs (e.g., changed bronchovascular pattern, increased pulmonary ventilation at rest) (Vertkin & Mamontov, 1970).

Pneumoconiosis due to inhalation of PVC dust was suggested following pulmonary biopsies in a male patient who had inhaled PVC dust for 1 year and exhibited granulomatous lesions due to foreign bodies. The severity of the disease was shown by the observed dyspnoea, secondary polyglobulia and reduced respiratory function (Szende *et al.*, 1970).

Fibrotic lung changes and altered pulmonary function tests have been reported in 96 workers exposed to PVC dust; the changes were more pronounced in those with long exposure (Lilis *et al.*, 1976; Waxweiler *et al.*, 1976).

Reduced pulmonary function and an enhancement of pulmonary function defects associated with an increased risk of respiratory impairment were noted in a number of nonsmoking workers exposed to an occupational environment contaminated with vinyl chloride fumes and PVC dust (Miller, 1975; Miller *et al.*, 1975).

In
tries',
mixed wi
reported
bromosul
in 1, sl
Raynaud'
of abnor
of 37 PV
abnormal
transami
abnormal

Cas
wrappers
sealed a
has not
Polakoff

Dur
distress
combusti
chloric
been rep
tance, a
thermal

The
be cause
1953).

3.3 Case

A pr
cates fro
cutting,
cating).
was found
[The stud
by the fa
vities d:

¹The
exposed t

In 15 polyvinyl production workers employed in 'PVC-processing industries', where stabilizers, colours and bulk materials were added to and mixed with the basic PVC powder, the following pathological findings were reported: in 7, slight to moderate thrombocytopenia; in 7, increased bromosulphalein retention; in 6, reticulocytosis; in 1, leucopenia; and in 1, slight splenomegaly. Neither scleroderma-like skin changes nor Raynaud's syndrome were observed (Lange *et al.*, 1975). A large proportion of abnormal liver function tests and platelet counts were found in a group of 37 PVC process workers: 20% had abnormal alkaline phosphatase, 60% abnormal lactic dehydrogenase, 30% abnormal serum glutamic-oxaloacetic transaminase, 10% abnormal serum glutamic-pyruvic transaminase, and 35% abnormal platelets (Wegman, 1975).

Cases of 'meat-wrappers' asthma have been reported, in which meat wrappers developed respiratory symptoms when exposed to fumes of PVC film sealed and cut with a hot wire. The identity of the causative agent(s) has not been established (Brooks & Vandervort, 1977; Falk & Portnoy, 1976; Polakoff *et al.*, 1975; Sokol *et al.*, 1973).

During the period 1970-1975, 175 fire-fighters experienced respiratory distress due to the toxicity of hydrogen chloride gas released from the combustion of PVC plastics (Dyer & Esch, 1976). Carbon monoxide, hydrochloric acid and phosgene (Cornish & Abar, 1969; Dyer & Esch, 1976) have been reported as the major PVC pyrolysis products of toxicological importance, although more than 75 components have been identified following the thermal degradation of PVC (Dyer & Esch, 1976).

The few cases of dermatitis that have been reported are believed to be caused by sensitivity to plasticizers in polyvinyl plastics (Morris, 1953).

3.3 Case reports and epidemiological studies¹

A proportional mortality study was carried out using death certificates from 1970-1972 of 707 male plastic workers (extruding, moulding, cutting, turning or otherwise machining plastics, and including PVC fabricating). A statistically significant excess of stomach cancer mortality was found (24 observed *versus* 16.4 expected, $P < 0.05$) (Baxter & Fox, 1976) [The study is limited by the use of proportionate mortality methodology and by the fact that not all deaths studied were among workers engaged in activities directly involving PVC].

¹The Working Group was aware of a study in progress on workers exposed to PVC (IARC, 1978c).

In a cross-sectional mortality study of 4341 deaths during 1964-1973 among current and former employees of 17 PVC fabricators, an excess in total cancer mortality, particularly that of the digestive system, was reported among both white males and white females. The risk of cancer of the breast and urinary organs was also reported to be in excess among white females (Chiazze *et al.*, 1977) [The study is limited by the use of proportionate mortality methodology and by the fact that not all deaths were among workers engaged in activities directly involving PVC].

Casterline *et al.* (1977) reported a case of a 22-year-old male who developed a squamous-cell carcinoma of the buccal mucosa as a result of a habit, acquired at the age of 8 years, of chewing plastic materials containing PVC. No prior history of mouth or lip lesions, of smoking tobacco or drinking alcohol, or of occupational exposure to vinyl chloride was noted.

Vinyl chloride-vinyl acetate copolymers

1. Chemical and Physical Data

1.1 Synonyms and trade names

Chem. Abstr. Services Reg. No.: 9003-22-9

Chem. Abstr. Name: Acetic acid ethenyl ester polymer with chloroethene

Acetic acid vinyl ester polymer with chloroethylene; chloroethylene-vinyl acetate polymer; polyvinyl chloride-polyvinyl acetate; vinyl acetate-vinyl chloride copolymer; vinyl acetate-vinyl chloride polymer; vinyl chloride-vinyl acetate polymer

A 15; A 15 (polymer); A 15-0; A 15S; Bakelite LP 70; Bakelite VLFV; Bakelite VMCC; Bakelite VSJD 10; Bakelite VYHD; Bakelite VYHH; Bakelite VYNS; Bakelite VYNW; Breon 351; Breon 425; Breon AS 60/41; Corvic 51/83; Corvic 236581; Corvic R 46/88; Denkalac 41M; Denka Vinyl MM 90; Diamond Shamrock 744; Diamond Shamrock 7401; Exon 450; Exon 454; Exon 470; Exon 481; Exon 760; Flovic; Geon 100x150; Geon 130x10; Geon 135; Geon 351; Geon 400x47; Geon 421; Geon 427; Geon 434; Geon 103EP-J; Geon 440L2; Geon 450x150PN; Geon 150XML; Geon 103ZX; Hostalit PVP; Leucovyl PA 1302; Lucovyl GA 8502; Lucovyl MA 6028; Lucovyl PA 1208; Marvinol VP 56; 50ME;

Nor
PVC
AX
PA
VAG
H 4
VYF
Vin

1.2 Str

1.3 Chem

(a)

(b)


1.4 Tech

Low
to obtain
vinyl ace
described
ves are u

No d
in the co

Norvinyl P 6; Opalon 400; Pevikon C 870; Pliovac AO; Pliovic AO; PVC Cordo; Resin 4301; Rhodopas 6000; Rhodopas AX; Rhodopas AX 30/10; Rhodopas AX 85/15; Sarpifan HP 1; Solvic 523KC; Solvic PA 513; Solvic 513PB; Sumilit PCX; Tennus 0565; VA 3 (copolymer); VAGD; VH 10/60; Vilit 40; Vinnol H 10/60; Vinnol H 15/45; Vinnol H 40/60; Vinylite VGHH; Vinylite VYDR; Vinylite VYDR 21; Vinylite VYFS; Vinylite VYHD; Vinylite VYHH; Vinylite VYNS; Vinylite VYNW; Vinyon; VLVF; VMCC; VYHH; VYNS; VYNW

1.2 Structural and molecular formulae and molecular weight



1.3 Chemical and physical properties of the copolymers

- (a) Description: White powder
- (b) Stability: Sensitive to excessive exposure to heat and light in the absence of added stabilizers. Hydrogen chloride gas is a decomposition product of degradation.

1.4 Technical products and impurities

Low levels of vinyl acetate are copolymerized with vinyl chloride to obtain resins with specific properties. Depending upon the use, the vinyl acetate level may vary from 2-20%, with an average of 11-12%. As described in the monograph on polyvinyl chloride, p. 405, various additives are used to aid in processing the resins.

No detailed information on the possible presence of unreacted monomers in the copolymers was available to the Working Group.