

DATE

March 4, 1969

cc R. E. Keller - Res. 1
E. P. Wheeler - GO

SUBJECT

AROCOR & WILDLIFE:
INCINERATION OF NCR PAPER

REFERENCE


TO

W. R. Richard - GO

It has been proposed from the observed distribution of polychlorinated biphenyls in tissue samples taken from marine organisms that global eco-distribution of these materials occurs via aerial transport (wind currents, airborne particulate matter, etc.).

A qualitative experiment has been performed with the objective of determining if Aroclor in N.C.R. paper is destroyed or volatilized when the paper is disposed of by burning.

The experiment was carried out in the following manner. Approximately 3g. (1 sheet) of paper (CB 15# white, 8-1/2 X 11, long grain, F-3468-4-24-64, The N.C.R. Company) was placed in the incinerator chamber of the apparatus shown in Figure I.



At this point the vacuum was turned on and the paper ignited. After the paper had completely burned, the hexane in each trap, and hexane extracts of the ash and the walls of the incinerator chamber were analyzed for the presence of Aroclor by GC/GC. Hexane samples from trap #1 were also analyzed after extraction with base and acid to remove any oxidized materials.

0631269

PCB-ARCH0745709

W. R. Richard
March 4, 1969
Page 2

Conclusions:

Under the conditions of this experiment:


1. Aroclor is easily volatilized when N.C.R. paper is burned.
2. Aroclor undergoes little, if any, decomposition when burned.
3. Unfortunately, it appears that significant air pollution can occur via burning of N.C.R. paper or other Aroclor containing materials even under more strenuous conditions.

E. S. Tucker

db

0631270

PCB-ARCH0745710



Material
Trapped
in Hexane

Neolon 1242
Standard

Material
Extracted
from Residue

Blank

FIG. 2

0631271




Fig 3

0631272




FIG. 1

0631273