

THE TOXICOLOGY OF PCB'S

**An Overview with Emphasis on Human Health
Effects and Occupational Exposures**

**Prepared by the
Hazard Evaluation System**

**Epidemiological Studies Section
State of California
Department of Health Services/Department of Industrial Relations
2151 Berkeley Way
Berkeley, California 94704
(415) 540-2115**

January, 1981

MONS 010655

THE TOXICOLOGY OF PCB'S

Table of Contents

I.	INTRODUCTION	1
II.	GENERAL BACKGROUND INFORMATION	3
III.	PHARMACOKINETICS	5
	Absorption	5
	Distribution, Accumulation (Mammals)	5
	Transplacental Exposure, Secretion in Milk	6
	Metabolism	7
IV.	ANIMAL TOXICOLOGY	10
	Acute	10
	Sub-acute, Chronic	11
	Reproductive Effects	12
	Other - Immunosuppressive, Endocrine	14
V.	CARCINOGENICITY/MUTAGENICITY	16
	Carcinogenicity.	16
	Test Results	16
	Mutagenicity	19
VI.	BIOCHEMICAL EFFECTS.	21
	Enzyme Induction	21
	Porphyria	22
VII.	HUMAN TOXICOLOGY AND EPIDEMIOLOGY	24
	Dermatologic Effects	24
	Systemic Symptoms	25
	Liver Damage	25
	Yusho	25
	Neurotoxicity	27
	Cancer	27
	Ongoing Occupational Studies	28
VIII.	MEDICAL SURVEILLANCE AND BIOLOGIC MONITORING	31
IX.	SUMMARY AND CONCLUSIONS	33

MQNS 010656

Table of Contents

REFERENCES	34
Table I. Dose-Response for Animal Toxicology	39
Table II. Occupational Exposure to PCBs.	40
Table III. Percent Distribution of Symptoms of Yusho Reported by 189 Patients Examined Before October 3, 1968	44
Figure I.	45
Figure II.	46
Figure III.	47
Figure IV.	48

MONS 010657

L INTRODUCTION

As a consequence of the EPA ban on further manufacture of PCB's^{*} in 1977, occupational exposures to these compounds have been drastically reduced. However, significant exposures may remain for particular occupational groups. Utility workers, for example, may experience sporadic but potentially massive exposures when cleaning up spills, or when servicing and dismantling transformers and capacitors that still contain PCB fluid. Electricians, appliance service workers and firefighters also may have continued occupational exposure. NIOSH estimates that 12,000 workers have potential exposure as a result of current uses of PCB's (NIOSH, 1977). Despite the vast scientific literature on the toxicology of PCB'S, the human health effects likely to result from such exposure remain ill-defined.

The Hazard Evaluation System has reviewed the literature on PCB toxicology in response to inquiries about worker health. Requests for information have come from unions and workers who handle PCB fluids in clean-up of spills, in maintenance work, and in transportation, storage and disposal of used equipment. Toxicology information has also been requested by medical professionals evaluating the clinical significance of PCB exposures, and public health officials who are attempting to set standards for occupational and environmental exposures. Our primary goal has been to review the data relevant to the human health effects of PCB's especially those resulting from occupational exposures. Since the published epidemiologic evidence is limited we have utilized animal toxicology studies where appropriate in anticipating potential biologic effects in humans. We have not attempted to summarize the extensive literature on PCB toxicology, but the reader is referred to a number of recent reviews (DHEW, 1978; IARC, 1978; Fishbein, 1974; Kimbrough, 1974; EPA, 1977; Nelson, 1972; NIOSH,

(1977).

• PCBs: Polychlorinated Biphenyls

MONS 010659

II. GENERAL BACKGROUND INFORMATION

In all animal species that have been studied PCB's have a very low acute toxicity. They are readily absorbed across biological membranes, poorly metabolized and only very slowly eliminated. Because PCB's persist in the environment and accumulate in living tissue, they are concentrated ("biomagnified") in the food chain in a similar manner to other organochlorine compounds like DDT. Concern about exposure to PCB's has, therefore, focused on their resistance to biodegradation with the consequent potential for long-term or delayed health effects.

A number of published reports have established "background" levels of PCB's in the blood and tissues of human populations with no previous history of exposure. Surveys in various geographical areas have found detectable residues in blood, fat and mothers' milk. Measurable levels of PCB's are typically found in greater than 50% of subjects tested with maximum blood levels generally less than 20 ppb (Finklea, 1972). The levels reported from adipose tissue are typically somewhat higher, in the range of 1-2 ppm (Kutz, 1975). Residues of PCB's in human milk have ranged from 40-100 ppb in whole milk (New York State Health Council, 1977).

Two facts complicate the documentation of the human and animal toxicology of PCB's:

1. Commercial products are rarely single agents, but rather are complex mixtures of chlorinated biphenyls with different numbers and arrangements of attached chlorine atoms (see Figure I). The metabolism and toxicology of PCB's seem to vary with the percent of chlorination and with the isomeric structure of the PCB molecule.
2. All commercial products are potentially contaminated with chlorinated naphthalenes and polychlorinated dibenzofurans (PCDF's). The degree of this contamination varies with different commercial mixtures (see Figure III).

MONS 010660

Contamination by dibenzofurans (PCDF's) is of particular concern because of the structural similarity of these compounds to the highly toxic dibenzodioxins (see Figure II). The pattern of observed effects in animals exposed to PCDF's closely resembles that seen following exposure to 2,3,7,8 tetrachlorodibenzodioxin (TCDD). In comparative animal studies the toxicity of PCDF's is much greater than the PCB's, particularly in the thymus, skin (acne), liver and hematopoietic system (Oishi et al., 1978; Moore et al., 1979). In addition, PCDF's are 1000 times more potent than PCB's as enzyme inducers (see Section VI).

Uncertainties in the analytic methods used for detection of PCB's must be considered when reviewing the published data on PCB toxicology. Monitoring PCB's in environmental or biological samples by gas-liquid chromatography/mass spectrometry (GC/MS) is made difficult by the presence of other chlorinated hydrocarbons (e.g., pesticides) which are commonly present at similar concentrations (Stalling et al., 1979). Because of the difficulties in the interpretation of GC/MS spectra, the PCB levels reported from different laboratories may show considerable variation.

MONS 010661

III. PHARMACOKINETICS

Absorption

There is relatively little information on the rate or degree of absorption of PCB's by any route for any species of animal. Since similar systemic toxicity has been observed in rodents after dermal, oral and inhalational administration of comparable doses, it is likely that PCB's are easily absorbed by all routes. The few quantitative measurements of relative absorption rates indicate that most, if not all, PCB's which contain six or fewer chlorine atoms are efficiently absorbed from the GI tract (Albro and Fishbein, 1972; Van Miller et al., 1973).

Distribution, Accumulation (Mammals)

As with other heavily chlorinated chemicals the major storage tissue for PCB's is body fat. The concentration in adipose tissue is 10-1000 times that found in other tissues, both following single oral doses (Grant et al., 1971) and after chronic administration. (Curley et al., 1971) The lowest concentrations are found in whole blood and plasma where levels are usually several fold lower than in other tissues examined. This preferential distribution of PCB's into fat has been well documented after i.v. dosing in the rat (Lutz et al., 1977). Results are consistent with the high distribution coefficient of PCB's in fat and low perfusion of adipose tissue compared to skin, liver, muscle and blood (Anderson et al., 1977).

Two important pharmacokinetic questions cannot now be resolved on the basis of available data:

MONS 010662

1. Does the concentration of PCB's reach steady state with constant exposure?
and
2. Will mobilization of adipose tissue after starvation or illness lead to a transient increase in PCB concentrations in blood and other tissues ?

Like most heavily chlorinated hydrocarbons the half-life of PCB's in animal tissue is quite long. In a chronic feeding study with Aroclor 1245 at 100 ppm in the diet of rats, a steady build-up of PCB's occurred in all tissues analyzed without a plateau level even after 240 days of treatment. By comparison, in a similar rat dietary study using DDT, plateaus in fat were attained after 90-140 days (Curley et al., 1971). In dairy cows a steady state was reached in 40-60 days (Fries, 1972), probably due to mobilization into fat micelles and secretion into milk. Thus, PCBs are unlikely to reach steady state levels in non-lactating animals; and fat mobilization or lactation may be expected to result in release of stored PCBs.

Transplacental Exposure, Secretion in Milk

Transplacental exposure of PCB's has been documented in mammals. Term fetuses taken from rats exposed to 10 mg/kg/day during days 7-15 gestation contained 0.63 ppm PCB, or about 1/60 the maternal dose. When the dose to the mother was increased five fold, the concentration in the fetuses increased two fold (Curley et al., 1973). The hyperpigmented babies observed in the Yusho incident (see Section VII) represent additional circumstantial evidence of transplacental passage of PCB's.

Secretion of PCB's into milk has also been observed. In mice, little passage of PCB's occurred across the placenta once PCB's had been sequestered into maternal adipose tissue, but they were readily transferred to suckling offspring through the milk (Vodichnik and Lech, 1980). These observations suggest that secretion of PCB's into milk may be quantitatively much more important as a source of exposure in newborns than is

MONS 010663

transplacental passage. This has recently been documented in a prospective study of Japanese mothers and their infants. (Hirokadzu and Ota, 1980)

Metabolism

PCB's are metabolized primarily by hydroxylation and conjugation with glucuronic acid. The primary site of biotransformation is assumed to be the liver, although no data is available currently on the possible role of peripheral metabolism, e.g., skin.

Many experimental feeding studies in both mammals and birds have shown an inverse relationship between percent chlorination and rates of metabolism. Less chlorinated PCB's are more readily metabolized than are more chlorinated ones, the rate of metabolism and excretion decreasing sharply as the number of chlorine atoms increases above five (EPA, 1977). The metabolism of the higher chlorinated biphenyls is also dependent on the position of chlorine atom substitution. The presence of two adjacent, unsubstituted carbon atoms is needed for the rapid enzymatic hydroxylation reaction (Jensen and Sundstrom, 1974). Since the more highly chlorinated biphenyls have a very much slower metabolic rate and longer half-life, they are more generally found as residues in human and animal tissues. This relationship alters with PCB's above 54% chlorination, presumably as a result of lower absorption from the gastrointestinal tract.

Arene oxide intermediates have been described in a major pathway of the metabolic transformation of PCB's by hepatic mixed function oxidases (Safe et al., 1975; Gardner et al., 1973). These intermediates are of particular concern since they are capable of direct interaction with DNA and may be the active form of carcinogenic polyacyclic hydrocarbons. (Jerina and Daly, 1974) The PCB molecules which are more readily metabolized and excreted also are more likely to form these arene oxides. It does not necessarily follow, however, that those compounds which persist in tissue and are more

likely to be measured in population sampling are less important in terms of their carcinogenic potential.

From the limited data available, it appears that significant differences exist between non-human primates and rodents in the metabolism and pharmacokinetics of PCB's. The marked variation observed in PCB toxicity between rodents and primates may be explained by such differences. Primates appear to be more susceptible to the toxic effects of PCB's than are rats or mice (see Section IV and Figure IV). When single doses of radiolabeled PCB were administered by gastric intubation to infant monkeys, the metabolites measured in urine, tissue (liver) and serum included hydroxylation products derived from arene oxide intermediates (Allen et al., 1976); while in the rat, direct hydroxylation is the rule (Hsu et al., 1975).

There is virtually no pharmacokinetic data in humans. A few generalizations can be made, however, based on studies reporting PCB blood levels: The higher the exposure levels, the higher blood concentration of PCB's (Hara et al., 1975; Inoue et al., 1975; Karppanen and Lotho, 1973; Baker et al., 1980); and the higher the environmental concentration and/or the longer the period of exposure, the longer the blood levels of PCB's remain elevated (Hara et al., 1975; Baker et al., 1980). However, there are a few reports which are inconsistent with this latter trend (Bumgarner et al., 1975; Hasegawa et al., 1972; Kitamura et al., 1973). PCB levels have also been correlated with race and geographic residence (NIOSH, 1977) and with age and dietary intake of fish (Kimbrough, 1980).

Comments

Certain generalizations can be made from the limited pharmacokinetic data that is available:

MONS 010665

1. Absorption occurs by all routes (skin, GI, inhalation).
2. Distribution is primarily into fat.
3. Metabolism and excretion are dependent on specific molecular structure, varying inversely with percent chlorination.
4. Excretion is in general quite slow so that bio-accumulation occurs even at low exposure levels.
5. Transplacental transfer occurs but may be quantitatively less significant than secretion into milk.
6. Arene oxide metabolites are found in the metabolic transformation of PCBs. These compounds are highly reactive and may represent the active carcinogens. (see Section V)
7. The relationship between percent chlorination and potency as carcinogen has not been established.
8. There are essentially no pharmacokinetic data in humans; it is not known, for example, if intermittent high doses are more or less hazardous than low level chronic exposures to the same total dose.

MONS 010666

IV. ANIMAL TOXICOLOGY

Acute Toxicity

When given as a single dose, the acute oral LD₅₀ of PCB's in rats, rabbits and mice ranges from 1-10 grams/kg of body weight. According to the American Industrial Hygiene Association classification system for acute toxicity, PCB's are classified as "slightly toxic" (0.5 - 5 g/kg), or "practically non-toxic" (5-15 g/kg). There is some evidence that young animals are more sensitive than adults, and that females are more susceptible than males to the acute effects of PCB's (Kimbrough et al., 1978). In rodents, the acute oral toxicity appears to decrease with increasing chlorine content of the administered PCB's. This may be secondary to decreased absorption of the higher chlorinated compounds or to the differences in metabolic transformation previously discussed.

Although few clinical signs of toxicity have been reported in experimental animals, pathologic findings are extensively documented. CNS depression (decreased pain response and diminished exploratory behavior), anorexia and oliguria followed by ataxia, coma and death have been observed in rats following acute administration of large doses of PCB's (Brackner et al., 1973). Consistent pathologic findings associated with death in rats, rabbits and guinea pigs include liver damage with fatty infiltration, centrilobular atrophy, and in some cases necrosis. Pathologic changes in other organs in these species are not often described, except for chloracne-like lesions which occur at the site of skin or intradermal application.

Comments

The low order of acute toxicity in experimental animals is consistent with the lack

of acute effects observed in workers exposed to PCB's. Reported symptoms after occupational exposures include mild irritation of the skin and eyes at levels above 0.1 mg/m³ with unbearable irritation occurring above 10 mg/m³ (ACGIH, 1976). Systemic symptoms of nausea and headache have been reported but may be secondary to the solvents (such as trichlorobenzene) in the PCB mixtures.

Sub-acute and Chronic Toxicity

In contrast to the low order of acute toxicity, effects from chronic exposures to relatively low doses of PCBs have been consistently observed and are of far greater concern. These sub-acute effects show appreciable variation among species, but liver damage is again the prominent finding.

The major changes in rats fed Aroclor 1248, 1254, and 1262 at 100 ppm in their diet for six weeks included liver hypertrophy, marked fatty infiltration and degeneration of parenchymal cells. As in acute toxicity studies PCB mixtures with lower chlorine content were more toxic (Allen and Abrahamson, 1973). In 8-12 month feeding studies increased serum lipids and focal areas of liver damage were observed (Allen et al., 1976; Kimbrough et al., 1972).

Non-human primates are more sensitive than rodents to the toxic effects of PCB's (see Table 1 and Figure IV). Adult female monkeys exposed to dietary levels of 2.5 ppm for 12 months (=0.08 mg/kg/day) developed facial edema, alopecia, acne, gastritis with ulceration, anemia, hypoproteinemia and bone marrow hypoplasia. At 100 ppm (=10 mg/kg/day) there was considerably more evidence of tissue damage than in rats, including marked hepatic hypertrophy with ultrastructural abnormalities (Allen, 1975). Based on extrapolation from the Yusho data, PCB's may cause symptoms to humans at levels (0.2 mg/kg/day) which are comparable to the lowest doses which produce effects in non-human primates (see Table 1).

MONS 010668

Most of the animal data are derived from feeding or oral intubation studies. There are relatively few reports of dermal or inhalation experiments. Inhalation studies again revealed liver damage to be the prominent finding in rodents. For summary of inhalation data see the NIOSH criteria document, page 125 (NIOSH, 1977). Dermal toxicity studies in rabbits have produced skin lesions at the site of application as well as systemic effects including liver and kidney damage, thymic atrophy, lymphopenia and increased fecal porphyrins (Vos and Beems, 1971).

Comments

Liver damage, documented histologically, is the consistent finding among various laboratory animals exposed to low chronic levels of PCB's. It has not been observed in the limited surveys of exposed workers. This may be an artifact, however, of the relative insensitivity of the standard liver function tests (such as serum levels of SGOT, SGPT) as compared to biopsy and histologic analysis.

Reproductive Effects

Adverse reproductive effects of PCB's have been noted in many mammalian and avian species. The pattern of reproductive effects include alterations in estrus cycles, failure of implantation, increased frequency of spontaneous abortions, low birth weight offspring and decreased post-natal survival. No specific teratogenic effects of PCB's have been observed in a variety of avian species. Transplacental effects, however, have been documented in both animals and humans (see Section VII).

PCB's given to mice for 10 weeks at dosage of 1.0 mg/kg/day lengthened the estrus cycle by more than two days and decreased the number of successfully implanted ova (Orberg and Kihlstrom, 1973). Similarly, mice that received PCB's as sucklings in a

MONS 010669

long-term transgenerational study showed subsequent alterations in estrus cycles, decrease in implantations, and when mated to each other (F1 studies), reduced number of offspring per litter (Kihlstrom et al., 1975).

In rats, studies suggest that reproductive effects of PCB's decrease as chlorination increases. No reproductive effects have been found with Aroclor 1260 (60 percent chlorination) at 1, 10, 100 ppm, but significant effects have been noted with Aroclor 1242 and 1254 (42 and 54 percent chlorination, respectively) at doses of 20 and 100 ppm. Aroclor 1260 began to exert toxic effects at doses of 500 ppm. Rats chronically fed from 20 to 100 ppm Aroclor 1242 and 1254 had reduced numbers of offspring. Surviving newborns showed increased mortality, with only 30 percent surviving to weaning. Five ppm of either Aroclor 1242 or 1254 produced no effects over two generations. Thus, the minimum effective doses ranged from 20 ppm for the lower chlorination mixtures to 100-500 ppm of the more highly chlorinated compounds. (Keplinger et al., 1971; Linder, et al., 1974)

Evidence of adverse reproductive effects is also available for non-human primates. Rhesus monkeys fed 2.5 and 5.0 ppm Aroclor 1248 for 18 months in the diet showed changes in menstrual cycles in addition to other systemic signs of toxicity. Evidence was also obtained for frequent resorptions and spontaneous abortions following breeding to normal males. In all, six infants were carried successfully to term out of 14 pregnancies. The offspring were of low birth weight and by two months began to show evidence of PCB toxicity, presumably from PCB's in the maternal milk; only three infants survived to six months. Behavioral tests in the three surviving animals showed marked deficits in several learning tasks, with increasing errors correlated with increasing body burdens of PCB's (Allen and Barsotti, 1976; Bowman et al., 1978; Barsotti et al., 1976).

MONS 010670

The effect of PCB's on the male reproductive system is not known. There is one report of four male Rhesus monkeys exposed to 5.0 ppm Aroclor 1248 in the diet for 18 months. After 12 months, one of four animals developed clinical signs of PCB intoxication, showed marked sperm count depression and was functionally sterile. A testicular biopsy revealed an absence of spermatogonia. A second biopsy one year after exposure showed complete recovery (Allen et al., 1979). PCB's are negative in the mouse sperm morphology assay (Heddlie and Bruce, 1977).

Comments

PCB's show significant effects on reproductive competence in a variety of species. These effects increase in intensity with increasing dosage and decrease with increasing chlorination of the PCB isomers. PCB's do not appear to be mammalian teratogens. A reasonable explanation for most of the reproductive effects of PCB's could be based on their estrogenic activity (see below).

PCB's have been detected in human semen (Dougherty et al., 1980), but there have been no studies of semen quality in relation to PCB exposure in humans. The effects of PCBs on the male reproductive system in animals or humans has not been adequately studied. The only other evidence to date on reproductive toxicity in humans come from the Yusho incident and is summarized in Section VII.

Other

1. Immunosuppressive Effects

A number of reports implicate PCB's as immunosuppressants (Fishbein, 1974). Lymphoid atrophy has been observed in rabbits, chickens and guinea pigs. Suppression of humoral immune responses to several antigens was observed in rabbits and guinea pigs, and decreased cell-mediated immune response

MONS 010671

followed PCB exposure in guinea pigs. A decreased tolerance to hepatitis virus was seen in ducklings without apparent intoxication. In monkeys exposed transplacentally and through contaminated milk, the lymph nodules of the spleen were extremely small and without germinal centers (Allen and Barsotti, 1976); morphologic changes were indicative of reduced immunologic competence.

2. Endocrine Effects

Subcutaneous administration of Aroclor compounds with lower chlorination produced an estrogenic effect on the rat uterus which was not shown with Aroclors of higher chlorination (Bitman and Ceal, 1970). Female primates fed Aroclor 1248 for six months showed an increase in concentration of urinary ketosteroids and a prolongation of their menstrual cycles with increased bleeding (Barsotti et al., 1976). Antiandrogenic effects have been described in birds although the mechanism is not clear. It may be secondary to an increased rate of androgen metabolism in the liver by induction of microsomal enzymes (see Section VI), or by virtue of PCB's exerting estrogenic effects.

Comments

The effect of PCB exposure on immune and endocrine system function has not been carefully studied in humans, so the relevance of these animal observations to human health remains unknown. There is one cross-sectional study of occupational exposure to PCB's which will include analysis of serum hormone levels and urinary metabolites, but results have not yet been published (Selikoff et al., in progress).

MONS 010672

V. CARCINOGENICITY/MUTAGENICITY

Carcinogenicity

Several PCB mixtures are clearly carcinogenic in rodent bioassays, producing liver tumors (hepatocellular carcinomas). Kanechlor 300 and Aroclor 1254 are carcinogenic in male mice (Ito et al., 1973; Kimbrough and Linder, 1974), and Aroclor 1260 is carcinogenic in separate studies in two strains of female rats (Kimbrough et al., 1975; Norback et al., 1980). In addition, a purified component of a PCB mixture, 2,4,5,2',4',5'-hexachlorobiphenyl, has recently been found to be carcinogenic in female rats, causing hepatocellular carcinomas (Norback et al., 1980).

*hepatocellular
carcinomas
in
liver
nodules*

Because high doses of PCB's are known to cause extensive injury to liver tissue it is important to consider the dose levels at which liver carcinomas were produced in the rodent bioassays. In two studies in rats, significant increases in hepatocellular carcinomas were present at doses which did not produce gross histologic changes. Hepatocytes were somewhat enlarged (probably due to microsomal enzyme induction), but no extensive fatty infiltration or necrosis occurred, as was characteristic of bioassays at higher dose levels (Kimbrough et al., 1975).

Test Results

1. Mice (Male)

- A. Kanechlor 300, 400, and 500 fed to groups of 12 eight-week-old male mice at 100, 250, and 500 ppm in the diet for 32 weeks produced hepatocellular carcinomas in 5 of 12 survivors in the high dose group fed Kanechlor 500. The remaining 7 mice in this group had nodular

MONS 010673

hyperplasia (neoplastic nodules). No metastases or other tumors were present in this or other dosed groups. The control group (6 mice) was likewise tumor-free (Ito et al., 1973).

- B. Aroclor 1254 administered to groups of 50 five to six-week-old male BALBc/J mice at dietary levels of 0 or 300 ppm (about 30 mg/kg body weight during the exposure) for 11 months produced neoplastic nodules (hepatomas or hyperplastic nodules) in 9 of 22 survivors in the dosed group. Other liver lesions (adenofibrosis) were present in all 22 survivors. Additional morphological changes in the livers of these animals included pleomorphism and areas of necrosis. Such changes and tumors were absent among survivors (24) in the control group (Kimbrough and Linder, 1974).

- A. Kanechlor 400 administered to ten-week-old Donryu rats (10 males and 10 females) at dietary levels which varied from 40-600 ppm during the 400-day study produced liver tumors (multiple adenomatous nodules) in 6/10 treated female rats. Such lesions were absent from the controls (5 males and 5 females) and the treated males (Kimura and Baba, 1973).

- B. Kanechlor 300, 400, or 500 administered to groups of 30 eight-week-old male Wistar rats at dietary levels of 0, 100, 500, or 1000 ppm produced increases in the incidence of cholangiofibrosis at the highest dose level of all Kanechlors (2/15, 2/10, and 4/13, respectively). All three compounds also produced hepatic nodular hyperplasia, the incidence of which increased with dose and extent of chlorination (Kanechlor 300 at 100 ppm: 1/22; Kanechlor 400 at 100 ppm: 2/16, and 1000 ppm: 3/10; Kanechlor 500 at 100 ppm: 3/25, at 500 ppm: 5/16, and at 1000 ppm: 5/13); (Ito et al., 1974).

MONS 010674

- C. Aroclor 1260 administered to groups of 200 three to four-week-old female Sherman rats at 0 and 100 ppm in the diet (varying between 5-10 mg/kg body weight during the 21 month exposure) produced at 23 months among the dosed survivors clearly significant increases of hepatocellular carcinomas (controls 0/173; dosed group 146/184) as well as neoplastic nodules (hyperplastic nodules: controls 0/173; dosed group 26/184). The incidences of non-hepatic tumors did not differ between the dosed and control groups (Kimbrough et al., 1973).
- D. Aroclor 1254 administered to groups of 24 eight-week-old Fisher 344 rats of either sex at dietary levels of 0, 25, 50, or 100 ppm for 105 weeks was not carcinogenic to any of the treated groups under the test conditions. It is important to note that two of the dose levels used were lower than those which produced a positive response in Sherman rats. Rare adenocarcinomas and carcinomas of the gastrointestinal tract appeared in both sexes and may be related to the administration of the PCBs (males: historical controls 6/600, dosed group 2/24). In addition a high incidence of non-neoplastic liver hyperplasia was present among the dosed groups (males: controls 0/24, low-dose 5/24, mid-dose 8/24, high-dose 12/24; females: controls 0/23, low-dose 6/24, mid-dose 9/22, and high-dose 17/24); (NCI 1978).
- E. Aroclor 1260 administered to groups of 50 male and female Sprague-Dawley rats at dietary levels of 0 and 100 ppm for 105 weeks was carcinogenic in female rats, causing significant increases in liver hepatocellular carcinomas (Norback and Weltman, 1980).

MONS 010675

F. A purified component of a PCB mixture, 2,4,3,2',4',5-hexachlorobiphenyl administered to groups of 50 male and female Sprague Dawley rats at dietary levels of 0 and 100 ppm for 105 weeks was carcinogenic in female rats, producing an increased incidence of liver hepatocellular carcinomas among the dosed animals (Norback and Weltman, 1980).

Mutagenicity

PCB mixtures have not been observed to have mutagenic activity nor to measurably affect chromosomes in repeated studies using a variety of in vitro or in vivo test systems. Evidence of genetic damage from PCB's in laboratory test systems including chromosomal aberrations, non-disjunction, loss of sex chromosomes or increased frequency of sister chromatid exchange has not been observed. Report of a weak effect of Aroclor 1221 and of a stronger effect of 4-chlorobiphenyl in Salmonella using PCB-induced rabbit liver homogenate as a liver activation system appears unfounded (Wyndham et al., 1976). Further attempts to repeat these results have been unsuccessful using a variety of Salmonella tester strains and liver activation systems (Katzenellenbogen and Ames, 1980; Safe, 1978).

However, PCB's belong to the class of heavily chlorinated animal carcinogens, most of which are not positive in short-term tests for mutagenicity. Examples in this class include dieldrin, chlordane, kepone, mirex, TCDD,* chloroform, and carbon tetrachloride. Whether this is because the in vitro metabolic activation systems do not produce the same spectrum of metabolites that occur in vivo or because heavily chlorinated compounds such as PCB's are carcinogenic by non-mutagenic mechanisms is not known at this time.

* TCDD: Tetrachlorodibenzodioxin

Validation of the carcinogenic effects in rodents is provided by a positive cell transformation assay using C3H10T1/2 clone eight mouse fibroblast cells in culture by two separate PCB mixtures (Aroclor 1254 and 1260) and a purified component 2,4,5,2',4',5'-hexachlorobiphenyl (Norback and Weltman, 1980).

Comments

A wide variety of PCB mixtures have been subjected to rodent cancer bioassays and to numerous in vitro and in vivo short-term tests for mutagenicity. Several of these PCB mixtures are carcinogenic. None of the PCB mixtures are active in short-term tests for mutagenicity, a finding that holds true for most heavily chlorinated carcinogens. However, substantial confirming evidence for carcinogenicity is provided by positive cell transformation assays using these same PCB mixtures. Thus, under OSHA published criteria, PCB mixtures should be considered Category I* carcinogens. Both IARC (IARC, 1978) and EPA (EPA, 1978) have concluded that based on available animal data PCB's should be considered as potential human carcinogens.

-
- * Category I: Human evidence or two positive mammalian bioassays or 1 positive mammalian bioassay with supporting results in short term tests.

Category II: One positive mammalian bioassay. (Source: Occupational Health and Safety Letter Vol. 9, No. 24 November 8, 1979)

MONS 010677

VL BIOCHEMICAL EFFECTS OF PCB'S

Enzyme Induction

The principal biochemical effect of PCB's is the stimulation and induction of certain enzyme systems. Enzyme induction occurs in both the microsomal monooxygenase or cytochrome P-450 system and the aryl hydrocarbon hydroxylase or cytochrome P-448 system, and it has been observed in both man and experimental animals. Induction is not restricted to the liver. It occurs in numerous other organs including kidney, adrenal, lung, gut, skin, and testes. Fetal enzyme induction may occur via transplacental exposure, and induction may also occur by exposure to contaminated milk. (ref)

Identification of structure-activity relationships for enzyme induction is difficult because of the large number of isomers in commercially prepared PCB's and because all commercial products contain trace amounts of polychlorinated dibenzofurans (PCDF's) which are orders of magnitude more potent as enzyme inducers than PCB's. (ref)

In early studies using commercial Aroclors, potency for enzyme induction was found to be dependent on chlorination of the PCB mixture. Later, when purified isomers were tested, potency was found to vary with the position of chlorine atom substitution (see Section III). (ref) Since rate of metabolism is also known to vary with isomeric configuration of the PCB molecule, it may be that potency for enzyme induction is simply a function of the relative rate of metabolism and excretion.

The enzyme induction properties of PCB's are utilized in the metabolic activation system of in vitro bioassays for mutagenicity. It is unlikely, however, that enzyme

induction would consistently enhance the effects of carcinogens or pro-carcinogens: It might function synergistically to activate a chemical, but they also might function to deactivate reactive carcinogens. Both phenomena have been observed in rodent cancer bioassays.

Porphyria

Porphyria cutanea tarda (PCT) in humans is an acquired defect in hepatic porphyrin metabolism characterized by uroporphorinuria, photosensitivity and mechanical fragility of the skin. PCT can be produced experimentally by a number of drugs, including tetrachlorodibenzodioxins and PCB's. All of these agents have the ability to stimulate the activity of 2-aminolevulinic acid (ALA) synthetase which is the initial enzyme in the heme synthetic pathway.

Experimental hepatic porphyria was observed in Sherman rats exposed to Aroclor 1254 in the diet. At doses of 100 ppm the animals became porphyric after a delay of approximately 2-4 months. The porphyria resembled hexachlorobenzene poisoning and human PCT (Goldstein et al., 1975).

In chronic feeding studies ALA-synthetase induction occurs after rats have become porphyric, although with large single doses the enzyme induction is seen almost immediately after dosing the animals (Goldstein et al., 1975).

It has not been established whether only certain isomers in the PCB mixtures or contamination with PCDF's is responsible for the production of hepatic PCT. Porphyria has not been reported in humans exposed to PCB's.

Comments

Enzyme induction has two important implications for human health

MONS 010679

1. The occurrence of disease secondary to the increased metabolism of endogenous or exogenous substances, and
2. The interference with medical therapy due to increased metabolism of administered drugs.

PCB's are more potent enzyme inducers than phenobarbital, a drug that occasionally causes clinical problems due to its enzyme inducing effects. While the effects of phenobarbital decline after administration ceases, enzyme induction from PCB's persists long after cessation of exposure.

MDNS 010680

VII. HUMAN TOXICOLOGY AND EPIDEMIOLOGY

Few good epidemiologic studies of the health effects of PCB's are available. Most studies reported in the literature have been characterized by one or more of the following shortcomings:

1. Small study populations.
2. Lack of accurate exposure data.
3. Simultaneous exposure of workers to other potentially harmful chemicals.
4. Lack of control for confounding variables, such as alcohol consumption.
5. Inability to separate PCB's from contaminants and/or difficulty in comparing PCB's manufacture by different firms.

In spite of these problems, some health effects have been consistently reported in studies of workers occupationally exposed to PCB's. In addition, a large-scale poisoning which resulted from ingestion of PCB-contaminated rice oil has been well documented and resulted in multiple signs and symptoms attributable to PCB's.

The health effects identified in a review of the epidemiologic literature are summarized below, and Table II briefly describes the major epidemiological studies from 1954 through 1980.

Dermatologic Changes

Chloracne, contact or allergic dermatitis, and brown chromodermatosis have been consistently reported in studies of workers exposed to PCB's (Hara et al., 1975; Hasegawa et al., 1972; Inoue et al., 1975; Kitamura et al., 1973; Baker et al., 1980; Meigs et al., 1954; Ouw et al., 1976; Schwartz, 1936).

MONS 010681

Systemic Symptoms

Nausea, digestive disturbances, headaches, upper respiratory problems, and persistent body odor have been reported as a result of occupational exposures (Ouw et al., 1976; Schwartz, 1936; Warshaw et al., 1979).

Liver Damage

This effect has been reported in some studies (Hasegawa et al., 1972; Higuchi, 1976; Meigs et al., 1954; Ouw et al., 1976). However, some investigators reporting abnormal liver function tests did not control for additional chemical exposures, previous medical problems, drinking patterns, etc. These confounding variables could explain some or all of the marginal differences encountered.

Yusho (Japanese word translated as "oil disease")

Both dermal and systemic health effects are well documented in the epidemiologic study of a poisoning epidemic in Japan caused by ingestion of contaminated rice oil in 1968 (Higuchi, 1976; Kuratsune et al., 1972)

It is not clear how much the health effects observed in Yusho victims can be extrapolated to occupational exposures for the following reasons:

1. The average amount of PCB (Kanechlor 400) ingested was estimated to be 2 grams and the minimum, 0.5 gram (Kuratsune et al., 1972). This is a higher dose than has been reported in most occupational exposures. In addition, the PCB's were ingested as opposed to inhaled or skin-absorbed as is the case with occupational exposures.
2. The contaminated oil contained "used" Kanechlor 400, the exact chemical composition of which is unknown.
3. Frying of foods with the rice oil could have produced new compounds which may have altered the toxicity of the PCB's or the toxicity of possible contaminants.

MONS 010682

4. Yusho oil was shown to contain high concentrations of dibenzofurans.
5. Reported concentration of PCB's in the oil may not have been accurate enough to permit a rigorous quantitative analysis since the methods for estimating PCB's in foods were not fully developed at the time.

Clinical features of the Yusho patients are listed in Table III. The Yusho incident is also important because it clearly documents the potential for reproductive and transplacental effects in humans:

A study was made of the thirteen infants of 11 mothers affected by Yusho and of 2 unaffected wives of patients: Two of the Yusho mothers had stillbirths; ten of the babies had transient greyish or dark-brown pigmentation of the skin, and 5 had similar pigmentation of the gingiva and/or nails; increased ocular discharge was present in 9; and 12 of the 13 infants were small when compared with the national average (Funatsu et al., 1972; Kikuchi et al., 1969; Kuratsune, 1976; Taki et al., 1969). Babies born to patients even 3 years after severe PCB exposure tended to show pigmentation of the skin on the back and the gingiva, although the degree of pigmentation was less than that of babies born to the same mothers up to one year after the poisoning (Kuratsune, 1976).

Congenital abnormalities have also been observed in PCB-intoxicated infants. In the population of 13 offspring of Yusho mothers, premature eruption of teeth was observed in 2 cases, and larger than normal frontal and occipital fontanelles, exophthalmos and the persistence of an abnormally wide sagittal suture were observed in 3 others. No other gross malformations were reported nor was any relationship between dose and outcome considered. (Funatsu et al., 1972).

MONS 010683

Mothers' milk contaminated with PCB's also appears to be a source of exposure for infants: one baby showed signs of poisoning even though the mother had ingested the contaminated rice oil only after the baby was delivered. The infant began to show signs of PCB intoxication after 3-4 months of breast feeding (Kuratsune, 1972; Yoshimura, 1974).

Neurotoxicity

Paresthesias were reported in over 30% of Yusho patients (see Table III). In the Yusho epidemic more detailed neurologic examinations were performed in 21 cases admitted to a University hospital in northern Japan. Ten of the patients complained of numbness or pain in the distal extremities, and in five cases decreased pain, touch and temperature sensation was observed. Sensory conduction velocity in sural and radial nerves was below normal in 6 of 10 individuals with neuropathic symptoms (Murai and Kuroiwa, 1971). Headache and peripheral nervous system symptoms were also reported in a poisoning episode which occurred in a Finnish paper company (Hakkinen et al., 1973).

A decrease in amplitude of muscle action potential evoked by nerve stimulation, and a decrease in sciatic nerve conduction velocity has been reported in rats intoxicated with tetrachlorobiphenyl. Thus, PCB's can affect peripheral nerve function in both humans and experimental animals, but these have been reported only at doses which cause other systemic signs of poisoning.

Cancer

There is too little epidemiological evidence available yet to evaluate the potential of PCB's as human carcinogens (Bahn et al., 1980; Brown and Jones, 1980). A follow-up of the Yusho patients through 1977 has reported 51 deaths (31 with cause of death

MONS 010684

confirmed) of the 1665 identified victims. There were 11 deaths from neoplasms, or 35.4% of the total. While this rate is higher than the 21.1% in the population of the same prefecture in 1977, these data were not age-adjusted. No particular site was elevated, and there were no deaths from malignant melanoma, a tumor previously suspected to be linked to PCB exposure (Bahn et al., 1976). Two liver cancers and two lung cancers were reported but smoking and drinking patterns were not available (Urabe et al., 1979).

A retrospective cohort mortality study of 2,567 workers in two capacitor manufacturing plants was recently completed by NIOSH. The reports, still in draft form, did not find any statistically significant excess mortality for any cause of death among the exposed workers. Deaths from liver cancer, cirrhosis of the liver, and rectal cancer were slightly higher than expected, but no information was available on medical histories, drinking patterns, etc. No correlation was observed between increased mortality and length of exposure, but the number of total deaths was small (163). NIOSH will continue to follow up the mortality experience of the cohort (Brown and Jones, 1980).

Ongoing Occupational Studies

Two additional cohort mortality studies are currently underway. The first is a mortality survey of the entire workforce employed between 1952-1957 at the largest U.S. facility that manufactured capacitors and transformers. There is detailed information available on exposure levels in the plant. While the duration from onset of exposure is shorter than optimal (only 25 years in some cases), the information will at least give data on the short-term mortality experience of a heavily exposed occupational group (Selikoff et al., in progress).

MONS 010685

The second is a similar occupational mortality study, also of workers exposed in capacitor and transformer manufacturing. Over 2,000 workers have been identified for this study but no further details are yet available (Bertazzi et al., in progress). One case control study is currently being conducted to assess whether there is excess risk of malignant melanoma among PCB-exposed workers. This data will not be available until March, 1981 (Bahn et al., 1976).

There is also one cross-sectional clinical field survey of 326 capacitor manufacturing workers at two sites, encompassing a total workforce of 800 (Fischbein et al., 1979). Exposures were classified as none, low, medium, and high based on job description at the time of the survey (1975). Researchers were able to identify the PCB's used and had some data on environmental air levels in the plants. A number of parameters were measured, including complete history and physical exams, SMA panels and pulmonary function tests. Results have been published on respiratory function and general signs/symptoms, and results of serum lipids, endocrine function and dermatologic findings are forthcoming. To date, the only positive association involves dermatologic signs and symptoms.

Further investigations of the effects of PCB exposure on serum lipids have been done in both occupational and general environmental exposure settings. Smith et al., (1978), reported some statistically significant differences between exposed and non-exposed workers at two sites. They reported higher serum triglycerides and lower levels of high density lipo-proteins in the exposed group. Whether the magnitude of the difference is biologically significant is not clear from this study. For example, the non-exposed group at site #1 compared to the non-exposed group at site #2 showed a greater difference than the exposed and non-exposed comparison at either site. In another

MONS 010686

study (Baker et al., 1980) workers and community residents with exposure to fertilizer made from sewage sludge contaminated with PCB's were studied. Plasma triglyceride levels were found to increase significantly with serum PCB concentration (both in drinkers and non-drinkers), and the authors concluded that PCB's may alter lipid metabolism at levels of exposure and bio-accumulation insufficient to produce other identifiable signs of toxicity.

Comments

Although many problems have been identified in the studies evaluating the health effects of PCB's, it is clear that occupational exposure, at a minimum, can produce dermatologic effects. The long half-life of PCB's and their bio-accumulation in various human tissues leaves open the possibility of substantial chronic and delayed effects analogous to those seen in animal bioassays. These effects have only recently begun to be studied in a rigorous manner, and although the epidemiological evidence is neither complete nor entirely consistent there can be no question of the necessity to protect the worker from exposure.

MONS 010687

VIII. MEDICAL SURVEILLANCE AND BIOLOGIC MONITORING

Medical surveillance and biologic monitoring are of limited usefulness in predicting health hazards if dose-response relationships are not known. This certainly is the case with PCB's. Based on animal toxicology, there are many suspected adverse effects of PCB's which might result from exposure in occupational settings, but very few have been documented well enough to give even rough estimates of "no-effect" or "safe" levels. A large percentage of non-occupationally exposed people have detectable PCB levels in body fat, blood and milk. However, any attempt to estimate an adverse health effect associated with increases above this background level necessarily involves extrapolation from animal data and therefore is subject to considerable error, especially when the marked variation in sensitivity of various animal species is appreciated. Furthermore, not enough is known regarding the relative dose-response characteristics of the various documented effects (e.g., liver damage, skin changes) to state that in the absence of a particular sign, symptom or laboratory abnormality, the risk of long-term effect (cancer, reproductive toxicity) will be negligible (see Table I).

For the clinician confronted with a worker who has a history of exposure to PCB's the approach to management cannot be easily outlined. Given the current analytic methodology, residues can be measured in blood or tissue in the ppb range and compared to background; but assigning a health risk to a given level is virtually impossible, especially given the lack of pharmacokinetic data. Often patients are being evaluated after a considerable lag period (years) since last exposure occurred and extrapolation to peak blood levels is not possible. In fact, it may be that residue levels bear little relationship to the health risk. For example, the lower chlorinated compounds may be more toxic but they are more rapidly metabolized and excreted and therefore less

MONS 010688

likely to persist in blood or fat. Further, with the possible exception of chloracne, the presence of specific signs, symptoms or laboratory abnormalities is very difficult to definitely relate to PCB exposure in any given patient.

Given these uncertainties and the potential for serious health effects, the approach to monitoring should emphasize environmental sampling and every attempt should be made to minimize exposure by engineering controls or personal protective measures in those settings where occupational exposure still occurs (e.g., utility repair workers). Biologic monitoring may be used to assess the effectiveness of environmental control, but it is really best utilized within a specific research protocol and probably has little value in the routine work-up of individual patients.

MONS 010689

SUMMARY AND CONCLUSIONS

PCB's have low acute toxicity but are of public health concern because of their persistence in the environment and in human tissues and their demonstrated potential for chronic or delayed toxicity. They are potent inhibitors of reproductive function in both rodents and non-human primates and are positive in animal cancer bioassays. As potent inducers of hepatic enzyme systems, PCB's may have additional unpredictable long-term health effects.

Some of the conflicting reports in the toxicology literature are undoubtedly related to the variable composition and trace chemical contamination of the tested mixtures. Occupational and environmental exposure is usually to those mixtures; but if we are to accurately assess the associated health hazards, further animal studies are needed which carefully define the toxicology of the individual agents.

Epidemiologic studies of occupational exposures to PCB's to date have failed to detect serious adverse effects but are considered insufficient, and further studies are clearly needed. Of particular interest is the continued exposure among utility workers. Because of the potential ability to cause cancer and other long-term adverse effects such as infertility and hepatic injury, human exposure to PCB's should be kept to the lowest level technically possible. The persistence of PCB's in the body and the irreversibility of some of its effects make it necessary to act now, rather than to wait until more definitive data are available.

MONS 010690

REFERENCES

- A.C.G.I.H. Chlorodiphenyl - 42% chlorine, Documentation of the TLV's for substances in Workroom Air. 3rd ed., pp51-2. Cincinnati, 1976.
- Albro, PW and Fishbein, L. Intestinal absorption of PCB's in Rats. Bull Environ Contam Toxicol 8:26, 1972.
- Allen, JR. and Abrahamson, L.J. Morphologic and biochemical changes in the liver of rats fed PCB's. Arch Environ Contam Toxicol 1:265, 1973.
- Allen, JR. Response of the non-human primate to PCB exposure. Fed Proc 34: 1675, 1975.
- Allen, JR and Barsotti, DA. The effects of transplacental and mammary movement of PCB's on infant rhesus monkeys. Toxicol 6:331, 1976.
- Allen, JR et al. Response of rats exposed to PCB's for 53 weeks. Arch Environ Toxicol 4:404, 1976.
- Allen, JR et al. Reproductive effects of halogenated aromatic hydrocarbons on non-human primates. Ann NYAS 320:419, 1979.
- Anderson, MW et al. The construction of a pharmacokinetic model for the disposition of PCB's in the rat. Clin Pharm Therap 22: 765, 1977.
- Bahn, AK et al. Melanoma after exposure to PCB's (letter to the editor) N Engl J Med 295: 450, 1976.
- Baker, E et al. Metabolic consequences of exposure to polychlorinated biphenyls (PCB) in sewage sludge. Amer J Epidem 112: 553, 1980.
- Barsotti, DA et al. Reproductive dysfunction in rhesus monkeys exposed to low levels of PCB's (Aroclor 1248) Food Cosmet Toxicol 14: 99, 1976.
- Bertazzi, PA et al. Mortality experience among PCB workers. Univ di Milano, Inst Occ Health, Milan, Italy, in progress.
- Bitman, J and Ceal, HC. Estrogenic activity of DDT analogs and PCBs. J Ag Food Chem 18: 1108, 1970.
- Bowman, R et al. Correlation of PCB body burden with behavioral toxicology in monkeys. Pharmacol Biochem Behav 9: 49-56, 1978.
- Brackner, JV et al. Biologic response of the rat to PCB's. Tox Appl Pharm 24: 434, 1973.
- Goldstein, JA et al. A comparative study of two PCB mixtures on induction of hepatic porphyria and drug metabolizing enzymes. Tox Appl Pharm 32:461, 1975.

MONS 010691

Brown, DP and Jones, M. Mortality and industrial hygiene study of workers exposed to polychlorinated biphenyls. Draft report, NIOSH Division of Surveillance, Hazard Evaluation and Field Studies 1980.

Bumgarner, JE et al. Polychlorinated biphenyl residues in refuse workers. National Institute of Environmental Health Sciences (as reported in NIOSH Criteria Document), 1973.

Burse, VW et al. PCB's-storage, distribution, excretion and recovery: Liver morphology after prolonged dietary ingestion. Arch Environ Health 29: 301, 1974.

Curley, A et al. PCBs - Distribution and Storage in body fluids and tissues of Sherman rats. Environ Res 4: 481 1971.

Curley, A et al. PCB's evidence of transplacental passage in the Sherman rat. Food Cosm Toxicol 11: 471, 1973.

DHEW: Subcommittee on health effects of PCB's and PBB's - series of articles appearing in Env Health Persp 24: 146-198, 1978.

Dougherty, RC et al. Sperm density and toxic substances: A potential key to environmental health hazard, submitted to Env Health Chem, 1980.

EPA: Halogenated polyaromatics, in: L Fishbein, Potential industrial carcinogens and mutagens, pub #360/5-77-005, Office of Toxic Substances EPA, Washington, DC pp 173-197, 1977.

Finklea, J et al. PCB residues in human plasma expose a major urban pollution problem. Amer J Public Health 62: 645, 1972.

Fishbein, A et al. Clinical findings among PCB-exposed capacitor manufacturing workers. Ann NYAS 320: 203, 1979.

Fries, GF. PCB residues in milk of environmentally and experimentally contaminated cows. Env Health Persp. 1: 55, April 1972.

Funatsu, I et al. Polychlorobiphenyls (PCB) induced fetopathy. I Clinical observation (Abstract No. 72-2360). Kurume Med J 19: 43-51.

Gardner, AM et al. PCB's Hydroxylated urinary metabolites of 2,5,2',5'-tetrachlorobiphenyl identified in rabbits. Biochem Biophys Res Comm 55: 1377, 1973.

Grant, DL et al. Metabolism of a PCB (Aroclor 1254) mixture in the rat. Bull Environ Contam Tox 6: 102, 1971.

*(Goldstein, JA et al. see bottom p. 34)

Hakkinen, J et al. Diphenyl poisoning in fruit paper production. Arch Environ Health 26: 70, 1973.

Hara, I et al. Follow-up study of condenser factory after use of PCB discontinued. (As reported NIOSH Criteria Document) Jpn J Ind Health 17: 371-372, 1975.

Hasegawa, H et al. Report on survey of work area environment where PCB is handled and of the health of workers handling PCB, in Special research report on prevention of environmental pollution by PCB-like substances. (As reported in NIOSH Criteria Document) Japan, Research Coordination Bureau, Science and Technology Agency, pp. 141-99, 1972.

MONS 010692

- Heddle, JA and Bruce, WR. Comparison of tests for mutagenicity or carcinogenicity using assays for sperm abnormalities, formation of micronuclei and mutations in *Salmonella*. In, *Origins of Human Cancer*, Cold Spring Harbor Lab., pp 1549, 1975.
- Higuchi, K ed. PCB poisoning and pollution. Academic Press, NY 1976.
- Hirokadzu, D and Ota, H. Transfer of PCBs to infants from their mothers. *Arch Environ Health* 35: (2) 95, 1980.
- Hsu, IC et al. Metabolic fate of ³H-2,5,2',5'-tetrachlorobiphenyl in infant non-human primates. *Bull Environ Contam Toxicol* 18:233, 1975.
- IARC: Working Group on the evaluation of the carcinogenic risk of chemicals to humans - Polychlorinated Biphenyls, Vol 18, 1978.
- Inoue, Y et al. Discovery of PCB pollution in textile factory—I. PCB level in blood serum of laborers and results of physical examination (As reported in NIOSH Criteria Document) *Jpn. J Public Health* 22: 1637, 1973.
- Ito, N et al. Histopathological studies on liver tumorigenesis in rats treated with PCB's *Gann* 65: 545, 1974.
- Jensen, S and Sundstrom, G. Structure and levels of most chlorobiphenyls in two technical PCB products and in human adipose tissue. *Ambio* 3: 70, 1974.
- Jerina, DM and Daly, JW. Arene Oxides: A New Aspect of drug metabolism. *Science* 185: 573 1974.
- Karppanen, E and Kolho, L. The concentration of PCB in human blood and adipose tissue in three different research groups. In, *PCB Conference II, Stockholm 1972*. (As reported in NIOSH Criteria Document) Solna, Sweden National Swedish Environment Protection/Publications, 4E, pp 124-28, 1973.
- Katzenellenbogen, J and Ames, BN. Personal Communication 1980.
- Keplinger, M et al. Toxicologic Studies with PCBs (abstract) *Tox Appl Pharmacol* 19: 402-403, 1971.
- Kihlstrom, JE et al. Sexual function of mice neonatally exposed to DDT or PCB. *Environ Phys Biochem* 5: 54, 1975.
- Kikuchi, M et al. An autopsy case of stillborn of chlorobiphenyls poisoning. *Fukuoka Acta Med* 60: 489, 1969.
- Kimbrough, RD et al. Morphologic changes in liver of rats fed PCB's. *Arch Ind Health* 25: 354, 1972.
- Kimbrough, RD. The toxicity of polychlorinated polycyclic compounds and related chemicals. *Crit Rev Toxicol* 2: 445, 1974.
- Kimbrough, RD and Linder, RE. Induction of adenofibrosis and hepatomas of the liver in BALB/cJ mice by PCB's (Aroclor 1254). *J Natl Cancer Inst* 53: 547, 1974.
- Kimbrough, RD et al. Induction of liver tumors in Sherman strain female rats by PCB (Aroclor 1260). *J Natl Cancer Inst* 55: 1453, 1975.

Kimbrough, R et al. Animal toxicology, in DHEW Subcommittee on Health Effects of PCB's and PBB's. Env Health Persp 28: 173, 1978.

Kimbrough, R Chronic toxicity of halogenated biphenyls and related compounds in animals and health effects in humans. CDC report, 1980.

Kimura, NT and Baba, T. Neoplastic changes in the rat liver induced by PCB. Gann 64:105, 1973.

Kitamura, M et al. PCB in blood of workers employed in an electrical parts manufacturing plant. (As reported in NIOSH Criteria Document) Jpn J Ind Health 13: 539, 1973 (Jap).

Kuratsune, M et al. Epidemiologic study on Yusho, A poisoning caused by ingestion of rice oil contaminated with a commercial brand of PCB's. Environ Health Persp 1: 119, 1972.

Kutz, FW and Strassman, SC. Residues of PCB's in the general population of the US. In: Proceedings of the natural conference on PCB, Chicago, EPA - 560/6-75-004, Washington DC, pp 139.

Linder, R et al. The effect of PCB's on rat reproduction. Food Cosmet Toxicol 12: 63, 1974.

Lutz, RJ et al. Preliminary pharmacokinetic model for several chlorinated biphenyls in the rat. Drug Metab Dis 3: 386, 1977.

Mathews, HB and Anderson, NW. Effect of chlorination on the distribution and excretion of PCB's. Drug Metab Dis 3: 371, 1975.

Meigs, JW et al. Chloracne from and unusual exposure to Aroclor. JAMA 154: 1417, 1954.

Miller, JW. Pathologic changes in animals exposed to a commercial chlorinated biphenyl. Public Health Rep 59: 1085, 1944.

Moore, JA et al. Comparative toxicity of three halogenated dibenzofurans in guinea pigs, mice and rhesus monkeys. Ann NYAS 320: 151, 1979.

Murai, Y and Yoshigoro, K. Peripheral neuropathy in chlorobiphenyl poisoning. Neurol 21: 1173, 1971.

NCI: Carcinogenesis technical report series #38, DHEW publication #(NIH) 78-838, 1978.

Nelson, N et al. PCB's - environmental impact. Environ Res 3: 249, 1972.

NIOSH: Criteria for a recommended standard - Occupational exposure to PCB's NIOSH, Cincinnati, 1977.

Norback, DH and Weltman, R. Personal communication 1980.

New York State Health Planning Commission. Report of the ad hoc committee on the health implications of PCB's in mothers' milk. Albany Health Advisory Council, 1977.

Oishi, S et al. Comparative toxicity of PCB's and dibenzofurans in rats. Tox Appl Pharm 43: 13, 1978.

Orberg, J and Kihlstrom, JE. Effects of long-term feeding of PCB, Clophen A-60 on length of estrus cycle and frequency of implanted ova in the mouse. *Environ Res* 6: 176, 1973.

Ouw, HK et al. The use and health effects of Aroclor 1242, a polychlorinated biphenyl, in an electrical industry. *Arch Environ Health* 31: 189, 1976.

Sato, M and Hasegawa, H. Amount of PCB in blood of laborers. (As reported in NIOSH Criteria Document) *Jpn J Ind Health* 16: 365, 1974 (Jpn).

Safe, S et al. The metabolism of 4-chlorobiphenyl in the pig. *Can J Phys Pharmacol* 53: 392, 1975.

Schwartz, L. Dermatitis from synthetic resins and waxes. *Dermatitis* 26: 586, 1936.

Selickoff, IJ et al. Mortality experience of factory workers exposed to PCB's in the manufacture of transformers and capacitors. Mt. Sinai School of Med, Env Sciences Lab, New York, NY, in progress, 1980.

Shiota, K. Postnatal behavioral effects of prenatal treatment with PCB's in rats. (As reported in IARC, 1978). *Okajimas Fol Anat Jpn* 53: 105, 1976.

Smith, AB et al. Lipid and Lipoprotein alterations: Occupational exposures to PCB. *Clinical Res* 26: 549, 1978.

Stalling, DL et al. An expanded approach to the study and measurement of PCB's and selected planar halogenated aromatic environmental pollutants. *Ann NYAS* 320: 48, 1979.

Taki, I et al. Report on Yusho (chlorobiphenyls poisoning): Pregnant women and their fetuses. *Fukuoka Acta Med* 60: 471, 1969.

Treon, JF et al. The toxicity of the vapors of Aroclor 1242 and Aroclor 1254. *Am Ind Hyg Q* 17: 204, 1956.

Urabe, H et al. Present state of Yusho patients. *Ann NYAS* 320: 273, 1979.

Vodjcnik, MJ and Lech, JJ. The transfer of 2,4,5,2',4',5',-hexachlorobiphenyl to fetuses and nursing offspring. *Tox Appl Pharm* 54: 293, 1980.

Van Miller, JP et al. Distribution and metabolism of ³H-2,5,2',5',-tetrachlorobiphenyl in rats. *Proc Soc Exp Biol Med* 148: 682, 1975.

Vos, JG and Beems, RB. Dermal toxicity studies of technical PCB's and fractions thereof in rabbits. *Tox Appl Pharm* 19: 617, 1971.

Warshaw, R et al. Decrease in vital capacity in PCB-exposed workers in a capacitor manufacturing facility. *Ann NYAS* 320: 277, 1979.

Wyndham, C et al. The *in vitro* metabolism, macromolecular binding and bacterial mutagenicity of 4-chlorobiphenyl, a model PCB substrate. *Res Commun Chem Pathol Pharmacol* 15: 363, 1976.

Yoshimura, T. Epidemiological study on Yusho babies born to mothers who had consumed oil contaminated by PCB. *Fukuoka Acta Med* 65: 74, 1974.

MONS 010695

TABLE I

DOSE-RESPONSE FOR ANIMAL TOXICOLOGY

Species	Age	Agent	Route	Dose mg/kg/day	Duration	Total Dose mg/kg	Effect	References
Rat	Adult	Aroclor 1254, 1:60	Oral	4,000 + 10,000	Single Exposure	-----	LD ₅₀	Kimbrough et al., 1978
Rat	Weanling	Aroclor 1254, 1:60	Oral	1,700 + 1,300	Single Exposure	-----	LD ₅₀	Kimbrough et al., 1978
Rat	Adult	Aroclor 1242, 1:254, 1:62	Oral	50	6 weeks	2100	Liver hypertrophy, fatty infiltration	Allen & Abrahamson, 1977
Rat	Weanling	Aroclor 1242, 1:616	Oral	5.9-6.6	6 months	700-1700	Fatty liver	Burke et al., 1976
Rat	Adult	Aroclor	Oral	100	7-15 day gestation	700	↓ survival - offspring no litters	Linden et al., 1976
Rat	Adult	Aroclor 1260	Oral	25.0	2 months prior to mating	1500	↓ litter size ↓ survival offspring	Linden et al., 1976
Rat	Adult	Aroclor 1254	Oral	5.0	2 months prior to mating	300	↑ mortality - offspring ↑ mating behavior - offspring	Linden et al., 1976
Rat	Adult	Remazol 500	Oral	1.0	Day 8-14 or 15-21 gestation	4.0	↓ various learning assays offspring	Shiota, 1976
Mice	Adult	"PCB's"	Oral	7.0	4 weeks starting at birth	200	↓ implantations, litter size-offspring; changes in estrous cycle-offspring	Kihlstrom et al., 1975
Mice	Adult	Claphen 660	Oral	1.0	"chronic"	Unknown	↑ prolonged estrous cycles ↓ implantations	Oberg & Kihl- strom, 1977
Monkey	Adult	Aroclor 1248	Oral	10	4 weeks	840	weight loss, alopecia, facial edema, eye discharge	Allen, 1975
Monkey	Adult	Aroclor 1248	Oral	2.5	1 month	100	liver abnormalities (histological)	Allen, 1975
Monkey	"young"	Aroclor 1242	Oral	0.1-10	9 months	77-2700	death, gastric ulceration, facial edema, thymus atrophy	Allen, 1975
Monkey	Adult	Aroclor 1248	Oral	0.08-0.17	1 year	29-62	facial edema, alopecia cone, infertility, irregular menses, poorly abortions, offspring: ↑ weight, head circumference, (detectable PCB's in tissues)	Allen, 1975
Pigs	Adult	Aroclors	Dermal	≈100	25 days	2500	slight Δ's - liver histology	Miller, 1946
Guinea pig	Adult	Aroclors	Dermal	≈70	Single Exposure	-----	death -- delayed up to 21 days with liver atrophy	Miller, 1946
Rabbit	Adult	Aroclor 1260	Dermal	≈20	38 days	760	skin, liver and kidney damage	Yao & Boone, 1971
Rat, Mouse, Guinea pig, and cat	Adult	Aroclor 1254	Inhal.	2.5 *	17 weeks	450	microscopic liver Δ's	Troen, 1956
	" "	" "	" "	0.7 *	31 weeks	217	reversible liver Δ's	Troen, 1956
Mice	8 weeks	Remazol 500	Oral	50	32 weeks	11,200 ^a	↑ hepatocellular carcinomas (control 0/6; dosed 5/12) ↑ neoplastic nodules (control 0/6; dosed 7/12)	Ito et al., 1973
Mice	5 weeks	Aroclor 1254	Oral	50	46 weeks	16,100 ^a	↑ neoplastic nodules (control 0/24; dosed 9/22)	Kimbrough & Linden, 1976
Rats	4 weeks	Aroclor 1260	Oral	10	21 months	6.3 P ^a	↑ hepatocellular carcinomas (control 1/173; dosed 146/184) ↑ neoplastic nodules (control 0/173; dosed 26/184)	Kimbrough et al., 1975
Rats	8 weeks	Aroclor 1260	Oral	10	104 weeks	7.2 P ^a	↑ hepatocellular carcinomas (unpublished, 1980)	Herbach et al., 1980
Rats	8 weeks	2,4,5,2',4',5'- hexachlorobi- phenyl	Oral	10	104 weeks	7.2 P ^a	↑ hepatocellular carcinomas (unpublished, 1980)	Herbach et al., 1980

* Animals were exposed to 1.4 or 1.5 mg/m³ of Aroclor 1254 for 7 hours/day, 5 days/week for 17 weeks. To approximate dose in mg/kg/day, the following corrections were applied: 0.2 (volume of air breathed in m³ by animal per day); 0.5 (weight of animal in kg); 24/7 (correction for 24 hour/day exposure); 7/5 (correction for 7 day/week exposure). Resulting value assumes 100% absorption.

so Total dose in gm/kg

TABLE II - OCCUPATIONAL EXPOSURE TO PCBs

Study	Exposure Level & Time	Study Population		Dermal Effects	Liver Function	Findings Blood Concentrations	Cancer/ Mortality	Comments
		Exposed	Controls					
Melga 1934 - Out- break of derma- titis in a chemical plant	0.1 mg/M ³ Arochlor - 5 to 19 months inter- mittent expo- sure through vapor leakage	14	0	7/14 mild to moderate chloracne	6 normal, 1 borderline (in chloracne cases)			Since blood concentra- tions of PCBs could not be measured, no cor- relation of individual dose and skin effects was possible.
Hasegawa et al, 1972 Study of 6 industrial plants includ- ing PCB manu- facture, cap- tactor, manu- facture, and biphenyl recovery	Vapors 13-965 $\mu\text{g}/\text{M}^3$ Particulates 4-250 (6,270 in a spill) <1 to 20 years	99	32	Various	Slightly abnormal (elevated liver enzymes)	Exposed: 370 ppm non-exposed: 20 ppb		Dermal ailments were unrelated to blood concentrations. Based on 3 plants, there was no rel- ationship of expo- sure to blood con- centration; fat metabolism was apparently affected
Hara et al 1973-1974 MONS 010697	Level of exposure not reported in NIOSH Criteria Document	118 (study concentr- ated on 17 immer- sion pro- cess workers)		45% black- heads, 37% acne, 13% irritation	not reported	Exposed 7-300 ppb		Blood concentrations closely related to years of exposure; Follow-up study after exposure ceased allowed calculation of serum half- life: The longer the duration of exposure, the longer the PCB half-life. (range 3 - 30 months)
Kitamura 1973 Study of workers medical exam in a capac- itor manu-act- uring plant	Exposure level not reported in NIOSH Criteria Document Time 2.5 years	13		Various: acne, seb- orrhea adiposa, folliculitis	Normal	820 ppb average (320- 2100 ppb)		No relationship was found between concentration in blood and expo- sure duration of expo- sure.

OCCUPATIONAL EXPOSURE TO PCBs

Study	Exposure		Study Population		Dermal Effects	Liver Function	Findings		Comments
	Level & Time		Exposed	Controls			Blood Concentrations	Cancer/Mortality	
Ow et al 1976 Study of capacitor manufacturer	Arochlor 1262 1.08-1.44 mg/M ³ (19 fillers) .32 mg/M ³ (15 assemblers) Workers wore no protective clothing Time 1 month to 23 years		34	30	Mild burn- ing, irri- tation of face, eyes and skin 5 had rashes, 1 chloracne, several dermatitis	Bromsul phot- halium tests elevated 4 of 7 fillers with blood levels > 500 ppb	Exposed 100- 602 ppb (mean 400 ppb) Not detected in non-exposed		Systemic effects reported such as nausea and per- sistent body odor. There was no adverse response at blood concentrations below 200 ppb
Bahn, et al 1976 Study of workers in a refinery	Arochlor 1254 over a 9 year period		51 researchers and development; 41 refinery workers					2 malignant melanomas observed, .04 expected (based on TNCS data)	These were pre- liminary results reported in a letter to the editor. Workers were also exposed to other chemicals. Study is in progress.
Brown and Jones 1980 cohort morbi- lity study of capacitor manufacturers (draft report)	Plant #1: 1.24 µg/M ³ 3.93 µg/M ³ Plant #2: 170 µg/M ³ 1260 µg/M ³		Plant #1-968 Plant #2-1599				All cause mortality was lower than expected (163 obs. vs 174 exp) All cancer mortality was lower than expected (39 obs. vs 40.6 exp.) Rectal and liver cancer were slightly elevated but not significantly.		Lower observed mortality may be attributable to the "health worker" effect. NIOSH will continue to follow- up mortality experience

OCCUPATIONAL EXPOSURE TO PCBs

Study	Exposure		Study Population		Dermal Effects	Liver Function	Findings		Cancer/Mortality	Comments
	Level & Time		Exposed	Controls			Blood Concentrations			
Roux et al 1975 Study of a family with thread processing operation	Not reported in NIOSH Criteria Document		Number studied not reported in NIOSH document (Latar)		Mild skin lesions including comedones	Unknown	130-520 ppb			Good correlation between degree of exposure and blood PCB levels.
	0.15-1.2 μM^3 0.13-4.4 $\mu\text{g}/\text{M}^3$		54 more persons were studied							
Jato and Hest-jawa 1974 Pressure sensitive carbonless paper manufacturer	measured 2 yrs. past PCB use		Not reported in NIOSH Criteria Document				Exposed 73 ppb non-exposed 20 ppb			
Carpien and Colino 1972 Study of 3 groups: 1) no exposure 2) analytical lab capacitor impregnation	High exposure (capacitor manufacturing); low exposure (analytical Lab)		High - 12 Low - 6	4 males 5 females			Unexposed 5.6-12 ppb Medium exposure 35-63 ppb High exposure 78-1,900 ppb			Unable to detect any "Biologic Effects"; type of monitoring not reported in NIOSH review
Bumgarner et al 1973 Study of refuse workers exposed to PCB in incineration of waste	Incinerated		37	36 lumber yard workers			Controls: max. 4.2 ppb Exposed: max. 14 ppb (4-14 ppb)			Concentrations not well correlated with duration of exposure, age or race

MONS

010699

OCCUPATIONAL EXPOSURE TO PCBs

Study	Exposure Level & Time	Study Population		Dermal Effects	Liver Function	Findings Blood Concentrations	Cancer/ Mortality	Comments
		Exposed	Controls					
Baker, Land- rigan et al 1980 Study of exposure to PCBs in sewage sludge	Liquid sewage entering plant 30-430 ppb. Up- stream sewage 1250-5500 ppb (Aroclor 1016) Concentrations in sludges (Aroclor 1242) were as high as 1700 ppm (mean 479.1 ppb) and 107.3 ppm in treated soil (mean 17.1 ppm)	89 sludge users, 18 workers exposed to PCBs, 19 members workers families	22 com- munity members	Acne, increased pigmenta- tion in 4 workers		Sludge users 17.4 ppb, workers 75.1 ppb Families 33.6 ppb, community 24.2 ppb		Plasma triglyceride levels increased significantly with serum PCB con- centrations. Data indicate that PCBs may alter lipid metabolism.

MONS 010700


TABLE III
PERCENT DISTRIBUTION OF SYMPTOMS OF YUSHO REPORTED
BY 189 PATIENTS EXAMINED BEFORE OCTOBER 31, 1962

Symptoms	Males (N-89)	Females (N- 100)
Dark brown pigmentation of nails	83.1	75.0
Distinctive hair follicles	64.0	56.0
Increased sweating at palms	50.6	55.0
Acnelike skin eruptions	87.6	82.0
Red plaques on limbs	20.2	16.0
Itching	42.7	52.0
Pigmentation of skin	75.3	72.0
Swelling of limbs	20.2	41.0
Stiffened soles in feet and palms of hands	24.7	29.0
Pigmented mucous membrane	56.2	47.0
Increased eye discharge	88.8	83.0
Hyperemia of conjunctiva	70.8	71.0
Transient visual disturbance	56.2	55.0
Jaundice	11.2	11.0
Swelling of upper eyelids	71.9	74.0
Feeling of weakness	58.4	52.0
Numbness in limbs	32.6	39.0
Fever	16.9	19.0
Hearing difficulties	18.0	19.0
Spasm of limbs	7.9	8.0
Headache	30.3	39.0
Vomiting	23.6	28.0
Diarrhea	19.1	17.0


Source: Kuratsune et al, 1972

MONS 010701


FIGURE 1



BIPHENYL MOLECULE AND RING NUMBERING SYSTEM



3-chlorobiphenyl



2,2',3,4',5-pentachlorobiphenyl

EXAMPLES OF NOMENCLATURE SYSTEM OF CHLOROBIPHENYL COMPOUNDS


NUMBER OF ISOMERS AND PERCENT CHLORINE
FOR THE 10 CHLOROBIPHENYL (PCB) CLASSES

Chlorobiphenyl	Empirical Formula	No. of Isomers	Weight % Cl
mono	C ₁₂ H ₉ Cl	3	18.79
di	C ₁₂ H ₈ Cl ₂	12	31.77
tri	C ₁₂ H ₇ Cl ₃	24	41.30
tetra	C ₁₂ H ₆ Cl ₄	42	48.56
penta	C ₁₂ H ₅ Cl ₅	46	54.30
hexa	C ₁₂ H ₄ Cl ₆	42	58.93
hepta	C ₁₂ H ₃ Cl ₇	24	62.77
octa	C ₁₂ H ₂ Cl ₈	12	65.98
nona	C ₁₂ HCl ₉	3	68.73
deca	C ₁₂ Cl ₁₀	1	71.18


Source: NIOSH, 1977

MONS 010702


FIGURE II




Polychlorinated Biphenyls (PCB's)




Polychlorinated Terphenyls



Chlorinated Dibenzofurans



Chlorinated Dibenzodioxin



Polychlorinated Naphthalenes

Source: Kimbrough, 1974

FIGURE III
CHLORODIBENZOFURAN TYPES AND CONCENTRATIONS ($\mu\text{g/g}$)
IN COMMERCIAL PCB PREPARATIONS

Mixture*	Chlorodibenzofurans						Total
	di	tri	tetra	penta	hexa	hepta	
(1) 1016	0.5						0.5
(1) 1016			<0.0001	<0.0001	<0.0001		
(1) 1248			0.5	1.2	0.3		2.0
(1) 1254			0.1	0.2	1.4		1.7
(1) 1254			0.2	0.4	0.9		1.5
(1) 1260			0.1	0.4	0.5		1.0
(1) 1260			0.2	0.3	0.3		0.8
(2) A-60			1.4	5.0	2.2		8.4
(3) DP-6			0.7	10.0	2.9		13.6
(4) K300			(a)	(a)			1-1.5
(4) K400	(c)**	(e)	(e)	(c)			17-18
(4) K500			(a)	(a)	(c)	(a)	2,5-4
(4) K600			(a)	(a)	(b)	(b)	3-5

* (1) Aroclor, (2) Clophen, (3) Phenoclor, (4) Kanechlor

** (a), (b), (c), (d), (e) represent relative amounts in increasing order

Source: NIOSH, 1977

MONS 010703

FIGURE IV
Responses of primates and rats to PCB's*

Response	Man	Monkey	Rat
Susceptibility to toxicity	High	High	Moderate
Acne	Yes	Yes	No
Hyperpigmentation of skin	Yes	Only infants	No
Alopecia	NA	Yes	No
Hyperactive Meibomian glands	Yes	Yes	No
Conjunctivitis	Yes	Yes	No
Oedema of eyelids	Yes	Yes	No
Subcutaneous oedema	Yes	Yes	No
Keratin cysts in hair follicles	Yes	Yes	No
Hyperplasia of hair follicle epithelium	Yes	Yes	No
Gastric hyperplasia	NA	Yes	No
Thymic atrophy	NA	NA	Yes
Hepatic hypertrophy	Yes	Yes	Yes
Liver enzyme change	NA	Yes	Yes
Decreased no. of red-blood cells	Yes	Yes	No
Decreased haemoglobin	Yes	Yes	No
Serum hyperlipidaemia	Yes	Hypolipidaemia	Yes
Leucocytosis	Yes	Yes	No

Source: IARC, 1978

* This table summarizes acute and subacute clinical effects but does not include chronic or delayed effects such as reproductive effects or cancer.

MONS 010704