

FILE NAME: Phenolic Resins (PHR)

DATE: 1936

DOC#: PHR031

DOCUMENT DESCRIPTION: Book Excerpt from Industrial Dust; from Eagle
Pitcher File

INDUSTRIAL DUST

*Hygienic Significance, Measurement
and Control*

BY

PHILIP DRINKER, S.B., Ch.E.

*Professor of Industrial Hygiene, Harvard
School of Public Health*

AND

THEODORE HATCH, B.S., S.M.

*Instructor in Industrial Sanitation, Harvard
School of Public Health and Harvard
Graduate School of Engineering*

FIRST EDITION
SECOND IMPRESSION

McGRAW-HILL BOOK COMPANY, Inc.

NEW YORK AND LONDON

1936

CONTENTS

	Page
CHAPTER I	
PHYSICAL PROPERTIES OF DUSTS, FUMES AND MISTS.	1
Definitions—Settling Velocities—Flocculation—Wetting—Electrical Properties—Optical Properties.	
CHAPTER II	
EFFECTS OF DUSTS AND FUMES UPON MAN.	20
Respiration and Pneumoconiosis—Silicosis—Asbestosis—Toxic Dusts—Metal-fume Fever—Allergic Reactions—The Dusty Trades.	
CHAPTER III	
PHYSICAL AND CHEMICAL FACTORS IN PNEUMOCONIOSIS	43
Quartz—Hardness—Solubility—Silicates, Asbestos, Sericite—The Silicosis Potency of Dusts—Particle Size—Differential Disintegration and Selective Settlement.	
CHAPTER IV	
PERMISSIBLE DUSTINESS.	64
The Retention of Inhaled Dust—Ash of Dusted Lungs—Standards of Dustiness: Quartz; Coal; Cement and Limestone; Lead; Zinc Oxide; Manganese—Limitations in Standards.	
CHAPTER V	
THE DUST SURVEY	80
Historical—Outline of Survey—Requirements and Limitations of Dust Determination.	
CHAPTER VI	
DUST DETERMINATION.	91
Settlement—Filtration—Impingement: Konimeter; Owens' Apparatus; Continuous Recorder.	
CHAPTER VII	
DUST DETERMINATION—(Continued)	110
Impinger: Lightfield and Darkfield Counting—Tyndallmeter—Electric Precipitator—Thermal Precipitator—Comparison of Sampling Instruments.	
CHAPTER VIII	
DETERMINATION OF PARTICLE SIZE	133
Collection of Sample—Measurement: Screen Analysis; Microscopic; Elutriation; Determining Size of Submicroscopic Particles—Graphical and Mathematical Analysis of Particle-size Data.	

CHAPTER IX

CHEMICAL AND MINERALOGICAL ANALYSES.	153
Collection and Treatment of Air-borne Dust—Recovery of Dust from Lungs—Determination of Free and Combined Silica—Combined Chemical and Petrographic Method—Immersion Method—Composition in Relation to Size of Particles—Microscopy of Sputum—Identification of Asbestosis Bodies.	

CHAPTER X

CONTROL OF THE DUST HAZARD—GENERAL.	167
Medical Control—Elimination of Sources of Dust—Preventing Dust Dispersion: Segregation; Enclosure; Wetting; Local Exhaust Ventilation—Personal Protection—Housekeeping.	

CHAPTER XI

DESIGN OF LOCAL EXHAUST SYSTEMS.	177
Dust Dispersion: by Dynamic Projection; by Air Currents—Control of Air Currents—Aerodynamic Characteristics of Exhaust Hoods—Hood Design—Determining Required Rate of Airflow into Hoods—Static Suction as an Index of Hood Operation—Exhaust Piping: Basic Equation; Future Expansion; Layout—Transporting Velocities—Pressure Losses—Power Consumption—Pipe Design—Pipe Construction—Source of Suction: Laws; Construction; Location of Fan.	

CHAPTER XII

AIR-CLEANING APPARATUS.	220
Amount of Dust Collected—Size of Particles—Air Cleaning Requirements—Methods of Air Cleaning—Discharge Stack—Gravitational Settling Chamber—Cyclone—Inertial Separator.	

CHAPTER XIII

AIR-CLEANING APPARATUS—FILTRATION.	233
Particle Size—Concentration—Air Velocity and Porosity—Physical Characteristics; Plugging—Filter Cloths—Construction and Operation of Dust Filters—Filters in Air Conditioning—Bacterial Filters—Production of Dust Clouds in the Laboratory—Testing Filter Materials, Resistance, and Filtering Efficiency.	

CHAPTER XIV

DUST RESPIRATORS AND AIR MASKS.	261
History—Testing Dust Respirators: Resistance; Facial Fit; Efficiency—Testing Air Masks—Second-grade Respirators. .	
BIBLIOGRAPHY.	277
NAME INDEX.	291
SUBJECT INDEX.	297

fully that, if he has silicosis, he got it in Canada and Canada will take care of him.

In the case of a first-stage silicotic without tubercular infection, there is no method, as yet, for determining his lessened efficiency. The lung change, if any, caused by his dust inhalation is too small to measure.

McCann and Hurtado (177) have estimated disability from lung fibroses of various origins but have not yet succeeded in developing a method that is simple enough for routine use by compensation boards, insurance examiners, and the like. Their procedures involve various simple measurements of respiratory function but they have not established any methods of detecting fibrosis which are more sensitive than the present X-ray examination.

Of course, industry would welcome a diagnostic routine for detecting early harm from dust or, better, for forecasting harm. No such method is available and it seems fruitless to expect such help. The pneumoconioses are not diagnosed until harm is demonstrable by X-ray.

Collis (42) has stressed the difference between the age groupings in pulmonary tuberculosis and in silicosis, with or without tubercular infection superimposed. Tuberculosis is most prevalent between the ages sixteen to twenty-four, while silicosis rarely becomes disabling until later in life. Inasmuch as a good many years of dust exposure are required before the average case of silicosis (or asbestosis) becomes evident, it is obvious that it is not likely to be acquired early in life. However, there is an epidemiological side to the question which is of great importance and it is that which Collis especially emphasized. The average healthy individual has acquired a fairly effective immunity or resistance to tuberculosis. His chest X-ray is likely to show one or more healed scars or fibrosed areas from tuberculosis. Such an individual, Cummings emphasizes (211), is a better silicosis risk than the man who shows no evidence of past exposure to pulmonary tuberculosis. Consequently it is wise to select for dusty jobs men who are past the age of forty and not young men just taking up a trade.

Asbestosis.—The X-ray picture of the typical asbestotic chest is confusing to the layman. The effect is described as a diffuse fibrosis. Characteristic silicotic nodules are absent and even

EFFECTS OF

the expert roentgen history is complete.

The pathology produces. The asbestos fibers irritate the lung tissue, causing what is called emphysema or transmigration of the blood vessels and no form of Fig. 13. As the at




FIG. 13.—Asbestosis b.

area causes serious suggests that the of second-stage a work of the heart work to pump blood normal lung.

In silicosis it is point, the victim's has ceased. But seen patients with

the expert roentgenologist withholds his diagnosis until the case history is complete.

The pathology produced by asbestos is not like that of silicosis. The asbestos fibers group about the neck of an alveolus and shut it off, causing what is known as *atelectasis*. There is no definite migration or transportation of the dust particles to the lymph nodes and no formation of the fibrous nodules as shown in Fig. 13. As the atelectatic areas increase, the reduction in lung




FIG. 15.—Asbestosis bodies in sputum. (After Ellman (75); courtesy J. Ind. Hyg.)

area causes serious dyspnea or labored breathing. Lanza (158) suggests that the enlarged hearts noted frequently in his cases of second-stage asbestosis may be the result of the increased work of the heart resulting from this condition; it takes more work to pump blood through the atelectatic than through the normal lung.

In silicosis it seems to be a general rule that, after a certain point, the victim's condition grows worse even if his dust exposure has ceased. But Wood and Gloyne (258) state that they have seen patients with asbestosis "whose condition appears to have

remained stationary since stopping work in the factory," but they advise definitely that the asbestotic individual be removed from his dusty job. Merewether (183) and Lanza are less certain on this point.

Asbestosis Bodies. In the lungs of patients who died after prolonged exposure to asbestos dust and in the sputum of men with considerable asbestos-dust exposure are found what first were called *curious bodies* and later *asbestosis bodies* (Fig. 15) (75). While somewhat similar bodies occur in the lungs of coal workers and even of normal persons, it is admitted that asbestosis bodies in sputum are characteristic of asbestosis. Stewart (223) gives considerable diagnostic weight to their presence.

Pneumoconiosis in Animals. Silicosis has been produced experimentally by quartz dusting guinea pigs, rabbits, mice, cats, and domestic fowl, thus emphasizing the specificity of quartz dust to biological tissue. In like manner asbestosis has been produced in experimental animals.

The Equidae, such as horses and mules, apparently are not susceptible to ordinary pulmonary tuberculosis but there is no reason to believe they have any special immunity or resistance to silicosis. The normal silica content of horses' or mules' lungs is not known but it would seem wholly reasonable to use the lungs of animals which have worked 5 to 15 years underground in mines as physiological dust samples. A brief report on this subject was published by Haynes (121) but data such as one needs are singularly lacking.

TOXIC DUSTS

Poisoning from inhaling toxic dusts is much more likely than poisoning from swallowing them. Dusts that reach the lungs may pass directly into the blood stream, thence to the heart, and immediately be pumped all over the body. Distribution to all the body tissue is thus brought about rapidly and effectively. But dust taken in with the food goes to the stomach and the major part passes out in the feces. Some is picked up by the portal blood circulation and moves on to the liver. That portion which causes poisoning must first pass through the liver, which is an effective filter and detoxifier; only then can it enter the general circulation.

The practi
between the t
(103) states,

A great dea
who sought to
grinders again
clean overalls
escape of lead
tunately this
must never be
enter the body
three times a
ten hours of h


Goodby (8
poisoning m
over ten tin
Drinker and
foreign dust
(188) emph
from inhaled
that the ab
rapid and m
dose by mo

All this
expressed by
now beginn
danger of be
them in foo

Metal-fur
ing of dust
malady tha
of metal fur
lead, probal

About 2 f
metallic or
followed by
following a
incomforta
day he feel

EXHIBIT K: 8e



613.63
D83

AFV	No.	8468
	Date	NOV 26 1951
	E.	CO.
		JOPPA, MO.

660.8
829

EXHIBIT "M"

82

Handbook of Dangerous Materials

By

N. IRVING SAX

Toxicologist, General Electric Co.
Schenectady, N. Y.

Assisted by

M. J. O'HERIN

Fungicide Laboratory
General Electric Co., Schenectady, N. Y.

and

W. W. SCHULTZ

General Engineering Laboratory
General Electric Co., Schenectady, N. Y.

BOOK DIVISION

REINHOLD PUBLISHING CORPORATION
330 West Forty-Second St., New York 18, U.S.A.

1951

Arsenical Mixtures or Compounds, N. O. S.*
Solid

Hazardous Properties: Poisonous materials. See Arsenic.

Shipping Regulations:†

Poison B 352, 354* Poison label 200 lbs.
(max. lot)

Arsenical Dip, Liquid (Sheep Dip)

Hazardous Properties: Poisonous liquid. See Arsenic.

Shipping Regulations:†

Poison B 338, 349* Poison label 55 gals.
(max. lot)

Arsenical Dust

Hazardous Properties: A poisonous material. See Arsenic.

Shipping Regulations:†

Poison B 352, 355* Poison label 200 lbs.
(max. lot)

Arsenical Flue Dust

Hazardous Properties: Poisonous material. See Arsenic.

Shipping Regulations:†

Poison B 352, 355* Poison label 200 lbs.
(max. lot)

Arsenous Acid, Solid

Hazardous Properties: A poison. See Arsenic.

Shipping Regulations:†

Poison B 352, 361* Poison label 200 lbs.
(max. lot)

Arsenous and Mercuric Iodide Solution, Liquid
Hazardous Properties: A poisonous liquid. See Arsenic and Mercury.

Shipping Regulations:†

Poison B 338, 349* Poison label 55 gals.
(max. lot)

Arsine

Maximum Allowable Concentration in Air:* 0.05 parts per million.

Hazardous Properties: See Arsenic. Caution! Arsine can be liberated from unusual places. For instance, in cleaning out steel tanks which contained concentrated sulfuric acid it is possible to liberate arsine; although concentrated sulfuric acid does not react with steel, dilute acid does react with it momentarily, liberating hydrogen, which reacts with the arsenic, forming arsine. This is but one of the very numerous possibilities for industrial exposure to this toxic material.

Synonyms: Arsenic hydride; arseniuretted hydrogen

Formula: AsH₃

M. Wt.: 77.9

M. P.: -113.5°C

B. P.: -55°C

Density: 3.484 g/liter

Description: Colorless gas

Shipping Regulations:† Poison label (1).

Arsphenamine

Hazardous Properties: A toxic material. See Arsenic.

Synonyms: 3-Diamino-4-dihydroxy-1-arsenobenzene hydrochloride; 606

Formula: C₆H₁₂AsN₂O₂·2HCl·2H₂O **M. Wt.:** 475.0

Description: Light yellow hygroscopic powder

Shipping Regulations:† None.

"Artic"

Trade mark of a refrigeration grade of methyl chloride. A proprietary material. See under component as listed.

Asbestos Particles or Asbestos Dust

Maximum Allowable Concentration in Air:* 5 million particles per cubic meter.

Hazardous Properties: Asbestos is a class name for various minerals of a fibrous nature. The commercial material is mainly fibrous serpentine, known as chrysotile. The implication is that this material merely constitutes a nuisance dust. The figure given for the M. A. C. is probably the safe limit. There is a specific lung disease known as asbestosis, which may be caused by this material; it can also cause chronic conjunctivitis with hypertrophy (6). Asbestosis is a comparatively new disease which was first described by English writers (1920 to 1925). Finely ground asbestos is said not to be dangerous, and this theory is supported clinically. The time necessary to develop asbestosis is from 7 to 9 years, if the exposure concentration is fairly high. Lower concentrations are likely to require 15 to 20 years of exposure. Once established, the disease is progressive, even though exposure has stopped. It would seem that the irritation due to this material is related to its physical nature rather than its chemical composition. The first sign is dyspnea; the cough is usually dry and auscultation reveals few if any physical signs; loss of weight, pallor and cyanosis are also early symptoms. Later, emaciation and chest symptoms appear. There is a distinction between this disease and silicosis, i.e., the chest appears emaciated and lacks the robust character noticed in silicosis. Occupational exposure occurs in the packing, insulating and fire-proofing industry or where this material is combined with textiles.

Treatment and Antidotes: Rest and removal from exposure are indicated. Consult a physician.

Storage and Handling: Personnel exposed to high concentrations or to unknown concentrations of asbestos particles should wear respiratory protection of an approved design.

Asphalt


Hazardous Properties: Asphalt causes skin eruptions which vary roughly in relation to the specific gravity and boiling point of the asphalt or bitumen. It can cause dermatitis. The fumes evolved are noxious and unpleasant with a strong odor, so that bottles of this material

being too
discomfor
evolved a
a fire ha
Furtherm
see naph
as a mixt
for use, t
as a duc
as a pa
coating f
roofing f
as a con.
Treatment c
hot asp
If this is
water a
removed
benzene
extensiv
Storage and
This is
sary to
fres. It
close to
bottles
itself th
insulating
should
It shou
should
the top
Shoes
no ope
the sh
down
should
not be
ot to
asphal
should
be ca
them
To
spray
chlori
Flask Pa
Asp. lin
Hazard
optic
Storage
mate
Oher
safer
skin.
Aspirin
Hazard
to c
* See 1

* See list of abbreviations and symbols, page vii. † Section 5 gives complete ICC Shipping Regulations. Numbers in parentheses refer to literature citations, page vi.

EXHIBIT "L"

86



613.6

J 64

Please xerox an
extra copy of all
this for BE.
These are
primary file mat'ls

A rectangular piece of paper with a double border, tilted to the right. It contains handwritten text in cursive. The text is arranged in five lines. The first line is "Please xerox an". The second line is "extra copy of all". The third line is "this for BE.". The fourth line is "These are". The fifth line is "primary file mat'ls".

OCCUPATIONAL MEDICINE AND INDUSTRIAL HYGIENE

By

RUTHERFORD T. JOHNSTONE, A.B., M.D.

Consultant in Industrial Health; Lecturer at the
University of California, Los Angeles

Formerly Assistant Professor of Medicine, University of
Pittsburgh School of Medicine; Formerly Director of
Department of Occupational Diseases,
Golden State Hospital

With One Hundred Seventeen Illustrations

Seven in Color

St. Louis

THE C. V. MOSBY COMPANY

1948

CONTENTS

PART ONE

CHAPTER I

Page

TOMORROW'S DOCTORS (THE BACKGROUND AND THE FUTURE OF INDUSTRIAL MEDICINE)	17
---	----

CHAPTER II

INDUSTRIAL MEDICINE AND WORKMEN'S COMPENSATION	25
Personal Injury, 29; By Accident, 29; Arising Out of and in the Course of Employment, 29; In the Course of Employment, 30.	

CHAPTER III

THE NEED OF SCIENTIFIC PROOF IN MEDICOLEGAL CONTROVERSY	54
---	----

CHAPTER IV

FUNCTIONS OF THE INDUSTRIAL PHYSICIAN	57
---	----

CHAPTER V

THE TEACHING OF INDUSTRIAL MEDICINE	69
---	----

CHAPTER VI

THE BASIC APPROACH TO A DIAGNOSIS OF OCCUPATIONAL DISEASE	75
---	----

CHAPTER VII

LABORATORY PROCEDURES OF VALUE IN THE DIAGNOSIS OF THE OCCUPATIONAL DISEASES	81
--	----

CONTENTS

PART TWO

CHAPTER VIII

	Page
THE INDUSTRIAL SOLVENTS.....	87
Saturated and Unsaturated Compounds, 90; Chains and Rings of Atoms, 90; The Aliphatic Compounds, 91; The Aromatic Compounds, 91; The Alcohols, 92; The Aldehydes, 92; Acids and Esters, 92.	

CHAPTER IX

THE ALIPHATIC HYDROCARBONS.....	99
Petroleum Series, 99; Gasoline, 100; Occupational Exposures, 100; Pathology, 100; Signs and Symptoms, 100; Laboratory Findings, 101; Disability, 101; Treatment, 102; Illustrative Cases, 103; Tetraethyl Lead, 106; Toxicity, 106; Pathology, 107; Symptoms and Signs, 107; Prodromal Period, 107; Physical Examination, 107; Complications, Sequelae, and Convalescence, 107.	

CHAPTER X

THE ALIPHATIC HYDROCARBONS—CONT'D.....	100
The Alcohols, 109; Methyl Alcohol, 109; Ethyl Alcohol, 110; Propyl Alcohol, 110; Butyl Alcohol, 110; Amyl Alcohol, 111; Treatment, 111; The Glycols, 112; The Aldehydes, 114; Occupational Exposures, 114; Prophylaxis of Formaldehyde Poisoning, 115; Treatment of Formaldehyde Poisoning, 115; The Ketones, 115; The Aliphatic Acids, 116.	

CHAPTER XI

THE ALIPHATIC HYDROCARBONS—CONT'D.....	118
The Aliphatic Oxides, 118; Carbon Monoxide, 118; Properties, 118; Occupational Hazards, 118; Pathology, 118; Diagnosis, 120; Differential Diagnosis, 124; Sequelae of Acute Carbon Monoxide Poisoning, 124; Treatment, 127; Carbon Dioxide, CO ₂ , 130; Carbon Oxychloride, COCl ₂ , 130; Aliphatic Nitrile Compounds, 130; Pathology and Physiology, 130; Signs and Symptoms, 131; Tests, 132; Medicolegal Aspects, 132; Treatment, 133; Discussion, 135.	

CHAPTER XII

THE HALOGENATED HYDROCARBONS.....	126
Methyl Bromide, 136; Properties, 136; Tests, 136; Pathology, 136; Signs and Symptoms, 139; Disability, 139; Prevention, 140; Treatment, 141; Case Histories, 141; Methyl Chloride (CH ₃ Cl), 142; Properties, 142; Pathology, 142; Signs and Symptoms, 142; Tests, 143; Medicolegal Aspects, 143; Treatment, 143; Illustrative Cases, 143; Dichlorodifluoromethane (CCl ₂ F ₂), 144; Dichloroethane (CH ₂ Cl-CH ₂ Cl), 144; Ethylene Chlorohydrin (CH ₂ Cl-CH ₂ OH), 144.	

CONTENTS

CHAPTER XIII

Page

THE CHLORINATED HYDROCARBONS..... 147

Carbon Tetrachloride (CCl_4) (Tetrachloromethane), 143; Occupational Uses, 143; Pathology, 143; Signs and Symptoms, 149; Laboratory Tests, 150; Possible Disability, 151; Treatment, 152; Preface to Case Histories, 154; Comment, 156; Trichlorethylene (CHClCHCl_2), 158; Pathology, 158; Signs and Symptoms, 159; Laboratory Tests, 160; Possible Disability, 160; Treatment, 160; Tetrachlorethane ($\text{CHCl}_2\text{CHCl}_2$), 162.

CHAPTER XIV

CHLORINATED NAPHTHALENES AND DIPHENYLS..... 165

Occupational Exposure, 165; Pathology, 165; Signs and Symptoms, 166; Laboratory Tests, 166; Disability, 166; Treatment, 166; Naphthalene, 169.

CHAPTER XV

MISCELLANEOUS ALIPHATIC COMPOUNDS..... 171

Carbon Disulfide, 171; Occupational Hazards, 171; Résumé of Group Study, 171; Pathology, 172; Signs and Symptoms, 174; Differential Diagnosis, 176; Medicolegal Aspects, 178; Treatment, 178; Case History, 179; Sulfur Compounds, 179; Hydrogen Sulfide, 179; Occupational Hazards, 179; Pathology, 180; Signs and Symptoms, 180; Medicolegal Aspects, 180; Treatment, 180; Sulfur Dioxide, 182; Occupational Exposure, 182; Signs and Symptoms, 182; Medicolegal Aspects, 182; Treatment, 183; Illustrative Cases, 183.

CHAPTER XVI

HALOGENS AND HALOGEN COMPOUNDS..... 185

Chlorine, 185; Signs and Symptoms, 185; Treatment, 185; Fluorine, 186; Hydrogen Chloride, 187; Hydrogen Fluoride, 187; Case Report, 189.

CHAPTER XVII

THE AROMATIC HYDROCARBONS..... 190

Benzol, 190; Occupational Exposure, 190; Pathology, 190; Blood Picture, 193; The "Latent Period" of Benzol Poisoning, 200; Summary, 201; Signs and Symptoms, 201; Laboratory Tests, 202; Possible Disability, 202; Treatment, 202; Case Histories, 204; Toluene, 205; Xylol, 208; Solvent Naphtha, 208; Trinitrotoluene, 208; Laboratory Tests, 210; Treatment, 211; Case History, 211; Tetryl, 214; Nitrobenzene, 214; Aniline, 215; Phenol, 215.

CHAPTER XVIII

THE OXIDES OF NITROGEN..... 218

Occupational Exposure, 218; Pathology, 218; Signs and Symptoms, 219; Treatment, 220; Case Histories, 221; Ammonia, 221; Signs and Symptoms, 222; Medicolegal Aspects, 222; Treatment, 222; Case History, 223.

CONTENTS

CHAPTER XIX

Page

DDT (2,2-Bis-(P-CHLOROPHENYL)-1,1,1-TRICHLOROETHANE), 2,4-DICHLOROPHENOXY ACETIC ACID, AND THIOLGLYCOLIC ACID AND THIOLGLYCOLATES.....	224
DDT or 2,2-bis-(P-Chlorophenyl)-1,1,1-Trichloroethane. 224; 2,4-Dichlorophenoxy Acetic Acid, 228; Thioglycolic Acid and Thioglycolates, 231.	

PART THREE

THE METALS

CHAPTER XX

LEAD	234
Occupational Exposure, 234; Absorption and Excretion of Lead, 234; The Fate of Lead in the Body, 235; Pathology, 236; Signs and Symptoms, 240; The Basis for Diagnosis, 249; Differential Diagnosis, 250; Sequelae of Lead Poisoning, 251; Treatment, 253; Treatment in Lead Palsy, 257; Treatment in Lead Encephalopathy, 257; Prophylaxis, 257; Case Histories, 258.	

CHAPTER XXI

CADMIUM	265
Occupational Exposure, 265; Pathology, 265; Signs and Symptoms, 266; Laboratory Tests, 266; Disability, 266; Treatment, 267; Inhalations, 267; Case Histories, 268.	

CHAPTER XXII

BERYLLIUM	277
Occupational Exposure, 277; Etiology, 278; Pathology, 278; X-Ray Findings, 281; Signs and Symptoms, 284; Differential Diagnosis, 284; Treatment, 284; Case History, 284.	

CHAPTER XXIII

ARSENIC	289
Occupational Hazards, 289; Pathology, 289; General Discussion, 289; Arsine or Hydrogen Arsenide, 291; Differential Diagnosis, 292; Tests, 292; Medicolegal Aspects, 292; Treatment, 292.	

CHAPTER XXIV

MANGANESE, MAGNESIUM, PHOSPHORUS, ANTIMONY, AND ZINC	295
Manganese, 295; Occupational Exposure, 295; Pathology, 295; Effect on Blood, 297; Absorption and Elimination of Manganese, 297; Signs and Symptoms, 298; Differential Diagnosis, 298; Tests, 298; Treatment, 299; Magnesium, 302; Occupational Exposure, 302; Pathology, 302; Treatment, 304; Phosphorus, 304; Antimony, 306; Occupational Exposure, 306; Pathology, 306; Zinc, 307; Chronic Poisoning, 307; Medicolegal Aspects, 308; Treatment, 308; Zinc Chloride Burns, 308.	

CONTENTS

Page
2,4-DICHLOROPHE-
NOLGLYCOLATES..... 224

224; 2,4-Dichlo-
loglycolates, 231.

..... 234

of Lead, 234; The
d Symptoms, 240;
Sequelae of Lead
y, 257; Treatment
Histories, 258.

..... 265

d Symptoms, 266;
Inhalations, 267;

..... 277

278; X-Ray Find-
ings, 284; Treat-

..... 289

Discussion, 289;
s, 292; Tests, 292;

..... 295

y, 295; Effect on
e, 297; Signs and
; Treatment, 299;
y, 302; Treatment,
posure, 306; Path-
ological Aspects, 308;

CHAPTER XXV

MERCURY, CHROMIUM, COPPER, AND ALUMINUM..... 310

Mercury, 310; Occupational Exposure, 310; Pathology, 310; Signs and
Symptoms, 310; Laboratory Tests, 311; Treatment, 311; Case Histories,
312; Chromium, 314; Occupational Hazards, 314; Signs and Symptoms,
314; Medicolegal Aspects, 315; Treatment, 315; Copper, 316; Aluminum,
316.

CHAPTER XXVI

SELENIUM, TELLURIUM, VANADIUM, AND RADIUM..... 317

Selenium, 317; Occupational Exposure, 317; Pathology, 317; Signs and
Symptoms, 317; Treatment, 319; Tellurium, 321; Occupational Exposure,
321; Pathology, 321; Vanadium, 322; Vanadium Pentoxide as Toxic Prod-
uct, 322; Signs and Symptoms, 323; Radium, 323; Alpha-Rays, 324; Beta-
Rays, 324; Gamma-Rays, 325; Pathology, 325; Signs and Symptoms, 325;
Treatment, 325.

CHAPTER XXVII

METAL FUME FEVER..... 329

Pathology, 329; Immunity, 329; Signs and Symptoms, 329; Medicolegal
Aspects, 330; Treatment, 330.

PART FOUR

THE DUSTS

CHAPTER XXVIII

SILICOSIS..... 333

Silica in Industries, 333; Asbestos and Talc, 334; Mica, 335; Sodium
Silicate, 335; Silica Gel, 335; Lime-Soda Glass, 335; Definition of Silicosis,
336; Size of the Dust Particle, 336; Permissible Concentrations, 336;
Length of Exposure, 337; Diagnosis, 337; Signs and Symptoms, 337; Path-
ology, 342; Spread of Fibrosis, 342; Alveolar Consolidation, 343; Produc-
tion of Dyspnea, 344; Silico-Tuberculosis, 344; Other Complications, 345;
Additional Aids in Diagnosis, 346; Disability, 346; Employability of the
Person With Silicosis, 351; Treatment, 352; Results of Treatment, 353;
Case Reports, 355; Chemistry of Minerals, 357.

CHAPTER XXIX

ANTHRACOSILICOSIS..... 361

Findings in Governmental Survey, 361; Symptoms, 361; Basis of Diagnosis,
361; Groups Studied, 362; Prevalence and Exposure, 362; Safe Limits of
Exposure, 363; Respiratory Disease, 363; Signs and Symptoms, 364; Path-
ology, 364; X-Ray Findings, 366; Medicolegal Aspects, 367.

CONTENTS

	Page
CHAPTER XXX	
ASBESTOSIS	368
Signs and Symptoms, 368; Pathology, 368; Action of Asbestos and Silica, 368; Microscopic Appearance, 372; X-Ray Findings, 372; Medicolegal Aspects, 372.	
CHAPTER XXXI	
SILICIOUS DUSTS OF UNDETERMINED PATHOLOGIC STATUS	373
Talc, 373; Bauxite, 375; Diatomaceous Earth, 382; Tripoli, 384.	
CHAPTER XXXII	
THE INERT DUSTS	385
Siderosis, 387; Byssinosis, 389; Cement Dust, 394; Calcium and Magnesium Carbonates, 396; Carbon Dust, 396; Tobacco Dust, 396; Mill Dust, 396; Discharge of Dust Into Air, 396; Grain or Malt Fever, 397; Mica Dust, 398.	
CHAPTER XXXIII	
TUBERCULOSIS AND PNEUMONIA IN INDUSTRY	402
Respiratory Irritants: Dusts, Fumes, Gases, 403; Lead Absorption Versus Tuberculosis, 405; Temperature and Humidity Versus Tuberculosis, 406; Tuberculosis Among Steel Plant Workers, 406; Tuberculosis in the Foundry Industry, 407; Tuberculosis in Granite and Marble Workers, 407; Tuberculosis in Hard Rock Miners, 408; Fatigue as a Factor in Industrial Tuberculosis, 408; Summary, 409; Industrial Environment and Pneumonia, 411.	
CHAPTER XXXIV	
THE DERMATOSES	413
Occupational Hazards, 413; Definition, 413; General Considerations of Etiology, 413; Diagnosis, 414; Epidermophytosis, 419; Infectious Eczematoid Dermatitis, 420; Turpentine Dermatitis, 421; Baker's Dermatitis, 421; Cosmetician's and Barber's Dermatitis, 423; Cement Dermatitis, 423; Dermatitis Artefacta, 424; Onychia, 426; Syphilis, 426; Medicolegal Aspects, 427; Treatment, 427; Infectious Eczematoid Dermatitis, 432.	
CHAPTER XXXV	
OXYGEN THERAPY IN OCCUPATIONAL MEDICINE	434

CONTENTS

Page	Page
..... 363	CHAPTER XXXVI 433
estros and Silica, 72; Medicolegal	SYNTHETICS
..... 373	Synthetic Rubber, 438; Synthetic Fibers, 446; Rayon, 446; Nylon, 448; Cellophane, Sealing Caps, 450; Cellulose Sponge, 450; The Plastics, 451; Thermosetting Resins, 452; Phenolic Resins, 452; Urea-Formaldehyde Resins, 453; Melamine Resins, 453; Alkyd Resins, 454; Thermoplastics Based on Cellulose, 454; Cellulose Ester Plastics, 454; Cellulose Acetate, 455; Ethyl Cellulose Plastics, 455; Thermoplastic Resins, 456; Acrylic Resins, 456; Vinyl Resins, 457; Vinyl Alcohol and Acetal Resins, 457; Vinylidene Chloride Resins, 457; Styrene Resins, 458; Protein Plastics, 458; Casein Plastics, 458.
..... 384	CHAPTER XXXVII
..... 386	SPECIAL INDUSTRIAL PROCESSES 461
and Magnesium Mill Dust, 396; Mica Dust, 398.	Electroplating, 461; Anodic Treatment, 465; Degreasing, 467; Recom- mendations, 471; Pickling, 472; Heat Treatment, 473; Sand Blasting, 475; Shot Blasting, 476; Tumbling, 476; Casting (Molding), 477; The Ceramic and Pottery Processes, 477; Diesel Engines, 478; The Fluorescent Lamp Process, 479; Storage Battery Manufacturing Operations, 479; Welding, 481; Oxyacetylene Welding, 481; The Oxyacetylene Flame, 482; Electric Arc Welding, 484; The Metallic Arc Process, 485; Shielded and Unshielded Arc Welding Processes, 485; Electric Resistance Welding, 486; Hazards of Welding, 488; Composition of Welding Rods, 489; Metallizing, 490.
..... 402	CHAPTER XXXVIII
bsorption Versus uberculosis, 406; osis in the Foun- e Workers, 407; eter in Industrial t and Pneumonia,	INDUSTRIAL HYGIENE—GENERAL 493
..... 413	General Plant Hygiene, 494; Plant Lighting, 495; Natural Lighting, 495; Artificial Lighting, 496; Functional Use of Color, 499; Explosion Hazards of Dusts, Gases, and Vapors, 500; Gases, Vapors, Liquids, 500; Respiratory Protection, 502; Gas Masks and Cartridge Respirators, 503; Filter Respira- tors, 504; Care of Equipment, 505; Respirators With Impervious Clothing, 505; Hose Masks, 505; Abrasive Blasting Protectors, 506; Oxygen Breathing Apparatus, 506; Air-Line Respirators, 507; Cleaning and Maintenance of Respiratory Devices, 508; Eye and Face Protection, 509; Impact Goggles, 509; Corrective Goggles, 510; Face Shields, 511; Acid Hoods, 511; Chem- ical Goggles, 512; Welding and Babbiting Masks, 512; Harmful Rays, 512; Preventing Skin Infections, 515; Skin Cleansers, 516; Cutting Oils, 516; Safety Clothing, 516.
Considerations of Infectious Ecze- ma's Dermatitis, t Dermatitis, 423; 426; Medicolegal Dermatitis, 432.	CHAPTER XXXIX
..... 434	INDUSTRIAL HYGIENE—GENERAL—CONT'D 517
	Control of Hazards, 517; Manner and Method of Exhausting, 513; Hoods, 520; Large Open Hoods, 520; Lateral Exhaust System, 521; Flexible Ex- haust System, 521; Spray Booths, 521; Protection Against Corrosion, 522; Application and Design of Duct Exhaust Systems, 523; Duct Sizes, 527; Transport Velocities, 529; Conveying Capacity of Ducts, 529; Construc- tion, 530.

CONTENTS

	Page
CHAPTER XL	
INDUSTRIAL HYGIENE—GENERAL—CONT'D.	531
Appraisal of Hazards, 531; Dust Collection and Analysis, 531.	

CHAPTER XLI	
PRE-EMPLOYMENT EXAMINATIONS AND THE PLACEMENT PROGRAM.	542
The Job Placement Program, 544; Elements in Successful Placement, 546; Evaluating the Worker, 547; Evaluating the Job, 548; Psychiatry and Job Placement, 552.	

APPENDIX	
THE CHEMICALS IN COMMON TRADE-NAME PRODUCTS.	555

ILLUSTRATIONS IN COLOR

Fig.	Page
23. Bone marrow (femoral) (Photomicrograph)	198
24. Spleen (Photomicrograph)	198
25. Spleen (Photomicrograph)	198
26. Spleen (Photomicrograph)	198
27. Liver (Photomicrograph)	198
28. Lymph node (Photomicrograph)	198
29. Blood cells of anemia of lead poisoning	233

CHAPTER XXX

ASBESTOSIS

Asbestosis is a comparatively new disease. The first comprehensive reports regarding asbestosis came from the English writers in the years between 1920 and 1925. In America in the early thirties studies were made by Lanza, Sayers, Bloomfield, and others.

Asbestos is a hydrated magnesium silicate. Occupational exposure occurs in those trades where it is used for packing, insulating, or fire-proofing, or where it is combined with cotton or other material in textile processes.

Signs and Symptoms.—As in silicosis,¹ the first sign is dyspnea, the cough is usually dry, and auscultation reveals few, if any, physical signs. Loss of weight is usually noticeable, the color is pale, and cyanosis is a fairly early sign. After some time, fine, crackling râles may be heard, and the chest appears emaciated and lacks the robust character noticed in silicosis. Early asbestosis must be differentiated from acute emphysema. If tuberculosis is present, the symptoms may take on the character of this disease, but this complication is far less frequent than in silicosis.

Pathology.—

PULMONARY CHANGES.—The pulmonary fibrosis of asbestosis is diffuse in character, peribronchial, and basal in location in contrast to silicosis in which the fibrosis is nodular in character and present in the upper part of the lungs. As the process continues, the fibrosis extends into almost all portions of the lung tissue. Bronchiolectasis and bronchiectasis are present within the substance of the fibroid areas, and bronchopneumonia and acute tracheobronchitis often cause the death of these patients. The pleurae, especially in the basal regions, are thickened and adherent. In some cases the pleural sac is completely obliterated; in others a fibrous or serous exudate may be present within it.

Action of Asbestos and Silica.—The difference between chrysotile asbestos and quartz silica in their mode of action has caused considerable speculation as well as experimentation. Gardner² particularly, at the Saranac Laboratory, carried on intensive study of these two materials. He felt that while the action of free silica is chemical in nature, the action of asbestos is mechanical. He stated:

"We have come to the conclusion that inhaled asbestos fibres are irritating not because they are silicates but because they are stiff fibres which mechanically irritate the lungs. Unlike the free silicas, these minerals will not stimulate fibroblasts in any part of the body; only those in the lungs are affected. It was inferred that these organs were

affected because and continuous if asbestos was that 2 microns. It is likely destroyed. The fibres were carried developed in spite of 125 million per contrast in a period of long fibre as two years."

If the effect decrease in size particles have it becomes more. With fibrous as

The histology. Even under the injection there is of leucocytes in. The connective the asbestos fibre contracts, the in animals, at least exposure to the reason is the of surface of the in bodies" (Figs. 1

Finally, it is minerals only they are fibrous. The pulmonary irritation group is greatly fact, chrysotile silicate, serpent would seem to composition of

One point variations; namely conclusion has known that although in form extremely fine. in American p

affected because the movements of respiration are so much more rapid and continuous than those of other viscera. Then it was discovered that if asbestos was ground very finely so that few of the fibres were longer than 2 microns in length the irritating property of the asbestos was practically destroyed. Inhalation experiments with such fine chrysotile asbestos were carried out over a period of several years. No fibrosis has developed in spite of the fact that an average atmospheric concentration of 125 million particles per cubic foot of air has been maintained. In contrast in a previously reported experiment one-third this concentration of long fibre asbestos dust produced well marked fibrosis after about two years."

If the effect of asbestos were chemical, one would expect that a decrease in size would accelerate tissue response. With free silica large particles have little effect, but as their size decreases cellular reaction becomes more vigorous and even constitutional symptoms may ensue. With fibrous asbestos the reverse is true.

The histology of early asbestosis does not suggest a chemical injury. Even under the most extreme conditions that can be created by artificial injection there is no preliminary phase of tissue necrosis with infiltration of leucocytes as occurs with high concentrations of very fine quartz. The connective tissue cells merely multiply very slowly in areas where the asbestos fibres are caught in the bronchioles. As collagen forms and contracts, the air spaces are obliterated by scar tissue. In experimental animals, at least, this change is not a progressive one after cessation of exposure to the dust as is the case in the response to quartz. Perhaps the reason is the deposition of the peculiar iron-containing coating on the surface of the inhaled fibres giving rise to the characteristic "asbestosis bodies" (Figs. 50 and 51).

Finally, it is most suggestive that among dozens of different silicate minerals only the five known as asbestos, which are unique because they are fibrous in structure, should be commonly recognized as pulmonary irritants. The variation in chemical composition within this group is greater than that between them and many other silicates. In fact, chrysotile asbestos has the same chemical formula as a nonfibrous silicate, serpentine, which is physiologically inert. Obviously irritation would seem to be associated with the physical rather than the chemical composition of these minerals.

One point of practical significance may be indicated by these observations: namely, that very finely ground asbestos is not dangerous. This conclusion has support in clinical observation, for it has long been known that at the Thetford Mills there was no clinical asbestosis even though in former years the atmosphere was very dusty and the dust was extremely fine. The fact that fabrication of fibres of the same mineral in American plants could produce disease was one of the puzzling fea-




Fig. 50.—Right lung, showing markedly thickened pleura with fusion of interlobar pleura. Dense fibrous tags over entire surface. (U. S. Public Health Service, Bulletin 241.)




Fig. 51.—Asbestosis bodies accompanied by giant cells. (U. S. Army Medical Museum, Specimen 52255.)



Fig. 32.—Section of a lung in asbestosis. (Gardner.)




Fig. 33.—X-ray film of a patient with asbestosis. (Fendergrass.)

tures of this disease. But these experiments offer a plausible explanation. The fine dust in the mills is composed of serpentine and extremely short chrysotile fibres; that in the spinning and weaving mills contains many more long fibres.

Microscopic Appearance.—In the early phases of the disease there is a thickening of the alveolar septa as a result of fibroblastic proliferation. The alveolar spaces contain numerous alveolar phagocytes. With progression of the disease, fibrosis becomes more marked and the alveolar structure gradually disappears, and in its place there is now dense fibrous tissue. The few remaining alveoli which lie in the area of fibrous tissue are lined by a low cuboidal epithelium and have a glandular appearance (Fig. 52).

Scattered throughout the lung in both the diseased and healthy parts are spindle-shaped structures (first described by McDonald²) known as asbestosis bodies. These bodies are slender in their center and bulbous at each extremity. Many are arranged like strings of graduated beads with the largest at the end of the chain.

X-Ray Findings.—Certain peculiarities exist here as contrasted to silicosis. In asbestosis, the lesions may be bilateral or largely unilateral. According to the experience of Pendergrass,³ the roentgenologic findings in asbestosis are largely limited to the lower half or two-thirds of the lung fields, while in silicosis the upper portions of the lungs are also involved. In moderately advanced asbestosis, x-rays will reveal lessened ventilation of the lung fields, the parietal pleura is thickened, and there is a "ground-glass" appearance to the picture. The vascular shadows lose their identity. Nodulation is absent. Pendergrass³ feels that a roentgenologic diagnosis of early asbestosis is unreliable and that the condition must be moderately or markedly advanced to make a differential diagnosis from the roentgenograms (Fig. 53).

Medicolegal Aspects.—It is felt that the time necessary to develop asbestosis is, on the average, from seven to nine years in a fairly high concentration; with a less severe concentration, from fifteen to twenty years. The allowable concentration is 10,000,000 particles per cubic foot of air, of a size between 0.5 and 5.

Once established, the disease is progressive, even after the cessation of exposure. The basis for diagnosis is similar to that given under Silicosis.

References

1. Lanza, A. J.: Asbestosis, J.A.M.A. 105: 368 (Feb.), 1936.
2. Gardner, L. U.: Recent Developments in Relation to Silicosis, Indust. Med. 9: 45 (Feb.), 1940.
3. Pendergrass, E. P.: Silicosis and Asbestosis, New York, 1938, Oxford University Press.

² McDonald, S.: Histology of Pulmonary Asbestosis, Brit. M. J., 2: 1025, 1927.

It has
(crystalline
disabling
been stud-
aily with
degree. I
of being c
As yet no
gation by
remain in

Male
be accom-
tremolite.
mineral,
dusting a

In 18
Valle² at-
noted the
stituted:
ers² tool
fibrosis;
morbidity
observat

"The
dyspnea.
The pro-
outlined
enlargem-
present
involved
cases the
areas an
product
numerous
present
and pne-
be pres-
therefor
Diffuse