

FOLLOW-UP STUDY OF CHRYSOTILE TEXTILE WORKERS: COHORT MORTALITY AND EXPOSURE-RESPONSE

Misty J. Hein ¹, Leslie T. Stayner ^{2,3}, Everett Lehman ⁴, John M. Dement ⁵

¹ Industrywide Studies Branch
Division of Surveillance, Hazard Evaluations and Field Studies
National Institute for Occupational Safety and Health
4676 Columbia Parkway, R-13
Cincinnati, Ohio 45226
USA
MHein@cdc.gov
513-841-4207 telephone
513-841-4486 fax

² Division of Epidemiology and Biostatistics
University of Illinois School of Public Health
Chicago, Illinois, USA

³ Risk Evaluation Branch
Education and Information Division
National Institute for Occupational Safety and Health
Cincinnati, Ohio, USA

⁴ Industrywide Studies Branch
Division of Surveillance, Hazard Evaluations and Field Studies
National Institute for Occupational Safety and Health
Cincinnati, Ohio, USA

⁵ Department of Community and Family Medicine
Division of Occupational and Environmental Medicine
Duke University Medical Center
Durham, North Carolina, USA

Keywords: asbestos; asbestos, serpentine; asbestosis; respiratory tract diseases; cohort studies

The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, an exclusive licence (or non exclusive for government employees) on a worldwide basis to the BMJ Publishing Group Ltd and its Licensees to permit this article to be published in OEM and any other BMJ PGL products to exploit all subsidiary rights, as set out in our licence (<http://oem.bmjournals.com/fora/licence.pdf>).

ABSTRACT

Objectives

This report provides an update of the mortality experience of a cohort of South Carolina asbestos textile workers.

Methods

A cohort of 3,072 workers exposed to chrysotile in a South Carolina asbestos textile plant (1916-1977) was followed up for mortality through 2001. Standardized mortality ratios (SMRs) were computed using U.S. and South Carolina mortality rates. A job exposure matrix provided calendar time dependent estimates of chrysotile exposure concentrations. Poisson regression models were fitted for lung cancer and asbestosis. Covariates considered included sex, race, age, calendar time, birth cohort, and time since first exposure. Cumulative exposure lags of 5 and 10 years were considered by disregarding exposure in the most recent 5 and 10 years, respectively.

Results

A majority of the cohort was deceased (64%) and 702 of the 1,961 deaths occurred since the previous update. Mortality was elevated based on U.S. referent rates for *a priori* causes of interest including all causes combined (SMR 1.33, 95% confidence interval (CI) 1.28-1.39); all cancers (SMR 1.27, 95% CI 1.16-1.39); esophageal cancer (SMR 1.87, 95% CI 1.09-2.99); lung cancer (SMR 1.95, 95% CI 1.68-2.24); ischemic heart disease (SMR 1.20, 95% CI 1.10-1.32); and pneumoconiosis and other respiratory diseases (SMR 4.81, 95% CI 3.84-5.94). Mortality remained elevated for these causes when South Carolina referent rates were used. Three cases of mesothelioma were observed among cohort members. Exposure-response modeling for lung cancer, using a linear relative risk model, produced a slope coefficient of $0.0198 \text{ (fiber-years/ml)}^{-1}$ (standard error 0.00496), when cumulative exposure was lagged 10 years. Poisson regression modeling confirmed significant positive relationships between estimated chrysotile exposure and lung cancer and asbestosis mortality observed in previous updates of this cohort.

Conclusions

This study confirms the findings from previous investigations of excess mortality from lung cancer and asbestosis and a strong exposure-response relationship between estimated exposure to chrysotile and mortality from lung cancer and asbestosis.

BACKGROUND

Asbestos is well recognized to be a cause of malignant and non-malignant respiratory diseases. However, a continuing debate exists over whether or not, and if so to what extent, the chrysotile form of asbestos is a cause of these diseases. Some have suggested that “pure” chrysotile may not in fact be carcinogenic and that respiratory cancer excesses that have been observed in studies of chrysotile exposed workers may be explained by trace tremolite contamination in commercially used chrysotile.[1] This speculation has been referred by some as the “amphibole hypothesis”. Others have argued against this hypothesis.[2, 3]

A recent study, which provides evidence against the amphibole hypothesis, is a 25-year longitudinal study of male workers ($n = 515$) at an asbestos plant in China, where exposure was to chrysotile with little tremolite contamination (< 0.001 percent).[4] An age- and smoking-adjusted relative risk of 8.1 (95% confidence interval (CI) 1.8-36.1) was observed for lung cancer among highly exposed workers (workers in poorly ventilated raw material and textile sections) relative to workers with low exposure to asbestos (office workers and workers in the well ventilated asbestos cement section) and two cases of malignant mesothelioma were observed.[4]

Another line of evidence against the amphibole hypothesis is the strong relationship between lung cancer and chrysotile observed in a cohort of textile workers in South Carolina.[5] The strong exposure-response relationships between chrysotile and lung cancer, in addition to pneumoconiosis and other respiratory diseases (including asbestosis), have persisted in updates of this cohort.[6-8] In addition, a case-control analysis of this cohort indicated that the relationship between chrysotile exposure and lung cancer was not confounded by exposures to mineral oil.[7] Predicted lifetime excess risks of lung cancer and asbestosis from exposure to chrysotile, based on extensive exposure-response modeling of this cohort, were estimated by Stayner et al.[8] Vital status follow-up was recently extended through 2001 for the cohort of South Carolina textile workers. The primary objectives of this study were to update mortality and exposure-response relations between chrysotile and lung cancer and asbestosis mortality. This study was approved by the National Institute for Occupational Safety and Health Human Subjects Review Board.

METHODS

The study plant, located in South Carolina, produced asbestos products beginning in 1896 and asbestos textile products beginning in 1909.[9] The plant exclusively used chrysotile fibers obtained from Quebec, British Columbia and Rhodesia; however, small amounts of crocidolite yarn were used to make woven tape or braided packing from the 1950s until 1975. The total quantity of crocidolite used was approximately 2,000 pounds compared to 6-8 million pounds per year of chrysotile during the same time period. Since the crocidolite was never carded, spun or twisted, and all weaving of crocidolite tapes was done wet on a single loom, the predominant exposure at the plant was to chrysotile. The cohort definition included all textile production departments including preparation, carding, spinning (ring and gang, mule), spooling, twisting, winding, braiding, rope, wick and cord, weaving, finishing, and quality control. The plant stopped using asbestos material by the end of 1977.

Cohort Description

The original cohort (n=1261) was defined as all white male workers employed in textile production operations for at least 1 month between 01/01/1940 and 12/31/1965, with vital status follow-up through 1975.[5] The cohort was expanded to include white and non-white males and white females (n=3022) and vital status follow-up was extended through 1990.[6, 7] For the current study, 21 non-white females were added to the cohort. Updated demographic information was available for 29 workers who were previously excluded from analysis. Therefore, the current study updates mortality for 3072 workers with vital status follow-up through 2001.

Cohort mortality

Follow-up methods in the original study and previous update of this cohort were applied to all workers, regardless of race or sex. For the present update, names of cohort members were submitted to the National Death Index (NDI) for determination of vital status from 01/01/1991 through 2001. NDI Plus provided underlying and contributing causes of death for deceased workers identified by the NDI. All deaths were coded according to the revision of the International Classification of Diseases (ICD) in effect at the time of death. Workers that were confirmed alive on 01/01/1979, with a valid (within assigned range) Social Security number and not shown to be deceased on the NDI were considered to be alive as of the study end date. Those lost to follow-up before 01/01/1979, with invalid Social Security numbers, and thus not able to be matched with the NDI, were considered "lost to follow-up".

The mortality experience of the cohort was analyzed using the National Institute for Occupational Safety and Health modified life table analysis system.[10, 11] Under this system, each cohort member accumulated person-years at risk (PYAR) for each year of life after January 1, 1940 or completion of the one month eligibility period (whichever was later) until the earliest of the following: the date of death for deceased cohort members, the date last observed for persons lost to follow-up or the study end date (12/31/2001).

Rate files created for the life table analysis system identify 92 cause of death categories for which rates begin in 1940 and extend through 2001.[12] The PYAR were stratified into five-year intervals by age and calendar time and then multiplied by the appropriate U.S. sex-, race-, and cause-specific mortality rates to calculate the expected number of deaths. The ratio of observed to expected number of deaths was expressed as the standardized mortality ratio (SMR). Expanded rate files that identify 119 cause of death categories, including asbestosis, were created for both U.S. and South Carolina referent populations.[12] In analyses using 119 cause of death categories, PYAR and observed deaths started to accumulate on January 1, 1960, when the rates were first available, or completion of the one month eligibility period, whichever was later.

We considered mortality from the following causes to be of *a priori* interest: mesothelioma; cancers of the digestive system, larynx, trachea/bronchus/lung, peritoneum, and pleura; non-malignant respiratory diseases including pneumoconiosis and asbestosis; and ischemic heart disease. Since a specific code for mesothelioma did not exist prior to the tenth revision of the ICD, we reviewed death certificates for any mention of mesothelioma in order to identify pleural and peritoneal mesothelioma deaths

during 1991-1998. Mesothelioma deaths occurring after 1998 were identified by ICD-10 code C45 in the NDI cause of death file.

Exposure assessment

Detailed work histories listing beginning and ending dates in departments and operations were available for each member of the cohort. A department-, operation-, and calendar year-specific job exposure matrix (JEM) was available to link with the detailed work histories to calculate cumulative exposure to chrysotile.[9] Chrysotile exposure concentrations (expressed as fibers longer than 5 micrometers per milliliter of air) were estimated using statistical modeling of nearly 6000 industrial hygiene sampling measurements taken over the period 1930-1975 and analyzed using phase contrast microscopy. Exposure concentrations were considerably higher prior to 1940, before engineering dust control measures were put into place. This JEM has been used in previous mortality studies and exposure-response analyses of this cohort.[5-8, 13]

Each day in the work history was assigned an exposure level based on the JEM and cumulative exposure was defined as the sum of the assigned exposure concentrations over all days worked. SMRs were calculated by cumulative exposure for cancer of the trachea, bronchus, and lung (hereafter referred to as lung cancer) and pneumoconiosis and other respiratory diseases. Cutpoints of cumulative exposure were selected to give six exposure strata with approximately equal numbers of deaths (cause-specific).

Internal exposure-response modeling

Since stratified SMRs are not directly comparable, rate ratios (RRs) for each cumulative exposure category relative to the lowest group, adjusted for sex, race, age (<50, 50-54...75-79, and ≥ 80 years), and calendar-year (<1970, 1970-1979, 1980-1989, and ≥ 1990), were obtained by Poisson regression. Exposure lags of 5 and 10 years were considered by disregarding exposures in the most recent 5 and 10 years, respectively.

More detailed Poisson regression analyses were conducted by treating cumulative exposure as a continuous variable. In these analyses, cumulative exposure was partitioned into 30 categories with approximately equal numbers of deaths (cause-specific) and modeled as a continuous variable using the mean exposure, weighted by PYAR, in each exposure category.[14] Birth cohort (<1900, 1900-1909, 1910-1919, 1920-1929, and ≥ 1930), was used, in the absence of smoking information for all cohort members, as a surrogate for smoking. Additional covariates considered included sex, race, age, calendar-year, and time since first exposure (TSFE; <20, 20-39, and ≥ 40 years).

For lung cancer, the underlying cause of death was used to define the response and the model form was based on a linear relative risk model that was similar to the 1986 U.S. Environmental Protection Agency lung cancer model.[15] In a previous analysis of this cohort, this model was found to best fit the lung cancer exposure-response.[8] Background incidence was modeled as a log (ln) linear function of the covariates (sex, race, birth cohort, age, and calendar year). Second-order interactions were assessed among the covariates in the baseline function. The predicted incidence rate (λ) was modeled as the product of the background incidence (λ_0) and a linear function of cumulative exposure, i.e., $\lambda = \lambda_0 \times (1 + \beta E_{10})$, where E_{10} represents cumulative exposure to chrysotile (fiber-years/ml, as measured by phase contrast microscopy) omitting any

exposure in the most recent 10 years (i.e., a lag of 10 years). Interactions between exposure and the baseline covariates were considered along with the interaction between exposure and TSFE.

For asbestosis, both underlying and contributing causes were used to define the response and background incidence was similarly modeled, but a power model ($\lambda = \lambda_0 \times (E + \alpha)^p$) was employed. In the power model, alpha (α) was solved iteratively by minimizing the model deviance. The power model was found to best fit the asbestosis exposure-response in a previous analysis of this cohort.[8] Cumulative exposure lags of 5 and 10 years were also considered. Interactions between exposure and the baseline covariates were considered along with the interaction between exposure and TSFE.

Restricted cubic spline and categorical models, the model deviance, and Akaike's Information Criterion were used to informally evaluate the continuous models.[16] Models with the smallest values of Akaike's Information Criterion were considered to be the best fitting models. Poisson regression models were fit using Epicure (version 2.11, Hirosoft Corporation).

RESULTS

Cohort mortality

The cohort of 3072 workers has been followed for 118,513 PYAR (Table 1). The additional eleven years of follow-up and the inclusion of non-white females to the cohort added 702 deaths and more than 13,000 PYAR. More than half of the cohort (63.8%) had died through 2001. Nearly 10% of the cohort, however, was considered lost to follow-up, mostly due to high rates of loss to follow-up among females. One hundred and twenty workers known to be dead as of December 31, 2001 were missing cause of death information. An additional 426 workers who died or were lost to follow-up prior to January 1, 1960 were excluded from analyses using South Carolina referent rates because rates were not available prior to 1960. Since the observed person-time and deaths among non-white females were too few to provide stable estimates for any cause of death, results for non-white females are only reported in analyses combining all females or all workers.

Table 1. Characteristics of the South Carolina asbestos textile workers cohort

	White males	White females	Non-white males	Non-white females	Total
Total number of workers	1256	1244	551	21	3072
Vital status through 12-31-2001					
Alive	373 (29.7%)	339 (27.3%)	129 (23.4%)	5 (23.8%)	846 (27.5%)
Dead, cause of death known	841 (67.0%)	647 (52.0%)	345 (62.6%)	8 (38.1%)	1841 (59.9%)
Dead, cause of death unknown	27 (2.1%)	50 (4.0%)	39 (7.1%)	4 (19.0%)	120 (3.9%)
Lost to follow-up	15 (1.2%)	208 (16.7%)	38 (6.9%)	4 (19.0%)	265 (8.6%)
Person-years at risk					
vital status through 2001 ^a	49,409.8	49,163.9	19,180.9	758.0	118,512.6
vital status through 1990 ^b	43,898.8	43,743.8	17,406.1	674.6	105,723.3
Age at death, years median (range)	64.5 (18.1 – 95.8)	73.4 (22.6 – 101.0)	61.0 (22.1 – 96.5)	66.2 (47.5 – 90.0)	67.1 (18.1 – 101.0)
Age at date last observed, years median (range)	66.6 (18.1 – 95.8)	74.1 (16.9 – 101.0)	63.3 (16.9 – 96.5)	72.8 (21.1 – 90.0)	68.8 (16.9 – 101.0)
Cumulative exposure, fiber-years/ml median (range)	4.4 (0.1 – 699.8)	4.2 (0.2 – 317.1)	14.5 (0.4 – 682.7)	5.9 (0.5 – 216.0)	5.5 (0.1 – 699.8)
Duration of employment, years median (range)	1.1 (0.1 – 46.8)	0.9 (0.1 – 43.7)	1.5 (0.1 – 43.8)	0.9 (0.1 – 30.9)	1.1 (0.1 – 46.8)

^a Length of follow-up through 2001. Person-years at risk was calculated from the later of (rate file begin date, date achieve 15 years of age, and date achieve 1 month of employment between January 1, 1940 and December 31, 1965) through the earlier of (date of death, study end date (31 December 2001), and date last observed to be alive).

^b Length of follow-up through 1990.

The distribution of duration of employment was strongly right skewed with approximately half of the workers employed for 1 year or less (i.e., “short-term workers”). Short-term workers were more likely to have been hired during the World War II years of 1939-1945. Most of the short-term and many of the long-term workers hired during these years did not continue working at the plant after the war.

Table 2 gives the results of the mortality analysis by race and sex based on the use of U.S. mortality rates. Mortality was elevated for all causes combined and all cancers. Among the *a priori* causes of interest, mortality from cancers of the digestive organs was not elevated, with the exception of excess mortality from esophageal cancer; mortality from cancers of the respiratory system was elevated, largely due to excess mortality from lung cancer; mortality from ischemic heart disease was elevated; and mortality from diseases of the respiratory system was elevated, particularly chronic obstructive pulmonary disease and pneumoconiosis and other respiratory diseases. These elevations persisted in subsequent analyses (not shown) in which South Carolina referent rates were used to generate expected numbers of deaths. When short-term workers were excluded, elevations in mortality from lung cancer (SMR 2.44, 95% CI 2.04-2.90) and pneumoconiosis and other respiratory diseases (SMR 7.07, 95% CI 5.49-8.97) persisted.

Table 2. Mortality through 2001 based on U.S. mortality rates for selected causes of death among workers in the South Carolina asbestos textile workers cohort

Underlying cause of death	White males		Non-white males		Females		All workers combined		
	O ^a SMR	95% CI	O SMR	95% CI	O SMR	95% CI	O	E SMR	95% CI
All deaths	868 1.52 **	1.42 - 1.62	384 1.10	0.99 - 1.22	709 1.29 **	1.20 - 1.39	1961	1469.9 1.33 **	1.28 - 1.39
All cancers	221 1.58 **	1.37 - 1.80	73 0.96	0.75 - 1.21	169 1.14	0.97 - 1.32	463	365.0 1.27 **	1.16 - 1.39
MN of buccal cavity and pharynx	4 1.14	0.31 - 2.90	2 0.83	0.10 - 3.00	2 1.08	0.13 - 3.90	8	7.8 1.03	0.44 - 2.03
MN of digestive organs and peritoneum	37 1.06	0.75 - 1.46	28 1.28	0.85 - 1.85	26 0.74	0.48 - 1.08	91	92.1 0.99	0.80 - 1.21
MN of esophagus	8 2.17	0.94 - 4.28	9 2.26 *	1.03 - 4.28	0		17	9.1 1.87 *	1.09 - 2.99
MN of stomach	4 0.79	0.22 - 2.02	9 1.94	0.88 - 3.67	0		13	13.3 0.97	0.52 - 1.67
MN of intestine except rectum	6 0.49	0.18 - 1.07	2 0.37	0.04 - 1.32	11 0.71	0.36 - 1.28	19	33.1 0.57 *	0.35 - 0.90
MN of rectum	2 0.66	0.08 - 2.38	2 1.54	0.19 - 5.55	5 1.77	0.57 - 4.13	9	7.2 1.26	0.57 - 2.38
MN of biliary passages and liver	7 2.10	0.84 - 4.32	1 0.40	0.01 - 2.23	2 0.55	0.07 - 1.98	10	9.5 1.05	0.51 - 1.94
MN of pancreas	9 1.28	0.58 - 2.43	5 1.34	0.43 - 3.12	7 0.93	0.37 - 1.91	21	18.3 1.15	0.71 - 1.75
MN of peritoneum and unspecified of digestive organs	1 1.79	0.05 - 9.94	0		1 1.38	0.03 - 7.69	2	1.6 1.28	0.15 - 4.60
MN of respiratory system	121 2.34 **	1.94 - 2.79	22 0.84	0.52 - 1.27	65 2.30 **	1.78 - 2.93	208	106.3 1.96 **	1.70 - 2.24
MN of larynx	4 2.20	0.60 - 5.63	1 0.79	0.02 - 4.36	1 2.08	0.05 - 11.6	6	3.6 1.68	0.61 - 3.66
MN of trachea, bronchus and lung	116 2.34 **	1.94 - 2.81	21 0.85	0.52 - 1.30	61 2.22 **	1.70 - 2.85	198	101.7 1.95 **	1.68 - 2.24
MN of pleura	0		0		0		0	0.3	
MN of other respiratory sites	1 3.25	0.08 - 18.1	0		3 12.9 **	2.65 - 37.6	4	0.7 5.49 *	1.49 - 14.0
MN of breast	0		0		28 0.99	0.66 - 1.43	28	28.6 0.98	0.65 - 1.42
MN of female genital organs	0		0		20 1.02	0.62 - 1.58	20	19.6 1.02	0.62 - 1.58
MN of male genital organs	12 1.10	0.57 - 1.92	8 0.79	0.34 - 1.56	0		20	21.0 0.95	0.58 - 1.47
MN of urinary organs	9 1.26	0.57 - 2.38	2 0.78	0.09 - 2.83	4 0.92	0.25 - 2.36	15	14.1 1.07	0.60 - 1.76
MN of other and unspecified sites	23 1.25	0.79 - 1.88	7 0.92	0.37 - 1.90	11 0.62	0.31 - 1.11	41	43.7 0.94	0.67 - 1.27
Neoplasms of lymphatic and hematopoietic tissue	15 1.11	0.62 - 1.84	4 0.78	0.21 - 2.00	13 0.98	0.52 - 1.67	32	31.9 1.00	0.69 - 1.42
Benign and unspecified neoplasms	1 0.54	0.01 - 3.01	2 2.09	0.25 - 7.53	4 1.46	0.40 - 3.74	7	5.5 1.26	0.51 - 2.60
Tuberculosis	6 1.62	0.59 - 3.52	6 0.75	0.27 - 1.62	0		12	14.7 0.82	0.42 - 1.43
Diabetes mellitus	18 1.79 *	1.06 - 2.84	8 1.08	0.47 - 2.14	18 1.24	0.74 - 1.97	44	31.9 1.38 *	1.00 - 1.85
Diseases of the blood and blood forming organs	0		2 1.84	0.22 - 6.64	2 0.86	0.10 - 3.10	4	5.2 0.77	0.21 - 1.97
Mental, psychoneurotic, and personality disorders	6 1.30	0.48 - 2.84	6 1.41	0.51 - 3.07	8 1.24	0.53 - 2.44	20	15.3 1.31	0.80 - 2.02
Disorders of the nervous system and sense organs	3 0.37	0.08 - 1.10	3 0.80	0.16 - 2.33	18 1.53	0.90 - 2.41	24	23.6 1.02	0.65 - 1.51
Diseases of the heart	295 1.38 **	1.23 - 1.55	101 0.95	0.77 - 1.15	198 1.12	0.97 - 1.29	594	495.7 1.20 **	1.10 - 1.30
Ischemic heart disease	250 1.39 **	1.22 - 1.58	70 0.92	0.72 - 1.17	149 1.11	0.94 - 1.30	469	390.1 1.20 **	1.10 - 1.32

Hypertension with heart disease	12	2.25 *	1.16 - 3.93	9	0.86	0.39 - 1.64	7	0.97	0.40 - 2.00	28	23.0	1.22	0.81 - 1.76
Other diseases of the heart	28	1.28	0.85 - 1.86	20	1.16	0.71 - 1.78	33	1.34	0.92 - 1.88	81	63.7	1.27 *	1.01 - 1.58
Other diseases of the circulatory system	65	1.46 **	1.12 - 1.86	42	1.11	0.80 - 1.50	78	1.25	0.99 - 1.56	185	145.0	1.28 **	1.10 - 1.47
Hypertension without heart disease	2	1.10	0.13 - 3.99	7	2.10	0.84 - 4.33	4	1.40	0.38 - 3.59	13	8.0	1.63	0.87 - 2.78
Cerebrovascular disease	49	1.70 **	1.25 - 2.24	28	1.03	0.69 - 1.49	54	1.19	0.90 - 1.56	131	101.3	1.29 **	1.08 - 1.53
Diseases of the respiratory system	84	2.00 **	1.60 - 2.48	33	1.42	0.98 - 2.00	72	1.67 **	1.30 - 2.10	189	108.4	1.74 **	1.50 - 2.01
Chronic obstructive pulmonary disease	29	1.41	0.94 - 2.02	6	0.90	0.33 - 1.96	26	1.36	0.89 - 1.99	61	46.4	1.31 *	1.01 - 1.69
Asthma	1	1.08	0.03 - 6.01	0			0			1	3.3	0.30	0.01 - 1.67
Pneumoconiosis and other respiratory diseases	41	5.67 **	4.07 - 7.69	12	3.02 **	1.56 - 5.28	32	4.94 **	3.38 - 6.98	85	17.7	4.81 **	3.84 - 5.94
Asbestosis ^b	20	172.5 **	105.3 - 266.4	4	129.5 **	35.3 - 331.2	12	1500 **	771.7 - 2611	36	0.2	232.5 **	162.8 - 321.9
Diseases of the digestive system	35	1.33	0.92 - 1.84	12	0.77	0.40 - 1.35	34	1.45 *	1.00 - 2.03	81	65.4	1.24	0.98 - 1.54
Cirrhosis and other chronic liver diseases	20	1.48	0.91 - 2.29	5	0.70	0.23 - 1.63	11	1.34	0.67 - 2.39	36	28.9	1.25	0.87 - 1.73
Diseases of the genitourinary system	13	1.66	0.88 - 2.84	8	0.89	0.38 - 1.75	16	1.46	0.84 - 2.38	37	27.8	1.33	0.94 - 1.84
Diseases of the skin and subcutaneous tissue	1	2.28	0.06 - 12.7	0			2	2.22	0.27 - 8.00	3	2.0	1.51	0.31 - 4.41
Diseases of the musculoskeletal system and connective tissue	0			0			3	1.05	0.22 - 3.08	3	4.8	0.63	0.13 - 1.84
Symptoms and ill-defined conditions	11	1.99	0.99 - 3.56	10	1.33	0.63 - 2.44	6	1.32	0.48 - 2.87	27	17.6	1.53 *	1.01 - 2.23
Accidents	52	1.45 *	1.08 - 1.90	25	1.11	0.72 - 1.64	13	0.88	0.47 - 1.51	90	73.1	1.23	0.99 - 1.51
Transportation accidents	28	1.43	0.95 - 2.06	10	0.99	0.48 - 1.83	7	1.10	0.44 - 2.27	45	36.0	1.25	0.91 - 1.67
Suicide	8	0.66	0.29 - 1.31	0			1	0.23	0.01 - 1.29	9	18.6	0.48 *	0.22 - 0.92
Homicide	5	1.47	0.48 - 3.43	7	0.55	0.22 - 1.13	1	0.87	0.02 - 4.83	13	17.3	0.75	0.40 - 1.28
Unknown cause	27			39			54			120			
Residual	17	1.70	0.99 - 2.72	7	0.78	0.31 - 1.61	12	0.85	0.44 - 1.49	36	33.0	1.09	0.76 - 1.51

^a O, observed number of deaths; SMR, standardized mortality ratio; CI, confidence interval; E, expected number of deaths (based on U.S. referent rates); MN, malignant neoplasm.

^b Observed deaths, SMRs and 95% CIs for asbestosis are for the time period 1960-2001 due to rate file restrictions.

* two-sided p-value < 0.05.

** two-sided p-value < 0.01.

Mortality was elevated from diabetes mellitus and diseases of the digestive system; these elevations persisted when South Carolina referent rates were used. Mortality was elevated based on U.S., but not South Carolina, referent rates for other diseases of the heart, other diseases of the circulatory system including cerebrovascular disease, and accidents. Reduced mortality was observed for cancer of the intestine (excluding the rectum) and suicide. Among white males, excess mortality was also observed for cirrhosis and other chronic liver diseases, particularly when South Carolina referent rates were used (SMR 1.88, 95% CI 1.15-2.91).

Mortality from asbestosis was highly elevated based on U.S. referent rates (observed 36, expected 0.15, SMR 232.5, 95% CI 162.8-321.9) and remained highly elevated based on South Carolina referent rates (expected 0.33, SMR 108.2, 95% CI 75.8-149.8). A manual review of the death certificates identified three mesothelioma deaths, all among white males. Two of these have been described in a previous update of this cohort (Dement et al., 1994). The third mesothelioma death occurred in 1995, nearly 50 years after the employee began working in the mule spinning department for approximately 2.5 years. None of the deaths under the tenth revision of the ICD were due to mesothelioma.

A mortality analysis among workers actively employed, which considered person-time and deaths from the person-year begin date through termination of employment at the plant, indicated reduced mortality overall (63 deaths, SMR 0.59, 95% CI 0.46-0.76), however, excess mortality was observed for pneumoconiosis and other respiratory diseases (7 deaths, SMR 8.14, 95% CI 3.26-16.8). Mortality in the first year post termination of employment was elevated overall (59 deaths, SMR 4.36, 95% CI 3.32-5.63) and for all cancers combined (18 deaths, SMR 7.87, 95% CI 4.66-12.4) with much of the elevation was due to cancers of the digestive organs (4 deaths, SMR 6.63, 95% CI 1.81-17.0), lung cancer (11 deaths, SMR 21.6, 95% CI 10.8-38.7), and pneumoconiosis and other respiratory diseases (6 deaths, SMR 46.4, 95% CI 17.0-101); all of these deaths were among workers with cumulative exposure to chrysotile in excess of 10,000 fiber-days/ml. All of the lung cancer and pneumoconiosis deaths in the first year post termination of employment occurred more than 10 years since first exposure and 3 (out of 4) digestive organ cancer deaths occurred more than 20 years since first exposure.

The results of the mortality analysis for lung cancer by cumulative exposure to chrysotile fibers are provided in Table 3. A majority (96%) of the 198 lung cancer deaths occurred 20 years or more after first exposure. Increasing trends in lung cancer SMRs with increasing cumulative exposure were observed for white males, non-white males, females and overall. Increasing and highly significant trends were also observed for lung cancer mortality in internal analyses using Poisson regression, which adjusted for sex, race, age and calendar-year. Results were similar and there was no improvement in model fit when a five- or ten-year lag was employed. RRs relative to the lowest cumulative exposure group were particularly elevated in the highest cumulative exposure category. The trend persisted when short-term workers were excluded from the analysis.

Table 3. Lung cancer mortality (underlying cause) through 2001 by cumulative exposure to chrysotile among workers in the South Carolina asbestos textile workers cohort

Group	Lag (years)	Cumulative exposure (fiber-years/ml) ^a						Trend P-value
		< 1.5	1.5 – < 5	5 – < 15	15 – < 60	60 – < 120	≥ 120	
White- males	0	O/E ^b	22 / 12.6	23 / 14.1	15 / 8.4	19 / 7.1	17 / 4.8	20 / 2.5
		SMR	1.74 (1.09 – 2.64)	1.63 (1.04 – 2.45)	1.79 (1.00 – 2.94)	2.69 (1.62 – 4.20)	3.56 (2.07 – 5.70)	7.85 (4.80 – 12.1)
		RR	1	0.94 (0.52 – 1.69)	1.03 (0.53 – 1.98)	1.52 (0.82 – 2.81)	2.00 (1.05 – 3.80)	4.43 (2.37 – 8.29)
Non-white- males	0	O/E	0 / 1.5	5 / 4.5	4 / 7.4	3 / 6.6	2 / 2.4	7 / 2.4
		SMR	0.00 (0.00 – 2.45)	1.11 (0.36 – 2.60)	0.54 (0.15 – 1.38)	0.45 (0.09 – 1.32)	0.84 (0.10 – 3.03)	2.96 (1.19 – 6.11)
		RR	1 ^c	1 ^c	0.63 (0.17 – 2.35)	0.54 (0.13 – 2.27)	1.08 (0.21 – 5.59)	3.99 (1.23 – 12.9)
Females	0	O/E	11 / 7.1	6 / 6.7	14 / 5.9	11 / 4.5	13 / 2.2	6 / 0.9
		SMR	1.54 (0.77 – 2.76)	0.89 (0.33 – 1.94)	2.35 (1.29 – 3.95)	2.43 (1.21 – 4.35)	5.82 (3.10 – 9.95)	6.64 (2.44 – 14.4)
		RR	1	0.59 (0.22 – 1.61)	1.51 (0.69 – 3.33)	1.52 (0.66 – 3.50)	3.40 (1.52 – 7.60)	3.84 (1.41 – 10.5)
Overall	0	O/E	33 / 21.3	34 / 25.3	33 / 21.8	33 / 18.2	32 / 9.4	33 / 5.8
		SMR	1.55 (1.07 – 2.18)	1.34 (0.93 – 1.88)	1.52 (1.04 – 2.13)	1.81 (1.25 – 2.54)	3.41 (2.33 – 4.81)	5.68 (3.91 – 7.98)
		RR	1	0.95 (0.59 – 1.54)	1.22 (0.75 – 1.99)	1.46 (0.90 – 2.37)	2.50 (1.53 – 4.08)	4.84 (2.94 – 7.97)
Overall	5	O/E	33 / 21.5	34 / 25.3	33 / 21.7	35 / 18.4	32 / 9.4	31 / 5.4
		SMR	1.53 (1.06 – 2.15)	1.34 (0.93 – 1.88)	1.52 (1.05 – 2.13)	1.90 (1.32 – 2.64)	3.41 (2.33 – 4.82)	5.78 (3.93 – 8.21)
		RR	1	1.00 (0.62 – 1.61)	1.28 (0.79 – 2.08)	1.59 (0.99 – 2.58)	2.61 (1.60 – 4.28)	5.14 (3.10 – 8.53)
Overall	10	O/E	34 / 22.1	33 / 25.3	34 / 21.7	35 / 18.8	37 / 9.2	25 / 4.7
		SMR	1.54 (1.07 – 2.15)	1.30 (0.90 – 1.83)	1.57 (1.08 – 2.19)	1.86 (1.30 – 2.59)	4.02 (2.83 – 5.53)	5.36 (3.47 – 7.92)
		RR	1	1.00 (0.62 – 1.62)	1.37 (0.85 – 2.21)	1.61 (1.00 – 2.59)	3.20 (1.99 – 5.14)	4.91 (2.88 – 8.35)
Long-term ^d	0	O/E	2 / 0.56	7 / 5.2	26 / 15.9	31 / 16.7	32 / 9.4	33 / 5.8
		SMR	3.56 (0.43 – 12.9)	1.34 (0.54 – 2.75)	1.64 (1.07 – 2.40)	1.85 (1.26 – 2.63)	3.41 (2.33 – 4.81)	5.68 (3.91 – 7.98)
		RR	1 ^e	1 ^e	1.14 (0.53 – 2.44)	1.39 (0.66 – 2.95)	2.39 (1.13 – 5.06)	4.62 (2.17 – 9.84)

^a Cutpoints of cumulative exposure were selected to approximately divide the deaths from lung cancer into six cumulative exposure groups. The numbers of workers in each cumulative exposure group were 705, 756, 628, 524, 264, and 195, respectively; the person-years at risk in each cumulative exposure group were 26667, 29188, 24449, 20561, 10295, and 7352, respectively.

^b O/E, ratio of observed to expected (based on U.S. referent rates) number of deaths; SMR, standardized mortality ratio from life-table analyses adjusted for sex, race, age and calendar year (95% confidence interval); RR, rate ratio relative to lowest cumulative exposure group from Poisson regression analyses adjusted for sex, race, age and calendar year (95% confidence interval).

- ^c Reference category for non-white males was cumulative exposure < 5 fiber-years/ml due to zero deaths in the < 1.5 fiber-years/ml category.
- ^d Workers with 1 year of more of exposure.
- ^e Reference category for long-term workers was cumulative exposure < 5 fiber-years/ml due to few deaths (n=2) in the < 1.5 fiber-years/ml category.

Results of the mortality analysis for the cause of death category 'pneumoconiosis and other respiratory diseases' (which includes asbestosis) by cumulative exposure to chrysotile fibers are provided in Table 4. Of the 85 observed deaths in this category, 42 were from asbestosis. Increasing trends in pneumoconiosis SMRs with increasing cumulative exposure were observed for white males, non-white males, females and overall. Increasing and highly significant trends were also observed for pneumoconiosis mortality in internal analyses using Poisson regression. Results were similar and there was no improvement in model fit when a five- or ten-year lag was employed. RRs relative to the lowest cumulative exposure group were particularly elevated in the highest cumulative exposure category. The trend persisted when short-term workers were excluded from the analysis.

Table 4. Pneumoconiosis and other respiratory diseases mortality (underlying cause) through 2001 by cumulative exposure to chrysotile among workers in the South Carolina asbestos textile workers cohort

Group	Lag (years)	Cumulative exposure (fiber-years/ml) ^a						Trend P-value
		< 3	3 – < 16	16 – < 60	60 – < 100	100 – < 150	≥ 150	
White- males	0	O/E ^b 9 / 3.0	4 / 2.1	6 / 1.0	7 / 0.6	4 / 0.3	11 / 0.3	
		SMR 3.05 (1.39 – 5.78)	1.92 (0.52 – 4.92)	6.10 (2.24 – 13.3)	11.5 (4.64 – 23.8)	12.3 (3.34 – 31.4)	38.9 (19.4 – 69.6)	< 0.0001
		RR 1	0.64 (0.20 – 2.07)	1.90 (0.68 – 5.36)	3.62 (1.32 – 9.93)	3.88 (1.17 – 12.9)	12.1 (4.86 – 30.4)	< 0.0001
Non-white- males	0	O/E 0 / 0.6	3 / 1.5	2 / 1.0	1 / 0.3	3 / 0.2	3 / 0.3	
		SMR 0.00 (0.00 – 5.96)	1.94 (0.40 – 5.68)	1.94 (0.23 – 7.00)	3.32 (0.08 – 18.5)	14.4 (2.97 – 42.1)	11.2 (2.30 – 32.6)	0.0003
Females	0	O/E 5 / 2.5	7 / 2.1	5 / 1.1	6 / 0.3	5 / 0.4	4 / 0.1	
		SMR 2.02 (0.66 – 4.72)	3.29 (1.32 – 6.78)	4.67 (1.52 – 10.9)	17.4 (6.39 – 37.9)	14.2 (4.61 – 33.1)	36.6 (9.97 – 93.7)	< 0.0001
		RR 1	1.69 (0.54 – 5.34)	2.47 (0.71 – 8.54)	9.02 (2.73 – 29.8)	8.21 (2.36 – 28.6)	21.4 (5.65 – 81.0)	< 0.0001
Overall	0	O/E 14 / 6.0	14 / 5.8	13 / 3.1	14 / 1.3	12 / 0.9	18 / 0.7	
		SMR 2.32 (1.27 – 3.89)	2.43 (1.33 – 4.08)	4.21 (2.24 – 7.20)	11.2 (6.11 – 18.8)	13.5 (6.99 – 23.7)	27.2 (16.1 – 43.1)	< 0.0001
		RR 1	1.20 (0.57 – 2.52)	2.14 (1.00 – 4.58)	5.61 (2.65 – 11.9)	6.89 (3.15 – 15.1)	15.6 (7.51 – 32.5)	< 0.0001
Overall	5	O/E 15 / 6.2	14 / 5.7	14 / 3.1	15 / 1.3	9 / 0.8	18 / 0.6	
		SMR 2.41 (1.35 – 3.97)	2.46 (1.35 – 4.16)	4.54 (2.48 – 7.62)	11.9 (6.68 – 19.7)	10.8 (4.93 – 20.5)	30.1 (17.8 – 47.6)	< 0.0001
		RR 1	1.22 (0.59 – 2.54)	2.29 (1.10 – 4.79)	5.99 (2.89 – 12.4)	5.68 (2.45 – 13.2)	17.4 (8.47 – 35.9)	< 0.0001
Overall	10	O/E 15 / 6.5	14 / 5.6	16 / 3.1	16 / 1.2	10 / 0.8	14 / 0.5	
		SMR 2.32 (1.30 – 3.83)	2.49 (1.36 – 4.18)	5.17 (2.96 – 8.40)	13.0 (7.42 – 21.1)	13.0 (6.21 – 23.8)	27.4 (15.0 – 45.9)	< 0.0001
		RR 1	1.38 (0.66 – 2.89)	2.88 (1.40 – 5.90)	7.29 (3.52 – 15.1)	7.84 (3.44 – 17.9)	17.5 (8.11 – 37.9)	< 0.0001
Long-term ^c	0	O/E 1 / 0.33	10 / 3.5	13 / 2.9	14 / 1.3	12 / 0.9	18 / 0.7	
		SMR 2.99 (0.08–16.7)	2.83 (1.36 – 5.21)	4.41 (2.35 – 7.54)	11.2 (6.11 – 18.8)	13.5 (6.99 – 23.7)	27.2 (16.1 – 43.1)	< 0.0001
		RR 1 ^d	1 ^d	1.75 (0.78 – 3.92)	4.54 (2.04 – 10.1)	5.55 (2.41 – 12.8)	13.1 (5.95 – 28.8)	< 0.0001

^a Cutpoints of cumulative exposure were selected to approximately divide the deaths from pneumoconiosis and other respiratory diseases into six cumulative exposure groups. The numbers of workers in each cumulative exposure group were 1125, 997, 491, 181, 155, and 123, respectively; the person-years at risk in each cumulative exposure group were 43524, 38133, 19209, 6959, 6097, and 4591, respectively.

^b O/E, ratio of observed to expected (based on U.S. referent rates) number of deaths; SMR, standardized mortality ratio from life-table analyses adjusted for sex, race, age and calendar year (95% confidence interval); RR, rate ratio relative to lowest cumulative exposure group from Poisson regression analyses adjusted for sex, race, age and calendar year (95% confidence interval).

^c Workers with 1 year of more of exposure.

^d Reference category for long-term workers was cumulative exposure < 16 fiber-years/ml due to few deaths (n=1) in the < 3 fiber-years/ml category.

Exposure-response modeling

Lung cancer mortality (198 observed deaths) was modeled as a continuous function of exposure using the linear relative risk model. When covariates were included one-at-a-time in the background function, mortality was lower for females and non-whites, increased with age and calendar time, but decreased with birth cohort (not shown). Model fit was improved when sex, race, and age were included in the baseline function, but not when either calendar time or birth cohort was additionally included in the model. Interactions among covariates in the baseline model did not improve model fit; consequently, the baseline function included terms for sex, race, and age. Cumulative exposure, lagged 10 years, was highly significant based on the likelihood ratio test. Interactions with cumulative exposure and the baseline covariates did not improve model fit; likewise, an interaction with cumulative exposure and TSFE also did not improve model fit. The model for lung cancer mortality is described in Table 5 and graphically displayed, along with the categorical and restricted cubic spline models, in Figure 1. Since the baseline function included categorical terms for sex, race, and age, the graph depicts the predicted mortality rate for, as an example, white males 60-64 years of age. Predicted curves for other groups can be constructed using the estimated parameters in Table 5. While the model was developed using the entire cohort, the graph only displays cumulative exposures through 200 fiber-years/ml, rather than the maximum of 700 fiber-years/ml since only 1% of the observed person-time is above 200 fiber-years/ml. With very little person-time and furthermore, very few people with person-time in the range of 200-700 fiber-years/ml, modeling cannot provide much information about the shape of the exposure-response in this range, particularly for a specific cause of death. The coefficient of cumulative exposure was $0.0198 \text{ (fiber-years/ml)}^{-1}$ based on a 10 year lag; coefficients were similar, but slightly reduced, when cumulative exposure was not lagged (0.0165) or lagged by only 5 years (0.0176). Cumulative exposure was positively and significantly associated with lung cancer mortality in all models tested.

Table 5. Model for lung cancer mortality as an underlying cause of death among workers in the South Carolina asbestos textile workers cohort ^a

Term in model			Estimate	Standard error	Likelihood ratio test ^b
Intercept		β_0	-8.72	0.358	---
Sex	Male	Ref	---	---	43.5, 1, <0.0001
	Female	β_1	-1.02	0.161	
Race	White	Ref	---	---	23.3, 1, <0.0001
	Non-white	β_2	-1.03	0.238	
Age	< 50 years	Ref	---	---	253.8, 7, <0.0001
	50-54 years	β_3	2.05	0.445	
	55-59 years	β_4	2.66	0.410	
	60-64 years	β_5	3.27	0.392	
	65-69 years	β_6	3.61	0.388	
	70-74 years	β_7	3.59	0.402	
	75-79 years	β_8	3.87	0.415	
	80+ years	β_9	4.20	0.432	
Exposure	Fiber-years/ml; 10 year lag	β_{10}	0.0198	0.00496	47.1, 1, <0.0001

^a Linear relative risk model for lung cancer mortality rate is given by $\lambda = \lambda_0 \times (1 + \beta_{10}E_{10})$ where $\lambda_0 = \exp[\beta_0 + \beta_1(\text{sex=female}) + \beta_2(\text{race=non-white}) + \beta_3(\text{age=50-54 years}) + \beta_4(\text{age=55-59 years}) + \beta_5(\text{age=60-64 years}) + \beta_6(\text{age=65-69 years}) + \beta_7(\text{age=70-74 years}) + \beta_8(\text{age=75-79 years}) + \beta_9(\text{age=80+ years})]$ and E_{10} is cumulative chrysotile exposure (in fiber-years/ml) lagged by 10 years.

^b Likelihood ratio chi-square statistic, degrees of freedom, p-value.

Asbestosis mortality (62 observed deaths) was modeled using the power model. Model fit was improved when sex, race, age and birth cohort were included in the baseline function, but not when calendar time was additionally included in the model. Age and birth cohort categories with similar estimates were combined resulting in four age categories (<50 years, 50-59 years, 60-74 years, and 75+ years) and two birth cohort categories (<1920, 1920+). Mortality was lower for females and non-whites, higher for workers born before 1920, and increased with age. With sex, race, age and birth cohort in the baseline function, the relative rate for cumulative exposure (E, fiber-years/ml) was given by $(E+0.9)^{1.07}/(0.9)^{1.07}$. Interactions among the baseline covariates did not improve the model, nor did interactions with cumulative exposure and TSFE; however, a statistically significant interaction between cumulative exposure and age category was observed (p-value = 0.026). Including this interaction term in the model results in estimates of the relative rate for cumulative exposure that depend upon age. This model for asbestosis mortality is described in Table 6 and graphically displayed, along with the categorical and restricted cubic spline models, in Figure 2 for, as an example, white males 60-74 years of age born in 1920 or later. Under this model, the relative rate for asbestosis as a function of cumulative exposure was given by $(E+0.9)^{1.52}/(0.9)^{1.52}$ for persons under 50 years of age, $(E+0.9)^{1.50}/(0.9)^{1.50}$ for workers 50-59 years of age, $(E+0.9)^{1.20}/(0.9)^{1.20}$ for persons 60-74 years of age, and $(E+0.9)^{0.62}/(0.9)^{0.62}$ for persons 75+ years of age. Similar results were obtained when cumulative exposure was lagged by 5 or 10 years (not shown). Cumulative exposure was positively and significantly associated with asbestosis mortality in all models tested.

Table 6. Model for asbestosis mortality as an underlying or contributing cause of death among workers in the South Carolina asbestos textile workers cohort ^a

Term in model			Estimate	Standard Error	Likelihood Ratio Test ^b
Intercept		β_0	-14.4	1.94	---
Sex	Male	Ref	---	---	10.5, 1, 0.0012
	Female	β_1	-0.938	0.299	
Race	White	Ref	---	---	11.9, 1, 0.0006
	Non-white	β_2	-1.20	0.390	
Age	< 50 years	Ref	---	---	20.4, 3, 0.0001
	50-59 years	β_3	1.06	2.59	
	60-74 years	β_4	3.32	2.14	
	75+ years	β_5	6.62	2.07	
Birth cohort	1920+	Ref	---	---	5.54, 1, 0.019
	< 1920	β_6	0.811	0.373	
Exposure	Fiber-years/ml; no lag	α	0.9	---	---
		β_7	1.52	0.417	95.5, 1, <0.0001
Exposure×Age	< 50 years	Ref	---	---	9.30, 3, 0.026
	50-59 years	β_8	-0.0175	0.540	
	60-74 years	β_9	-0.314	0.456	
	75+ years	β_{10}	-0.897	0.451	

^a Power model for asbestosis mortality rate is given by $\lambda = \lambda_0 \times (E + \alpha)^\beta$ where $\lambda_0 = \exp[\beta_0 + \beta_1(\text{sex=female}) + \beta_2(\text{race=non-white}) + \beta_3(\text{age=50-59 years}) + \beta_4(\text{age=60-74 years}) + \beta_5(\text{age=75+ years}) + \beta_6(\text{birth cohort=prior to 1920})]$, E is cumulative chrysotile exposure (in fiber-years/ml), $\alpha = 0.9$ (estimated iteratively), and $\beta = (\beta_7 + \beta_8(\text{age=50-59 years}) + \beta_9(\text{age=60-74 years}) + \beta_{10}(\text{age=75+ years}))$.

^b Likelihood ratio chi-square statistic, degrees of freedom, p-value.

DISCUSSION

The primary focus of this update was to re-examine both mortality and the exposure-response relationships between cumulative exposure to chrysotile and mortality from lung cancer and asbestosis in the South Carolina asbestos textile workers cohort with an additional eleven years of follow-up. All cause mortality remained elevated, as was mortality from all cancers including esophageal and lung cancer, ischemic heart disease, pneumoconiosis and other respiratory diseases, diabetes, chronic obstructive pulmonary disease, and diseases of the digestive system. The number of deaths from mesothelioma is consistent with the results of other cohort mortality studies where workers were predominantly exposed to the chrysotile form of asbestos.[2,17]

Increasing and highly statistically significant trends with increasing cumulative exposure were observed for both lung cancer and asbestosis, with 72 and 17 additional deaths since the last update, respectively. Standardized mortality ratios were elevated for lung cancer and pneumoconiosis and other respiratory diseases even among the lowest exposure group. Previous analyses of the exposure-response relation between chrysotile and lung cancer mortality in this cohort selected the linear relative risk model over other model forms and estimated the relative rate for cumulative chrysotile exposure (E , fiber-years/ml) to be $(1 + 0.022 E)$ for a worker with 15-29 years of TSFE, $(1 + 0.037 E)$ for a worker with 30-39 years of TSFE and $(1 + 0.011 E)$ for a worker with 40 or more years of TSFE.[8] Using the updated data, and employing a 10-year lag on cumulative exposure, the relative rate was estimated to be $(1 + 0.020 E_{10})$; however, there was no evidence of an interaction between TSFE and exposure.

Previous analysis of the exposure-response relation between chrysotile and mortality from asbestosis in this cohort selected the power model over other model forms and estimated the relative rate for cumulative chrysotile exposure (E , fibers-years/ml) to be equal to $(E + 0.5)^{1.3}/(0.5)^{1.3}$. [8] The power model, which fit the data better than other models tested (results not shown), is still a reasonable model for asbestosis mortality in this cohort, but was improved by adding an interaction with age.

Smoking information on the cohort is limited. The U.S. Public Health Service administered surveys to active workers in 1964 and again to active workers in 1971. These surveys indicated that, compared to the U.S. population in 1965, smoking prevalence among white males ($n = 292$) was similar to the prevalence among U.S. white males; however, prevalence among non-white males ($n = 113$) was lower than the prevalence among U.S. non-white males and prevalence among white females ($n = 124$) was higher than the prevalence among U.S. white females.[7] The strong exposure-response patterns observed for lung cancer is unlikely to be fully explained by uncontrolled confounding by smoking since these analyses were conducted within the cohort and smoking is unlikely to vary by level of asbestos exposure. In a study examining the potential for tobacco and alcohol to confound the relationship between laryngeal cancer and metal working fluids, Kriebel et al. concluded that, for large studies, systematic or chance differences in smoking and drinking habits among the exposure groups are unlikely to cause more than a 17% change in the relative risk.[18] Smoking prevalence in the U.S. is related to sex, race, education and birth cohort.[19, 20] In the absence of smoking information, others have used birth year or birth cohort as a surrogate for smoking.[21] Birth cohort, however, was neither an effect modifier nor a confounder in our models of lung cancer with cumulative exposure. In our models of asbestosis

mortality, the addition of a term to identify workers born prior to 1920 in the baseline function significantly improved model fit, but an interaction term between birth cohort and cumulative chrysotile exposure did not.

There is some evidence of the healthy worker survival effect in that reduced mortality was observed among actively employed workers and highly elevated mortality, mostly lung cancer and pneumoconiosis and other respiratory diseases, was observed in the first year following termination of employment. Time since termination of employment has been shown to confound exposure-response relationships in occupational cohorts.[22] In supplemental analyses (not shown), time since last employment at the plant (used as a surrogate for time since termination of overall employment) did not confound the exposure-response relationship between chrysotile and lung cancer.

The cohort included many “short-term” workers with nearly half of the workers employed for less than one year at the plant. Since higher than expected death rates among short-term employees have been observed in a number of occupational mortality studies, hypotheses have been developed that short-term workers might have certain characteristics that distinguish them from long-term workers.[23,24] In this cohort, a high percentage (71%) of the short-term workforce was only employed during the World War II years and the nature of their short-term employment was more likely related to the reduction of war-time demand as opposed to possible socioeconomic, lifestyle or occupational factors. Furthermore, mortality from lung cancer and pneumoconiosis and other respiratory diseases was still elevated and positively associated with estimated cumulative exposure to chrysotile in supplemental analyses excluding short-term workers.

In addition to a lack of smoking histories for all of the cohort members, the findings reported herein are subject to additional limitations including incomplete life-time work histories and high rates of loss to follow-up, especially among female workers. The detailed exposure-response modeling is not without limitations in that estimates of risk are based on a single model and are somewhat dependent on the form of the model, the cutpoints used to stratify exposure and the lag period selected. Cumulative exposure, however, was positively and highly statistically significantly associated with lung cancer and asbestosis mortality when additional model forms were considered (not shown). The use of a large number of cutpoints and the treatment of exposure as a continuous variable in these models lessens the dependence of the results on the selected cutpoints.[14] Exposure lag periods of 5 and 10 years were considered in addition to un-lagged analyses. Results were similar in lagged and un-lagged analyses, however, in this cohort, many years have passed since exposure ceased, so this was not too surprising.

In spite of these limitations, there are many strengths of the analysis, including a long follow-up time, a large number of observed deaths due to lung cancer and asbestosis, detailed work histories and a well-defined JEM. The JEM used here characterizes exposure to chrysotile in terms of the number of fibers longer than 5 micrometers per milliliter of air. Exposure concentrations were estimated using statistical modeling of nearly 6000 industrial hygiene sampling measurements taken over the period 1930-1975 and analyzed using phase contrast microscopy. There is evidence that fiber size (i.e., diameter and length) may play an important role in characterizing the exposure-response relationship between chrysotile exposure and mortality from lung cancer and

asbestosis.[25] Plans are underway for the re-analysis of this cohort using an updated JEM based on a transmission electron microscopy analysis of the air samples that considers the size distribution of the fibers in addition to the number of fibers.

In conclusion, this study confirms the findings from previous investigations of excess mortality from lung cancer and asbestosis and a strong exposure-response relationship between exposure to chrysotile and mortality from lung cancer and asbestosis.

DISCLAIMER

The findings and conclusions in this report are those of the authors and do not necessarily represent the views of the National Institute for Occupational Safety and Health.

ACKNOWLEDGEMENTS

The authors express their appreciation to Ms. Christine M. Gersic for her assistance with the data files.

Figure 1. Estimated lung cancer mortality for white males, 60-64 years of age as a function of cumulative exposure to chrysotile (10 year lag) based on the model described in Table 5 (linear relative risk model – solid curve; restricted cubic spline model – dashed curve; categorical model – step function).

Figure 2. Estimated asbestosis mortality for white males, 60-74 years of age, and born in 1920 or later as a function of cumulative exposure to chrysotile based on the model described in Table 6 (power model – solid curve; restricted cubic spline model – dashed curve; categorical model – step function).

REFERENCES

- 1 McDonald JC, McDonald AD. Chrysotile, tremolite and carcinogenicity. *Ann Occup Hyg* 1997;**41**(6):699-705.
- 2 Stayner LT, Dankovic DA, Lemen RA. Occupational exposure to chrysotile asbestos and cancer risk: a review of the amphibole hypothesis. *Am J Public Health* 1996;**86**(2):179-86.
- 3 Landrigan PJ, Nicholson WJ, Suzuki Y, *et al.* The hazards of chrysotile asbestos: a critical review. *Ind Health* 1999;**37**(3):271-80.
- 4 Yano E, Wang Z, Wang X, *et al.* Cancer mortality among workers exposed to amphibole-free chrysotile asbestos. *Am J Epidemiol* 2001;**154**(6):538-43.
- 5 Dement JM, Harris RL Jr, Symons MJ, *et al.* Exposures and mortality among chrysotile asbestos workers. Part II: Mortality. *Am J Ind Med* 1983;**4**(3):421-33.
- 6 Brown DP, Dement JM, Okun A. Mortality patterns among female and male chrysotile asbestos textile workers. *J Occup Med* 1994;**36**(8):882-8.
- 7 Dement JM, Brown DP, Okun A. Follow-up study of chrysotile asbestos textile workers: cohort mortality and case-control analyses. *Am J Ind Med* 1994;**26**(4):431-47.

- 8 Stayner L, Smith R, Bailer J, *et al.* Exposure-response analysis of risk of respiratory disease associated with occupational exposure to chrysotile asbestos. *Occup Environ Med* 1997;**54**(9):646-52.
- 9 Dement JM, Harris RL Jr, Symons MJ, *et al.* Exposures and mortality among chrysotile asbestos workers. Part I: Exposure estimates. *Am J Ind Med* 1983;**4**(3):399-419.
- 10 Waxweiler RJ, Beaumont JJ, Henry JA, *et al.* A modified life-table analysis system for cohort studies. *J Occup Med* 1983;**25**(2):115-24.
- 11 Steenland K, Beaumont J, Spaeth S, *et al.* New developments in the Life Table Analysis System of the National Institute for Occupational Safety and Health. *J Occup Med* 1990;**32**(11):1091-8.
- 12 Robinson CF, Schnorr TM, Cassinelli RT II, *et al.* Tenth revision U.S. mortality rates for use with the NIOSH Life Table Analysis System. *J Occup Environ Med* 2006;**48**(7):662-7.
- 13 Dement JM, Brown DP. Cohort mortality and case-control studies of white male chrysotile asbestos textile workers. *J Occup Med Toxicol* 1993;**2**(4):355-63.
- 14 Richardson DB, Loomis D. The impact of exposure categorisation for grouped analyses of cohort data. *Occup Environ Med* 2004;**61**(11):930-5.
- 15 U.S. Environmental Protection Agency. Airborne Asbestos Health Assessment Update. 1986. Report 600/8-84-003F, U.S. Environmental Protection Agency.
- 16 Herndon JE II, Harrell FE Jr. The restricted cubic spline as baseline hazard in the proportional hazards model with step function time-dependent covariables. *Stat Med* 1995;**14**(19):2119-29.
- 17 Hodgson JT, Darnton A. The quantitative risks of mesothelioma and lung cancer in relation to asbestos exposure. *Ann Occup Hyg* 2000;**44**(8):565-601.
- 18 Kriebel D, Zeka A, Eisen EA, *et al.* Quantitative evaluation of the effects of uncontrolled confounding by alcohol and tobacco in occupational cancer studies. *Int J Epidemiol* 2004;**33**(5):1040-5.
- 19 Escobedo LG, Peddicord JP. Smoking prevalence in US birth cohorts: the influence of gender and education. *Am J Public Health* 1996;**86**(2):231-6.
- 20 Freedman DM, Tarone RE, Doody MM, *et al.* Trends in reproductive, smoking, and other chronic disease risk factors by birth cohort and race in a large occupational study population. *Ann Epidemiol* 2002;**12**(6):363-9.
- 21 Luebeck EG, Heidenreich WF, Hazelton WD, *et al.* Biologically based analysis of the data for the Colorado uranium miners cohort: age, dose and dose-rate effects. *Radiat Res* 1999;**152**(4):339-51.
- 22 Richardson D, Wing S, Steenland K, *et al.* Time-related aspects of the healthy worker survivor effect. *Ann Epidemiol* 2004;**14**(9):633-9.
- 23 Gilbert ES. Some confounding factors in the study of mortality and occupational exposures. *Am J Epidemiol* 1982;**116**(1):177-88.
- 24 Kolstad HA, Olsen J. Why do short term workers have high mortality? *Am J Epidemiol* 1999;**149**(4):347-52.
- 25 Berman DW, Crump KS, Chatfield EJ, *et al.* The sizes, shapes, and mineralogy of asbestos structures that induce lung tumors or mesothelioma in AF/HAN rats following inhalation. *Risk Anal* 1995;**15**(2):181-95.




Figure 1




Figure 2