

The Honorable Lisa P. Jackson
Administrator
U.S. Environmental Protection Agency
1200 Pennsylvania Avenue, N.W.
Washington, D.C. 20460

Subject: SAB Review of EPA's Reanalysis of Key Issues Related to Dioxin
Toxicity and Response to NAS Comments

Dear Administrator Jackson:

EPA's Office of Research and Development (ORD) requested that the Science Advisory Board (SAB) review the Agency's draft report entitled *EPA's Reanalysis of Key Issues Related to Dioxin Toxicity and Response to NAS Comments* ("Report"). The Report contains EPA's technical response to key comments in the 2006 National Academy of Sciences (NAS) report, *Health Risks from Dioxin and Related Compounds: Evaluation of the EPA Reassessment*. The NAS reviewed EPA's 2003 exposure and human health reassessment of dioxin and recommended that the Agency should: more thoroughly justify and communicate its approaches to dose-response modeling for the health effects of dioxin, taking into consideration both nonlinear and linear methods for characterizing cancer risk; improve the transparency and clarity of the selection of key data sets for the dioxin dose-response analysis; reevaluate its cancer weight-of-evidence determination for dioxin based on the Agency's 2005 Cancer Guidelines; consider using physiologically-based pharmacokinetic (PBPK) modeling in the dioxin risk assessment; and improve transparency, thoroughness and clarity in quantitative uncertainty analysis. The NAS also encouraged EPA to calculate a reference dose (RfD), which had not been derived in the 2003 reassessment.

In response to EPA's request, the SAB convened an expert panel to review the Agency's Report. The SAB Panel was asked to comment on the scientific soundness of EPA's responses to the NAS recommendations. The enclosed SAB report provides the consensus advice and recommendations of the Panel, with the exception of one member who offered a dissenting opinion mainly on 2,3,7,8 Tetrachlorodibenzo-p-dioxin (TCDD) carcinogenicity.

The SAB finds that EPA's Report is clear, logical, and responsive to many but not all of the recommendations of the NAS. We have provided recommendations to further enhance the transparency, clarity, and scientific integrity of the Report. The SAB has identified deficiencies in EPA's Report with respect to the completeness of its consideration of two critical elements of the TCDD assessment: 1) nonlinear dose-response for TCDD carcinogenicity, and 2) uncertainty analysis of TCDD toxicity. Our major comments and recommendations are provided below:

- 1
2 • The SAB commends EPA for the comprehensive and rigorous process that was
3 used to identify, review, and evaluate the TCDD literature. The criteria for study
4 selection have been clearly articulated, well justified, and applied in a
5 scientifically sound manner. To further improve clarity and transparency of the
6 Report, we recommend that EPA include a better means of tracking and
7 describing which studies did not satisfy inclusion criteria. Similarly, we
8 recommend that EPA strengthen its justification for excluding studies of dioxin-
9 like compounds. The Report can be enhanced by incorporating information from
10 studies with dioxin-like compounds into a qualitative discussion of the weight-of-
11 evidence for cancer and noncancer endpoints.
12
- 13 • EPA used the Emond physiologically-based pharmacokinetic model to evaluate
14 the internal dose of TCDD in human and rodent tissue, and to estimate the
15 continuous daily TCDD intake over the relevant period of exposure. The SAB
16 agrees with EPA that this model provides the best available basis for the dose
17 metric calculations. We also support EPA's use of blood TCDD concentrations
18 as the relevant dose metric. However, we recommend that EPA expand the
19 discussion of other published models, evaluate the impact of model selection on
20 dose metric prediction, provide a more quantitative uncertainty analysis, and
21 conduct an external peer review of the mouse model because it has not been
22 published in the peer-reviewed literature.
23
- 24 • The SAB agrees with EPA's classification of TCDD as carcinogenic to humans in
25 accordance with EPA's *2005 Guidelines for Carcinogen Risk Assessment*. The
26 SAB recommends that in the weight-of-evidence characterization EPA build upon
27 all available data to support its decision and clearly indicate how different types
28 of data support each other. One Panel member, however, indicated that at best,
29 there is equivocal evidence for TCDD classification as a human carcinogen.
30
- 31 • The SAB agrees with EPA's selection of the Cheng et al. (2006) study, which
32 analyzed the National Institute for Occupational Safety and Health (NIOSH)
33 occupational cohort, as the critical study for the quantitative cancer assessment.
34 The SAB also agrees that it is appropriate to use all-cancer mortality as the basis
35 of the oral slope factor because of the extensive dose-response information.
36
- 37 • The SAB finds that the Report did not respond adequately to the NAS
38 recommendation to adopt both linear and nonlinear methods of extrapolation in
39 order to account for the uncertainty of the dose-response curve for TCDD. The
40 Report states that only a linear approach could be justified. We recommend that
41 EPA revise the Report to provide a discussion of evidence of possible modes of
42 action that include both linear and nonlinear alternatives for the cancer endpoint.
43 In the absence of a definitive nonlinear mode of action, estimates based on the
44 linear option can serve as the baseline for comparison with other estimates.

- The SAB supports EPA's selection and use of two co-critical epidemiologic studies for the derivation of the RfD for TCDD. These studies evaluated the effects of human exposure to TCDD following accidental release at a chemical plant near Seveso, Italy. The SAB finds that the study endpoints used by EPA to determine the RfD (decrease in sperm count and motility and increased thyroid stimulating hormone in blood) are relevant to public health. The selection of these endpoints also resolves the critical issue of differing windows of susceptibility to environmental toxic agents over the course of the life cycle, with pre- and periconceptional exposures comprising the window of greatest susceptibility. We recommend, however, that EPA provide a discussion of the strengths and weaknesses of the studies and an indication of whether these weaknesses affect the RfD determination.
- The SAB also agrees with the benchmark dose modeling approaches used by EPA in the Report and the decision to use human data as preferred to animal data for the RfD determination.
- EPA's Report discusses a broad range of philosophical and methodological issues to be considered in conducting an uncertainty analysis for TCDD toxicity. Although the SAB acknowledges the challenges of a unified quantitative uncertainty analysis, we do not agree with the position taken in the Report that such an analysis is unfeasible and we have suggested a number of methods that could be used for this purpose.
- Finally, EPA's Report could be improved by editing and restructuring to better integrate the material presented in various sections, eliminate redundancies, and move some material into appendices to provide more succinct responses to NAS concerns. In addition, we recommend including a glossary in the Report to help minimize confusion and misinterpretation among diverse users.

The SAB appreciates the opportunity to provide EPA with advice on this important subject. We support EPA in its effort to move in a proficient and expeditious manner to finalize the IRIS document for dioxin and look forward to receiving the Agency's response.

Sincerely,

Dr. Deborah L. Swackhamer, Chair
EPA Science Advisory Board

Dr. Timothy J. Buckley, Chair
SAB Dioxin Review Panel

NOTICE

This report has been written as part of the activities of the EPA Science Advisory Board, a public advisory committee providing extramural scientific information and advice to the Administrator and other officials of the Environmental Protection Agency. The Board is structured to provide balanced, expert assessment of scientific matters related to problems facing the Agency. This report has not been reviewed for approval by the Agency and, hence, the contents of this report do not necessarily represent the views and policies of the Environmental Protection Agency, nor of other agencies in the Executive Branch of the Federal government, nor does mention of trade names or commercial products constitute a recommendation for use. Reports of the EPA Science Advisory Board are posted on the EPA Web site at: <http://www.epa.gov/sab>

U.S. Environmental Protection Agency
Science Advisory Board
Dioxin Review Panel

CHAIR

Dr. Timothy Buckley, Associate Professor and Chair, Division of Environmental Health
Sciences, College of Public Health, The Ohio State University, Columbus, OH

MEMBERS

Dr. Harvey Clewell, Director of the Center for Human Health Assessment, The Hamner
Institutes for Health Sciences, Research Triangle Park, NC

Dr. Louis Anthony (Tony) Cox, Jr., President, Cox Associates, Denver, CO

Dr. Elaine Faustman, Professor and Director, Institute for Risk Analysis and Risk
Communication, School of Public Health, University of Washington, Seattle, WA

Dr. Scott Ferson, Senior Scientist, Applied Biomathematics, Setauket, NY

Dr. Jeffrey Fisher, Research Toxicologist, National Center for Toxicological Research,
U.S. Food and Drug Administration, Jefferson, AR

Dr. Helen Håkansson, Professor of Toxicology, Unit of Environmental Health Risk
Assessment, Institute of Environmental Medicine, Karolinska Institutet, Stockholm,
Sweden

Dr. Russ Hauser, Frederick Lee Hisaw Professor, Department of Environmental Health,
Harvard School of Public Health, Boston, MA

Dr. B. Paige Lawrence, Associate Professor, Departments of Environmental Medicine
and Microbiology and Immunology, School of Medicine and Dentistry, University of
Rochester School of Medicine and Dentistry, Rochester, NY

Dr. Michael I. Luster, Professor, Department of Community Medicine, West Virginia
University Health Sciences Center, Morgantown, WV

Dr. Paolo Mocarelli, Professor of Clinical Biochemistry, Department of Clinical
Laboratory, Hospital of Desio-Nuovo Monoblous, University of Milano Bicocca, Desio-
Milano, Italy

Dr. Victoria Persky, Professor, Epidemiology and Biostatistics Program, School of
Public Health, University of Illinois at Chicago, Chicago, IL

Dr. Sandra L. Petersen, Professor, Associate Graduate Dean, Department of Veterinary
and Animal Sciences, College of Natural Sciences, University of Massachusetts-
Amherst, Amherst, MA

Dr. Karl Rozman^{*}, Professor, Pharmacology, Toxicology and Therapeutics, The
University of Kansas Medical Center, Kansas City, KS

Dr. Arnold Schecter, Professor, Environmental and Occupational Health Sciences,
School of Public Health-Dallas Campus, University of Texas, Dallas, TX

Dr. Allen E. Silverstone, Professor, Department of Microbiology and Immunology,
Health Science Center, SUNY Upstate Medical University, Syracuse, NY and Adjunct
Professor of Environmental Medicine, University of Rochester School of Medicine and
Dentistry, Rochester, NY.

Dr. Mitchell J. Small, The H. John Heinz III Professor of Environmental Engineering,
Department of Civil and Environmental Engineering and Engineering and Public Policy,
Carnegie Mellon University, Pittsburgh, PA

Dr. Anne Sweeney, Professor of Epidemiology, Department of Epidemiology and
Biostatistics, School of Rural Public Health, Texas A&M Health Science Center, College
Station, TX

Dr. Mary K. Walker, Professor, Division of Pharmaceutical Sciences, College of
Pharmacy, University of New Mexico, Albuquerque, NM

SCIENCE ADVISORY BOARD STAFF

Dr. Thomas Armitage, Designated Federal Officer, U.S. Environmental Protection
Agency, Washington, DC

Dr. Diana Wong, Designated Federal Officer, U.S. Environmental Protection Agency,
Washington, DC

^{*}Dissenting opinion in Appendix A

**U.S. Environmental Protection Agency
Science Advisory Board
BOARD**

CHAIR

Dr. Deborah L. Swackhamer, Professor and Charles M. Denny, Jr. Chair in Science,
Technology and Public Policy - Hubert H. Humphrey School of Public Affairs and Co-
Director of the Water Resources Center, University of Minnesota, St. Paul, MN

SAB MEMBERS

Dr. David T. Allen, Professor, Department of Chemical Engineering, University of
Texas, Austin, TX

Dr. Claudia Benitez-Nelson, Full Professor and Director of the Marine Science
Program, Department of Earth and Ocean Sciences, University of South Carolina,
Columbia, SC

Dr. Timothy J. Buckley, Associate Professor and Chair, Division of Environmental
Health Sciences, College of Public Health, The Ohio State University, Columbus, OH

Dr. Patricia Buffler, Professor of Epidemiology and Dean Emerita, Department of
Epidemiology, School of Public Health, University of California, Berkeley, CA

Dr. Ingrid Burke, Director, Haub School and Ruckelshaus Institute of Environment and
Natural Resources, University of Wyoming, Laramie, WY

Dr. Thomas Burke, Professor, Department of Health Policy and Management, Johns
Hopkins Bloomberg School of Public Health, Johns Hopkins University, Baltimore, MD

Dr. Terry Daniel, Professor of Psychology and Natural Resources, Department of
Psychology, School of Natural Resources, University of Arizona, Tucson, AZ

Dr. George Daston, Victor Mills Society Research Fellow, Product Safety and
Regulatory Affairs, Procter & Gamble, Cincinnati, OH

Dr. Costel Denson, Managing Member, Costech Technologies, LLC, Newark, DE

Dr. Otto C. Doering III, Professor, Department of Agricultural Economics, Purdue
University, W. Lafayette, IN

1 **Dr. David A. Dzombak**, Walter J. Blenko, Sr. Professor of Environmental Engineering,
2 Department of Civil and Environmental Engineering, College of Engineering, Carnegie
3 Mellon University, Pittsburgh, PA
4

5 **Dr. T. Taylor Eighmy**, Vice President for Research, Office of the Vice President for
6 Research, Texas Tech University, Lubbock, TX
7

8 **Dr. Elaine Faustman**, Professor and Director, Institute for Risk Analysis and Risk
9 Communication, School of Public Health, University of Washington, Seattle, WA
10

11 **Dr. John P. Giesy**, Professor and Canada Research Chair, Veterinary Biomedical
12 Sciences and Toxicology Centre, University of Saskatchewan, Saskatoon, Saskatchewan,
13 Canada
14

15 **Dr. Jeffrey K. Griffiths**, Associate Professor, Department of Public Health and
16 Community Medicine, School of Medicine, Tufts University, Boston, MA
17

18 **Dr. James K. Hammitt**, Professor, Center for Risk Analysis, Harvard University,
19 Boston, MA
20

21 **Dr. Bernd Kahn**, Professor Emeritus and Associate Director, Environmental Radiation
22 Center, Georgia Institute of Technology, Atlanta, GA
23

24 **Dr. Agnes Kane**, Professor and Chair, Department of Pathology and Laboratory
25 Medicine, Brown University, Providence, RI
26

27 **Dr. Madhu Khanna**, Professor, Department of Agricultural and Consumer Economics,
28 University of Illinois at Urbana-Champaign, Urbana, IL
29

30 **Dr. Nancy K. Kim**, Senior Executive, Health Research, Inc., Troy, NY
31

32 **Dr. Catherine Kling**, Professor, Department of Economics, Iowa State University,
33 Ames, IA
34

35 **Dr. Kai Lee**, Program Officer, Conservation and Science Program, David & Lucile
36 Packard Foundation, Los Altos, CA (affiliation listed for identification purposes only)
37

38 **Dr. Cecil Lue-Hing**, President, Cecil Lue-Hing & Assoc. Inc., Burr Ridge, IL
39

40 **Dr. Floyd Malveaux**, Executive Director, Merck Childhood Asthma Network, Inc.,
41 Washington, DC
42

43 **Dr. Lee D. McMullen**, Water Resources Practice Leader, Snyder & Associates, Inc.,
44 Ankeny, IA
45

1 **Dr. Judith L. Meyer**, Professor Emeritus, Odum School of Ecology, University of
2 Georgia, Lopez Island, WA
3

4 **Dr. James R. Mihelcic**, Professor, Civil and Environmental Engineering, University of
5 South Florida, Tampa, FL
6

7 **Dr. Jana Milford**, Professor, Department of Mechanical Engineering, University of
8 Colorado, Boulder, CO
9

10 **Dr. Christine Moe**, Eugene J. Gangarosa Professor, Hubert Department of Global
11 Health, Rollins School of Public Health, Emory University, Atlanta, GA
12

13 **Dr. Horace Moo-Young**, Dean and Professor, College of Engineering, Computer
14 Science, and Technology, California State University, Los Angeles, CA
15

16 **Dr. Eileen Murphy**, Grants Facilitator, Ernest Mario School of Pharmacy, Rutgers
17 University, Piscataway, NJ
18

19 **Dr. Duncan Patten**, Research Professor, Hydroecology Research Program, Department
20 of Land Resources and Environmental Sciences, Montana State University, Bozeman,
21 MT
22

23 **Dr. Stephen Polasky**, Fesler-Lampert Professor of Ecological/Environmental
24 Economics, Department of Applied Economics, University of Minnesota, St. Paul, MN
25

26 **Dr. Arden Pope**, Professor, Department of Economics, Brigham Young University,
27 Provo, UT
28

29 **Dr. Stephen M. Roberts**, Professor, Department of Physiological Sciences, Director,
30 Center for Environmental and Human Toxicology, University of Florida, Gainesville, FL
31

32 **Dr. Amanda Rodewald**, Professor of Wildlife Ecology, School of Environment and
33 Natural Resources, The Ohio State University, Columbus, OH
34

35 **Dr. Jonathan M. Samet**, Professor and Flora L. Thornton Chair, Department of
36 Preventive Medicine, University of Southern California, Los Angeles, CA
37

38 **Dr. James Sanders**, Director and Professor, Skidaway Institute of Oceanography,
39 Savannah, GA
40

41 **Dr. Jerald Schnoor**, Allen S. Henry Chair Professor, Department of Civil and
42 Environmental Engineering, Co-Director, Center for Global and Regional Environmental
43 Research, University of Iowa, Iowa City, IA
44
45

1 **Dr. Kathleen Segerson**, Philip E. Austin Professor of Economics, Department of
2 Economics, University of Connecticut, Storrs, CT

3
4 **Dr. Herman Taylor**, Director, Principal Investigator, Jackson Heart Study, University of
5 Mississippi Medical Center, Jackson, MS

6
7 **Dr. Barton H. (Buzz) Thompson, Jr.**, Robert E. Paradise Professor of Natural
8 Resources Law at the Stanford Law School and Perry L. McCarty Director, Woods
9 Institute for the Environment, Stanford University, Stanford, CA

10
11 **Dr. Paige Tolbert**, Professor and Chair, Department of Environmental Health, Rollins
12 School of Public Health, Emory University, Atlanta, GA

13
14 **Dr. John Vena**, Professor and Department Head, Department of Epidemiology and
15 Biostatistics, College of Public Health, University of Georgia, Athens, GA

16
17 **Dr. Thomas S. Wallsten**, Professor and Chair, Department of Psychology, University of
18 Maryland, College Park, MD

19
20 **Dr. Robert Watts**, Professor of Mechanical Engineering Emeritus, Tulane University,
21 Annapolis, MD

22
23 **Dr. R. Thomas Zoeller**, Professor, Department of Biology, University of Massachusetts,
24 Amherst, MA

25
26
27 **SCIENCE ADVISORY BOARD STAFF**

28 **Dr. Angela Nugent**, Designated Federal Officer, U.S. Environmental Protection Agency,
29 Washington, DC

30
31 **Ms. Stephanie Sanzone**, Designated Federal Officer, U.S. Environmental Protection
32 Agency, Washington, DC

TABLE OF CONTENTS

1		
2		
3		
4		
5	ABBREVIATIONS AND ACRONYMS.....	xii
6	EXECUTIVE SUMMARY	1
7	INTRODUCTION.....	9
8	RESPONSES TO EPA’S CHARGE QUESTIONS.....	11
9	Charge Question 1. General Charge Questions	11
10	Charge Question 2. Transparency and Clarity in the Selection of Key Data Sets for Dose-Response	
11	Analysis.....	13
12	Charge Question 3. The Use of Toxicokinetics in the Dose-Response Modeling for Cancer	
13	and Noncancer Endpoints.....	19
14	Charge Question 4. Reference dose.....	25
15	Charge Question 5. Cancer assessment	33
16	Charge Question 6. Feasibility of Quantitative Uncertainty Analysis.....	40
17	REFERENCES.....	50
18	APPENDIX A: DISSENTING OPINION FROM KARL ROZMAN, PH.D.	A-1
19	APPENDIX B: VALUE OF INFORMATION.....	B-1
20	APPENDIX C: EDITORIAL COMMENTS AND CORRECTIONS	C-1
21	APPENDIX D: EPA’S CHARGE QUESTIONS	D-1
22		
23		
24		
25		
26		
27		
28		

ABBREVIATIONS AND ACRONYMS

1		
2		
3		
4	AhR	aryl hydrocarbon receptor
5	BMD	benchmark dose
6	BMDL	benchmark dose lower bound
7	BMR	benchmark response level
8	CYP	cytochrome P450
9	DLC	dioxin-like compound
10	ED	effective dose
11	EPA	U.S. Environmental Protection Agency
12	HED	human equivalent dose
13	IRIS	integrated risk information system
14	LASC	lipid-adjusted serum concentrations
15	LOAEL	lowest-observed-adverse-effect level
16	MOA	mode of action
17	NAS	National Academy of Sciences
18	NHEERL	National Health and Environmental Effects Research Laboratory
19	NIOSH	National Institute for Occupational Safety and Health
20	NOAEL	no-observed-adverse-effect level
21	NRC	National Research Council
22	OSF	oral slope factor
23	PBPK	physiologically-based pharmacokinetic
24	PCDDs	polychlorinated dibenzo- <i>p</i> -dioxin
25	PCDFs	polychlorinated dibenzofuran
26	POD	point of departure
27	RfD	reference dose
28	RR	relative risk
29	SAB	Science Advisory Board
30	T3	triiodothyronine
31	T4	thyroxine
32	TCDD	2,3,7,8-Tetrachlorodibenzo- <i>p</i> -dioxin
33	TEF	toxicity equivalence factor
34	TEQ	toxicity equivalence
35	TSH	thyroid stimulating hormone
36	UF	uncertainty factor
37	WHO	World Health Organization

EXECUTIVE SUMMARY

In 2003, EPA, along with other federal agencies, asked the National Academy of Sciences (NAS) to review aspects of the science in EPA's draft dioxin reassessment entitled, *Exposure and Human Health Reassessment of 2,3,7,8-Tetrachlorodibenzo-p-Dioxin (TCDD) and Related Compounds* and, in 2004, EPA sent the 2003 draft dioxin reassessment to the NAS for review. In 2006, the NAS released the report of its review entitled, *Health Risks from Dioxin and Related Compounds: Evaluation of the EPA Reassessment*. The NAS recommended that EPA should: more thoroughly justify and communicate its approaches to dose-response modeling for the health effects of dioxin, taking into consideration both nonlinear and linear methods for characterizing cancer risk; improve the transparency and clarity of the selection of key data sets for the dioxin dose-response analysis; reevaluate its cancer weight-of-evidence determination for dioxin based on the Agency's 2005 Cancer Guidelines; consider using physiologically-based pharmacokinetic (PBPK) modeling in the dioxin risk assessment; and improve transparency, thoroughness and clarity in quantitative uncertainty analysis. The NAS also encouraged EPA to calculate a reference dose (RfD), which had not been derived in the 2003 reassessment.

EPA's Office of Research and Development (ORD) prepared the draft report, entitled *EPA's Reanalysis of Key Issues Related to Dioxin Toxicity and Response to NAS Comments* (EPA, 2010) (hereafter referred to as the *Report*), and requested that the EPA Science Advisory Board (SAB) conduct an independent external peer review of the *Report*. This Executive Summary highlights the findings and recommendations of the SAB Dioxin Review Panel (the "Panel") in response to charge questions concerning each of the six sections of the *Report*.

General Charge

The SAB Panel was asked to comment on: whether the *Report* was clear and logical, whether the Agency had objectively and clearly presented the key National Academy of Sciences (NAS) recommendations, and whether there were other critical studies that would make a significant impact on the conclusions of the hazard characterization or dose-response assessment of the chronic noncancer and cancer health effects of TCDD.

As further discussed in the responses to Charge Question 1, the Panel found that EPA was effective in developing a report that was clear, logical, and responsive to many but not all of the recommendations of the NAS. The Panel has provided recommendations to further improve the clarity, organization, and responsiveness of various parts of the *Report*. The Panel was impressed with the process that EPA used to identify, review, and evaluate the relevant literature. The Panel found that EPA's process was comprehensive and rigorous and included public participation. However, the Panel recommends that the *Report* be improved by: incorporating text to better integrate the material presented in the individual chapters, providing greater clarity and transparency

1 in indicating which studies did not satisfy criteria for inclusion in EPA's assessment of
2 TCDD, and editing the *Report* to provide greater clarity in writing and make it more
3 concise by moving some material into appendices.
4

5 During the course of its discussion, the Panel did not identify any additional
6 studies that would make a significant impact on the conclusions of the hazard
7 characterization and dose-response assessment. The Panel recommends that EPA
8 provide an assessment of both the null studies and positive studies with more discussion
9 and clarity concerning the exclusion of null epidemiologic studies. In addition, as further
10 discussed in the responses to the relevant charge questions, the Panel has identified
11 deficiencies in the *Report* with respect to the completeness of its consideration of two
12 critical elements: 1) nonlinear dose-response for TCDD carcinogenicity, and 2)
13 uncertainty analysis. As discussed below, the Panel has provided recommendations to
14 improve the *Report* in these areas.
15

16 **Transparency and Clarity in the Selection of Key Data Sets for Dose-Response** 17 **Analyses** 18

19 The NAS proposed that EPA develop a clear and readily understandable
20 methodology for evaluating and including epidemiologic and animal bioassay data sets in
21 dose-response evaluations. The SAB Panel was asked to comment on: whether EPA had
22 been responsive to NAS concerns about transparency and clarity in data set selection,
23 whether the epidemiology and animal bioassay study criteria and considerations had been
24 scientifically justified and clearly described, and whether EPA had applied the
25 epidemiology and animal bioassay study criteria considerations in a scientifically sound
26 manner.
27

28 Section 2 of the *Report* contains a clear presentation of the process EPA used to
29 select key data sets for dose-response analysis and is thus responsive to NAS
30 recommendations in this area. The *Report* also clearly identifies the studies that were
31 used for dose-response analysis. However, the Panel has provided recommendations to
32 further enhance the overall clarity and transparency of Section 2 of the *Report*. The Panel
33 recommends careful and extensive editing to revise and consolidate Section 2.
34 Specifically, editing should include aspects of grammar and syntax, minimizing
35 redundancies, and use of more succinct language in responses to NAS concerns. The
36 Panel also recommends restructuring Section 2 to improve its integration into the overall
37 document and make it easier to follow the studies used by EPA from one section of the
38 *Report* to another. In this regard, the Panel suggests that Section 2 could be used as the
39 foundation for the entire document.
40

41 The Panel also found that EPA's epidemiology and animal bioassay study criteria
42 and considerations were scientifically justified, clearly described, and applied in a
43 scientifically sound manner. The Panel has provided recommendations to improve and
44 strengthen the scientific justification and clarity of description of EPA's study criteria and
45 considerations. The Panel recommends that EPA better justify the rationale for using

only studies where the exposure was primarily to TCDD for derivation of the reference dose. This justification should include both scientific and practical reasons. The Panel also recommends that EPA incorporate information from studies with dioxin-like compounds (DLCs) into a qualitative discussion of the weight-of-evidence for cancer and noncancer endpoints. In addition, the Panel has provided a number of specific recommendations to further clarify the justifications for some of the study inclusion and exclusion criteria.

Use of Toxicokinetics in Dose-Response Modeling for Cancer and Noncancer Endpoints

In the Section 3 of the *Report*, EPA discussed the use of a physiologically-based pharmacokinetic (PBPK) model (Emond et al., 2004, 2005, 2006) with blood concentration as the dose metric rather than first-order body burden. The Panel was asked to comment on the scientific justification for EPA's application of this model, the model modifications that EPA implemented, and EPA's characterization of uncertainty in the model. EPA also developed a PBPK model to estimate TCDD concentration in mouse tissues. The Panel was asked to comment on the scientific rationale for development of the mouse model, the performance of the mouse model, and whether model uncertainty had been adequately characterized. In addition, the Panel was asked to comment on the use of the Emond PBPK model to estimate human intake based on internal exposure measures, EPA's sensitivity analysis of the kinetic modeling, and EPA's estimate of lifetime average daily dose.

The Panel agrees with EPA's use of blood TCDD concentration as a surrogate for tissue TCDD exposure. Blood TCDD concentration is a better choice than using body burden (as in the 2003 Reassessment) because it is more closely related to the biologically relevant dose metric: the free concentration of dioxin in the target tissues. The Panel further agrees that the PBPK model developed by Emond et al. (2004, 2005, 2006) provides the best available basis for the dose metric calculations in the assessment. However, the Panel recommends that EPA clarify how the model was used in studies that reported the concentrations of dioxin in plasma, serum, blood, or blood fat:blood measurements. The Panel also recommends additional discussion of: other published models, the intended use of the Emond model in the assessment, and the basis for selection of the Emond model. The Panel found that the EPA modifications to the published Emond model were minor and appropriate. However, the Panel notes that the use of 0.6 as the Hill coefficient in the Emond model for CYP1a2 induction is well outside the confidence interval of 0.78 and 1.14 reported by Walker et al. (1999). The use of a Hill coefficient value well below unity would lead to a nonlinear model behavior that is biologically implausible. As a result, when the human model was used for extrapolation to lower doses (in the calculation of risk-specific doses), the model would estimate a lower exposure level for a given blood concentration. The Panel suggests repeating the human Emond model calculations with multiple values for the Hill coefficient to characterize the resulting uncertainty in the exposure estimates.

1 The Panel also recommends that a more quantitative uncertainty analysis be
2 conducted for the PBPK model. Methods that could be useful and informative for such
3 analysis are suggested in the response to Charge Question 6.2. The sensitivity analysis in
4 the *Report* left out the Hill coefficient, which is one of the most important parameters in
5 the model for low-dose extrapolation. Model sensitivities are species, dose, and dose-
6 scenario dependent, so they need to be determined under the same exposure conditions
7 that dose metrics are calculated.

8
9 The Panel found that the mouse model developed by EPA based on the published
10 rat model (Emond et al., 2004, 2005, 2006) was appropriate, but it is recommended that
11 an external peer review of the mouse model be performed. The Panel agrees with the
12 average daily dose calculation approaches described in the *Report*. However, the Panel
13 recommends that EPA carefully explain how the early life stage internal doses were
14 calculated because serum thyroid stimulating hormone (TSH) levels in newborns are used
15 as a critical effect.

16 17 **Reference Dose**

18
19 In Section 4 of the *Report* EPA discussed the use of two co-critical studies
20 (Mocarelli et al., 2008 and Baccarelli et al., 2008) for development of the reference dose
21 for TCDD. The Panel was asked to comment on the scientific justification for selection
22 and use of these studies to develop the reference dose.

23 24 *a. Selection of Critical Studies and Effects*

25
26 The Panel supports EPA's selection of the Mocarelli et al. (2008) and Baccarelli
27 et al. (2008) studies for identifying "co-critical" effects for the derivation of the reference
28 dose (RfD). The Panel found that these two human epidemiological studies were well
29 designed. The studies provided sufficient exposure information, including biological
30 concentrations that could be used to establish acceptable lifetime daily exposure levels.
31 The rationale for selecting these two studies over numerous other available studies for
32 determining the RfD was clearly described but study weaknesses were not clearly
33 delineated. The Panel recommends that EPA provide a discussion of the strengths and
34 weaknesses of these studies with an indication of whether the weaknesses affect
35 determination of the RfD. In addition, the Panel recommends that the comprehensive
36 data base of both animal and human epidemiological studies be used to demonstrate a
37 consistent and integrative signal of toxicity across species and endpoints for TCDD. The
38 collective impact of the studies should be made stronger in the *Report* by including
39 discussion of both human and experimental animal studies that have examined the effects
40 of dioxin and DLCs on other reproductive and endocrine endpoints. In this regard, dose-
41 response relationships as well as comparisons of no-observed-adverse-effect levels
42 (NOAELs) and lowest-observed-adverse-effect levels (LOAELs) should be discussed.

43
44 The Panel agrees with EPA's assertion that traditional (e.g., immune, endocrine,
45 reproductive) endpoints are more appropriate than biochemical endpoints for establishing

1 points of departure (PODs). The associations of traditional endpoints with health
2 outcomes have been well studied and they are more tightly associated with adverse
3 outcomes than biochemical endpoints. However, EPA should discuss biochemical
4 endpoints, particularly P450s, relevant to establishing and strengthening the proposed
5 reference dose.

6
7 *b. Estimation of Continuous Exposure for Mocarelli et al. (2008)*
8

9 Mocarelli et al. (2008) reported male reproductive effects (decrease in sperm
10 count and motility) observed later in life for boys with high acute exposure to TCDD
11 between the ages of 1 and 9 (average age 5 years), followed by low level background
12 dietary exposure. EPA identified a 10 year critical exposure window and estimated the
13 continuous TCDD intake as the average of the high acute exposure and the 5 year
14 average exposure during the critical exposure window. The Panel found that the pattern
15 of exposure from Seveso posed some extrapolation issues for the EPA, particularly
16 whether the same endpoints and or dose-response from high acute exposures would be
17 expected when extrapolating to low-dose chronic exposures. It would be useful for EPA
18 to provide a discussion of published examples in which dioxin studies were conducted
19 using both high-dose acute and low-dose chronic exposures in animals for the same
20 endpoint and how the outcomes compare both qualitatively and quantitatively. The life
21 stage-specific approach to hazard and dose-response characterization for children's health
22 risk assessment in EPA's *Framework for Assessing Health Risks of Environmental*
23 *Exposures to Children* (EPA, 2006) is also relevant to addressing this issue and should be
24 discussed.

25
26 *c. Designation of a 20% Decrease in Sperm Count as a LOAEL for Mocarelli et al.*
27 *(2008)*
28

29 The Panel supports the use of the change from normal sperm counts and sperm
30 motility for determining an RfD. While the shifts observed in sperm counts may or may
31 not pose a significant health effect in a single individual, such shifts on a population basis
32 could presumably lead to an increased incidence of adverse health outcomes. The Panel
33 recommends that World Health Organization (WHO) reference values for male
34 reproductive parameters and life stage differences in sperm counts in humans be
35 discussed in the *Report*.

36
37 *d. Determination of Effective Exposure Estimate for the Baccarelli et al. (2008)*
38 *Study*
39

40 EPA determined the maternal intake at the LOAEL from the maternal serum-
41 TCDD vs. neonatal TSH regression model by finding the maternal TCDD lipid-adjusted
42 serum concentrations (LASC) at which neonatal TSH exceeded 5 µU/ml. EPA then used
43 the Emond PBPK model under the human gestational scenario to estimate the continuous
44 daily oral TCDD intake that would result in a TCDD LASC corresponding to a neonatal
45 TSH of 5 µU/ml at the end of gestation. EPA estimated the effective maternal intake as

0.024 ng/kg-day. The Panel supports EPA's decision to use the Baccarelli et al. (2008) estimates of the relevant effective doses. The Panel also suggests that since the bulk of the calculations were based on zonal averages of exposed individuals in Baccarelli et al. (2008), EPA should clarify how these measurements relate to ranges and variations in exposure *in utero*.

e. Designation of 5μ-units TSH per ml blood as a LOAEL for Baccarelli et al. (2008)

EPA selected a LOAEL of 5μ-units TSH per ml blood in neonates. The Panel supports EPA's designation of the TSH endpoint within the context of the broader dioxin literature. While the shift observed in TSH levels may or may not pose a significant health effect in a single individual, such a shift on a population basis could presumably lead to an increased incidence of adverse health outcomes. There is a need to better describe the potential adverse health outcomes related to altered neonatal TSH levels. For example, in addition to effects on growth, both cognitive and motor deficits have been found in young adults with congenital hypothyroidism. The *Report* could better describe the consequences of transient hypothyroidism on reproductive outcomes.

f. Selection of Uncertainty Factors

A composite uncertainty factor of 30 (an uncertainty factor of 10 for the lack of a NOAEL, and an uncertainty factor of 3 for human interindividual variability) was applied to the LOAEL of 0.020 ng/kg-day from Mocarelli et al. (2008) to obtain the RfD. The Panel agrees that EPA has used the appropriate uncertainty factors for the derivation of the RfD. However, a short discussion of the decision not to include an uncertainty factor for data quality is needed.

g. Benchmark Dose (BMD) Modeling of animal bioassay data and EPA's Choice of POD from These Studies

The Panel agrees with the BMD modeling approaches used in the *Report*. In addition, the Panel agrees that the animal data have sufficient limitations that preclude their use to establish a RfD.

Cancer Assessment

In Section 5 of the *Report* EPA has provided: a weight-of-evidence characterization of TCDD as a known human carcinogen, conclusions regarding the mode of carcinogenic action for TCDD, EPA's selection of data sets for cancer dose-response modeling, and consideration of approaches for assessment of TCDD carcinogenicity. The Panel was asked to comment on the scientific soundness of these aspects of EPA's cancer assessment.

1 a. Weight-of-Evidence Cancer Descriptor

2
3 The Panel agrees with EPA's conclusion that TCDD is "Carcinogenic to
4 Humans." The Panel recommends that the Agency provide more discussion of the
5 power of the studies used and the difficulties involved when assessing rare tumors. The
6 Panel also recommends that EPA consider including studies with substantial DLC
7 exposure where toxicity equivalence factors (TEFs) can be calculated in the weight-of-
8 evidence discussion. EPA should also attempt to characterize the uncertainty regarding
9 the carcinogenicity of TCDD at low human exposures, since the minimum dose at which
10 carcinogenic effects would be expected to occur cannot be clearly delineated from the
11 current epidemiological human data.

12
13 b. Mode of Action

14
15 The Panel believes the mode of action for TCDD toxicity should be "reasonably
16 well known" rather than "largely unknown," although the Panel agrees that the exact
17 mechanism of action has not been fully delineated for any distinct TCDD toxicity
18 endpoint. The Panel recommends that EPA provide a discussion of the evidence for
19 possible modes of action that include both linear and nonlinear alternatives; and that the
20 description of the nature of a receptor mediated dose-response be expanded by including
21 more evidence regarding the nonlinearity of the receptor mediated dose-response for
22 dioxin.

23
24 c. Selection of Critical Study for Cancer Endpoint

25
26 The Panel agrees with the inclusion of the Cheng et al. (2006) study in the cancer
27 assessment. This study incorporated information on gradation of exposure. However,
28 expanded discussion of several other studies would support the weight-of-evidence for
29 carcinogenicity in less common cancers such as lymphomas and soft tissue sarcoma. The
30 Panel agrees that Cheng et al. (2006) was the appropriate study for quantitative cancer
31 assessment, and that it was appropriate to use all-cancer mortality in this case, because of
32 the extensive dose-response information. The Panel also agrees that the use of the
33 Emond model to estimate risk-specific doses from Cheng et al. (2006) dose-response
34 modeling results was scientifically justified and clearly described but, as previously
35 discussed, the value of the Hill coefficient used in the model is problematic. The Panel
36 found that Cheng et al. (2006) study did not provide completely clear information
37 regarding risks below current background exposure levels. The Panel therefore suggests
38 that EPA expand the discussion to consider the possibility that mode of action
39 considerations could help indicate whether linear extrapolation of the Cheng et al. (2006)
40 data is appropriate to obtain risk estimates in this range of exposures.

41
42 d. Nonlinear Approach for Assessment of TCDD Carcinogenicity

43
44 The Panel found that the *Report* did not respond adequately to the NAS
45 recommendation to adopt "both linear and nonlinear methods of risk characterization to

1 account for the uncertainty of dose-response relationship shape below the ED01.” The
2 Panel recommends that EPA present both linear and nonlinear risk assessment
3 approaches. The nonlinear examples in the document should be formalized and
4 extended. In the absence of a definitive nonlinear mode of action, the linear option
5 results can serve as the baseline for comparison with these other estimates.
6

7 **Quantitative Uncertainty Analysis**

8

9 Section 6 of the *Report* discusses a broad range of philosophical and
10 methodological issues to be considered in conducting an uncertainty analysis for TCDD
11 toxicity. The Panel was asked to comment on: whether the discussion in this section of
12 the *Report* was clearly presented and scientifically justified, the conclusion that a
13 comprehensive quantitative uncertainty analysis (QUA) is not feasible, the discussion
14 regarding volitional uncertainty and how it limits the ability to conduct a QUA, and
15 approaches that EPA used to conduct sensitivity analyses.
16

17 The Panel found that Section 6 of EPA’s *Report* was clearly presented and
18 provided many useful insights for EPA’s dioxin reassessment, but it was not scientifically
19 justified. As further discussed in the responses to Charge Question 6, the Panel does not
20 agree with EPA’s argument that conducting a unified QUA for TCDD toxicity is
21 unfeasible. EPA’s decision to not conduct an integrated QUA may be based primarily on
22 grounds of practicality or timeliness. In particular, EPA argues that a complete
23 quantitative uncertainty analysis would require data and resources not available. We
24 disagree with this logic. More limited evaluations can, and should, be implemented to
25 inform critical issues in the dioxin reassessment. In the response to Charge Question 6.2
26 we suggest a number of methods that could be used. The Panel recommends that EPA
27 revise its argument that QUA for dioxin toxicity is unfeasible.
28

29 EPA’s document contrasted volitional uncertainty with cognitive uncertainty.
30 The Panel recommends that the term “volitional uncertainty,” which might also have
31 been called “decisional uncertainty,” be dropped from the Agency’s document. The
32 Panel recommends that EPA focus on uncertainties about the state of the world and
33 display different modeling choices and the consequences of making them. The Panel
34 recommends that EPA apply standard tools and techniques for analysis of model
35 uncertainty.
36

37 In addition, the Panel found that the sensitivity studies EPA has already
38 completed are useful. The Panel is mindful of the need to minimize further delay of the
39 finalization of EPA’s already protracted dioxin assessment and we recommend that
40 sensitivity studies that EPA has already completed be integrated into whatever overall
41 uncertainty analysis the Agency elects to undertake.
42

INTRODUCTION

EPA has been preparing an assessment of the potential health impacts of 2,3,7,8-Tetrachlorodibenzo-p-Dioxin (TCDD) for many years. In 2003, EPA released an external review draft report entitled, *Exposure and Human Health Reassessment of 2,3,7,8-Tetrachlorodibenzo-p-Dioxin (TCDD) and Related Compounds* (U.S. EPA, 2003) (hereafter referred to as the *2003 Reassessment*) that was reviewed by the EPA Science Advisory Board (SAB), and then by the National Academy of Sciences (NAS). In 2006, the National Research Council (NRC) of the National Academies published their evaluation of EPA's reassessment, *Health Risks from Dioxin and Related Compounds: Evaluation of the EPA Reassessment* (NRC, 2006).

The NAS identified key recommendations that they believed would result in substantial improvement to the *2003 Reassessment* and thus support a scientifically robust characterization of human responses to exposures to TCDD. The NAS recommended that EPA should: more thoroughly justify and communicate its approaches to dose-response modeling for the health effects of dioxin, taking into consideration both nonlinear and linear methods for characterizing cancer risk; improve the transparency and clarity of the selection of key data sets for the dioxin dose-response analysis; reevaluate its cancer weight-of-evidence determination for dioxin based on the Agency's 2005 Cancer Guidelines; consider using physiologically-based pharmacokinetic (PBPK) modeling in the dioxin risk assessment; and improve transparency, thoroughness and clarity in quantitative uncertainty analysis. The NAS also encouraged EPA to calculate a reference dose (RfD), which had not been derived in the *2003 Reassessment*.

In 2010, EPA's Office of Research and Development (ORD) prepared the draft report, *EPA's Reanalysis of Key Issues Related to Dioxin Toxicity and Responses to NAS Comments* (EPA, 2010) (hereafter referred to as the *Report*). The *Report* includes new analyses completed in response to the NAS recommendations and recently published literature, as well as a discussion of topics where EPA's views differed from those of the NAS. The *Report* is not an assessment per se; it was designed to supplement the information provided in EPA's *2003 Reassessment*. However, the *Report* provides a noncancer reference dose and updated cancer values. Detailed discussions of many of the issues addressed in the *Report* are available in the *2003 Reassessment* and were not reproduced in the *Report*.

ORD requested that the EPA Science Advisory Board (SAB) conduct an independent external peer review of the *Report*. In its review, the SAB was asked to consider the accuracy, objectivity, and transparency of EPA's reanalysis and responses.

In response to ORD's request, the SAB convened an expert panel to conduct the review. The Panel held an initial public teleconference on June 24, 2010 to receive an orientation to EPA's *Report*. The Panel then held two public face-to-face meetings (July 13 – 15, 2010 and October 27 – 29, 2010) to deliberate on the charge questions (see Appendix D) and two public teleconferences (March 1 and 2, 2011) to discuss its report.

1 Public comments were provided in oral and written form to the Panel at the
2 teleconferences and face-to-face meetings. There were charge questions on the 6 sections
3 of EPA's *Report*. The questions focused on: transparency and clarity in the selection of
4 key data sets for dose-response analysis, the use of physiologically-based
5 pharmacokinetic (PBPK) modeling in dose-response modeling for cancer and noncancer
6 endpoints, derivation of a proposed oral reference dose (RfD) for noncancer endpoints,
7 cancer weight-of-evidence classification, mode of action of dioxin carcinogenicity,
8 derivation of oral slope factor (OSF) for dioxin, and quantitative uncertainty analysis.
9 This report provides the consensus advice and recommendations of the Panel, with the
10 exception of one member who offered a dissenting opinion mainly on the TCDD
11 carcinogenicity (see Appendix A).
12

RESPONSES TO EPA'S CHARGE QUESTIONS

Charge Question 1. General Charge Questions

The Panel was asked to comment on: whether the EPA's *Report* was clear and logical, whether the Agency had objectively and clearly presented the key National Academy of Sciences (NAS) recommendations, and whether there were other critical studies that would make a significant impact on the conclusions of the hazard characterization or dose-response assessment of the chronic noncancer and cancer health effects of TCDD.

1.1. Is the draft Response to Comments clear and logical? Has EPA objectively and clearly presented the three key NRC recommendations?

Response:

The Panel found that EPA has developed a report that is clear, logical and responsive to many but not all of the recommendations of the NAS. While we provide a general assessment of opportunities for improvement in the context of this overview charge question, most of the issues related to clarity, organization, and responsiveness are addressed more completely and specifically in the context of the subsequent, more specific charge questions.

With respect to the first question, the Panel found that the EPA was effective in developing a clear, transparent, and logical response. The Panel was particularly impressed with the process that EPA used for identifying, reviewing, and evaluating the relevant literature. EPA's process was comprehensive, rigorous, and included public participation. The Agency's report, *EPA's Reanalysis of Key Issues Related to Dioxin Toxicity and Response to NAS Comments*, consists of two volumes. The first volume contains the main text of the *Report* and is 690 pages long (including 37 pages of references). The second volume contains appendices and is 1,159 pages long. Because of the size and complexity of EPA's *Report*, the Panel found the Executive Summary to be particularly important and valuable in providing a concise and accurate summary. As described in detail in the response to Charge Question 2, the Panel identified the need for better integration across chapters and greater clarity and transparency in indicating which studies did not satisfy inclusion criteria, and therefore were not carried forward for further consideration in subsequent chapters of the *Report* to meet particular needs. In contrast, the Panel was satisfied that the inclusion criteria and the associated retained studies were well described and transparent. The Panel acknowledges that, given the enormity of the dioxin published literature, it is not a trivial matter to characterize what has been "left on the cutting room floor." We therefore suggest that EPA do this in a way that provides only a general consideration of this issue.

The *Report* is long and dense with a considerable amount of jargon and in some places it is quite repetitive. These features, while a necessity of this type of document, at times detract from clarity or make the EPA's logic difficult to discern. The Panel found some instances where the *Report* would benefit from greater clarity in writing. For example, topic sentences are sometimes not easily connected to paragraph content. A specific example of this is in the second paragraph on page xxvii of volume 1 where the text does not clearly identify separate EPA activities to address NAS comments. Another example of a *Report* section that could be edited to improve clarity is the qualitative discussion of the uncertainty in the RfD (Section 4.4 of volume 1). The clarity of this section could be improved by including bullet points to highlight and separate key points and/or provide links to information in other sections of the document (e.g., Section 6 – Feasibility of Quantitative Uncertainty Analysis). The Panel suggests a careful review by a qualified technical editor. Similarly, the Panel suggests that the clarity and accessibility of the *Report* could be enhanced by the inclusion of a glossary to help minimize confusion and misinterpretation among the diverse users of the document. At 690 pages, volume 1 is a formidable report. The Panel appreciates the dilemma of preparing a report that is both complete and rigorous and at the same time succinct and efficient, but it is suggested that EPA find additional efficiencies (e.g., greater use of appendices and elimination of redundancies) that would yield a more approachable document.

With respect to the second part of Charge Question 1.1 (i.e., objectivity and clarity of presentation of the three key NAS recommendations), the Panel found that EPA has been successful. The Panel found EPA's *Report* to be clear in presentation of the key NAS recommendations. However, as described more fully in responses to the relevant specific charge questions below, the Panel identified deficiencies in the *Report* with respect to the completeness of its consideration of two critical elements: 1) nonlinear dose-response for TCDD carcinogenicity and 2) uncertainty analysis.

Recommendations

- As further discussed in the response to Charge Question 2, the Panel recommends that the *Report* be revised to provide greater clarity and transparency in the discussion of studies that did not satisfy inclusion criteria for use in the dioxin assessment. Given the enormity of the dioxin published literature, the Panel recognizes that it is not a trivial matter to characterize the studies that were not considered, and therefore the Panel suggests that the *Report* be revised to generally indicate how this issue was considered.
- The *Report* is long and dense and contains a considerable amount of jargon. It would benefit from greater clarity in writing. The Panel therefore recommends that the *Report* be carefully reviewed by a qualified technical editor and revised to incorporate such improvements as better integration across chapters, better

connection between topic sentences and paragraph content, and elimination of repetition.

- The Panel recommends that the clarity and accessibility of EPA's *Report* be enhanced by the inclusion of a glossary to help minimize confusion and misinterpretation among the diverse users of the document.
- The Panel recommends that EPA find additional efficiencies (e.g., greater use of appendices and elimination of redundancies) to yield more succinct and approachable document.
- As discussed in the responses to other charge questions in this report, the Panel identified deficiencies in EPA's *Report* with respect to the completeness of its consideration of two critical elements: 1) nonlinear dose-response for TCDD carcinogenicity, and 2) uncertainty analysis. In the relevant charge question responses below, the Panel has provided recommendations to improve the *Report* in these areas.

1.2. Are there other critical studies that would make a significant impact on the conclusions of the hazard characterization and the dose-response assessment of the chronic noncancer and cancer health effects of TCDD?

Response:

During the course of its discussion, the Panel did not identify any additional studies that would impact the hazard characterization or the dose-response assessment. However, the Panel found that EPA's *Report* should provide more clarity on the exclusion of null epidemiologic studies.

Recommendations

- The Panel recommends that EPA's *Report* provide more discussion and clarity on the exclusion of null epidemiologic studies.

Charge Question 2. Transparency and Clarity in the Selection of Key Data Sets for Dose-Response Analysis

General Comments:

The NAS committee proposed that EPA develop a clear and readily understandable methodology for evaluating and including epidemiologic and animal bioassay data sets in dose-response evaluations. Section 2 of EPA's *Report* describes the Agency's approach to ensuring transparency and clarity in the selection of the studies for

dose-response analyses. The Panel was asked to comment on: whether EPA had been responsive to NAS concerns about transparency and clarity in data set selection, whether the epidemiology and animal bioassay study criteria and considerations had been scientifically justified and clearly described, and whether EPA had applied the epidemiology and animal bioassay study criteria considerations in a scientifically sound manner.

EPA developed and applied two sets of criteria for the animal bioassays and epidemiologic data. The Agency collected and evaluated these studies, including studies from the *2003 Reassessment* and newer studies found through literature searches and through public submissions. The Panel viewed with favor all of the efforts made by EPA to develop this section of the document. The Panel compliments the Agency for its efforts to present the nuanced differences and complicating issues surrounding this subject in a comprehensive and logical manner. The intention of the comments and recommendations provided below is to assist the EPA in further improvement of Section 2.

2.1. Is this section responsive to the NAS concerns about transparency and clarity in data set selection for dose-response analysis?

Response:

Members of the Panel found that Section 2 of the *Report* was responsive to NAS concerns about transparency and clarity. Moreover, it was perceived and appreciated that, in addressing these concerns, EPA had improved the approach in the original *2003 Reassessment*. The EPA's collaboration with Argonne National Laboratory, and invitation to the public to engage in updating the literature search to identify all appropriate studies for evaluation, as well as the conduct of the Dioxin Workshop in February of 2009, were instrumental in enhancing the transparency and clarity regarding the process of selection of studies for the dose-response analysis. The development of clear criteria for study evaluation and inclusion was crucial in addressing the concerns raised by the NAS.

EPA's *Report* presents a clear identification of the study selection process and the studies that were used for dose-response analysis. For example, the process and criteria used to select key data sets for dose-response analyses is described in Section 2.3 of the *Report* and in the Executive Summary. Flow diagrams (e.g., ES-1 and ES-2) clearly demonstrate how studies were chosen for inclusion. Likewise, Appendix B, which includes a point-by-point evaluation of which epidemiological studies were included and excluded, was useful and provides a detailed rationale explaining why the EPA used the particular studies selected in the *Report*. In addition, the results of the literature search performed by EPA are available online. Clarity could be improved by providing search words used for the MedLine searches. A clear case for including high-quality human studies over animal studies is also made.

While Section 2 of the *Report* is deemed responsive to NAS concerns, the Panel found that overall clarity and transparency regarding dataset selection would be further and markedly enhanced if EPA were to make Section 2 (and the document as a whole) more concise. In its present form, Section 2 was viewed by the Panel as overly verbose, to the detriment of overall clarity and we provide the following recommendations to improve the *Report*.

Recommendations

- The Panel strongly recommends careful and extensive editing to revise and consolidate Section 2 and the *Report* as a whole. Specifically, editing should include aspects of English grammar and syntax, minimizing redundancies, and efforts to provide more succinct responses to NAS concerns.
- The Panel recommends restructuring Section 2 to make it easier to follow a study used by EPA from one section of the *Report* to another. In other words, EPA should improve overall document integration using Section 2 as the foundation for this integration.

Charge Questions 2.2 and 2.3

- 2.2. *Are the epidemiology and animal bioassay study criteria/considerations scientifically justified and clearly described?*
- 2.3. *Has EPA applied the epidemiology and animal bioassay study criteria/considerations in a scientifically sound manner? If not, please identify and provide a rationale for alternative approaches.*

Response:

The Panel's discussion of Charge Questions 2.2 and 2.3 was highly integrated. Therefore, comments and specific recommendations that stem from these two questions are presented together.

The Panel found that EPA's study criteria and considerations were scientifically justified and clearly described, and that these were presented in a scientifically sound manner. Thus, Section 2 was deemed responsive to NAS concerns regarding the scientific justification and clarity of description for epidemiology and animal bioassay study criteria/considerations. However, several concerns were discussed by the Panel, and are summarized here.

The Panel's major concern pertains to improving clarity with regard to the decision to include or exclude particular studies and groups of studies from the data sets to be used. The rationale for distinct criteria for epidemiological and animal studies should be made stronger, and data set selection for noncancer and cancer endpoints has

room for further clarification and justification. There was discussion, with differences of opinion among members of the Panel, regarding EPA's scientific justification and clarity of description concerning the Agency's decision to exclude dioxin-like compounds. There was consensus among Panel members that the following recommended improvements would strengthen this section, and thus the document as a whole.

Recommendations

Rationale for excluding dioxin-like compounds

- EPA should better justify the rationale for using studies where the exposure is primarily to TCDD (or for animal studies only to TCDD) to calculate the reference dose. This justification should include scientific and practical reasons.
- EPA should incorporate information from studies with dioxin-like chemicals into a qualitative discussion of the weight-of-evidence for cancer and noncancer endpoints.

Study inclusion and exclusion criteria and considerations

- EPA should further clarify the justifications for study inclusion and exclusion criteria/considerations. To be clear, this recommendation does not indicate that the Panel suggests that a different approach to data set selection is needed. However, the approach used should be explained more effectively and clearly. In this regard, the following specific recommendations are provided to address points of concern raised by Panel members about the study inclusion and exclusion criteria:
 - o EPA should remove the criterion that studies must contain an explicit statement of TCDD purity. For research purposes, TCDD is available from a limited set of vendors, and all sell it as a highly purified compound. Thus, for the animal studies, it is highly unlikely that any study would be conducted using impure TCDD. Therefore, excluding a study simply due to absence of statements regarding TCDD purity runs the risk of excluding high quality studies because the author or journal editorial staff did not elect to include this piece of information.
 - o EPA should revise the explanation of the in vivo mammalian bioassay evaluation indicating that the "study design is consistent with standard toxicological practices." This is too vague as it likely has different meaning to readers from different backgrounds. In addition to defining this more clearly, it is recommended that, if possible, a reference should be provided to an EPA document in which these practices are described in detail.

- o EPA should consider eliminating use of the phrase “outside the range of normal variability,” especially when discussing animal studies.
- o EPA should define the phrase “common practices,” and if possible cite appropriate Agency documents to which the reader can refer for further detail. To provide further context, this recommendation refers specifically to statements such as the following one on page 2-5: “The study criteria shown below and in Figure 2-3 for animal bioassay data reflect EPA’s preferences for TCDD-specific study inclusion, some of which are based on common practices and guidance for POD selection and RfD and OSF derivation.”
- o EPA should provide a more thorough (albeit concise) discussion of data set limitations to educate the reader regarding Agency decisions about study inclusion/exclusion criteria. For instance, consider adding an expanded discussion on suitability of studies of immunological effects and/or thyroid and diabetes (e.g., Baccarelli et al., 2002, 2004; Calvert, 1999; Steenland, 2001).

Considerations concerning selection of epidemiology studies

- The Panel recommends that EPA better justify and explain considerations relating to the selection of epidemiology studies. The following specific recommendations are provided. Many of these specifically address the use of more standard epidemiology vocabulary and descriptors.
 - o EPA evaluated the available epidemiologic cohorts and studies based on five considerations presented on pages 2-6 and 2-7 of the *Report*. The Panel found that Consideration #2 (page 2-6) was worded awkwardly and that epidemiologic terms are misspecified. The Panel therefore recommends that EPA revise Consideration #2 as follows:
 - Define “susceptible to important biases.” This is a non-specific term and the biases should be explained.
 - Clarify what is meant by “control for potential confounding exposures.” Does this refer to only exposure to dioxin-like compound exposures or was it meant to more broadly refer to other exposures as well (NIOSH cohort studies)? Does the text “bias arising from study design” refer to selection bias or is this phrase used more broadly to describe how exposure and outcome are measured and covariate data collected?
 - Define what is meant by the phrase “bias arising from statistical analyses.” It is unclear if bias is the correct term, rather this may refer to model misspecification.

- o With regard to scientific justification and application of Consideration #3 (listed on page 2-7), the Panel recommends that EPA provide more discussion and clarity on the exclusion of null epidemiologic studies.
- o In Exclusion Criterion #3 (listed on page 2-7) EPA should define “reported dose.”
- o The Panel recommends that the discussion in Section 2 of the consideration of “confounding and other potential sources of bias” be clarified. The differences between males and females with regard to TCDD half-life are discussed, but the description of the number of males and females in each study population were often missing or very difficult to determine. Also, in the occupational cohort studies, the possibility of men and women performing different job tasks also increased the possibility that the men and women were exposed at different levels. However, when the job categories with assigned TCDD exposure levels were presented, there was often no discussion of the numbers by gender in the categories. For example, the Manz et al. study (1991) of the Hamburg cohort (1,583 men and 399 women) does not describe the TCDD categories by gender. In addition, the validity of the TCDD exposure levels assigned to the categories was examined “in a group of 48 workers who provided adipose tissue samples” (Page 2-41, lines 18-19). How were these workers selected? How many were approached but refused to provide a sample? Assessment of selection bias in this and other similar circumstances was lacking in some of the studies. This is particularly notable in the lack of overall response rates reported for several of these studies. Inclusion of these factors in the study review would be very helpful.
- o The Panel recommends that discussion of the consideration that “statistical precision, power, and study follow-up are sufficient” be clarified. These metrics can be difficult to determine with the smaller sample size populations, but there are studies that can be very useful even given the small samples. For example, the relative risks calculated for increasing TCDD exposure and risk of breast cancer in the Seveso study were greatly increased in the 3rd and 4th highest exposure categories, but the relative risks were not statistically significant (page 2-56, lines 1-8).

Charge Question 3. The Use of Toxicokinetics in the Dose-Response Modeling for Cancer and Noncancer Endpoints

In its *Report*, EPA used a physiologically-based pharmacokinetic (PBPK) model (Emond et al., 2004, 2005, 2006) with blood concentration as the dose metric rather than first-order body burden. The Panel was asked to comment on the scientific justification for EPA’s application of this model, the model modifications that EPA implemented, and

EPA's characterization of uncertainty in the model. EPA also developed a PBPK model to estimate TCDD concentration in mouse tissues. The Panel was asked to comment on the scientific rationale for development of the mouse model, the performance of the mouse model, and whether model uncertainty had been adequately characterized. In addition, the Panel was asked to comment on the use of the Emond PBPK model to estimate human intake based on internal exposure measures, EPA's sensitivity analysis of the kinetic modeling, and EPA's estimate of lifetime average daily dose.

3.1. The 2003 Reassessment utilized first-order body burden as the dose metric. In the draft Response to Comments document, EPA used a physiologically-based pharmacokinetic (PBPK) model (Emond et al., 2004, 2005, 2006) with whole blood concentration as the dose metric rather than first-order body burden. This PBPK model was chosen, in part, because it includes a biological description of the dose-dependent elimination rate of TCDD. EPA made specific modifications to the published model based on more recent data. Although lipid-adjusted serum concentrations (LASC) for TCDD are commonly used as a dose metric in the literature, EPA chose whole blood TCDD concentrations as the relevant dose metric because serum and serum lipid are not true compartments in the Emond PBPK models (LASC is a side calculation proportional to blood concentration).

Please comment on the following:

3.1.a. Please comment on the justification of applying a PBPK model with whole blood TCDD concentration as a surrogate for tissue TCDD exposure in lieu of using first-order body burden for the dose-response assessment of TCDD.

Response:

The use of body burden in the 2003 Reassessment represents an improvement over the usual default metric of administered dose (mg/kg/d) because the default metric would not properly reflect the accumulation of dioxin in the tissues over time. However, because the accumulation of dioxin in liver is dose-dependent, body burden would not serve as a direct surrogate for tissue exposure. The use of blood concentration is a better choice than body burden because it is more closely related to the biologically relevant dose metric: the free concentration of dioxin in the target tissues (liver, fetus, etc.). Blood concentrations are routinely used to estimate biologically effective exposures for pharmaceuticals.

The rationale for the use of blood concentration rather than lipid adjusted serum concentration (LASC) should not be based on the Emond model structure. It would be trivial to change the model so that LASC could be predicted. Indeed, the model is apparently used to estimate LASCs in the RfD calculations (e.g., page xli, line 21 in the Executive Summary of the *Report*). The question that should be addressed is only whether blood concentrations or LASCs provide better surrogates for cross-species and

cross-study comparisons of free dioxin concentration in the target tissues. LASC is the preferred measure for reporting dioxin biomonitoring data, and is the measurement reported in most of the human epidemiological studies. A metric that considers blood lipid content is also more likely to reflect free dioxin concentration in the plasma, and hence free concentration in the target tissue. The EPA pointed out (page xxxiv in the Executive Summary of the *Report*) that the LASC was related to the blood concentration by a scalar; however, EPA incorrectly concluded that the metrics are equivalent and later (page 3-511, line 6 of the *Report*) discussed the fact that the relationship between them was subject to inter-individual and inter-species variation. If the LASC were used to drive the distribution of TCDD to tissues, the pharmacokinetic outcome would be different from using blood as the driver because the tissue:blood ratio would differ. If the blood fat:blood and tissue:blood values were accounted for in the model, the use of blood and LASC would be similar. It's not clear at this point how this issue was addressed in the dose metric calculations. Consideration of this issue is unlikely to drastically affect the outcome of the risk calculations, but it would be important for a quantitative uncertainty analysis.

Recommendations

- The use of the blood metric is acceptable for the PBPK model. EPA should clarify how the model deals with studies that report the concentration of dioxin in plasma, serum, blood or blood fat:blood measurements.

3.1.b. Please comment on the scientific justification for using the Emond et al. model as opposed to other available TCDD kinetic models.

Response:

The Emond model provided the best available basis for the dose metric calculations in the assessment. It is the product of a high-caliber, multi-year research effort at EPA's National Health and Environmental Effects Research Laboratory, and represents a significant effort in terms of data collection. This model builds on prior PBPK modeling efforts conducted by Andersen et al. (1997). However, additional discussion of other published models and quantitative evaluation of the impact of model selection on dose metric predictions should also be provided.

Recommendations

- The *Report* should discuss how the model was intended to be used in the assessment, which would then dictate why a particular model was selected. That is, for the intended purposes, was the Emond model more robust and/or simpler than other models, and did it contain sufficient details for biological determinants deemed important by the Agency?

1 3.1.c. *Please comment on the modifications implemented by EPA to the published*
2 *Emond et al. model.*
3

4 **Response:**
5

6 The EPA modifications to the published Emond model (modifications described
7 on page 3-44 of the *Report* account for volume of plasma and describe urinary clearance
8 using blood concentration and not a lumped compartment) are minor and appropriate.
9 The model changes are scientifically appropriate and well supported.
10

11 3.1.d. *Please comment on whether EPA adequately characterized the uncertainty in the*
12 *kinetic models.*
13

14 **Response:**
15

16 The *Report* presents a reasonably thorough qualitative characterization of the
17 uncertainty in the kinetic models that is sufficient to support their use in the assessment.
18 A more quantitative uncertainty analysis is needed. Methods that could be useful and
19 informative for such an analysis are suggested in the response to Charge Question 6.2. It
20 is critical to demonstrate the dependence of human equivalent dose (HED) and risk
21 predictions on uncertainty and variability in the model parameters, particularly those with
22 high sensitivity (Evans and Andersen, 2000). Moreover, dose metric uncertainty needs to
23 be determined under the same exposure conditions that dose metrics are calculated: both
24 for the various studies that serve as the basis for the dose-response assessments and for
25 human exposures at the corresponding HEDs and risk specific doses.
26

27 The Hill coefficients for CYP1a1 and CYP1a2 induction used in the Emond
28 model were 1.0 and 0.6, respectively, based on fitting of kinetic data from single doses of
29 dioxin (Wang et al., 1997; Santostefano et al., 1998). However, Walker et al. (1999)
30 subsequently estimated a Hill coefficient of 0.94 for both CYP1a1 and CYP1a2 induction
31 using chronic exposures which were more relevant to the use of the Emond model in the
32 dioxin risk assessment. The value of 0.6 used in the Emond model was well outside the
33 confidence interval of 0.78 to 1.14 reported by Walker et al. (1999). The use of a Hill
34 coefficient value well below unity would lead to a nonlinear model behavior that is
35 biologically implausible (hypersensitivity to induction at doses near zero). As a result,
36 when the human model was used for extrapolation to lower doses (as in the calculation of
37 risk-specific doses) the model would tend to estimate a lower exposure level for a given
38 blood concentration. This effect could be seen in Table ES-1 of the *Report*, where a 5
39 order-of-magnitude change in risk was associated with a 6 order-of-magnitude change in
40 risk specific dose. That is, the model-estimated risk specific doses in the vicinity of 10^{-6}
41 risk were about a factor of 10 lower (more conservative) than linear extrapolation. The
42 evidence for this parameter needs to be carefully reviewed and the reasonable range of
43 values determined. At the least, the human Emond model calculations will need to be
44 repeated with multiple values to characterize the resulting uncertainty in the estimates.

When this is done, the Agency should also consider increasing the fat:blood partition in the human model from 100 to 200 to be more consistent with the human data (Patterson et al., 1988; Schechter and Ryan, 1989; Schechter et al., 1989; Iida et al., 1999; Maruyama et al., 2002). The Hill coefficient is not likely to have as significant an effect on calculations with the animal models, since low-dose extrapolation was not performed in the animals, but this should also be verified by sensitivity/uncertainty analysis of the animal models. Public comments were submitted to the Panel recommending consideration of a Hill coefficient value of 1.0 and pointing out why lower values are inappropriate (comments from Drs. Thomas Starr, July 7, 2010 and October 26, 2010 and Melvin E. Andersen, November 4, 2010).

Recommendations

- The Panel recommends additional efforts to fully characterize the uncertainty in the models with special consideration of the Hill coefficient value.

3.2. *Several of the critical studies for both noncancer and cancer dose-response assessment were conducted in mice. A mouse PBPK model was developed from an existing rat model in order to estimate TCDD concentrations in mouse tissues, including whole blood.*

Please comment on the following:

3.2.a. *Please comment on the scientific rationale for the development of EPA's mouse model based on the published rat model (Emond et al., 2004, 2005, 2006).*

Response:

The Panel agrees that an appropriate approach was used to develop the mouse model on the basis of the published rat model and the available mouse kinetic data. It should be noted that the NAS recommendation to use human data for dose metric could be accomplished because dose-dependent elimination of TCDD has been described in humans, albeit in just a few cases. Dose-dependent elimination has been reported repeatedly in animals and the PBPK model reflected this dose-dependence. Using CYP1A2 data from humans (caffeine metabolism) and mice would offer an opportunity to validate and/or adjust the mouse model.

Recommendations

- An external peer review of the mouse model should be conducted because this model has not been published in the peer-reviewed literature. This is typically a requirement for models to be used by the Agency.

1 3.2.b. *Please comment on the performance of the mouse model in reference to the*
2 *available data.*

3
4 **Response:**

5
6 The Panel found that the mouse model performed reasonably well, apart from
7 under-prediction of urinary excretion data. The urinary excretion data can be improved
8 by taking into account the fact that urine contains metabolites only, which partition
9 differently from the parent compound. The model appeared to be adequate for use in
10 estimating dose metrics for the assessment, but with greater uncertainty than the rat and
11 human models. This was considered a reasonable approach to solve a deficiency in
12 published PPBK models to meet the needs of this assessment.

13
14 The EPA's suggestion in the RfD chapter that the clustering of mouse points of
15 departure (PODs) at the lowest doses was due to mouse model failure was inappropriate
16 and should be rewritten.

17
18 **Recommendations**

- 19
20 • EPA should use the mouse model. The scientific credibility of the model will be
21 enhanced by its publication in an appropriate peer reviewed journal.

22
23 3.2.c. *Please comment on whether EPA adequately characterized the uncertainty in the*
24 *mouse and rat kinetic models. Please comment specifically on the scientific*
25 *justification of the kinetic extrapolation factor from rodents to humans.*

26
27 **Response:**

28
29 EPA provided an adequate characterization of the qualitative uncertainty in the
30 mouse and rat kinetic models sufficient to justify their use, together with the human
31 model, to estimate rodent-to-human extrapolation factors. On the other hand, formal
32 recalibration of the PBPK model parameters using a Hierarchical Bayesian approach such
33 as Markov chain Monte Carlo analysis was not considered necessary or particularly
34 useful. However, a more quantitative uncertainty analysis is needed.

35
36 **Recommendations**

- 37
38 • A more quantitative uncertainty analysis is recommended. Methods that could be
39 useful and informative for such an analysis are suggested in the response to
40 charge question 6.2 in this report.

41
42 3.3. *Please comment on the use of Emond et al. PBPK model to estimate human*
43 *intakes based on internal exposure measures.*
44

1 **Response:**

2
3 The modified Emond model is the best available approach for estimating
4 exposures on the basis of internal exposure measurements. Nevertheless, there is
5 considerable uncertainty associated with attempting to reconstruct prior exposures in a
6 human population (e.g., Seveso).
7

8 **Recommendations**

- 9
10 • The modeling of the Cheng et al. (2006), Moccarelli et al. (2008), and Bacarelli et
11 al. (2008) studies needs to be described in more detail and the impact of model
12 parameter uncertainty and exposure uncertainty in these studies should be
13 evaluated quantitatively.
14

15 3.4. *Please comment on the sensitivity analysis of the kinetic modeling (see Section*
16 *3.3.5).*
17

18 **Response:**

19
20 The *Report* only presented the sensitivity analysis published by Emond et al.
21 (2006), which was not entirely adequate for the purposes of this assessment. The analysis
22 left out the Hill coefficient, which was one of the most important parameters in the model
23 for low dose extrapolation (Evans and Andersen, 2000). Moreover, model sensitivities
24 were species, dose, and dose-scenario dependent, so they need to be determined under the
25 same exposure conditions as those for which dose metrics were calculated: both for the
26 various studies that serve as the basis for the dose-response assessments and for human
27 exposures at the corresponding HEDs and risk specific doses. This represents the most
28 pragmatic path forward for an evaluation of model sensitivity as it relates to potential
29 environmental regulation.
30

31 **Recommendations**

- 32
33 • EPA should provide a sensitivity analysis of the model to authenticate the model
34 for its intended purpose.
35

36 3.5. *Both EPA's noncancer and cancer dose-response assessments are based on a*
37 *lifetime average daily dose. Did EPA appropriately estimate lifetime average*
38 *daily dose? If not, please suggest alternative approaches that could be readily*
39 *developed based on existing data.*
40

41 **Response:**

42
43 The Panel agrees with the average daily dose calculation approaches described in
44 the *Report*. It was not clear to some Panel members how the computational estimates of

internal dose for newborns were carried out since a lactation model was not used. This is important because of the use of TSH in newborns as a critical effect. EPA, and Baccarelli et al. (2008), developed an empirical description of the relationship between maternal TCDD levels (lipid adjusted) in serum at birth of neonate and the measured serum TSH in the newborns up to 3 days of age. The Emond et al. model was run in an iterative fashion by adjusting chronic daily intake (ng/kg/day) in the human gestation model to predict maternal serum level of TCDD at term that was associated with infant serum thyroid stimulating hormone (TSH) concentration of 5 uU/ml (by using the regression equation). The result was 0.024 ng/kg bw/day.

Recommendations

- EPA should carefully explain how the early life stage internal doses are calculated.

Charge Question 4. Reference dose

EPA selected two co-critical studies (Mocarelli et al., 2008 and Baccarelli et al., 2008) for development of the reference dose for TCDD. The Panel was asked to comment on the scientific justification for selection and use of these studies to develop the reference dose.

4.1. *The Mocarelli et al. (2008) and Baccarelli et al. (2008) studies were selected as co-critical studies for the derivation of the RfD. Is the rationale for the choice of Mocarelli and Baccarelli scientifically justified and clearly described? Please identify and provide the rationale for any other studies that should be selected, including the rationale for why the study would be considered a superior candidate for the derivation of the RfD. Also comment on whether the selection of male reproductive effects and changes in neonatal thyroid hormone levels was scientifically justified and clearly described.*

Response:

The Panel found that use of the Mocarelli et al. (2008) and Baccarelli et al. (2008) studies was appropriate for identifying “co-critical” effects for the RfD calculation. These are human epidemiological studies that were well designed and executed. The studies provided sufficient exposure information, including biological concentrations that could be used to help establish acceptable life-time daily exposure levels. Some of the strengths of the human studies included the use of a well-characterized human cohort, conducted by dioxin epidemiology experts, and the fact that similar PODs were found across a broad spectrum of other reported dioxin toxicities in multiple species. The rationale for selecting these two studies over numerous other available studies was clearly described and the Panel believed that, overall, EPA provided a well-considered and rational discussion of why these two human studies were selected for determining the

1 RfD. However, one issue discussed by the Panel was that, while the strengths of the two
2 human studies were well-described, the study weaknesses were not always clearly
3 delineated. For example, in the Baccarelli (2008) study there was limited discussion of
4 how the presence of polychlorinated dibenzo-*p*-dioxins (PCDDs), polychlorinated
5 dibenzofurans (PCDFs) and coplanar polychlorinated biphenyls (PCBs) that were also
6 found in the blood might confound the interpretation of TCDD association with elevated
7 TSH levels. In addition there was no discussion of the potential impact of residential
8 histories (e.g., individuals who may have moved in and out of Zone A after the accident).
9 The Panel believes that more discussion of the strengths and weaknesses of these two
10 studies is needed.

11
12 As indicated above, the Panel agreed that the major strengths of the human studies
13 were the use of a well-characterized dioxin-exposed human cohort, conducted by dioxin
14 epidemiology experts, and the fact that similar PODs were found across a broad spectrum
15 of other reported dioxin toxicities in multiple species. However, in isolation from each
16 other, and lacking a description of supportive animal and epidemiological studies, the
17 studies were less useful for setting the RfD. The Panel emphasizes the need to consider
18 these other supportive studies within the context of the weight of the dioxin and dioxin-
19 like compound (DLC) database. The strength of the RfD should not be based solely on
20 these two human epidemiology studies, but rather should be supported by integration
21 with other similar supporting dioxin and DLC studies. A strong voice from the
22 committee was given for looking at the comprehensive data base of both animal and
23 human epidemiological studies together to demonstrate a consistent and integrative signal
24 of toxicity across species and endpoints for TCDD. It was suggested that similar studies
25 with DLCs should also be included as these would be supportive, at least for a semi-
26 quantitative comparative analysis. This “collective” impact of the studies was stated in
27 the *Report* but needs to be made stronger as it represents the contextual framing for
28 understanding dioxin health impacts. This response would include discussions of both
29 human and experimental animal studies that have examined the effects of dioxin or DLCs
30 on other reproductive and endocrine endpoints and should, for example, include
31 discussion of dose-response relationships as well as comparisons of no-observed-adverse
32 effect levels (NOAELs) and lowest-observed-adverse effect levels (LOAELs).

33
34 The Panel notes that Figures 4.3 and 4.4 in the *Report* show quantitative
35 comparisons across the RfDs and benchmark dose lower bounds (BMDLs) calculated
36 from the animal and epidemiological studies. These figures are useful in understanding
37 the quantitative similarities (to the PODs in the chosen studies) in these calculations. The
38 Panel also notes that since the figures did not have an indication of endpoints being
39 measured, just the reference to the publications, the consistency in signal (i.e., the
40 similarities in PODs determined) was not as readily apparent as it could be.

41
42 Although it has been addressed in the *Report*, the Panel recommends expanding
43 the discussion of the known human age-specific variability in endpoints such as sperm
44 counts, though the data from Moccarelli et al. (2008) do show ranges and variance (in
45 Figure 3/Table 2), and neonatal TSH levels.

Recommendations

- EPA should provide a discussion of the strengths and weaknesses of the Mocarelli et al. (2008) and Baccarelli et al. (2008) studies with an indication of whether the weaknesses affect determination of the RfD.
- EPA should label the endpoints for studies included in Figures 4.3 and 4.4.
- The comprehensive data base of both animal and human epidemiological studies, including studies with DLCs (e.g., studies cited in Goodman et al., 2010), should be discussed together to demonstrate a consistent and integrative signal of toxicity across species and endpoints for TCDD.

4.2. *In the Seveso cohort, the pattern of exposure to TCDD is different from the average daily exposure experienced by the general population. The explosion in Seveso created a high dose pulse of TCDD followed by low level background dietary exposure in the exposed population. In the population, this high dose pulse of TCDD was slowly eliminated from body tissues over time. There is uncertainty regarding the influence of the high-dose pulse exposure on the effects observed later in life.*

4.2.a. *Mocarelli et al. (2008) reported male reproductive effects observed later in life for boys exposed to the high dose pulse of TCDD between the ages of 1 and 10. EPA identified a 10 year critical exposure window. In the development of the candidate RfD, EPA used an exposure averaging approach that differs from the typical approach utilized for animal bioassays. EPA determined that the relevant exposure should be calculated as the mean of the pulse exposure and the 10-year critical exposure window average. Please comment on the following:*

4.2.a.i. *Please comment on EPA's approach for identifying the exposure window and calculating average exposure for this study*

Response:

The Panel discussed extensively, both as part of the deliberations on Section 4 of the *Report* and also as part of the discussion on Section 3, extrapolation issues posed by the pattern of exposure from Seveso. Issues raised included the question of whether the same endpoints and or dose-response would be expected from such exposure scenarios with high acute exposures when extrapolating to low-dose chronic exposures. It would be useful for EPA to provide a discussion of published examples in which dioxin studies were conducted using both high-dose acute and low-dose chronic exposures in animals for the same endpoint and how the outcomes compare both qualitatively and quantitatively. It would be important to determine whether similar results were observed

for similar endpoints. Several Panel members thought there were sufficient data in the immunological or reproductive areas that may allow such a comparison. Several chronic dioxin animal studies may be useful in this regard (Yoshizawa et al., 2009; Sand et al., 2010; Yoshizawa et al., 2010). The life stage-specific approach to hazard and dose-response characterization for children's health risk assessment found in EPA's *Framework for Assessing Health Risks of Environmental Exposures to Children* (EPA, 2006), is also relevant to addressing this issue and should be discussed. The Panel also recommends that the publication of Bell et al., (2010), which summarized and presented data on some differences about chronic vs. acute exposure in maternal transfer, be considered in this discussion.

4.2a.ii. Please comment on EPA's designation of a 20% decrease in sperm count (and an 11% decrease in sperm motility) as a LOAEL for Mocarelli et al. (2008).

Response:

The Panel found that changes from normal sperm counts and sperm motility are of public health relevance and therefore of interest for determining an RfD. Collectively, there was support for these endpoints within the context of the broader dioxin literature. The Panel discussed whether the magnitude of these changes would represent an adverse health effect. While the shifts observed in sperm counts may or may not pose a significant health effect in a single individual, such shifts on a population basis could presumably lead to an increased incidence of adverse health outcomes. Although there was concern expressed about the sample size used for sperm number and known variability in the biological endpoint, the Panel found that sample collection was conducted consistently across subjects and the differences in groups were apparent.

The Panel supports EPA's approach of using the World Health Organization (WHO) reference value for determining relevant TSH levels, and the Panel strongly suggests that further discussion of WHO reference values for male reproductive parameters be included in the *Report*. Several references were available which provided background information and current values recommended by WHO regarding sperm counts (e.g., Skakkebaek, 2010). The Panel suggests that the standard deviations or range of changes from Mocarelli et al. (2008) be discussed in the *Report* because this provides a better understanding of the potential magnitude of effect.

Life stage differences in sperm counts were discussed by the Panel. Members of the public also provided comments on this issue. It would be appropriate to indicate in the *Report* that life stage differences clearly exist in sperm counts in humans and to cite and discuss the EPA life stage document (EPA, 2006).

1 **Recommendations**

- 2
- 3 • Discussion on WHO reference values for male reproductive parameters should be
 - 4 included in the *Report* (e.g., Skakkebaek, 2010).
 - 5
 - 6 • The standard deviations or range of changes from the Mocarelli (2008) study should
 - 7 be discussed in the *Report* to provide a better understanding of the potential
 - 8 magnitude of effect.
 - 9

10 4.2.b. *For Baccarelli et al. (2008), the critical exposure window occurs long after the*

11 *high-dose pulse exposure. Therefore, the variability in the exposure over the*

12 *critical exposure window is likely to be less than the variability in the Mocarelli et*

13 *al. subjects. EPA concluded that the reported maternal exposures from the*

14 *regression model developed by Baccarelli et al. provide an appropriate estimate*

15 *of the relevant effective dose as opposed to extrapolating from the measured*

16 *infant TCDD concentrations to maternal exposure. Additionally, EPA selected a*

17 *LOAEL of 5 μ -units TSH per ml blood in neonates; as this was established by*

18 *World Health Organization (WHO) as a level above which there was concern*

19 *about abnormal thyroid development later in life. Please comment on the*

20 *following:*

21

22 4.2.b.i. *Please comment on EPA's decision to use the reported maternal levels and the*

23 *appropriateness of this exposure estimate for the Baccarelli et al. study.*

24

25 **Response:**

26

27 The Panel discussed and supports EPA's decision to use the Baccarelli et al.

28 (2008) estimates of the relevant effective doses. Since the bulk of the calculations were

29 based on zonal averages, it should be made clearer how these measurements relate to

30 ranges and variations in exposure *in utero*.

31

32 4.2.b.ii. *Please comment on EPA's designation of 5 u-units TSH per ml blood as a*

33 *LOAEL for Baccarelli et al., (2008.)*

34

35 **Response:**

36

37 The change in TSH levels reported by Baccarelli et al. (2008) was of public health

38 relevance and therefore of interest for determining an RfD. Collectively, there was

39 support for this endpoint within the context of the broader dioxin literature. There was

40 discussion on whether the magnitude of these changes would represent an adverse health

41 effect. The Panel notes that the shift observed in TSH levels may or may not pose a

42 significant health effect in a single individual, but such a shift on a population basis could

43 presumably lead to an increased incidence of adverse health outcomes. The Panel also

44 discussed the variability in neonatal TSH levels but concerns about this issue were

1 minimized by the fact that samples were all collected on the same postnatal day. The
2 Panel suggests that if any follow-up data on thyroid hormone levels, such as T3, T4 or
3 TSH levels, are available from the population studied, then these results should be
4 discussed in the *Report*. The Panel discussed several studies describing health effects
5 associated with elevated neonatal TSH levels not always recognized as associated with
6 congenital hyperthyroidism (CH). There is a need to better describe the potential adverse
7 health outcomes related to altered neonatal TSH levels. For example, in addition to
8 effects on growth, both cognitive and motor deficits have been found in young adults
9 with congenital hypothyroidism (Oerbeck et al., 2003; Oerbeck et al., 2007). The *Report*
10 could better describe the consequences of transient hypothyroidism on reproductive
11 outcomes e.g., see Anbalagan et al. (2010). Other references that relate to this question
12 include: Chevrier et al. (2007), Dimitropoulos et al. (2009), and Yr (2008).

13 14 **Recommendations**

- 15
16 • EPA should better describe the potential adverse health outcomes related to altered
17 neonatal TSH levels (e.g., effects on both cognitive and motor deficits).

18
19 4.3. *Please comment on the rationale for the selection of the uncertainty factors (UFs)*
20 *for the RfD. If changes to the selected UFs are proposed, please identify and*
21 *provide a rationale.*

22 23 **Response:**

24
25 A composite uncertainty factor of 30 (an uncertainty factor of 10 for the lack of a
26 NOAEL, and an uncertainty factor of 3 for human interindividual variability) was applied
27 to the LOAEL of 0.020 ng/kg-day from Mocarelli et al. (2008) to obtain the RfD. The
28 Panel agrees that the appropriate uncertainty factors (UFs) were included. The exclusion
29 or inclusion of the UFs in the *Report* is obvious, clearly discussed, and adequately
30 rationalized. The *Report* would be more transparent if EPA included a short discussion
31 of the basis for the decision not to include a UF for data quality.

32
33 4.4. *EPA did not consider biochemical endpoints (such as CYP induction, oxidative*
34 *stress, etc.) as potential critical effects for derivation of the RfD for TCDD due to*
35 *the uncertainties in the qualitative determination of adversity associated with*
36 *such endpoints and quantitative determination of adversity associated with such*
37 *endpoints and quantitative determination of appropriate response levels for these*
38 *types of endpoints in relation to TCDD exposure. Please comment on whether the*
39 *decision not to consider biochemical endpoints is scientifically justified and*
40 *clearly described.*

41 42 **Response:**

43
44 Biochemical endpoints such as P450 activation, increased oxidative stress, etc.
45 may be acceptable endpoints to establish PODs, particularly when the quantitative

1 relationship between the biochemical endpoint and an adverse health outcome is clearly
2 evident. However, with respect to TCDD, the Panel agrees that more traditional
3 endpoints (e.g., immune, endocrine, reproductive) are more appropriate because
4 associations of these endpoints with health outcomes are well studied and provide a
5 stronger association to an adverse outcome than biochemical endpoints. However,
6 because of the wealth of data on P450s and their importance in disease development,
7 normal development, and chemical response to exogenous agents, EPA should discuss
8 biochemical endpoints, particularly P450s, relevant to establishing and strengthening the
9 proposed reference dose.

10
11 4.5. *In using the animal bioassays, EPA averaged internal blood TCDD*
12 *concentrations over the entire dosing period, including 24 hours following the*
13 *last exposure. Please comment on EPA's approach for averaging exposures*
14 *including intermittent and one-day gestation exposure protocols.*
15

16 **Response:**

17
18 For animal studies it has been shown that for some effects acute exposure could
19 give different results than chronic exposure. For TCDD, however, its persistence might
20 suggest that such differences would be partly negated. In Baccarelli et al., (2008), there
21 was extensive discussion regarding the use of the exposure average time for the TCDD
22 concentrations. This is of biological significance as several papers have indicated the
23 unique aspects of high peak exposure of TCDD as occurred in Seveso and in several of
24 the animal studies. The endpoints affected as a result of these peaks do not always
25 translate to impacts from lower chronic exposures. As stated earlier in this section, it
26 would be helpful to discuss any available animal studies comparing high-dose acute vs.
27 low-dose chronic effects on similar endpoints for dioxin or DLCs. By returning to the
28 broader animal literature and using time and dose-response studies from the dioxin and
29 DLC studies, biological support for the two critical endpoints might be found.
30

31 4.6. *Please comment on the benchmark dose (BMD) modeling conducted by EPA to*
32 *analyze the animal bioassay data and EPA's choice of points of departure*
33 *(PODs) from these studies.*
34

35 **Response:**

36
37 The Panel agrees with the BMD modeling approaches used in Section 4 of the
38 *Report*. EPA conclusions that the animal data had sufficient limitations that precluded
39 their use to establish a RfD are adequately justified. The reasons provided, however, are
40 quite diverse, (e.g., no NOAEL, not considered an adverse effect, the effect at the
41 LOAEL is too divergent from the control group, insufficient dose groups at the low-end
42 of the dose-response curve, monotonic responses) and there is no way for the reader to
43 determine which study has particular deficiencies without going back to the original
44 paper. To help address this gap, the Panel suggests that several of the best animal studies
45 be discussed in some detail so these limitations are more apparent to the reader. As

indicated previously, the EPA authors need to better cite the endpoint guidance that is present within EPA documents for defending these approaches and application of BMD models for the critical effects. This is especially necessary given public comments that EPA was not following its own guidelines

4.7. *For the animal bioassay modeling, EPA applied the kinetic extrapolation at the level of the POD prior to applying the uncertainty factors because EPA has less confidence in the kinetic model output at lower doses reflective of the RfD. Please comment on whether the kinetic extrapolation at the level of the POD prior to applying the uncertainty factors was scientifically justified and clearly described.*

Response:

The EPA approach of applying the kinetics on the actual data present at the POD is preferred in this assessment (see additional discussion in the response to Charge Question 3 - The use of toxicokinetics in the dose-response modeling for cancer and noncancer endpoints).

4.8. *Please comment as to whether EPA's qualitative discussion of uncertainty in the RfD is justified and clearly described.*

Response:

The Panel agrees that EPA provided a clear and justified discussion of the uncertainties in deriving the RfD using the Seveso cohort. Section 4 of the *Report* discussed study limitations regarding the need to adjust from acute exposure to average daily dose, the issue of critical windows, co-exposure to DLCs, and the strength/weaknesses of the animal data. The Panel agrees with EPA that the major limitation of the Seveso cohort is the uncertainty arising from how well the effects resulting from high-dose acute exposure translate to low-dose daily exposures. Again, it might be useful to re-review the animal studies to identify whether there are any studies where dioxin or DLCs were administered by acute as well as chronic (or even subchronic) exposure and comparable endpoints were examined. If so, the information can be used to help confirm or refute the accuracy of the "average daily dose" adjustment. This is of particular concern in the Mocarelli study as "time periods of susceptibility" appear in male reproductive development and these periods (windows) may be very short. Again, animal studies, particularly those involving male reproduction, may be helpful.

It would also be useful to include a discussion of potential uncertainty in the exposure estimates from the Baccarelli study. Serum dioxin levels were only established in a subset of the cohort (approximately 51) at the time of the study while dioxin levels from the main cohort were estimated from data collected from zone of residence (A or B) at a much earlier time.

The discussion in the *Report* of whether the background DLC exposure may have a significant impact, particularly at the lower TCDD exposure levels, is important. While the Panel agrees that the true DLC impact can't be determined, it might be helpful to provide some general estimates of the variability that may occur at the proposed RfD.

Charge Question 5. Cancer assessment

In the *Report* EPA has provided: a weight-of-evidence characterization of TCDD as a known human carcinogen, conclusions regarding the mode of carcinogenic action for TCDD, EPA's selection of data sets for cancer dose-response modeling, and consideration of approaches for assessment of TCDD carcinogenicity. The Panel was asked to comment on the scientific soundness of these aspects of EPA's cancer assessment.

General Comment:

The Panel was impressed by the extensive work performed by EPA in its response to the NAS comments on cancer assessment. The comments below are intended to support the Agency in further developing Section 5 of the *Report*.

5.1. Weight-of-Evidence Cancer Descriptor: The 2003 Reassessment concluded that TCDD is a "known human carcinogen." In the current draft Response to Comments document, EPA concluded that under the 2005 Guidelines for Carcinogen Risk Assessment (U.S. EPA, 2005) TCDD is "carcinogenic to humans." Is the weight-of-evidence characterization scientifically justified and clearly described?

Response:

The Panel agrees on the classification that "TCDD is carcinogenic to humans" under EPA's 2005 *Guidelines for Carcinogen Risk Assessment*. Available occupational epidemiologic studies provide convincing evidence of an association between TCDD and human cancer that cannot be reasonably attributed to chance or confounding and other types of bias, and with a demonstration of temporality, strength of association, consistency, biological plausibility, and a biological gradient. Additional evidence from animal studies and from mechanistic studies provides additional support for the classification of TCDD as carcinogenic to humans. A dissenting opinion (see Appendix A of this report) was expressed by one Panel member who indicated that at best, there is equivocal evidence for the carcinogenicity of TCDD in the occupational setting where body burdens were much higher than current or previous background levels. The Panel provides the following recommendations to strengthen the *Report*.

Recommendations

- EPA should provide more discussion of the power of studies used and the difficulties involved when assessing rare tumors. Thoroughly addressing these aspects will make the weight-of-evidence characterization in Section 5 of the *Report* more clear and transparent.
- In the weight-of-evidence characterization, the Agency should build on all the available data to support the decision. It needs to be made clear how different types of data (in vitro, in vivo, human) support each other; or not.
- EPA should consider including studies with substantial DLC exposure where TEFs can be calculated. Specific experimental studies include Li and Rozman (1995), Rozman et al. (1993, 2005), and Viluksela et al. (1994, 1997a,b, 1998a,b).
- EPA should attempt to characterize the uncertainty regarding the carcinogenicity of TCDD at low human exposures, since the minimum dose at which carcinogenic effects would be expected to occur cannot be clearly delineated from the current epidemiological human data. The agency has concluded that AhR activation is a necessary but not sufficient precursor event in the carcinogenic activity of TCDD. Therefore, it would be beneficial if the Agency could evaluate available data on AhR activation and related effects in human cells and animal models to help inform the doses at which these precursor events are observed for comparison with the epidemiological data.

5.2. *Mode of Action: The mode of action of a carcinogen can inform identification of hazards and approaches used for a dose-response assessment. The mode of carcinogenic action for TCDD has not been elucidated for any tumor type. EPA concluded that, while interaction with the Ah receptor is likely to be a necessary early event in TCDD carcinogenicity in experimental animals, the downstream events involved are unknown.*

5.2.a. *Are the available data related to mode(s) of action for the carcinogenicity of TCDD appropriately characterized and clearly presented?*

Response:

The Panel appreciates the attempts by the Agency to further develop cancer mode of action concepts based on available dioxin liver, lung, and thyroid toxicity data. Such innovative and explorative work is clearly fundamental to the continued need to further develop risk assessment sciences and make more detailed and integrated use of already existing and published data.

The Panel complements the Agency for providing an up-to-date dioxin cancer mode of action section in its response to NAS comments. It could, however, be improved by incorporating additional data on linear and nonlinear modes of action in different target tissues and life stages. A large amount of data related to the mode of action for the carcinogenicity of TCDD is described in the *Report*, but the focus appears to be on presenting evidence that supports the use of a default linear approach rather than providing a balanced evaluation of alternative mode of action hypotheses.

The discussion of the likely dose-response for receptor mediated processes focuses only on the first step, binding of the agonist to the receptor, which is ultimately linear at low concentrations. However, no discussion is given to the nature of the dose-response for the down-stream sequelae of receptor activation, for which there is evidence of nonlinearity. It is, in fact, the fundamentally nonlinear nature of the dose-response for receptor mediated processes that underlies the conviction of a large segment of the scientific community that a nonlinear approach should be preferred for the risk assessment for dioxin.

Recommendations

- EPA should further expand the discussion of mode of action data available to delineate linear versus nonlinear modes of action and effects in different target tissues at different life stages.

5.2.b. Do the available data support EPA's conclusion that the overall mode(s) of action for TCDD-induced carcinogenesis is largely unknown? Please comment on whether this evaluation is clearly described.

Response:

The Panel notes that much is known about TCDD toxicity and mode of action. Some Panel members indicated that the characterization of the mode of action should be "reasonably well known" rather than "largely unknown." Nevertheless, the Panel agrees that the exact mechanism-of-action has not been fully delineated for any distinct TCDD-toxicity end-point. For example, it was pointed out that most TCDD toxicities are mediated by activation of the AhR. Many studies have demonstrated that TCDD can activate or interfere with the activity of estrogen receptors, as well as other steroid receptors. Such interference can disrupt the regulation of cell proliferation, cell death and tissue differentiation. By disrupting these cell functions, TCDD can have profound and lasting effects as demonstrated by studies showing that TCDD exposure during development produces adult neural dysfunctions.

1 **Recommendations**

- 2
- 3 • EPA should provide a discussion of the evidence for possible modes of action that
 - 4 include both linear and nonlinear alternatives.
 - 5
 - 6 • EPA should describe the receptor mediated nonlinear mode of action for dioxin
 - 7 (e.g., Van den Heuvel et al., 1994; Li and Rozman 1995; Andersen et al., 1997;
 - 8 Bhattacharya et al 2010; Gim et al., 2010) and DLCs (e.g., Rozman et al., 1993;
 - 9 2005, Stahl et al., 1994; Viluksela et al., 1994, 1997a,b, 1998a,b), as well as
 - 10 evidence regarding the fundamentally nonlinear nature of receptor mediated
 - 11 cellular responses (e.g., Andersen et al., 1999; Louis and Becskei, 2002; Zhang et
 - 12 al., 2010).
 - 13

14 5.3. *Is EPA's approach for selecting data sets from the key epidemiologic studies and*

15 *animal bioassays identified for cancer dose response modeling scientifically*

16 *justified and clearly described?*

17

18 **Response:**

19

20 The Panel agrees with the inclusion of the Cheng study in the cancer assessment.

21 The study incorporated information on gradation of exposure. Expanded discussion of

22 several other studies would support the weight-of-evidence for carcinogenicity in less

23 common cancers such as lymphomas and soft tissue sarcoma. The Panel discussed the

24 possible value of including studies with DLCs in the evaluation of the weight-of-

25 evidence, in light of the small number of studies involving primarily exposure to TCDD.

26 There are a numerous studies in the literature involving the health effects of DLCs.

27 These include rice oil poisoning incidents in Japan and Taiwan. These incidents have

28 been described, and additional references have been provided, in Schecter and Gasiewicz

29 (2003).

30

31 **Recommendations**

- 32
- 33 • EPA should present in a clear and visible format, for example in a table, which
 - 34 studies were carried forward or not, and the reasons for the decisions made. The
 - 35 weight-of-evidence discussion should be expanded to include evidence from
 - 36 studies of individual cancers for which precise gradation of exposure data is
 - 37 lacking.
 - 38

39 5.4. *For the animal bioassay data, potential cancer oral slope factors (OSFs) were*

40 *calculated by linear extrapolation (using a linear, non threshold cancer*

41 *approach) from the point of departure (POD). EPA also estimated the composite*

42 *risk of the occurrence of several tumor types from the animal cancer bioassay*

43 *data.*

44

1 *5.4.a. Please comment on whether the approach for estimating cancer risk, including*
2 *the use of tumor modeling of the TCDD animal cancer bioassay data, is*
3 *scientifically justified and clearly described.*
4

5 **Response:**
6

7 The Panel agrees that the approach for estimating cancer risk from animal studies
8 was scientifically justified and clearly described.
9

10 *5.4.b. Please comment on the choice of using a BMDL01 as the POD for the*
11 *development of candidate oral slope factors derived from the TCDD animal*
12 *cancer bioassays.*
13

14 **Response:**
15

16 The Panel notes the consistency of the selection of the BMDL01 as the POD with
17 Agency guidelines and has no further comments.
18

19 *5.5. EPA selected Cheng et al. (2006) – an analysis of the NIOSH occupational*
20 *cohort – as the critical study for oral slope factor (OSF) development. This study*
21 *was chosen because it considers dose-dependent elimination of TCDD rather than*
22 *first-order kinetics.*
23

24 *5.5.a. Please comment on whether the rationale for this selection is scientifically*
25 *justified and clearly described. Please identify and provide the rationale for any*
26 *other studies that should be considered and provide a critical evaluation of the*
27 *study and of its suitability for meeting the goals of a quantitative cancer*
28 *assessment.*
29

30 **Response:**
31

32 The Panel agrees that Cheng et al (2006) is the appropriate study for OSF
33 development. The selection of this study is well described.
34

35 *5.5.b. Cheng et al. (2006) analyzed all-cancer mortality. Please comment on the use of*
36 *all-cancer mortality as the basis of the OSF.*
37

38 **Response:**
39

40 The Panel agrees that it is appropriate to use all-cancer mortality in this case
41 because of the extensive dose-response information.
42

43 *5.5.c. Please comment on whether the use of the Emond PBPK model in the estimation*
44 *of risk-specific doses from the Cheng et al. dose-response modeling results is*
45 *scientifically justified and clearly described.*

Response:

The Panel agrees that the use of the Emond model to estimate risk-specific doses from the Cheng et al. (2006) dose-response modeling results is scientifically justified and clearly described. This is because the “concentration-and-age-dependent elimination model” (CADM) used in Cheng et al. (2006) did not facilitate this process. Also, the dose conversions were consistent with those used in the derivation of the RfD. However, as discussed in the response to charge question 3.1.d, the Panel is concerned about the value of the Hill coefficient used.

5.5.d. EPA elected to use the log linear relationship of fat concentration and rate ratio to estimate risk-specific doses at all risk levels. EPA could have estimated a POD for cancer risk itself at a single risk level (BMR) for extrapolation to the origin. Please comment on EPA’s choice of extrapolation approach.

Response:

Since the fat concentrations generated by CADM were not linear with the oral exposure at higher doses, a single oral slope factor to be used for all risk levels could not be obtained. EPA used the upper 95% bound on the slope (from Cheng et al., 2006) of the linear relationship between the natural logarithm of the rate ratio and the cumulative fat TCDD concentration (fat-AUC) to estimate risk-specific doses for TCDD at all risk levels. The Panel agrees that the Agency has chosen the appropriate extrapolation approach.

5.5.e. The slope factor derived from Cheng et al. (2006) was extrapolated below the background TCDD exposure levels experienced by the NIOSH cohort. Please comment on this extrapolation.

Response:

The ability of the Cheng study to be informative regarding risks below current background exposure levels is not completely clear.

Recommendations

- EPA should expand the discussion in the *Report* to consider the possibility that mode of action considerations could help to inform whether linear extrapolation of the Cheng data to obtain risk estimates in this range of exposures is appropriate.

5.6. Please comment on whether EPA has clearly described the major qualitative uncertainties in the derivation of the OSF.

1 **Response:**

2
3 The Panel found the description of qualitative uncertainties in the derivation of
4 the OSF to be clear and adequate.
5

6 5.7. *EPA did not consider dioxin-like compounds (DLCs) in the cancer dose-response*
7 *modeling because the occupational exposures in the available cohorts were*
8 *primarily to TCDD. Background DLC exposures were not incorporated in the*
9 *dose-response modeling because EPA judged that it was not possible to*
10 *disaggregate the responses from background exposure to DLCs and occupational*
11 *exposure to TCDD. Please comment on whether this approach is scientifically*
12 *justified and clearly described.*
13

14 **Response:**

15
16 While the Panel found that it was important to include DLC studies in the weight-
17 of-evidence analysis, we are conflicted on their use as a source of dose-response
18 estimates for TCDD. The Panel notes the scientific importance and regulatory relevance
19 of including a coordinated TEQ/DLC discussion in the *Report*. Including TEQ/DLC
20 aspects in the evaluation would allow for the use of additional studies with dose-response
21 information that more closely mirror environmental exposures. On the other hand, the
22 Panel recognizes the complications associated with developing a TCDD risk estimate that
23 is dependent on current TEF values.
24

25 **Recommendations**

- 26
27 • DLC studies should be considered in the weight-of-evidence discussion.
28

29 5.8. *The NRC suggested that EPA consider nonlinear approaches for the assessment*
30 *of TCDD carcinogenicity. In the Response to Comments, EPA presents two*
31 *illustrative nonlinear approaches for cancer, but considers both inappropriate to*
32 *use because lack of MOA information.*
33

34 5.8.a. *Please comment on these two illustrative nonlinear approaches including EPA's*
35 *conclusions regarding the limitations of these approaches.*
36

37 **Response:**

38
39 EPA's *Report* did not respond adequately to the NAS recommendation to adopt
40 "both linear and nonlinear methods of risk characterization to account for the uncertainty
41 of dose-response relationship shape below the ED01." Instead of adopting both linear
42 and nonlinear methods, the EPA argued that only a linear approach could be justified, and
43 derived two examples of RfD development using a nonlinear approach that they
44 characterized as an illustrative exercise only. The choice not to include both linear and

1 nonlinear risk assessment approaches for TCDD was inconsistent with the EPA (2005)
2 cancer guidelines (page 3-23/24):
3

4 “Nonlinear extrapolation having a significant biological support may be presented
5 in addition to a linear approach when the available data and a weight-of-evidence
6 evaluation support a nonlinear approach, but the data are not strong enough to
7 ascertain the mode of action applying the Agency’s mode of action framework.”
8

9 “In the absence of data supporting a biologically based model for extrapolation
10 outside of the observed range, the choice of approach is based on the view of
11 mode of action of the agent arrived at in the hazard assessment. If more than one
12 approach (e.g., both a nonlinear and linear approach) are supported by the data,
13 they should be used and presented to the decision maker.”
14

15 ***Recommendations***

- 16
17 • EPA should present both linear and nonlinear risk assessment approaches. In the
18 absence of a definitive nonlinear mode of action, the linear option results can
19 serve as the baseline for comparison with other estimates. The examples in the
20 current document should be formalized and extended to allow for such a
21 comparison.
22

23 5.8.b. *Are there other nonlinear approaches that could be readily developed based on*
24 *existing data for the assessment of TCDD carcinogenicity? If so, please suggest*
25 *alternative approaches and describe their utility and suitability for meeting the*
26 *goals of a quantitative cancer assessment.*
27

28 ***Recommendations***

- 29
30 • Since the EPA nonlinear analysis only used studies in S-D rats that were
31 identified in Section 2 of the *Report* for potential noncancer dose-response
32 modeling, additional alternative PODs should be added. For example, Simon et
33 al. (2010), which was cited in EPA’s *Report*, provided a number of alternative
34 PODs for a nonlinear approach that should be included in the EPA risk
35 assessment.
36
37

38 **Charge Question 6. Feasibility of Quantitative Uncertainty Analysis**

39
40 In its evaluation of EPA’s 2003 *Reassessment*, the NAS committee recommended
41 that EPA improve the transparency, thoroughness, and clarity in quantitative uncertainty
42 analysis (QUA). Section 6 of EPA’s Response to NAS Comments document addresses
43 NAS comments regarding QUA. The Panel was asked to comment on: whether Section 6
44 of EPA’s *Report* was clearly presented and scientifically justified; EPA’s conclusion that

1 a QUA is not feasible; the discussion of volitional uncertainty, and the utility of the
2 limited sensitivity studies presented by EPA.

3
4 6.1. *Please comment on the discussion in this Section. Is the response clearly*
5 *presented and scientifically justified?*
6

7 **Response:**
8

9 As discussed below, the Panel found that Section 6 of EPA's *Report* is clearly
10 presented, but it is not scientifically justified. In particular, the Panel disagrees with
11 EPA's argument that, since the most detailed and complete available methods for
12 conducting a quantitative uncertainty analysis (QUA) are unfeasible, no analysis can be
13 conducted at all. There are a number of approaches for conducting a QUA, some of
14 which are feasible given current knowledge and data. Specific methods that can be
15 implemented in a timely manner using available data and knowledge are suggested.
16

17 *Clarity of the EPA response to the NAS presented in Section 6 of the Report*
18

19 The EPA response is clearly presented. The *Report* addresses a broad range of
20 philosophical and methodological issues in conducting an uncertainty analysis for TCDD
21 toxicity, specifically for estimates of cancer oral slope factors and noncancer reference
22 doses. Section 6 is successful in identifying the challenges involved in assessing
23 uncertainty in toxicity estimates based on a small set of available models for
24 toxicokinetics, dose-response relationships, and low dose extrapolation, with limited
25 application, testing, and verification; and a small set of animal bioassay, epidemiological
26 or clinical/case studies, many with differing endpoints, dose metrics, and (in the case of
27 the human studies) uncertain exposure and subject data.
28

29 Section 6 of the *Report* provides many useful insights for the Agency's dioxin
30 reassessment. However, in its discussion of available methods, the *Report* is somewhat
31 biased in its treatment of certain statistical methods (discussed below) which could
32 address some of these issues (though the *Report* does note the potential contribution of
33 the methods at the end of Section 6, as part of ongoing or future studies) and overly
34 pessimistic regarding our ability provide improved quantitative estimates for certain
35 portions of the toxicity assessment.
36

37 Some Panel members indicated that the whole section should be rewritten to make
38 it more accessible to non-statisticians. As further discussed in the editorial comments on
39 Section 6 in Appendix C of this report, some phrasing and word choices in the text
40 should be reconsidered, in particular "exotic methods," "volitional uncertainty," and
41 "epistemic uncertainty." The Panel found that the definition of "quantitative uncertainty
42 analysis" was overly narrow and should be expanded to embrace other common and
43 useful methods discussed below. In a few other places, the *Report's* wording in Section 6
44 is strongly at variance with the literature on uncertainty analysis (see editorial comments
45 in Appendix C of this report).

1
2 *Scientific justification of the arguments presented in Section 6*
3

4 The Panel found that the arguments in Section 6 are not scientifically justified. In
5 Section 6, EPA's decision to not do an integrated quantitative uncertainty analysis is
6 presented and a variety of theoretical issues are discussed, but EPA's decision may be
7 based primarily on grounds of practicality or timeliness. EPA indicates that a complete
8 quantitative uncertainty analysis would require data and resources not available. We
9 disagree with this logic. More limited evaluations can, and should, still be implemented
10 to inform critical issues in the dioxin reassessment. EPA should be methodical in
11 considering what variables and components of the assessment would be included in the
12 analysis. The Panel found that the uncertainty narratives and sensitivity analyses already
13 in the document are an excellent beginning and may constitute the lion's share of the
14 work necessary to implement quantitative uncertain analysis based on simple bounding.
15

16 The Panel does not concur with the specific argument EPA used to justify not
17 doing a unified QUA. If the answer to the question of why EPA did not undertake one is
18 that it was not possible to specify precise marginal distributions and dependence
19 functions from existing data, then the conclusion would be that EPA has not been
20 responsive to the NAS criticism, because there are many possible approaches that could
21 be used that do not depend on such specifications. If the argument is that EPA guidance
22 doesn't require a QUA, then one might agree that the NAS criticism is perhaps itself
23 unreasonable. If EPA had asserted that it actually had done an uncertainty analysis in the
24 form of uncertainty factors (UFs) and the limited sensitivity studies that were performed,
25 then that might be understandable, though not consistent with the current state-of-the-art
26 in risk and uncertainty analysis. Even if the argument had been that mounting a QUA is a
27 significant and controversial undertaking itself and that doing one shouldn't delay the
28 finalization of the *Report*, then such a practicality argument would be understandable
29 given the protracted delay in completing the dioxin reassessment.
30

31 Instead, EPA asserts that "Data are the ultimate arbiter of whether quantitative
32 uncertainty analysis ... has sufficient evidentiary support." This flies in the face of how
33 uncertainty analyses are normally conceived. Of course, the absence of data is never a
34 substantive reason *not* to conduct an uncertainty analysis; it is the reason *to* do one.
35

36 In its *Report*, EPA indicates that it needs an "underlying distribution from which
37 to sample" in order to conduct a quantitative uncertainty analysis. The Panel notes that
38 this is not necessarily true, and it is facile to shrug off a call to characterize and account
39 for important uncertainties in the assessment process on these grounds alone. If one can
40 *estimate* the value of a quantity, then one should be able to express the uncertainty about
41 the value, otherwise one does not really have a scientific measurement in the first place.
42 One is not forced to identify precise probability distributions and dependence functions
43 for everything that is to be characterized as uncertain. Even when the uncertainty is
44 volitional (or decisional or just model uncertainty), there can be relevant ranges that are
45 interesting to decision makers and stakeholders. In some cases, the analysis may be

1 formally closer to a sensitivity analysis, but some appropriate response is usually
2 possible, if not always practicable. To its credit, EPA has acknowledged the legitimacy
3 of the call for QUA by NAS and undertaken some efforts in this direction.
4

5 In the *Report*, EPA calls uncertainty analysis an “emerging area in science” and
6 this is inarguably true, but it does not seem reasonable to hold that methodological
7 research is necessary for EPA to do anything more comprehensive to respond to NAS’s
8 criticism, even if we disallow the use of expert elicitation. Under a commitment to the
9 idea that analyses be *data-driven*, it is possible to do something that’s useful, even if it is
10 not predicated on precise distributions. There are a variety of ways to conduct a
11 quantitative uncertainty analysis, even an entirely probabilistic one that obeys the
12 Kolmogorov axioms (Gillies, 2000) that require neither extensive data nor expert
13 elicitation. The response to Charge Question 6.2 below provides a list of various ways
14 (with references) to accomplish this. The list includes probability trees or model choice
15 trees that articulate the structure of the model and dependencies, sensitivity analyses,
16 simple interval analysis that just propagates the plausible ranges, and the supervaluation
17 approach that uses nested inner and outer intervals (with the inner range representing the
18 values that most everyone considers to be plausible values and the outer range
19 representing conservatively broad ranges). There is also a continuous and unbounded
20 version of nesting intervals in an approach known as info-gap analysis that would be
21 useful if one cannot develop finite bounds on some of the inputs. One can also propagate
22 *bounds* on distribution functions, so whatever imperfect information about each input
23 variable’s distribution is available, one can fashion bounds on distribution functions and
24 propagate them through the calculations, with or without assumptions or information
25 about the dependencies among variables.
26

27 The Panel notes that the approaches mentioned above require EPA to make
28 certain modeling judgments, in the same way that developing any analysis requires
29 judgments. However, this does not mean that analysts would be required to make up
30 numbers or elicit any expert opinion. Such an analysis does not necessarily require a lot
31 of extra work by EPA. These methods can be simple to develop, and they are mostly
32 computationally trivial. Of course, the more comprehensive the analysis is, the harder it
33 is to complete. But the analysis does not have to be fully comprehensive to provide
34 useful insights.
35

36 We note that there was not perfect consensus among Panel members about the
37 value of a quantitative uncertainty analysis. Some on the Panel agree that an uncertainty
38 analysis is not an absolute good. For instance, if the final answer is already clear, an
39 uncertainty analysis can be a waste of time and resources. It would not be reasonable to
40 insist on another analysis which would merely waste time and resources. Likewise, if the
41 analysis is done poorly, or without appeal to available evidence from the real world, it
42 can be misleading. For instance, the idea, mentioned in footnote 66 on page 6-20 of
43 EPA’s *Report*, of arbitrarily converting uncertainty factors to independent lognormal
44 random variables in a scattered attempt to mount a QUA would entail a suite of
45 unjustified and probably untenable assumptions rendering the exercise nearly pointless.

Finally, if the analysis is used *strategically* to avoid rendering or finalizing a decision that is proper, it can be counterproductive. However, most members of the Panel felt that quantitative uncertainty analysis is an integral part of any good assessment, and that one is essential to address the many empirically unresolved questions and issues that have arisen in this assessment which beg for explicit consideration in the context of an uncertainty analysis. In its discussion of the other charge questions, the Panel has identified a number of important issues that should be addressed in an eventual uncertainty analysis.

Other methods to be considered

The Panel found that relevant Bayesian methods have been inadequately addressed and improperly dismissed in Section 6. In particular, methods that should be given a more extensive and balanced discussion with more citations to the literature include: 1) Bayesian hierarchical modeling (Axelrad et al., 2007; Choi et al., 2010; Coull et al., 2003; Ryan, 2008) which is used for combining information from multiple studies, and 2) Bayesian model averaging (Morales et al., 2006; Viallefont et al., 2011; Wheeler and Bailer 2007, 2009) which would be useful for considering more than one dose-response equation, while allowing the data to weight their *relative* likelihood and contribution to the estimate. These Bayesian methods should not be referred to as “exotic.” For example, in agreeing with the Section 6 authors that these methods should be pursued in ongoing and future case studies, White et al. (2009) refer to them as “advanced,” rather than exotic. Specifically, they recommend that health scientists should explore statistical approaches to model selection and suggest that “improvements to statistical approaches for model selection, such as model averaging, should be pursued. Case study applications of these advanced statistical approaches will identify potential strengths and weaknesses of the approaches and their significance for risk characterization” (White et al., 2009).

Recommendations

- The Panel recommends that EPA revise Section 6 of the *Report* because, as discussed above, the arguments in this section are not scientifically justified. In particular, EPA should consider revising its argument that quantitative uncertainty analysis is unfeasible for the dioxin assessment. Specific suggestions regarding feasible methods for quantitative uncertainty analysis are provided herein.

6.2. *Please comment on EPA’s overall conclusion that a comprehensive quantitative uncertainty analysis is not feasible.*

Response:

The Panel rejects EPA’s argument that a quantitative uncertainty analysis is unfeasible. Although a quantitative uncertainty analysis is challenging, the Panel does not agree that it is impossible or even impractical to undertake one. While it may well be

1 true that we lack an adequate empirical basis for full Monte-Carlo propagation of input
2 distributions, there are many other options available. Many on the Panel indicated that
3 the present circumstances warrant a compromise approach that would be simple and
4 achievable with modest effort by the Agency. Various bounding approaches, sensitivity
5 studies, uncertainty set analyses, and event trees (probability trees without the
6 probabilities) are suggested as possible approaches that could be used. With such
7 methods, legitimate and comprehensive uncertainty analyses (including even fully
8 probabilistic analyses) are possible. They would be useful and sufficient to respond to
9 NAS' criticism.

10
11 The Panel agrees with EPA's assertion that expert elicitation would be
12 problematic and should be "off the table." However, many on the Panel further
13 suggested that value-of-information methods would also be very useful, although
14 feedback from EPA included reservations about this idea. A discussion of value of
15 information methods is provided in Appendix B of this report.

16
17 The Panel considered the use of bounding approaches for quantitative uncertainty
18 analysis and asked EPA to provide information about the limitations of bounding
19 approaches. In response, EPA asked Dr. Roger Cooke to send the Panel a document on
20 bounding analysis. The short bounding analysis document provided to the Panel by Dr.
21 Cooke focused on the features of interval analysis, although this is not by any means the
22 only approach that might be useful in the context of the dioxin assessment. The bounding
23 analysis document mentions one issue that could be construed as a disadvantage of this
24 simplest bounding approach. It is the idea that the ranges are supposed to be absolute
25 bounds on the possible values of each input variable. So, for instance, the only thing one
26 can say about a percentage is that it is between zero and 100%, or the only thing one can
27 say about a dispersal distance is that it is between zero and the circumference of the Earth
28 (these are Dr. Cooke's examples). But the Panel finds that this criticism seems to
29 represent a misunderstanding of the word "absolute." Vacuous (e.g., physically limiting)
30 bounds are not the only bounds that can be used in interval analysis. In fact, they are
31 meant to be informed by observed study results. Furthermore, one is not necessarily
32 limited to interval ranges and interval analysis.

33
34 The Panel suggests that there are in fact a variety of methods that, with proper
35 application, could be useful and informative, including:

- 36
37 • **Sensitivity analysis studies** (even if not completely comprehensive) (Saltelli et
38 al., 2000a,b; Frey and Patil, 2002),
39 • **Interval analysis** (Moore 1966; Neumaier, 1990) which has been widely used for
40 decades and can be applied to complex models and even blackbox models (Trejo
41 and Kreinovich, 2001),
42 • **Nesting of intervals**, e.g., two levels, wide and narrow can give conservative and
43 optimistic characterizations of overall uncertainty (van Frassen, 1966, 1980),

- 1 • **Probability bounds analysis** (Ferson and Long, 1995; Ferson et al., 2003)
2 including Bayesian p-boxes (Montgomery, 2009), which has been used in a
3 variety of applications (Aughenbaugh and Paredis, 2007; Dixon, 2007; Karanki et
4 al., 2009; Minnery et al., 2009; Regan et al., 2002a,b), including assessments at
5 two Superfund sites (EPA, 2007, 2002-2005),
- 6 • **Info-gap decision theory** (Ben-Haim, 2006) which has been used in several
7 applications, (Davidovitch et al., 2009; Hall and Harvey, 2009; Regan et al., 2005;
8 Rout et al., 2009; Yokomizo, 2009),
- 9 • **Robust optimization** (Bertsimas and Brown, 2009; Bertsimas et al., 2009;
10 Bertsimas et al., 2010; Ben-Tal et al., 2010), and
- 11 • **Probability trees**, which are distributional methods for considering alternative
12 assumptions and models at various stages of the toxicity assessment. Small
13 (2008) explains that the distributional approach for characterizing uncertainty in
14 cancer risk assessment was developed by Evans, Sielken, and co-workers
15 beginning in the 1990s (Holland and Sielken, 1993; Evans et al., 1994a,b, 1995;
16 Sielken, 1990, 1993; Sielken and Valdez-Flores, 1996, 1999; Sielken et al., 1995)
17 and has also been referred to as information analysis, weight-of-evidence analysis,
18 the comprehensive methodology, and comprehensive realism (Sielken, 1990;
19 Sielken et al., 1995, 1996). The method has since been acknowledged in a number
20 of reviews of cancer risk assessment practice and research needs (Boyce, 1998;
21 Moschandreas and Karuchit, 2002; Zeise et al., 2002), and applied in various
22 forms for risk assessment of different chemical compounds (Humphreys et al.,
23 2001; Rai et al., 2002; Kirman et al., 2004; Starr et al., 2006; David et al., 2006;
24 Crump, 1994). The distributional approach enables consideration of a “portfolio-
25 of-mechanisms” that may contribute to carcinogenesis (Cox, 2006).

26
27 These methods are nontrivial and potentially valuable alternatives to traditional
28 probabilistic uncertainty analysis, and they are able to provide insights on critical
29 uncertainties in the assessment endpoints and the ongoing and future research needed to
30 achieve their resolution. The motivation for all of these approaches is the recognition
31 that the use of a single set of assumptions for the components of a cancer risk assessment,
32 whether default, conservative, or otherwise, fails to capture the full range of plausible or
33 likely relationships, how these relationships depend upon our current state of knowledge,
34 the implications for computed values of potency or unit risk, and the opportunities for
35 improved estimates. The methods require modeling judgment as any analysis does, but
36 they can provide a basis for ongoing integration and value of information assessment as
37 new studies and knowledge accumulate over time (Brusick et al., 2008). These methods
38 can at least provide useful bounds on the plausible risks and on the value of information
39 (VOI) of reducing uncertainties further (especially, perhaps, on whether the dose-
40 response relation has a threshold).

41
42 There are, of course, many significant benefits to undertaking a quantitative
43 uncertainty analysis. Although a completely comprehensive analysis might indeed be too
44 much to expect, it is possible and practical to provide readers with much more useful

information about uncertainty. A policy maker might reasonably expect the *Report* to provide insight into major uncertainties and questions such as the following:

- How likely is it that TCDD is not a human carcinogen at current exposure levels? Full discussion of this uncertainty may help to overcome probability neglect and action bias (Patt and Zeckhauser, 2000).
- How likely is it that TCDD at current exposure levels has health effects that have not yet been identified in the toxicological or epidemiological literature (Diamanti-Kandarakis et al., 2009; Soto and Sonnenschein, 2010)?
- What is the probability that reducing TCDD exposures would not reduce cancer risk at all, or only by amounts that would not be measurable, based on recent epidemiological studies and updates such as Pesatori et al. (2009)?
- What is the probability that reducing TCDD exposures would reduce cancer risk in the whole U.S. population, or targeted subpopulations, by amounts significantly greater than a prediction derived from the cancer slope factor estimated by EPA?
- What is the probability that reducing TCDD exposures would increase cancer risk (e.g., if the dose-response relation is J-shaped or U-shaped)?
- What is the decision-analytic value of information (VoI) from collecting more information on AhR kinetics and dose-response before making risk management decisions?
- What is the probability that TCDD interacts with other compounds to which U.S. or targeted subpopulations are exposed, increasing cumulative risk for cancer or other health effects (Carpenter et al., 2002)?

Although many members of the public believe that it is imprudent or even morally wrong to delay tighter regulation of TCDD exposures (perhaps reflecting beliefs that TCDD is a potent carcinogen, developmental toxin, etc.) many on the Panel felt that EPA should provide a thorough quantitative decision analysis that makes explicit the current uncertainties and trade-offs and that shows the conditions under which acting now or postponing action are the optimal actions. Without such quantitative analysis, risk management decisions for TCDD will not be adequately informed, and principles other than those of rational decision making (e.g., the biases discussed in Sunstein and Zeckhauser, 2010) may dominate risk management decisions for TCDD. EPA's uncertainty analysis should provide the scientific basis for improved decision making. The current decision, in effect, to "punt" on quantitative uncertainty analysis is not adequate for informing responsible risk management decision and policy-making, and is not justified.

Recommendations

- The Panel recommends that EPA reconsider the argument for not doing a quantitative uncertainty analysis, or undertake one. EPA could follow the recommendation of the NAS on this point by using one or more of the techniques suggested above.

6.2a. *Please comment on the discussion in Section 6 regarding volitional uncertainty and how this type of uncertainty limits the ability to conduct a quantitative uncertainty analysis.*

Response:

In the *Report*, EPA contrasts volitional uncertainty with cognitive uncertainty. The Panel recommends that the term “volitional uncertainty,” which might also have been called “decisional uncertainty,” should be dropped from the *Report*. EPA should focus instead on uncertainties about the state of world and display the different modeling choices and the consequences of making them. The decisions mentioned in the discussion in Section 6 of volitional uncertainty are modeling choices, and they should be dealt with using techniques for model uncertainty. Standard tools and techniques for analysis of model uncertainty can be applied.

Recommendations

- The Panel recommends that EPA delete from the *Report* the notion of “volitional uncertainty.” EPA should display the different modeling choices and the consequences of making them.

6.3. *Throughout the document (including the Appendices), EPA presents a number of limited sensitivity analyses (e.g., toxicokinetic modeling, RfD ranges, cancer OSF ranges, cancer RfD development). Please comment on the approaches used, and the utility of these sensitivity analyses in clarifying potential significant uncertainties.*

Response:

The Panel congratulates EPA on the sensitivity studies that it has already done and considers them to be very useful. The Panel felt these studies should be integrated and unified in an overall uncertainty analysis. The Panel emphasizes that EPA has already done the lion’s share of the effort needed in their considerations described in the uncertainty narratives. The Panel feels the agency should take credit for this hard work and extend the sensitivity studies to respond fully to the NAS criticism.

The Panel is mindful of the need to minimize further delay of the finalization of this already protracted dioxin assessment. The work the EPA has already done in the sensitivity studies should be leveraged to hasten the completion of whatever uncertainty analysis EPA elects to undertake.

1 ***Recommendations***
2

- 3 • The Panel recommends that sensitivity studies that have already been completed
4 be integrated into whatever overall uncertainty analysis EPA elects to undertake.
5
6
7
8
9

REFERENCES

- Abbaspour, K. C., R. Schulin., E. Schlappi and H. Fluhler 1996. A Bayesian approach for incorporating uncertainty and data worth in environmental projects. *Environmental Modeling and Assessment* 1: 151-158.
- Anbalagan, J, A.M. Sashi, G. Vengatesh, J.A. Stanley, R. Neelamohan, and M.M. Aruldas. 2010. Mechanism underlying transient gestational-onset hypothyroidism-induced impairment of posttesticular sperm maturation in adult rats. *Fertility and Sterility* 93(8): 2491-2497.
- Andersen, M.E., and H.A. Barton. 1999. Biological regulation of receptor-hormone complex concentrations in relation to dose-response assessments for endocrine-active compounds. *Toxicological Sciences* 48(1):38-50.
- Andersen, M.E., L.S. Birnbaum, H.A. Barton, and C.R. Eklund. 1997. Regional hepatic CYP1A1 and CYP1A2 induction with 2,3,7,8-tetrachlorodibenzo-p-dioxin evaluated with a multicompartiment geometric model of hepatic zonation. *Toxicology and Applied Pharmacology* 144(1):145-155.
- Aughenbaugh, J. M., and C.J.J. Paredis. 2007. Probability bounds analysis as a general approach to sensitivity analysis in decision making under uncertainty. *SAE 2007 Transactions Journal of Passenger Cars: Mechanical Systems, (Section 6)* 116:1325-1339, SAE International, Warrendale, Pennsylvania. [Available online at: http://www.srl.gatech.edu/Members/jaugenbaugh/papers_presentations/2007-01-1480.pdf]
- Axelrad D.A., D.C. Bellinger, L.M. Ryan, and T.J. Woodruff. 2007. Dose-response relationship of prenatal mercury exposure and IQ: an integrative analysis of epidemiologic data. *Environmental Health Perspectives* 115:609–615.
- Baccarelli, A, P. Mocarelli, D.G. Patterson, Jr., M. Bonzini, A.C. Pesatori, N. Caporaso, and M.T. Landi. 2002. Immunologic effects of dioxin: new results from Seveso and comparison with other studies. *Environmental Health Perspectives* 110:1169-1173.
- Baccarelli, A., A.C. Pesatori, S.A. Masten, D.G. Patterson, Jr., L.L. Needham, P. Mocarelli, N.E. Caporaso, D. Consonni, J.A. Grassman, P.A. Bertazzi, and M.T. Landi. 2004. Aryl-hydrocarbon receptor-dependent pathway and toxic effects of TCDD in humans: a population-based study in Seveso, Italy. *Toxicology Letters* 149:287-292.
- Baccarelli, A., S.M. Giacomini, C. Corbetta,, M.T. Landi, M. Bonzini, D. Consonni, P. Grillo, D.G. Patterson, Jr., A.C. Pesatori, and P.A. Bertazzi. 2008. Neonatal thyroid function in Seveso 25 years after maternal exposure to dioxin. *PLoS Med* 5: e161.

- 1 Bell, D.R., S. Clode, M.Q. Fan, A. Fernandez, P.M. foster, T. Jiang, G. Loizou, A.
- 2 MacNicol, B.G. miller, M. Rose, L. and S. White. 2010. Interpretation of studies on the
- 3 developmental reproductive toxicology of 2,3,7,8-tetrachlorodibenzo-p-dioxin in male
- 4 offspring. *Food and Chemical Toxicology* 48:1439-1447.
- 5
- 6 Ben-Haim, Y. 2006. *Info-Gap Decision Theory: Decisions Under Severe Uncertainty*.
- 7 2nd edition, Academic Press, London.
- 8
- 9 Ben-Tal, A., W. Davidson, D. Bertsimas, and D.B. Brown. 2010. A soft robust model for
- 10 optimization under ambiguity. *Operations Research* 58 (4): 1220-1234. [Available online
- 11 at:
- 12 [http://www.mit.edu/~dbertsim/papers/Robust%20Optimization/A%20Soft%20Robust%20](http://www.mit.edu/~dbertsim/papers/Robust%20Optimization/A%20Soft%20Robust%20Model%20for%20Optimization%20Under%20Ambiguity.pdf)
- 13 [Model%20for%20Optimization%20Under%20Ambiguity.pdf](http://www.mit.edu/~dbertsim/papers/Robust%20Optimization/A%20Soft%20Robust%20Model%20for%20Optimization%20Under%20Ambiguity.pdf)]
- 14
- 15 Bertsimas, D., and D.B. Brown. 2009. Constructing uncertainty sets for robust linear
- 16 optimization. *Operations Research* 57(6):1483-1495.
- 17
- 18 Bertsimas, D., O. Nohadani, and K.M. Teo. 2009. Nonconvex robust optimization for
- 19 problems with constraints. *INFORMS Journal on Computing*. Articles in Advance, 1-15.
- 20 [Available online at:
- 21 [http://www.mit.edu/~dbertsim/papers/Robust%20Optimization/Nonconvex%20Robust%20](http://www.mit.edu/~dbertsim/papers/Robust%20Optimization/Nonconvex%20Robust%20Optimization%20for%20Problems%20with%20Constraints.pdf)
- 22 [Optimization%20for%20Problems%20with%20Constraints.pdf](http://www.mit.edu/~dbertsim/papers/Robust%20Optimization/Nonconvex%20Robust%20Optimization%20for%20Problems%20with%20Constraints.pdf)]
- 23
- 24 Bertsimas, D., D.A. Iancu, and P.A. Parrilo. 2010. Optimality of affine policies in
- 25 multistage robust optimization. *Mathematics of Operations Research* 35(2):363-394.
- 26 [Available online at:
- 27 [http://www.mit.edu/~dbertsim/papers/Robust%20Optimization/Optimality%20of%20Aff](http://www.mit.edu/~dbertsim/papers/Robust%20Optimization/Optimality%20of%20Affine%20Policies%20in%20Multistage%20Robust%20Optimization.pdf)
- 28 [ine%20Policies%20in%20Multistage%20Robust%20Optimization.pdf](http://www.mit.edu/~dbertsim/papers/Robust%20Optimization/Optimality%20of%20Affine%20Policies%20in%20Multistage%20Robust%20Optimization.pdf)
- 29
- 30 Bhattacharya S, R.B. Conolly, N.E. Kaminski, R.S. Thomas, M.E. Andersen, and Q.
- 31 Zhang. 2010. A bistable switch underlying B-cell differentiation and its disruption by
- 32 the environmental contaminant 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Toxicological*
- 33 *Sciences* 115(1):51-65.
- 34
- 35 Bosgra, S., P.M. Bos, T.G. Vermeire, R.J. Luit, and W. Slob. 2005. Probabilistic risk
- 36 characterization: an example with di(2-ethylhexyl) phthalate. *Regulatory Toxicology and*
- 37 *Pharmacology* 43:104-113.
- 38
- 39 Boyce, C.P. 1998. Comparison of approaches for developing distributions for
- 40 carcinogenic slope factors. *Human and Ecological Risk Assessment* 4 (2):527-577.
- 41
- 42 Brand, K.P., and M.J. Small. 1995. Updating uncertainty in an integrated risk assessment:
- 43 Conceptual framework and methods. *Risk Analysis* 15(6):719-731.
- 44

- 1 Brusick, D., M.J. Small, E.L. Cavalieri, D. Chakravarti, X. Ding, D.G. Longfellow, J.
- 2 Nakamura, E.C. Rogan, and J.S. Swenberg. 2008. Possible genotoxic modes of action for
- 3 naphthalene. *Regulatory Toxicology and Pharmacology* 51 (2):43-50.
- 4
- 5 Calvert, G.M., M.H. Sweney, J. Deddens, and D.K. Wall. 1999. Evaluation of diabetes
- 6 mellitus, serum glucose, and thyroid function among United States workers exposed to
- 7 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Occupational and Environmental Medicine* 56:270-
- 8 276.
- 9
- 10 Carpenter, D.O., K. Arcaro, and D.C. Spink. 2002. Understanding the human health
- 11 effects of chemical mixtures. *Environmental Health Perspectives* 110(suppl. 1):25-42.
- 12
- 13 Chao, P.T., and B.F. Hobbs. 1997. Decision analysis of shoreline protection under
- 14 climate change uncertainty, *Water Resources Research* 33(4):817-830.
- 15
- 16 Cheng, H, L. Aylward, C. Beall, T.B. Starr, R.C. Brunet, G. Carrier, and E. Delzell. 2006.
- 17 TCDD exposure-response analysis and risk assessment. *Risk Analysis* 26: 1059-1071.
- 18
- 19 Chevrier, J, B. Eskenazi, A. Bradman, L. Fenster, and D.B. Barr. 2007. Associations
- 20 between prenatal exposure to polychlorinated biphenyls and neonatal thyroid-stimulating
- 21 hormone levels in a Mexican-American population, Salinas Valley, California.
- 22 *Environmental Health Perspectives* 115(10):1490-1496.
- 23
- 24 Choi, T., M.J. Schervish, K.A. Schmitt and M.J. Small. 2010. Bayesian hierarchical
- 25 analysis for multiple health endpoints in a toxicity study. *Journal of Agricultural,*
- 26 *Biological, and Environmental Statistics* 15(3):290-307 [Available online at:
- 27 <http://www.springerlink.com/content/2h416p2581210773/fulltext.pdf>]
- 28
- 29 Clemen, R.T. 1996. *Making Hard Decisions: An Introduction to Decision Analysis* (2nd
- 30 ed.), Duxbury Press, Belmont, CA.
- 31
- 32 Cooke, R.M. 2009. (ed) *Uncertainty Modeling in Dose Response, Bench-testing*
- 33 *Environmental Models*. Wiley, New York.
- 34
- 35 Coull, B., M. Menzetti, and L. Ryan. 2003. A Bayesian hierarchical model for risk
- 36 assessment of methylmercury, *Journal of Agricultural, Biological and Environmental*
- 37 *Statistics* 8(3):253-270.
- 38
- 39 Cox, L.A. 2006. Quantifying potential health impacts of cadmium in cigarettes on smoker
- 40 risk of lung cancer: A portfolio-of-mechanisms approach. *Risk Analysis* 26(6):1581-1599.
- 41
- 42 Crump, K.S. 1994. Risk of benzene-induced leukemia: A sensitivity analysis of the
- 43 Pliofilm cohort with additional follow-up and new exposure estimates. *Journal of*
- 44 *Toxicology and Environmental Health* 42:219-242.
- 45

- 1 David, R.M., H.J. Clewell, P.R. Gentry, T.R. Covington, D.A. Morgott, and D.J. Marino.
2 2006. Revised assessment of cancer risk to dichloromethane II. Application of
3 probabilistic methods to cancer risk determinations. *Regulatory Toxicology and*
4 *Pharmacology* 45(1):55–65.
- 5
6 Davidovitch, L., R. Stoklosa, J. Majer, A. Nietrzeba, P. Whittle, K. Mengersen, and Y.
7 Ben-Haim. 2009. Info-gap theory and robust design of surveillance for invasive species:
8 The case study of Barrow Island. *Journal of Environmental Management* 90(8):2785-
9 2793.
- 10
11 Diamante-Kandarakis, E., J-P. Bourguignon, L.C. Giudice, R. Hauser, G.S. Prins, A.M.
12 Soto, R. T. Zoeller, and A.C. Gore. 2009. Endocrine disrupting chemicals: an Endocrine
13 Society scientific statement. *Endocrine Reviews* 30:293-342.
- 14
15 Dimitropoulos, A, L. Molinari, K. Etter, M. Lang-Muritano, O.G. Jenni, R.H. Largo, and
16 B. Latal. 2009. Children with congenital hypothyroidism: long-term intellectual outcome
17 after early high-dose treatment. *Pediatric Research* 65(2):242-248.
- 18
19 Dixon, W.J. 2007. *The Use of Probability Bounds Analysis for Characterizing and*
20 *Propagating Uncertainty in Species Sensitivity Distributions*. Arthur Rylah Institute for
21 Environmental Research, Technical Report Series No. 163, Department of Sustainability
22 and Environment. Victoria, Australia [Available online at:
23 [http://www.land.vic.gov.au/CA256F310024B628/0/F11A9B711262E4C3CA257384001](http://www.land.vic.gov.au/CA256F310024B628/0/F11A9B711262E4C3CA2573840014660E/$File/ARI+Technical+Report+163+-+The+use+of+probability+bounds+analysis+for+characterising+and+propagating+uncertainty+in+species+sensitivity+distributions.pdf)
24 [4660E/\\$File/ARI+Technical+Report+163+-](http://www.land.vic.gov.au/CA256F310024B628/0/F11A9B711262E4C3CA2573840014660E/$File/ARI+Technical+Report+163+-+The+use+of+probability+bounds+analysis+for+characterising+and+propagating+uncertainty+in+species+sensitivity+distributions.pdf)
25 [+The+use+of+probability+bounds+analysis+for+characterising+and+propagating+uncert](http://www.land.vic.gov.au/CA256F310024B628/0/F11A9B711262E4C3CA2573840014660E/$File/ARI+Technical+Report+163+-+The+use+of+probability+bounds+analysis+for+characterising+and+propagating+uncertainty+in+species+sensitivity+distributions.pdf)
26 [ainty+in+species+sensitivity+distributions.pdf\]](http://www.land.vic.gov.au/CA256F310024B628/0/F11A9B711262E4C3CA2573840014660E/$File/ARI+Technical+Report+163+-+The+use+of+probability+bounds+analysis+for+characterising+and+propagating+uncertainty+in+species+sensitivity+distributions.pdf)
27
- 28 Emond, C, L.S. Birnbaum, and M.J. DeVito 2004. Physiologically based
29 pharmacokinetic model for developmental exposures to TCDD in the rat. *Toxicological*
30 *Sciences* 80:115-133.
- 31
32 Emond, C, J.E. Michalek, and L.S. Birnbaum. 2005. Comparison of the use of a
33 physiologically based pharmacokinetic model and a classical pharmacokinetic model for
34 dioxin exposure assessments. *Environmental Health Perspectives* 113:1666-1668.
- 35
36 Emond, C, L.S. Birnbaum, and M.J. DeVito. 2006. Use of a physiologically based
37 pharmacokinetic model for rats to study the influence of body fat mass and induction of
38 CYP1A2 on the pharmacokinetics of TCDD. *Environmental Health Perspectives* 114:
39 1394-1400.
- 40
41
42
43
44

- 1 EPA (U.S. Environmental Protection Agency) 2002-2005. *GE/Housatonic River Site in*
2 *New England*. Several documents available online at
3 <http://www.epa.gov/region1/ge/thesite/restofriver-reports.html> describing the U.S. EPA's
4 use of probability bounds analysis in the "rest of river" component of the assessment for
5 the GE/Housatonic River Site in New England, including both the Ecological Risk
6 Assessment and the Human Health Risk Assessment.
7
8 EPA (U.S. Environmental Protection Agency). 2006. *A framework for assessing health*
9 *risks of environmental exposures to children*. EPA/600/R-05/093F, National Center for
10 Environmental Assessment, Office of Research and Development, Washington, D.C.
11
12 EPA (U. S. Environmental Protection Agency). 2007. *Region 6 Superfund Program:*
13 *Calcasieu Estuary Initiative*. <http://www.epa.gov/earth1r6/6sf/sfsites/calcsinit.htm>,
14 [accessed July 2009]
15
16 EPA (U.S. Environmental Protection Agency). 2010. *EPA's Reanalysis of key issues*
17 *related to dioxin toxicity and response to NAS comments*. EPA/600/R-10/038A.
18 (May 2010 draft)
19
20 Evans, M.V. and Andersen, M.E. 2000. Sensitivity analysis of a physiological model for
21 2,3,7,8,-tetrachlorodibenzo-p-dioxin (TCDD): assessing the impact of specific model
22 parameters on sequestration in liver and fat in the rat. *Toxicological Sciences* 54:71-80.
23
24 Evans, J.S., J.D. Graham, G.M. Gray, and R.L. Sielken Jr. 1994a. A distributional
25 approach to characterizing low-dose cancer risk. *Risk Analysis* 14(1):25-34.
26
27 Evans, J.S., G.M. Gray, R.L. Sielken Jr., A.E. Smith, C. Valdez-Flores, and J.D. Graham.
28 1994b. Use of probabilistic expert judgment in distributional analysis of carcinogenic
29 potency. *Regulatory Toxicology and Pharmacology* 20(1):15-36.
30
31 Evans, J.S., J.D. Graham, G.M. Gray, R.L., and Sielken Jr. 1995. A distributional
32 approach to characterizing low-dose cancer risk. In *Low-Dose Extrapolation of Cancer*
33 *Risks*, S. Olin, W. Farland, C. Park, L. Rhomberg, R. Scheuplein, T. Starr, J. Wilson
34 (eds.), 253-274, ILSI Press Washington, DC.
35
36 Ferson, S., and T.F. Long. 1995. Conservative uncertainty propagation in environmental
37 risk assessments. *Environmental Toxicology and Risk Assessment, Third Volume, ASTM*
38 *STP 1218*, J.S. Hughes, G.R. Biddinger and E. Mones (eds.) 97-110, American Society
39 for Testing and Materials, Philadelphia, .
40
41 Ferson, S., V. Kreinovich, L. Ginzburg, D.S. Myers, and K. Sentz. 2003. *Constructing*
42 *Probability Boxes and Dempster-Shafer Structures*. SAND2002-4015. Sandia National
43 Laboratories, Albuquerque, NM.
44

- 1 Ferson, S., C.A. Joslyn, J.C. Helton, W.L. Oberkampf, and K. Sentz. 2004. Summary
2 from the epistemic uncertainty workshop: consensus amid diversity. *Reliability and*
3 *Engineering and System Safety* 85:355-370.
- 4
- 5 Finkel, A.M., and Evans, J.S. 1987. Evaluating the benefits of uncertainty reduction in
6 environmental health risk management, *Journal Air Pollution Control Association*,
7 37:1164-1171.
- 8
- 9 Gillies, D. 2000. *Philosophical Theories of Probability*. Routledge, New York, NY.
- 10
- 11 Gim, J., H.S. Kim, J. Kim, M. Choi, J.R. Kim, Y.J. Chung, and K.H. Cho. 2010. A
12 system-level investigation into the cellular toxic response mechanism mediated by AhR
13 signal transduction pathway. *Bioinformatics* 26(17):2169-2175.
- 14
- 15 Glimcher, P.W. 2003. *Decisions, Uncertainty, and the Brain: The Science of*
16 *Neuroeconomics*. The MIT Press, Cambridge, MA.
- 17
- 18 Goodman, J.E., L.E., Kerper, C.P. Boyce, R.L. Prueitt, and L.R. Rhomberg. 2010.
19 Weight-of-evidence analysis of human exposures to dioxins and dioxin-like compounds
20 and associations with thyroid hormone levels during early development. *Regulatory*
21 *Toxicology and Pharmacology* 58(1):79-99.
- 22
- 23 Hall, J.W., and H. Harvey 2009. Decision making under severe uncertainty for flood risk
24 management: a case study of info-gap robustness analysis. *Proceedings of the Eighth*
25 *International Conference on Hydroinformatics*, January 12-16, 2009, Concepcion, Chile.
26 [Available online at: [http://www.floodsite.net/html/partner_area/project_docs/T20-08-05-](http://www.floodsite.net/html/partner_area/project_docs/T20-08-05-hic-infogap.pdf)
27 [hic-infogap.pdf](http://www.floodsite.net/html/partner_area/project_docs/T20-08-05-hic-infogap.pdf)]
- 28
- 29 Holland, C.D., and R.L. Sielken, Jr. 1993. *Quantitative Cancer Modeling and Risk*
30 *Assessment* (Chapter 7). Prentice Hall, Englewood Cliffs, NJ.
- 31
- 32 Hornberger, G. M., and R. C. Spear. 1983. An approach to the analysis of behavior and
33 sensitivity in environmental systems. in *Uncertainty and Forecasting of Water Quality*,
34 M.B. Beck and G. van Stratten (eds.), 101-116, Springer-Verlag.
- 35
- 36 Hsu, M., M. Bhatt, R. Adolphs, D. Tranel, and C.F. Camerer. 2005. Neural systems
37 responding to degrees of uncertainty in human decision-making. *Science* 310
38 (5754):1680-1683. [Available online at:
39 <http://www.sciencemag.org/content/310/5754/1680.abstract>]
- 40
- 41 Humphreys, S.H., C. Carrington, and M. Bolger. 2001. A quantitative risk assessment for
42 fumonisins B1 and B2 in US corn. *Food Additives and Contaminants* 18 (3):211-220.
- 43

- 1 Iida, T., H. Hirakawa, T. Matsueda, J. Magayama, and T. Nagata. 1999. Polychlorinated
2 dibenzo-*p*-dioxins and related compounds: correlations of levels in human tissues and in
3 blood. *Chemosphere* 38:2767-2774.
- 4
- 5 Karanki, D.R., H.S. Kushwaha, A.K. Verma, and S. Ajit. 2009. Uncertainty analysis
6 based on probability bounds (p-box) approach in probabilistic safety assessment. *Risk*
7 *Analysis* 29(5):662-75.
- 8
- 9 Keeney, R.L. 1982. Decision analysis: an overview, *Operations Research* 39: 803-838.
- 10
- 11 Kirman, C. R., L.M. Sweeney, M.J. Teta, R.L. Sielken, C. Valdez-Flores, R.J. Albertini,
12 and M.L. Gargas. 2004. Addressing nonlinearity in the exposure-response relationship for
13 a genotoxic carcinogen: Cancer potency estimates for ethylene oxide. *Risk Analysis*
14 24(5):1165-1183.
- 15
- 16 Li, X., and K.K. Rozman. 1995. Subchronic effects of 2,3,7,8-tetrachlorodibenzo-*p*-
17 dioxin (TCDD) and their reversibility in male Sprague-Dawley rats. *Toxicology* 97:133-
18 140.
- 19
- 20 Louis M., and A. Becskei. 2002. Binary and Graded Responses in Gene Networks.
21 *Sciences Signal Transduction Knowledge Environment* (143):pe33.
- 22
- 23 Manz, A., J. Berger, J.H. Dwyer, D. Flesch-Janys, S. Nagel, and H. Waltsgott. 1991.
24 Cancer mortality among workers in chemical plant contaminated with dioxin. *Lancet*
25 338: 959-964.
- 26
- 27 Merz, R., J. Parajka, and G. Blöschl. 2009. Scale effects in conceptual hydrological
28 modeling *Water Resources Research* 45:W09405.
- 29
- 30 Minnery, J.G., J.G. Jacangelo, L.I. Boden, D.J. Vorhees and W. Heiger-Bernays. 2009.
31 Sensitivity analysis of the pressure-based direct integrity test for membranes used in
32 drinking water treatment. *Environmental Science and Technology* 43(24):9419-9424.
- 33
- 34 Mocarelli, P., P.M. Gerthoux, D.G. Patterson, Jr., S. Milani, G. Limonta, M. Bertona, S.
35 Signorini, P. Tramacere, L. Colombo, C. Crespi, P. Brambilla, C. Sarto, V. Carreri, E.
36 Sampson, W.E. Turner, and L. Needham. 2008. Dioxin exposure, from infancy through
37 puberty, produces endocrine disruption and affects human semen quality. *Environmental*
38 *Health Perspectives* 116:70-77.
- 39
- 40 Montgomery, V. 2009. *New Statistical Methods in Risk Assessment by Probability*
41 *Bounds*. Ph.D. dissertation, Durham University, UK. [Available online at:
42 <http://maths.dur.ac.uk/stats/people/fc/thesis-VM.pdf>]
- 43
- 44 Moore, R.E. 1966. *Interval Analysis*. Prentice Hall, Englewood Cliffs, NJ.
- 45

- 1 Morales, K.H., J.G. Ibrahim, C.-J. Chen, and L.M. Ryan. 2006. Bayesian model
2 averaging with applications to benchmark dose estimation for arsenic in drinking water.
3 *Journal of the American Statistical Association* 101(473):9-17.
4
- 5 Moschandreas, D.J., and S. Karuchit, 2002. Scenario-model-parameter - A new method
6 of cumulative risk uncertainty analysis. *Environment International* 28 (4):247-261.
7
- 8 NRC (National Research Council). 2006. *Health risks from Dioxin and Related*
9 *Compounds: Evaluation of the EPA Reassessment*. National Academies Press,
10 Washington, D.C. [Available online at:
11 http://www.nap.edu/catalog.php?record_id=11688]
12
- 13 Neumaier, A. 1990. *Interval Methods for Systems of Equations*. Cambridge University
14 Press, Cambridge, UK.
15
- 16 Oberkampf, W.L., J.C. Helton, C.A. Joslyn, S.F. Wojtkiewicz, and S. Ferson. 2004.
17 Challenge problems: uncertainty in system response given uncertain parameters.
18 *Reliability and Engineering and System Safety* 85:11-20.
19
- 20 Oerbeck, B., K. Sundet, B.F. Kase, and S. Heyerdahl. 2003 Congenital hypothyroidism:
21 influence of disease severity and L-thyroxine treatment on intellectual, motor, and
22 school-associated outcomes in young adults. *Pediatrics* 112:923-930.
23
- 24 Oerbeck, B., I. Reinvang, K. Sundet, and S. Heyerdahl. 2007. Young adults with severe
25 congenital hypothyroidism: cognitive event related potentials (ERPs) and the significance
26 of an early start of thyroxine treatment. *Scandinavian Journal of Psychology* 48:61-67.
27
- 28 Paterson, S, and D. Mackay. 1987. A steady-state fugacity-based pharmacokinetic model
29 with simultaneous multiple exposure routes. *Chemosphere* 6:395-408.
30
- 31 Patt, A., and R. Zeckhauser. 2000. Action bias and environmental decisions. *Journal of*
32 *Risk and Uncertainty* 21(1):45-72. [Available online at:
33 <http://www.springerlink.com/content/k47064873365w720/>]
34
- 35 Patwardhan, A., and M.J. Small. 1992. Bayesian methods for model uncertainty analysis
36 with application to future sea level rise. *Risk Analysis* 4:513-523.
37
- 38 Pesatori, A.C., D. Consonni, M. Rubagotti, P. Grillo, and P.A. Bertazzi. 2009. Cancer
39 incidence in the population exposed to dioxin after the "Seveso accident": twenty years of
40 follow-up. *Environmental Health* 8:39. [Available online at:
41 <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2754980/?tool=pmcentrez&report=abstract>]
42
43

- 1 Rai, S.N., S. Bartlett, D. Krewski, and J. Paterson. 2002. The use of probabilistic risk
2 assessment in establishing drinking water quality objectives. *Human and Ecological Risk*
3 *Assessment* 8(3):493-509.
- 4
- 5 Raiffa, H. 1968. *Decision Analysis: Introductory Lectures on Choice under Uncertainty*.
6 Addison-Wesley, Reading, MA.
- 7
- 8 Regan, H.M., B.K. Hope, and S. Ferson. 2002a. Analysis and portrayal of uncertainty in a
9 food-web exposure model. *Human and Ecological Risk Assessment* 8:1757-1777.
- 10
- 11 Regan, H.M., B.E. Sample, and S. Ferson. 2002b. Comparison of deterministic and
12 probabilistic calculation of ecological soil screening levels. *Environmental Toxicology*
13 *and Chemistry* 21:882-890.
- 14
- 15 Regan, H.M., Y. Ben-Haim, B. Langford, W.G. Wilson, P. Lundberg, S.J. Andelman, and
16 M.A. Burgman. 2005. Robust decision-making under severe uncertainty for conservation
17 management. *Ecological Applications* 15(4):1471-1477.
- 18
- 19 Rout, T.M., C.J. Thompson, and M.A. McCarthy. 2009. Robust decisions for declaring
20 eradication of invasive species. *Journal of Applied Ecology* 46:782-786.
- 21
- 22 Rozman, K.K., B.U. Stahl, M. Viluksela, and L.S. Birnbaum. 1993. Multiple dose
23 (subchronic) toxicity of heptachlorodibenzo-p-dioxin. In *Dioxin '93*, H. Fiedler, H. Frank,
24 O. Hutzinger, W. Parzefall, A. Riss and S. Safe (eds.) 13:133-136.
- 25
- 26 Rozman, K.K., M. Lebofsky, and D.M. Pinson. 2005. Chronic toxicity and
27 carcinogenicity of 1,2,3,4,6,7,8-heptachlorodibenzo-p-dioxin displays a distinct dose/time
28 threshold ($c \times t = k$) and a profound subthreshold hormetic effect: prolongation of life.
29 *Food and Chemical Toxicology* 43:729-40.
- 30
- 31 Ryan, L. 2008. Combining data from multiple sources, with applications to
32 environmental risk assessment. *Statistics in Medicine* 27(5):698-710.
- 33
- 34 Santostefano, M.J., X. Wang, V.M. Richardson, D.G. Ross, M.J. DeVito, and L.S.
35 Birnbaum 1998. A pharmacodynamic analysis of TCDD-induced cytochrome P450 gene
36 expression in multiple tissues: dose- and time-dependent effects. *Toxicology and Applied*
37 *Pharmacology* 151:294-310.
- 38
- 39 Sand, S., N. Fletcher, D. von Rosen, F. Kalantari, M. Viluksela, J.T. Tuomisto, J.
40 Tuomisto, A. Falk-Filipsson, and H. Håkansson, 2010. Quantitative and statistical
41 analysis of differences in sensitivity between Long-Evans and Han/Wistar rats following
42 long-term exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Regulatory Toxicology and*
43 *Pharmacology* 57(2-3):136-145.
- 44

- 1 Schechter, A.J., and Ryan, J.J. 1989. Blood and adipose tissue levels of PCDDs/PCDFs
2 over three years in a patient after exposure to polychlorinated dioxins and dibenzofurans.
3 *Chemosphere* 18(1-6):635-642.
4
5 Schechter, A., and T. Gasiewicz (eds). 2003. *Dioxins and Health*, 2nd Edition. Wiley
6 Interscience, Hoboken, NJ.
7
8 Schechter, A.J., J. Mes, and D. Davies. 1989. Polychlorinated biphenyl (PCB), DDT, DDE
9 and hexachlorobenzene (HCB) and PCDD/F isomer levels in various organs in autopsy
10 tissue from North American patients. *Chemosphere* 18(1-6):811-818.
11
12 Sielken, R.L., Jr. 1990. A weight-of-evidence approach to quantitative cancer risk
13 assessment: Information analysis. In *Risk Assessment in Chemical Carcinogenesis*. G.
14 Schettler, D. Schmähl, and T. Klenner (eds). Springer-Verlag, New York. Proceedings of
15 the Satellite Symposium on Risk Assessment in Chemical Carcinogenesis, Heidelberg,
16 Germany August 24-25, 1990.
17
18 Sielken, R.L., Jr., and C. Valdez-Flores. 1996. Comprehensive realism's weight-of-
19 evidence based distributional dose-response characterization. Special Issue of the *Human*
20 *and Ecological Risk Assessment* on Theoretical, Toxicological and Biostatistical
21 Foundations for Deriving Probability Distribution Functions for Reference Doses and
22 Benchmark Doses with Application to Carcinogens and Noncarcinogens 2(1):175-193.
23
24 Sielken, R. L., Jr., and C. Valdez-Flores. 1999. Probabilistic risk assessment's use of trees
25 and distributions to reflect uncertainty and variability and to overcome the limitations of
26 default assumptions. Special Issue of *Environmental International on Modeling and*
27 *Simulation* 25:755-772.
28
29 Sielken, R.L., Jr., R.S. Bretzlaff, and D.E. Stevenson. 1995. Challenges to default
30 assumptions stimulate comprehensive realism as a new tier in quantitative cancer risk
31 assessment. *Regulatory Toxicology and Pharmacology* 21:270-280.
32
33 Skakkeback, N.E. 2010. Normal reference ranges for semen quality and their relations to
34 fecundity. *Asian Journal of Andrology* 12(1):95-98.
35
36 Small, M.J. 2008. Methods for assessing uncertainty in fundamental assumptions and
37 associated models for cancer risk assessment. *Risk Analysis* 28(5):1289-1307.
38
39 Sohn, M.D., M.J. Small, and M. Pantazidou. 2000. Reducing uncertainty in groundwater
40 site characterization using Bayes Monte Carlo methods. *Journal of Environmental*
41 *Engineering* 126(10): 893-902.
42
43 Soto, A.M. and C. Sonnenschein. 2010. Environmental causes of cancer: endocrine
44 disrupters as carcinogens. *Nature Reviews Endocrinology* 6:364-371.
45

- 1 Stahl, B.U., M. Viluksela, J. Deliberto, L.S. Birnbaum, and K.K. Rozman. 1994.
- 2 Subchronic (13 week) toxicity of a mixture of four chlorinated dibenzo-p-dioxins in
- 3 Sprague-Dawley rats. In *Dioxin '94* (H. Fiedler, O. Hutzinger, L. Birnbaum, G. Lambert,
- 4 L. Needham and S. Safe, (eds.) 21:341-345.
- 5
- 6 Starr, T.B., G. Matanoski, M.W. Anders, and M.E. Andersen. 2006. Workshop overview:
- 7 Reassessment of the cancer risk of dichloromethane in humans. *Toxicological Sciences*
- 8 91(1): 20-28.
- 9
- 10 Steenland, K., G. Calvert, N. Ketchum, and J. Michalek. 2001. Dioxin and diabetes
- 11 mellitus: an analysis of the combined NIOSH and Ranch Hand data. *Occupational and*
- 12 *Environmental Medicine* 58:641-648.
- 13
- 14 Sunstein, C.R., and R. Zeckhauser. 2010. Dreadful possibilities, neglected probabilities.
- 15 *The Irrational Economist: Making Decisions in a Dangerous World*, E. Michel-Kerjan
- 16 and P. Slovic (eds.), Public Affairs Books, NY: New York. [Available online at:
- 17 <http://www.hks.harvard.edu/fs/rzeckhau/Sunstein4-6-09.pdf>]
- 18
- 19 Taylor, A.C., J. Evans, and T. McKone. 1993. The value of animal test information in
- 20 environmental control decisions. *Risk Analysis* 13:403-412.
- 21
- 22 Trejo, R., and V. Kreinovich. 2001. Error estimations for indirect measurements:
- 23 randomized vs. deterministic algorithms for 'black-box' programs. In *Handbook of*
- 24 *Randomized Computing*, S. Rajasekaran, P. Pardalos, J. Reif, and J. Rolim, Kluwer.
- 25 (eds), 673-729, Springer, New York, NY
- 26
- 27 Vanden Heuvel, J.P., G.C. Clark, M.C. Kohn, A.M. Tritscher, W.F. Greenlee, G.W.
- 28 Lucier, and D.A. Bell. 1994. Dioxin-responsive genes: examination of dose-response
- 29 relationships using quantitative reverse transcriptase-polymerase chain reaction. *Cancer*
- 30 *Research* 54(1):62-68.
- 31
- 32 Viallefont, V., A.E. Raftery, and S. Richardson. 2001. Variable selection and Bayesian
- 33 model averaging in case-control studies. *Statistics in Medicine* 20:3215-3230.
- 34
- 35 Viluksela, M., B.U. Stahl, and K.K. Rozman. 1994. Subchronic (13-week) toxicity of
- 36 heptachlorodibenzo-p-dioxin in male Sprague-Dawley rats. *Chemosphere* 29:2381-2393.
- 37
- 38 Viluksela, M., B.U. Stahl, L.S. Birnbaum, K.W. Schramm, A. Kettrup, and K.K.
- 39 Rozman. 1997a. Subchronic/chronic toxicity of heptachlorodibenzo-p-dioxin (HpCDD)
- 40 in rats. Part 1. Design, general observations, hematology and liver concentrations.
- 41 *Toxicology and Applied Pharmacology* 46:207-216.
- 42
- 43 Viluksela, M., B.U. Stahl, L. Birnbaum, and K.K. Rozman. 1997b. Subchronic/chronic
- 44 toxicity of heptachlorodibenzo-p-dioxin (HpCDD) in rats. Part 2. Biochemical effects.
- 45 *Toxicology and Applied Pharmacology* 46:217-226.

- 1
2 Viluksela, M., B.U. Stahl, L. Birnbaum, K.W. Schramm, A. Kettrup, and K.K. Rozman.
3 1998a. Subchronic/chronic toxicity of a mixture of four chlorinated dibenzo-p-dioxins
4 (CDDs) in rats. Part I. Design, general observations, hematology and liver concentrations.
5 *Toxicology and Applied Pharmacology* 151:57-69.
6
7 Viluksela, M., BU.. Stahl, L. Birnbaum, and K.K. Rozman. 1998b. Subchronic/chronic
8 toxicity of a mixture of four chlorinated dibenzo-p-dioxin (CDDs) in rats. Part II.
9 Biochemical effects. *Toxicology Applied Pharmacology* 151:70-78.
10
11 Walker, N.J., C.J. Portier, and S.F. Lax 1999. Characterization of the dose-response of
12 CYP1B1, CYP1A1, and CYP1A2 in the liver of female Sprague-Dawley rats following
13 chronic exposure to 2,3,7,8-tetrachlorodibenzo-*p*-dioxin. *Toxicology and Applied*
14 *Pharmacology* 154:279-286.
15
16 Walley, P. 1991. *Statistical Reasoning with Imprecise Probabilities*. Chapman and
17 Hall/CRC Press.
18
19 Wang, X, M.J. Santostefano, M.V. Evans, V.M. Richardson, J.J. Diliberto, and L.S.
20 Birnbaum. 1997. Determination of parameters responsible for pharmacokinetic behavior
21 of TCDD in female Sprague-Dawley rats. *Toxicology and Applied Pharmacology*
22 147:151-168.
23
24 Wheeler, M.W., and A.J. Bailer. 2007. Properties of model-averaged BMDLs: a study of
25 model averaging in dichotomous risk estimation. *Risk Analysis* 27:659-670.
26
27 Wheeler, M.W., and A.J. Bailer. 2009. Comparing model averaging with other model
28 selection strategies for benchmark dose estimation. *Environmental and Ecological*
29 *Statistics* 16(1):37-51.
30
31 White, R.H., I. Cote, L. Zeise, M. Fox, F. Dominici, T.A. Burke, P.D. White, D. Hattis,
32 and J.M. Samet. 2009. State-of-the-science workshop report: issues and approaches in
33 low dose-response extrapolation for environmental health risk assessment. *Environmental*
34 *Health Perspectives* 117:283-287.
35
36 Winkler, R.L., and A.H. Murphy. 1985. Decision analysis. In *Probability, Statistics, and*
37 *Decision Making in the Atmospheric Sciences*, A.H. Murphy and R.W. Katz (eds.), 493-
38 524, Westview Press, Boulder, CO,
39
40 Yokomizo, H., H.P. Possingham, M.B. Thomas, and Y.M. Buckley. 2009. Managing the
41 impact of invasive species: the value of knowing the density-impact curve. *Ecological*
42 *Applications* 19(2):376-386.
43
44 Yr, J. 2008. Neuro-developmental deficits in early-treated congenital hypothyroidism.
45 *Annals of the Academy of Medicine, Singapore* 37,Supp. 12:42-43.

- 1
2 Yoshizawa, K., A.E. Brix, D.M. Sells, M.P. Jokinen, M. Wyde, D.P. Orzech, G.E.
3 Kissling, N.J. Walker, and A. Nyska. 2009. Reproductive lesions in female Harlan
4 Sprague-Dawley rats following two-year oral treatment with dioxin and dioxin-like
5 compounds. *Toxicologic Pathology* 37(7):921-937.
6
7 Yoshizawa, K., N.J. Walker, A. Nyska, G.E. Kissling, M.P. Jokinen, A.E. Brix, D.M.
8 Sells, and M.E. Wyde. 2010. Thyroid follicular lesions induced by oral treatment for 2
9 years with 2,3,7,8-tetrachlorodibenzo-p-dioxin and dioxin-like compounds in female
10 Harlan Sprague-Dawley rats. *Toxicologic Pathology* 38(7):1037-1050.
11
12 Zeise, L., D. Hattis, , M. Andersen, A.J. Bailer, S. Bayard, C. Chen, H. Clewell, R.
13 Conolly, K. Crump, D. Dunson, A. Finkel, L. Haber, A.M. Jarabek, R. Kodell, D.
14 Krewski, D. Thomas, T. Thorslund, and J. Wassell. 2002. Improving risk assessment:
15 Research opportunities in dose response modeling to improve risk assessment. *Human*
16 *and Ecological Risk Assessment* 8(6):1421–1444.
17
18 Zhang Q., S. Bhattacharya, M.E. Andersen, and R.B. Conolly. 2010. Computational
19 systems biology and dose-response modeling in relation to new directions in toxicity
20 testing. *Journal of Toxicology and Environmental Health B* 13:253–276.

1 **Appendix A: Dissenting Opinion from Karl Rozman, Ph.D.**
2

The University of Kansas Medical Center

Karl K. Rozman, Ph.D.
Professor
School of Medicine
Department of Pharmacology,
Toxicology and Therapeutics

December 9, 2010

Thomas Armitage, Ph.D.
Designated Federal Officer
USEPA Science Advisory Board (1400R)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

RE: A Dissenting Opinion

Dear Tom,

As I have indicated in my previous written and oral opinions to this panel, I disagree with the panel conclusions regarding the carcinogenicity of TCDD and the adequacy of the EPA response to the criticisms of the NAS report.

There is at best equivocal evidence (statistically not significant) for the carcinogenicity of TCDD (or DLCs) in the occupational setting where the body burdens were at least 100 or 1000 times higher than the current or previous background levels. Therefore, the consideration of a practical threshold for any defined population requires acceptance of the compelling scientific conclusion that there is negligible (essentially zero) carcinogenic risk at current background levels which are much lower than past levels. Any other conclusion is incompatible with sound science and no amount of modeling or data manipulation will transform a non-existing effect at occupational exposure levels into a risk at current background levels other than the non-scientific, policy-driven, non-threshold extrapolation by EPA.

Further, it is my opinion that the EPA document (2010 Reanalysis of Key Issues Related to Dioxin Toxicity and Response to NAS Comments, 600/-10/038A) is deliberately non-responsive to the recommendations of the NAS report.

Respectfully,



Karl K. Rozman, Ph.D., D.A.B.T.
Professor

3901 Rainbow Blvd., MS1018, Kansas City, KS 66160 • (913) 588-7717 • FAX (913) 588-7501
krozman@kumc.edu

Appendix B: Value of Information


When human health risk assessments include an explicit representation of uncertainty, the potential value of new information (VOI) can be estimated by its ability to reduce uncertainties that matter most to the assessment target. While methods for determining VOI are most commonly associated with the decision analysis literature in the context of informing management or regulatory decisions (Raiffa, 1968; Keeney, 1982; Winkler and Murphy, 1985; Finkel and Evans, 1987; Taylor et al., 1993; Clemen, 1996; Chao and Hobbs, 1997), there are many steps in a scientific assessment well before (or even without subsequent) decision support and decision making where VOI evaluations can be of benefit in characterizing current scientific knowledge and the potential for its improvement. EPA should integrate these methods into their current and ongoing assessments of dioxin toxicity.

When uncertainty in a scientific assessment is measured by the variance of model predictions, a first measure of VOI is the extent to which this variance might be reduced by new or additional data (e.g., Patwardhan and Small, 1992; Brand and Small, 1995; Abbaspour et al., 1996; Chao and Hobbs, 1997; Sohn et al., 2000; Bosgra et al., 2005; Cooke, 2009). The relative contribution of different model assumptions and parameter uncertainties to the variance of the estimated effect (e.g., the BMD), or the cancer slope factor) provides an indication of which of these uncertainties would be most beneficial to address. In addition, a VOI assessment considers the potential for the component uncertainties to be reduced, based on the feasibility, resource requirements (time and funding), and likelihood of success of the studies that would be needed to achieve the necessary improvement in scientific knowledge.

A scientific VOI study may also target a key classification inference that results from a risk assessment, for example, whether a compound is genotoxic. Assuming the current assessment leads one to assign an inconclusive probability to this outcome (e.g., between 10% and 90%, so that neither inference can be rejected with a high degree of confidence), then potentially valuable studies are those able to shift subsequent probabilities to high values (e.g., above 90, 95, or 99%) with a positive result (e.g., providing support for genotoxicity) and/or to low values (below 10, 5, or 1%) with a negative result.

To illustrate, Small (2008) presents a simple probability tree model (a “distributional approach”) for assessing genotoxicity based on studies of DNA damage response caused by naphthalene and its metabolites. In the proposed studies a series of isogenic cell lines deficient in various DNA metabolism pathways are used to characterize the DNA damage responses caused by the targeted compounds. Following results from the cultured cells, mice deficient in the specific DNA damage responses would be exposed to naphthalene. Possible inferences are identified based on the assessed sensitivity and selectivity of study results to the genotoxicity of naphthalene. Study outcomes considered include: i) DNA damage responses in the isogenic cells; ii) increased numbers of stable DNA adducts in the DNA repair deficient mouse lung; and

iii) heightened Clara cell toxicity in the DNA repair deficient mouse lung. Illustrative results using Netica are presented as follows:



As noted, the results shown above are intended solely to demonstrate the way in which study results can be combined to support or refute targeted inferences.

Even when the uncertainty tree method is only used to delineate the set of possible outcomes and relationships among steps and assumptions in the risk assessment (i.e., mode of action; dosimetry measures for exposure; the mathematical form of the dose-response relationship; the experimental data set(s) used to fit the relationship; and the procedure used for interspecies extrapolation) *without the assignment of probabilities to the tree branches*, key assumptions and the experiments needed to support or refute them can still be identified. These will typically involve elements of the assessment that, depending on their resolution, effectively restrict the set of possible outcomes to either a positive or a negative inference regarding the endpoint of the risk assessment. Establishing a procedure of this type will allow the Agency to put in place a more formal mechanism for identifying, conducting, and integrating the results of key studies for future assessments.

Appendix C: Editorial Comments and Corrections

a. Section 2

Minor suggestions to further improve clarity regarding data set inclusion/exclusion criteria

Page xxxvii, Lines 16-19. The sentence needs clarification. It currently gives the impression that studies that were eliminated for further analysis would have NOAELs available.

Pages 2-234 – 2-247. EPA should consider adding information to the appendices and/or tables to provide readers with clarification regarding the exclusion of particular studies. For example, an extra column in Table 2-7 listing, by numbered reference, the criteria that were or were not met for each study would be helpful.

b. Section 6

Page 6-2. Add NRC (1996).

Page 6-3, bottom: The word “margins” should be “marginals.”

Page 6-3, line 26: If EPA wants to use the adverb “always”, the phrase “as a joint distribution” should be “as some characterization of a joint distribution” to be correct.

Page 6-4, lines 9-12: This text is strange and off-putting. A reader might ask who wrote this and why. It seems opinionated and unnecessary.

Page 6-4, line 9: The tone is too pedagogical (“This is not the place . . .”).

Footnote 54: The discussion of alternatives to strict, single-measure probability theory is ham-handed. Neither interval probabilities nor imprecise probabilities (*sensu* Walley, 1991) depart from probability theory; they follow the Kolmogorov axioms. They are motivationally and essentially equivalent to sensitivity analyses, except they do not make use of sampling strategies and can be more comprehensive.

Lines 29-30: It is simply untrue that sensitivity analyses have to be systematic. The word “systematic” might better be “comprehensive” and the word “essential” should be weakened, perhaps to “advantageous”.

Page 6-5, lines 4-7 and footnote 55: There seem to be only two axioms mentioned in the text, but Kolmogorov needs three to make probability theory.

Page 6-5: The meaning of the phrase “epistemic uncertainty” given on this page is plainly incorrect. Epistemic uncertainty is the uncertainty that arises from imperfect

1 knowledge such as from limitations on the amount or quality of data available or
2 deficiencies in our causal understanding about a system. It is not true that a quantity
3 about which there is epistemic uncertainty is necessarily fixed. Although it is perhaps
4 clear how one might come to this mistaken impression, no researchers use the phrase to
5 imply that the underlying quantity has no variability (although all would admit that this
6 could be the case given our ignorance about it). Indeed, a variable can have both forms
7 of uncertainty. For example, when body weight varies across a population, but with a
8 distribution that is unknown, the variable has both aleatory and epistemic uncertainty.
9 This mistake echoes in a couple of other places throughout this section.

10
11 It is not clear what the authors take to be the difference between epistemic uncertainty
12 and what they call “cognitive uncertainty.” It seems that the latter phrase was introduced
13 because the meaning of “epistemic uncertainty” had been misunderstood. Normally, the
14 phrase “cognitive uncertainty” would refer to an individual person’s uncertainty about the
15 validity of the results of his or her own information processing. The assertion that
16 cognitive uncertainty may be represented by probability (i.e., by precise probability
17 measures) is unnecessary and may be misleading. In fact, researchers in human cognition
18 and neuroscience have shown that humans process this kind of uncertainty (which they
19 often call “ambiguity”) separately and differently from what we think of as probability or
20 frequentist risk (Hsu et al., 2005; Glimcher, 2003). The section can omit the phrase
21 “cognitive uncertainty” altogether and use in its place “epistemic uncertainty.” There are
22 slight differences between the two ideas (e.g., epistemic uncertainty could be shared by
23 members in a group, whereas cognitive uncertainty is always personal), but these appear
24 to be unimportant in this context.

25
26 Page 6-5: The words “aleatoric” and “aleatory” are both used on this page as
27 (synonymous) adjectives of uncertainty. Actually, in the engineering literature, only
28 “aleatory” is preferred for this use. In any case, please pick one to use.

29
30 Page 6-5, line 10: The assertion that the frequentist and Bayesian interpretations are not
31 mutually exclusive may be misleading. They are mutually exclusive in the sense that it
32 would be improper to mix and match components of each into an analysis. It would be
33 appropriate to omit the clause with the phrase “mutually exclusive,” although it is surely
34 fair to say that subjective probabilities can and do track relative frequencies.

35
36 Page 6-5, lines 30-32: The text on the subject of dependence is strange here, and also in
37 section 6.1.3.3. It is incorrect that the “[i]ssues involving...epistemic and aleatory
38 uncertainty translate into issues of dependence.” This is just wrong (even under their
39 unusual definition of “epistemic”).

40
41 Page 6-6: Section 6.1.3.2 starting on this page discusses a way to address uncertainty for
42 sample data. This Spartan treatment does not mention that sampling uncertainty is not
43 the only kind of uncertainty that can be associated with data, nor that it may not even be
44 the largest kind of uncertainty. Mensurational uncertainty (including the plus-minus part
45 of a measurement, and censoring) may be more important. In some cases, the family or

1 shape of the marginal distribution may be unknown, which is a kind of model
2 uncertainty. As suggested on page 6-35, such uncertainties can be significant. The
3 section suggests only a resampling approach to expressing the uncertainty, but fails to
4 mention the often severe limitations of such approaches, and says nothing about what one
5 might do if there is no relevant sample data.

6
7 Page 6-6, line 20: Maybe the last word of the header should be plural.

8
9 Line 21: Modern practice has replaced “error” with “uncertainty” in this context.

10
11 Footnote 56: EPA could add “or subtracting” after “adding.”

12
13 Page 6-7, line 14: “The role of dependence modeling” should be replaced with
14 “Dependence among variables.”

15
16 Page 6-7. More examples of use of expert judgment for health assessment are available
17 and should be cited.

18
19 Page 6-7, last paragraph: This paragraph extending onto the next page should be
20 rewritten. The example is reasonable and important, but the discussion about it is
21 confused. The first sentence is incorrect. The uncertainty mentioned in the second
22 sentence may be epistemic, but the sentence is erroneous in its claim. In the following
23 sentences, the words “variable” and “fixed” (or “constant”) should be used rather than
24 “aleatoric” and “epistemic.” It is nonsense to say that a kinetic constant is “completely
25 correlated across individuals.” It’s not correlated; it is invariant. This case is not an
26 example of a dependence issue. There is no correlation between a distribution and a
27 fixed quantity (even if it’s uncertain). Correlation is defined between *varying* quantities.
28 If the number is fixed, whether or not we know what it is, then one cannot say it is
29 correlated with anything. The authors may have come to this twisted language because
30 they’re thinking of the uncertainties in terms of how they might plan to quantitatively
31 characterize them in a Monte Carlo simulation (repeatedly selecting a random deviate for
32 the kinetic constant but assigning it to every individual). Of course, variables such as
33 body fat, age, and smoking, on the other hand, can and do exhibit correlations that
34 definitely should be accounted for in the quantitative assessments. Likewise, it is also
35 important to keep track of the constancy of particular quantities about which we may not
36 know the precise value. These two issues should be untangled and discussed in a less
37 confusing way.

38
39 Page 6-8, line 12: The first paragraph of section 6.1.3.4 seems to be saying that one can
40 sometimes express model uncertainty as parametric uncertainty, which simplifies its
41 handling. This could be said more plainly. It would be helpful to mention that this trick
42 cannot always be used (as when the possible models cannot be listed). It might also be
43 especially helpful to mention that this trick is not so much a way to propagate model
44 uncertainty as a way to sweep it under the rug. Model averaging, including Bayesian
45 model averaging (which is mentioned in several places, including 6-36, lines 3ff), erases

1 model uncertainty in the same way that averaging variable quantities erases their
2 variation.

3
4 Page 6-8, line 13: Omit the unnecessary fancy after the semicolon.

5
6 Lines 15-17: This sentence is nonsense, if we understand what a linear low-dose model
7 is. Parsing the sentence, it seems to say “uncertainty over a...slope...may be quantified,
8 but uncertainty...in slope...cannot be captured” which is self-contradictory. Perhaps
9 what the authors mean to say is that the linearity assumption is not itself subject to
10 uncertainty quantification.

11
12 Page 6-9, line 1: The mathematical symbol x should be italicized, as should all Roman
13 letters throughout the document that represent unknown quantities, i.e., are symbols
14 representing something else rather than names like “e” the base of the natural logarithms.

15
16 Lines 14 and 16: The prefixes “pseudo” and “quasi” are not words. Hyphens are needed.

17
18 Page 6-9, line 18: Provide citations for dependence modeling.

19
20 Page 6-9: Section 6.1.3.6 might also mention *graphs*, and other traditional
21 communication tools other than correlation indices.

22
23 Page 6-10, line 4: Add mention of methods that identify uncertain assumptions or
24 parameters that are *important* for determining whether the model is consistent with
25 observed data (Hornberger and Spear, 1983) and for affecting a decision that is made as a
26 result of the model (Merz et al., 2009).

27
28 Page 6-10, lines 29-30: Do the authors mean “*this* probabilistic language,” referring to
29 the word “likely” in the quoted text?

30
31 Page 6-11, line 19: Of course there is no guarantee that linear will be protective.

32
33 Page 6-13, line 18: Of course it isn’t really apodictic knowledge at all, but rather only an
34 opinion or an assumption. We see the authors’ point and agree with it entirely, but
35 perhaps they should use a word other than “apodictic” here since it’s not technically
36 correct.

37
38 Page 6-14, lines 33-34: The parenthetical phrase “volitional uncertainty” should be
39 expanded into a sentence that says what the authors mean to express. The phrase
40 “cognitive uncertainty” does not mean anything in this context. Perhaps if the authors
41 expanded it into a sentence too, maybe making it “epistemic uncertainty” along the way,
42 it would be possible to understand what they are trying to say here.

43
44 Footnote 62: “Effective” is misspelled, as is “cancer.”
45

1 Page 6-16, line 5: We note that it's not really a guarantee of course.

2
3 Line 8: The word "common" should be "predominant".

4
5 Page 6-16, line 20: Perhaps we can say that variability (and uncertainty) in the factors
6 that are used to determine a particular UF can be considered in choosing the particular
7 value of the UF.

8
9 Page 6-17, lines 3-14: This problem can be addressed using a Bayesian analysis with a
10 beta conjugate for the uncertain response probability, p , with uninformative (uniform)
11 prior for p . The probability that "an experiment with a null response might have yielded
12 a positive response" can be estimated from the predictive distribution (which will depend
13 on the number of test animals in the original study that yielded zero responses) for the
14 next experiment (with any number of exposed animals).

15
16 Page 6-17, line 28: The word "band" should be "limit".

17
18 Page 6-20, footnote 66: The text starting "each have an error factor" should be followed
19 by "of" rather than "or".

20
21 Page 6-21, line 6: It would be helpful to say something about what the concerns are.

22
23 Page 6-21, lines 12-14: NAS was not suggesting that EPA use the *uncertainty factors*
24 approach to mount an uncertainty analysis, but rather a more modern approach.

25
26 Page 6-22, line 19: Would it be appropriate to note "and establishes a concomitant
27 reduction in some UFs?"

28
29 Line 29: The word "invokes" should perhaps be "would require".

30
31 Page 6-23, line 33 and passim: The word "exotic" is a poor choice that is unnecessarily
32 and transparently loaded.

33
34 Page 6-25, line 29: This sentence is ungrammatical.

35
36 Page 6-26, line 24 and Figure 6-1: Would it be helpful to draw the 45-degree line on the
37 graph?

38
39 Page 6-27, line 10: The word "epistemic" here is acceptable.

40
41 Line 14: The word "epistemic" here should be replaced by "fixed across individuals,"
42 and "is estimated from" should be replaced by "varies with." How does half life's
43 estimability from data imply that it is variable?

44
45 Page 6-28, lines 1-2: One would need the dependence between the variables to proceed.

Line 9: We suggest that “and” should be “although.”

Page 6-29, line 1-2: There are bounding techniques based on the classical Fréchet inequality that do not require any knowledge of or any assumptions about dependencies.

Line 32: Omit “to.”

Page 6-31, line 24: The pessimistic conclusion is a bit strong. Any *estimate* made from data is amenable to a quantitative uncertainty analysis so, if one is measuring anything, one can propagate uncertainties such as mensurational uncertainty, sampling uncertainty, and perhaps even surrogacy uncertainty. It’s not quite as hard to get quantitative models as the text here seems to suggest.

Page 6-32, lines 13-14: The dour conclusion is confusing. One could do a sensitivity analysis in this case, couldn’t one? If so, it seems that some kind of uncertainty analysis is clearly possible.

Page 6-33: The example in the text box is great, but the second table seems to say the log-likelihood for LLD is 2.46 and for Hill is 2.16, which would make LLD’s larger than Hill’s, which contradicts what’s said in the text.

Page 6-34, line 4: Shouldn’t “*Delivered dose*” be a new bullet?

Line 8: We don’t think this statement is true. Perhaps “statistically more powerful” should be “typically yield more sensitive”.

Lines 24-25: We don’t think it is necessary or helpful to persist with Box’s platitude. Model uncertainty is the uncertainty about a model’s predictions that arises from doubt about the relevance of that model for making such predictions.

Page 6-37, line 29: The caveat is overwrought. Exploring relevant alternative values in a sensitivity analysis could constitute a quantitative uncertainty analysis, even if the exploration is limited.

Page 6-37, line 30: This sentence is false. Analytical methods of propagation (convolution) don’t “sample” anything, and analyses based on intervals or imprecise probabilities don’t depend on uncertainty “distributions” (i.e., precise probability distributions).

It is important to keep in mind that, in general, we are not necessarily limited to identifying precise probability distributions for everything that is to be characterized as uncertain (as seems to be suggested here). Simple intervals about uncertain quantities can support a straightforward, albeit crude, interval analysis that propagates uncertainty about parameters and other model choices to statements about the range of possible

1 results. Similarly, an approach based on interval probabilities, probability boxes, or
2 general imprecise probabilities (Walley, 1991) can combine such intervals with precise
3 distributions if they are known for some other inputs, and with structures that are
4 intermediate between coarse intervals and delicate probability distributions when some
5 but incomplete knowledge is available. If the inputs are profoundly uncertain, the results
6 from such analyses are likely to be wide in reflection of these uncertainties. In almost all
7 cases, it is possible to be entirely rigorous without necessarily being precise and without
8 completely specifying each probability distribution.

9
10 Page 6-37, line 31: There does not need to be a specified “underlying distribution from
11 which to sample” in order to conduct a quantitative uncertainty analysis. It is facile to
12 shrug off a call to characterize and account for important uncertainties in the assessment
13 process on these grounds alone. Even when the uncertainty is volitional, there can be
14 relevant ranges that are interesting to decision makers and stakeholders. In such cases,
15 the analysis may be formally closer to a sensitivity analysis, but some appropriate
16 response is usually possible, if not always practicable. To their credit, EPA has
17 acknowledged the legitimacy of the call and undertaken some efforts in this direction,
18 notably Tables 5-18 and 5-19 (although some kind of graphical summary of the results
19 might have been better).

20
21 Page 6-38, line 30 and passim: The adjective “data driven” needs a hyphen, as it has
22 elsewhere in the document.

23
24 Line 23-24: We think this sentence is true, but, again, sampling from a distribution is not
25 the only way to conduct a quantitative uncertainty analysis.

26
27 Line 26: What is “(2.a)”?

28
29 Page 6-41, line 23: Omitting the word “extra” would make the sentence more easily
30 understandable.

31
32 Line 31: What does “How Forward?” mean? Is this idiomatic?

33
34 Section 6.5.2: The assertions in this section are rather surprising and questionable. EPA
35 says that uncertainty quantification is an “emerging area in science” and that it is “an area
36 where research could be focused” because “the requisite knowledge does not yet exist” to
37 apply quantitative uncertainty analysis in assessments such as this one for dioxin. The
38 document peremptorily dismisses the utility of “convening a blue-ribbon panel” to
39 identify the proper approach and suggests instead that “multiple approaches should be
40 encouraged.” Is the inference that the present review panel shouldn’t try to say what the
41 proper approaches to uncertainty quantification are, even if we think the area is more
42 mature than emerging? It is hard to understand what these statements are suggesting.
43 Will the Agency support intramural and extramural research efforts in this direction? If
44 not, what do the statements mean? Is it impossible that EPA could benefit from some
45 tech transfer efforts as well as basic research on uncertainty quantification? The

- 1 paragraph beginning on page 6-42 (line 3) mentions a European idea of bench-test
- 2 exercises to compare different approaches. It may be worth mentioning that this idea has
- 3 been implemented in the United States as well (Oberkamp et al., 2004; Ferson et al.,
- 4 2004).
- 5
- 6 The document's reference list is alphabetically arranged, but seems to go from Z back to
- 7 A again on page R-33.

Science Advisory Board (SAB) 5/4/11 Draft. Do not cite or quote.
This draft SAB Panel report has been prepared for quality review and approval by the chartered SAB. This draft does not represent EPA policy.

1 Appendix D: EPA's Charge Questions



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
NATIONAL CENTER FOR ENVIRONMENTAL ASSESSMENT
WASHINGTON, DC 20460

May 27, 2010

OFFICE OF
RESEARCH AND DEVELOPMENT

MEMORANDUM

SUBJECT: Request for Science Advisory Board Review of the Draft Report, "EPA's Reanalysis of Key Issues Related to Dioxin Toxicity and Response to NAS Comments"

FROM: Becki Clark, Deputy Director
National Center for Environmental Assessment (8601P)
Office of Research and Development

TO: Vanessa T. Vu, Ph.D., Director
EPA Science Advisory Board (1400F)

This is to request a review by the Science Advisory Board of the draft report entitled "EPA's Reanalysis of Key Issues Related to Dioxin Toxicity and Response to NAS Comments" (EPA/600/R-10/038A). This draft report details the Environmental Protection Agency's (EPA) response to key comments and recommendations included in the 2006 NAS report ("Health Risks from Dioxin and Related Compounds: Evaluation of the EPA Reassessment") on their review of the (EPA) 2003 draft dioxin reassessment. This draft report also includes significant new analyses on both the potential cancer and noncancer human health effects that may result from chronic exposures to dioxins.

Attached is the Charge that provides background information as well as questions that are to be the focus of the Science Advisory Board review of this draft report.

Please let me know if you have any questions. Thank you.

Attachment: Charge for EPA's Science Advisory Board – Review of the Draft Report, "EPA's Reanalysis of Key Issues Related to Dioxin Toxicity and Response to NAS Comments"

cc: Peter W. Preuss
Annette Gatchett
Glenn Rice
Cheryl Itkin

**Proposed Charge to the Science Advisory Board for Peer Review Of Draft Report
“EPA’s Reanalysis of Key Issues Related to Dioxin Toxicity
and Response to NAS Comments”**

May, 2010

EPA has been preparing an assessment of the potential health impacts of 2,3,7,8-Tetrachlorodibenzo-p-Dioxin (TCDD) for many years. In 2003, EPA released an external review draft report entitled, *Exposure and Human Health Reassessment of 2,3,7,8-Tetrachlorodibenzo-p-Dioxin (TCDD) and Related Compounds* (U.S. EPA, 2003) (herein referred to as “2003 Reassessment”) that was reviewed by the EPA Science Advisory Board (SAB), and then by the National Academy of Sciences (NAS). In 2006, the National Research Council (NRC) of the National Academies published their report of EPA’s reassessment, *Health Risks from Dioxin and Related Compounds: Evaluation of the EPA Reassessment* (NRC, 2006).

The current Report *EPA’s Reanalysis of Key Issues Related to Dioxin Toxicity and Response to NAS Comments* (“*Response to Comments*”) before the SAB is a response to the review by the NRC, and includes new analyses completed in response to the NRC recommendations and recently published literature, as well as a discussion of topics where our views differed. The draft *Response to Comments* document is not an assessment per se; it is designed to supplement the information provided in the 2003 *Reassessment*. However, the draft *Response to Comments* provides a noncancer reference dose and updated cancer values. Detailed discussions of many of the issues addressed in the draft *Response to Comments* are available in the 2003 *Reassessment* and have not been reproduced in the current Report – whenever appropriate; the reader is directed to the pertinent chapters of the 2003 *Reassessment*.

The NRC identified three key recommendations that they believed would result in substantial improvement to the EPA 2003 *Reassessment* and thus support a scientifically robust characterization of human responses to exposures to TCDD. These three key areas are (1) improved transparency and clarity in the selection of key data sets for dose-response analysis, (2) further justification of approaches to dose-response modeling for cancer and noncancer endpoints, and (3) improved transparency, thoroughness, and clarity in quantitative uncertainty analysis. The NRC Report also encouraged EPA to calculate a reference dose (RfD), which had not been derived in the 2003 *Reassessment*. The draft *Response to Comments* document addresses each of these issues. Please consider the accuracy, objectivity, and transparency of EPA’s reanalysis and responses in your review.

General Charge Questions

- 1.1 Is the draft *Response to Comments* clear and logical? Has EPA objectively and clearly presented the three key NRC recommendations?

- 1.2 Are there other critical studies that would make a significant impact on the conclusions of the hazard characterization or dose-response assessment of the chronic noncancer and cancer health effects of TCDD?

Specific Charge Questions

Section 2. Transparency and Clarity in the Selection of Key Data Sets for Dose-Response Analysis

- 2.1. Is this Section responsive to the NAS concern about transparency and clarity in data-set selection for dose-response analysis?
- 2.2. Are the epidemiology and animal bioassay study criteria/considerations scientifically justified and clearly described?
- 2.3. Has EPA applied the epidemiology and animal bioassay study criteria/considerations in a scientifically sound manner? If not, please identify and provide a rationale for alternative approaches.

Section 3. The Use of Toxicokinetics in Dose-Response Modeling for Cancer and Noncancer Endpoints

- 3.1 The *2003 Reassessment* utilized first-order body burden as the dose metric. In the draft *Response to Comments* document, EPA used a physiologically-based pharmacokinetic (PBPK) model (Emond et al., 2004, 2005, 2006) with whole blood concentration as the dose metric rather than first-order body burden. This PBPK model was chosen, in part, because it includes a biological description of the dose-dependent elimination rate of TCDD. EPA made specific modifications to the published model based on more recent data. Although lipid-adjusted serum concentrations (LASC) for TCDD are commonly used as a dose metric in the literature, EPA chose whole blood TCDD concentrations as the relevant dose metric because serum and serum lipid are not true compartments in the Emond PBPK models (LASC is a side calculation proportional to blood concentration).

Please comment on:

- 3.1.a. The justification of applying a PBPK model with whole blood TCDD concentration as a surrogate for tissue TCDD exposure in lieu of using first-order body burden for the dose-response assessment of TCDD.
- 3.1.b. The scientific justification for using the Emond et al. model as opposed to other available TCDD kinetic models.
- 3.1.c. The modifications implemented by EPA to the published Emond et al. model.

- 1
2 3.1.d. Whether EPA adequately characterized the uncertainty in the kinetic
3 models.
4
5 3.2. Several of the critical studies for both noncancer and cancer dose-response
6 assessment were conducted in mice. A mouse PBPK model was developed from an
7 existing rat model in order to estimate TCDD concentrations in mouse tissues,
8 including whole blood.
9
10 Please comment on:
11
12 3.2.a. The scientific rationale for the development of EPA's mouse model based
13 on the published rat model (Emond et al., 2004, 2005, 2006).
14
15 3.2.b. The performance of the mouse model in reference to the available data.
16
17 3.2.c. Whether EPA adequately characterized the uncertainty in the mouse and
18 rat kinetic models. Please comment specifically on the scientific
19 justification of the kinetic extrapolation factor from rodents to humans.
20
21 3.3 Please comment on the use of the Emond et al. PBPK model to estimate human
22 intakes based on internal exposure measures.
23
24 3.4. Please comment on the sensitivity analysis of the kinetic modeling (see Section
25 3.3.5).
26
27 3.5. Both EPA's noncancer and cancer dose-response assessments are based on a
28 lifetime average daily dose. Did EPA appropriately estimate lifetime average daily
29 dose? If not, please suggest alternative approaches that could be readily developed
30 based on existing data.
31

32 **Section 4. Reference Dose**

- 33
34 4.1. The Mocarelli et al. (2008) and Baccarelli et al. (2008) studies were selected as co-
35 critical studies for the derivation of the RfD. Is the rationale for this selection
36 scientifically justified and clearly described? Please identify and provide the
37 rationale for any other studies that should be selected, including the rationale for
38 why the study would be considered a superior candidate for the derivation of the
39 RfD. In addition, male reproductive effects and changes in neonatal thyroid
40 hormone levels, respectively, were selected as the co-critical effects for the RfD.
41 Please comment on whether the selection of these critical effects is scientifically
42 justified and clearly described. Please identify and provide the rationale for any
43 other endpoints that should be selected as the critical effect.
44
45 4.2. In the Seveso cohort, the pattern of exposure to TCDD is different from the average

1 daily exposure experienced by the general population. The explosion in Seveso
2 created a high dose pulse of TCDD followed by low level background dietary
3 exposure in the exposed population. In the population, this high dose pulse of
4 TCDD was slowly eliminated from body tissues over time. There is uncertainty
5 regarding the influence of the high-dose pulse exposure on the effects observed
6 later in life.

7
8 4.2.a. Mocarelli et al. (2008), reported male reproductive effects observed later
9 in life for boys exposed to the high dose pulse of TCDD between the ages
10 of 1 and 10. EPA identified a 10 year critical exposure window. In the
11 development of the candidate RfD, EPA used an exposure averaging
12 approach that differs from the typical approach utilized for animal
13 bioassays. EPA determined that the relevant exposure should be
14 calculated as the mean of the pulse exposure and the 10-year critical
15 exposure window average. Please comment on the following:

16
17 4.2.a.i. EPA's approach for identifying the exposure window and
18 calculating average exposure for this study.

19
20 4.2.a.ii. EPA's designation of a 20% decrease in sperm count (and an
21 11% decrease in sperm motility) as a LOAEL for Mocarelli et al.
22 (2008).

23
24 4.2.b. For Baccarelli et al. (2008), the critical exposure window occurs long
25 after the high-dose pulse exposure. Therefore, the variability in the
26 exposure over the critical exposure window is likely to be less than the
27 variability in the Mocarelli et al. subjects. EPA concluded that the
28 reported maternal exposures from the regression model developed by
29 Baccarelli et al. provide an appropriate estimate of the relevant effective
30 dose as opposed to extrapolating from the measured infant TCDD
31 concentrations to maternal exposure. Additionally, EPA selected a
32 LOAEL of 5 μ -units TSH per ml blood in neonates; as this was
33 established by World Health Organization (WHO) as a level above which
34 there was concern about abnormal thyroid development later in life.
35 Please comment on the following:

36
37 4.2.b.i. EPA's decision to use the reported maternal levels and the
38 appropriateness of this exposure estimate for the Baccarelli et al.
39 study.

40
41
42 4.2.b.ii. EPA's designation of 5 μ -units TSH per ml blood as a LOAEL
43 for Baccarelli et al. (2008).

44
45 4.3. Please comment on the rationale for the selection of the uncertainty factors (UFs)

for the RfD. If changes to the selected UFs are proposed, please identify and provide a rationale.

4.4. EPA did not consider biochemical endpoints (such as CYP induction, oxidative stress, etc.) as potential critical effects for derivation of the RfD for TCDD due to the uncertainties in the qualitative determination of adversity associated with such endpoints and quantitative determination of appropriate response levels for these types of endpoints in relation to TCDD exposure. Please comment on whether this decision is scientifically justified and clearly described.

4.5. In using the animal bioassays, EPA averaged internal blood TCDD concentrations over the entire dosing period, including 24 hours following the last exposure. Please comment on EPA's approach for averaging exposures including intermittent and one day gestation exposure protocols.

4.6. Please comment on the benchmark dose (BMD) modeling conducted by EPA to analyze the animal bioassay data and EPA's choice of points of departure (PODs) from these studies.

4.7. For the animal bioassay modeling, EPA applied the kinetic extrapolation at the level of the POD prior to applying the uncertainty factors because EPA has less confidence in the kinetic model output at lower doses reflective of the RfD. Please comment on whether this approach was scientifically justified and clearly described.

4.8. Please comment as to whether EPA's qualitative discussion of uncertainty in the RfD is justified and clearly described.

Section 5. Cancer Assessment

5.1. Weight of Evidence Cancer Descriptor: The *2003 Reassessment* concluded that TCDD is a "known human carcinogen." In the current draft *Response to Comments* document, EPA concluded that under the 2005 Guidelines for Carcinogen Risk Assessment (U.S. EPA, 2005) TCDD is "carcinogenic to humans." Is the weight-of-evidence characterization scientifically justified and clearly described?

5.2. Mode of Action: The mode of action of a carcinogen can inform identification of hazards and approaches used for a dose-response assessment. The mode of carcinogenic action for TCDD has not been elucidated for any tumor type. EPA concluded that, while interaction with the Ah receptor is likely to be a necessary early event in TCDD carcinogenicity in experimental animals, the downstream events involved are unknown.

5.2.a. Are the available data related to mode(s) of action for the carcinogenicity of TCDD appropriately characterized and clearly presented?

- 1
2 5.2.b. Do the available data support EPA's conclusion that the overall mode(s)
3 of action for TCDD-induced carcinogenesis is largely unknown? Please
4 comment on whether this evaluation is clearly described.
5
- 6 5.3. Is EPA's approach for selecting data sets from the key epidemiologic studies and
7 animal bioassays identified for cancer dose response modeling scientifically
8 justified and clearly described?
9
- 10 5.4. For the animal bioassay data, potential cancer oral slope factors (OSFs) were
11 calculated by linear extrapolation (using a linear, nonthreshold cancer approach)
12 from the point of departure (POD). EPA also estimated the composite risk of the
13 occurrence of several tumor types from the animal cancer bioassay data.
14
- 15 5.4.a. Please comment on whether the approach for estimating cancer risk,
16 including the use of tumor modeling of the TCDD animal cancer
17 bioassay data, is scientifically justified and clearly described.
18
- 19 5.4.b. Please comment on the choice of using a BMDL₀₁ as the POD for the
20 development of candidate oral slope factors derived from the TCDD
21 animal cancer bioassays.
22
- 23 5.5. EPA selected Cheng et al. (2006) – an analysis of the NIOSH occupational cohort –
24 as the critical study for oral slope factor (OSF) development. This study was
25 chosen because it considers dose-dependent elimination of TCDD rather than first-
26 order kinetics.
27
- 28 5.5.a. Please comment on whether the rationale for this selection is scientifically
29 justified and clearly described. Please identify and provide the rationale
30 for any other studies that should be considered and provide a critical
31 evaluation of the study and of its suitability for meeting the goals of a
32 quantitatively cancer assessment.
33
- 34 5.5.b. Cheng et al. (2006) analyzed all-cancer mortality. Please comment on the
35 use of all-cancer mortality as the basis of the OSF.
36
- 37 5.5.c. Please comment on whether the use of the Emond PBPK model in the
38 estimation of risk-specific doses from the Cheng et al. dose-response
39 modeling results is scientifically justified and clearly described.
40
- 41 5.5.d. EPA elected to use the log linear relationship of fat concentration and rate
42 ratio to estimate risk-specific doses at all risk levels. EPA could have
43 estimated a POD for cancer risk itself at a single risk level (BMR) for
44 extrapolation to the origin. Please comment on EPA's choice of
45 extrapolation approach.

- 5.5.e. The slope factor derived from Cheng et al. (2006) was extrapolated below the background TCDD exposure levels experienced by the NIOSH cohort. Please comment on this extrapolation.
- 5.6. Please comment on whether EPA has clearly described the major qualitative uncertainties in the derivation of the OSF.
- 5.7. EPA did not consider dioxin-like compounds (DLCs) in the cancer dose-response modeling because the occupational exposures in the available cohorts were primarily to TCDD. Background DLC exposures were not incorporated in the dose-response modeling because EPA judged that it was not possible to disaggregate the responses from background exposure to DLCs and occupational exposure to TCDD. Please comment on whether this approach is scientifically justified and clearly described.
- 5.8. The NRC suggested that EPA consider nonlinear approaches for the assessment of TCDD carcinogenicity. In the *Response to Comments*, EPA presents two illustrative nonlinear approaches for cancer, but considers both inappropriate to use because of the lack of MOA information.
- 5.8.a. Please comment on these two illustrative nonlinear approaches including EPA's conclusions regarding the limitations of these approaches.
- 5.8.b. Are there other nonlinear approaches that could be readily developed based on existing data for the assessment of TCDD carcinogenicity? If so, please suggest alternative approaches and describe their utility and suitability for meeting the goals of a quantitative cancer assessment.

Section 6. Feasibility of Quantitative Uncertainty Analysis from NAS Evaluation of the 2003 Reassessment

- 6.1. Please comment on the discussion in this Section. Is the response clearly presented and scientifically justified?
- 6.2. Please comment on EPA's overall conclusion that a comprehensive quantitative uncertainty analysis is not feasible.
- 6.2.a. Please comment on the discussion in Section 6 regarding volitional uncertainty and how this type of uncertainty limits the ability to conduct a quantitative uncertainty analysis.
- 6.3. Throughout the document (including the Appendices), EPA presents a number of limited sensitivity analyses (e.g., toxicokinetic modeling, RfD ranges, cancer OSF ranges, cancer RfD development). Please comment on the approaches used, and the

- 1 utility of these sensitivity analyses in clarifying potential significant uncertainties.